

1
2 **The impact of agricultural practices on soil organisms: lessons learnt from**
3 **market-gardens**
4
5
6

7 **Abstract**

8 The aim of this study was (i) to establish a typology of farming practices in vegetables
9 cropping systems in Guadeloupe and (ii) to determinate a relationship between these cropping
10 systems and soil fauna. Based on the analysis of cropping systems of an initial set on the
11 whole territory, we selected a representative subset of 18 farms located on vertisols in
12 Grande-Terre. On these 18 farms, we performed a PCA and a HCA. These methods allowed
13 us to build a typology in which farms were distributed between two types. In type A, farmers
14 are using conventional agricultural practices while in type B, farmers are using alternative
15 farming practices. In a second step, we collected soil fauna, from December 2016 to January
16 2017 in type A and type B farms. The results showed no significant difference between soil
17 fauna abundance in both types. However, the number of species richness was higher in type
18 B. Our results also showed that the abundance of litter transformers was significantly higher
19 in type B. Soil fauna activity in type A was probably affected by the use of synthetic
20 fertilizers and herbicides. Taxonomic richness and soil fauna functional diversity thus
21 strongly depend on agricultural practices in vegetables cropping systems in Guadeloupe.
22

23 **Keywords:** Vegetables cropping systems, Agroecology, Survey, Soil fauna, Functional
24 diversity.
25

26 **1. Introduction**

27 Intensive agriculture relied heavily on the use of synthetic inputs and low genetic diversity
28 [1,2,3]. It is well recognized that conventional intensive agriculture had negative impact on
29 natural resources such as soil (soil pollution, erosion), water quality (pollution of rivers, lakes
30 and streams), biodiversity loss and human health (inadequate use of pesticides) [4,5,6,7,8,9].
31 Therefore, such unsustainable models need to be modified to agroecosystems that can
32 optimize ecological functions while maintaining high productivity [9]. Since 1990s, there
33 was a growing interest in developing alternative sustainable farming strategies. All of these
34 strategies share the same objective in term of minimizing the use of synthetic inputs (or even
35 non-use at all), enhancing organic matter recycling and improving agroecosystems health,
36 while maintaining a high production level [10,11,12]. These strategies are part of the field of
37 agroecology as they promote the development of practices based on the mobilization of
38 natural regulations. According to Pretty (2008) [13], sustainable agriculture jointly produce
39 food and goods for farmers and the environment.

40 In 2017, worldwide agricultural production of vegetables was 182 million metric of tomatoes,
41 97 million metric of onions, 83 million metric of cucumbers and gherkins, 71 million metric
42 of cabbages and other brassicas and 52 million metric of eggplants [14]. China, India and the
43 United States of America were the main producers in 2017 [14]. Market-gardening has a
44 major place in agriculture production and in human health due to the providing of compounds
45 such as vitamin A and C, minerals, folic acid and fibers [15,16].

46 In Guadeloupe, agriculture is one of the most important economic sectors. It is a major source
47 of exported goods mainly based on the agroindustrial models developed with banana and
48 sugarcane. The surface of agriculture land decline mainly due to urban construction (e.g. from
49 57 385 ha in 1981 to 30 965 ha in 2013 [17]. However, it still occupies one third of the area of

50 the archipelago. In 2016, the island's main crops were sugarcane (590 299 tones) and banana
51 (66 208 tones). The others crops were vegetables (28 841 tones) and tubers (4 370 tones) [18].
52 Sugarcane and banana were the main studied cropping systems in Guadeloupe [19,20] as they
53 represent agricultural dominant systems, because of the engagement of farmers in market
54 channels and professional and public organizations. Sugarcane and banana also benefit from
55 major public subsidies, which helped farmers to invest and maximize their production. By
56 contrast, we have few informations on vegetables cropping practices though they are models
57 of alternative diversified systems, assumed less dependent on chemical inputs. Therefore, the
58 study concentrated on identifying agricultural practices in vegetables farming systems in
59 Guadeloupe. As we know agricultural practices have an influence on soil fauna activity;
60 however, we wanted to know what kind of alternative practices are used in vegetable cropping
61 systems and in what extent those practices affect soil biota. We hypothesized that there was a
62 positive correlation between practices quality developed in vegetables cropping systems and
63 soil organisms. Soil is then considered as a indicator of the quality of the practices.
64 Considering the lack of scientific knowledge on vegetables cropping systems influence on
65 soil organisms, this article intends to fill this gap by providing consistent informations on the
66 functioning of such agroecosystems. Thus, this paper aims at: (i) identifying the practices
67 developed in vegetables cropping systems and explaining their degree of ecologization. (ii)
68 On this basis, a typology of cropping practices in these agrosystems in Guadeloupe was
69 established. (iii) Using this typology, we demonstrate the relationship between cropping
70 systems and the quality of soils proxied by biological indicators (abundance and diversity of
71 soil fauna).

72

73 **2. Materials and Methods**

74 *2.1 Research area*

75 The study was carried out in Guadeloupe (French West Indies), which is a part of the
76 Winward islands, in the eastern Caribbean Sea. This archipelago includes two main islands
77 with distinct environment. Basse-Terre (848 km²) is dominated by a mountain chain oriented
78 North-West to South-East. The annual temperature is comprised between 20.1 and 31.9 °C
79 (France Meteorological Service, <http://www.meteo.gp>). This island is characterized by a
80 humid tropical climate and a variety of soil types: ferralsols, nitisols, andosols and vertisols
81 [20]. The mean annual rainfall in Basse-Terre is comprised between 1400 mm and 3500 mm
82 (France Meteorological Service, <http://www.meteo.gp>). At the opposite, Grande-Terre (586
83 km²) is characterized by a slightly undulating surface, and the relief rarely exceed 40 m [20].
84 The climate is tropical, with a mean annual rainfall between 1300 mm and 1600 mm, and
85 soils are mostly vertisols.

86

87 *2.2 Farm surveys and typology*

88 To collect data on the practices declined in vegetables cropping systems, a survey was carried
89 out between September and November 2016. 49 farms were randomly sampled: 21 in Grande-
90 Terre and 28 in Basse-Terre. We only targeted farms, which have all or a part of their
91 productions devoted to vegetables cropping systems. We visited those farmers to observe their
92 practices. In the survey, we used variables that best described and discriminated farms. Some
93 variables are intangible (i.e. soil type) while others depend on farmers strategies: crops
94 rotation, soil tillage, irrigation, use of pesticides, weed control, use of synthetic fertilizer or
95 organic amendment, mulch, and management of crop residues. Based on cropping systems of
96 the initial set of 49 farms on the whole territory, we selected a representative subset of 18
97 farms developed on vertisols in Grande-Terre. This selection was due to the fact that in
98 Guadeloupe, vegetables cropping systems are mostly concentrated on vertisol [21]. Indeed,
99 these soils are rich in calcium, magnesium, potassium and they maintain a pH neutral to

100 slightly basic [22] . In addition, the large diversity of soils in Basse-Terre makes it difficult to
101 build a typology.

102 On the 18 farms, we performed a PCA and a HCA. These methods allowed us to build a
103 typology, by gathering farms based on their characteristics and practices. This analysis was
104 realized by using the following variables: (i) soil tillage separated farms into 3 classes: deep,
105 superficial and manual tillage ; (ii) the type of pesticides divided farms in 3 classes: chemical
106 pesticides, pesticides used in biological agriculture or no pesticides; (iii) the use of synthetic
107 herbicides distributed farms in 3 classes: intensive, intermediate and occasional; (iv) weed
108 control separated farms in two classes: mechanical or manual; (v) amendment divided farms
109 in 4 classes: application of synthetic fertilizer, application of organic matter, application of
110 both, and no fertilization; (vi) the use of the mulch practice separated farms in 2 classes:
111 presence or absence; (vii) the management of crop residues divided farms in three classes:
112 removed from the field, incorporated into the soil, and left in the plot; (viii) the application of
113 slash-and-burn practices distributed farms in two classes: with or without slash-and-burn
114 practices; (ix) finally, the observation of soil biodiversity on the surface separated farms in
115 four classes: high, medium, low and no activity.

116

117 *2.3 Soil fauna*

118 From December 2016 to January 2017, in each selected farms on Vertisol, five soil samples of
119 25 cm (length) × 25 cm (width) × 20 cm (deep) were taken for soil macrofauna extraction
120 using TSBF method [23] . Each sample was separated at least 200 m from the others, and was
121 collected 1 km far away from any road and walking path. Animals were collected in alcohol,
122 counted and identified at the taxonomic level under a dissecting microscope. The following
123 taxonomic groups of soil fauna were identified: Oligochaeta, Formicidae, Isoptera, Isopoda,
124 Diplopoda, Dictyoptera, Coleoptera, Diptera, Lepidoptera, Gasteropoda, Homoptera,
125 Orthoptera, Heteroptera, Arenidae, Chilopoda, Dermaptera, Turbellaria, Insects larvae, and
126 Other insects. They were gathered in different functional groups: litter transformers, predators
127 and ecosystem engineers, and we calculated the taxonomic richness. This functional approach
128 can provide information on soil framework and vegetation quality [24,25].

129

130 *2.4 Data analysis methods*

131 To establish a typology of farming practices in vegetables cropping systems, a principal
132 component analysis (PCA) was performed. PCA is a multivariate data analysis that based on
133 projection methods. It is a useful technique for reducing the dimensionality of such datasets,
134 increasing interpretability but at the same time minimizing information loss [26]. Base on the
135 PCA, a hierarchical cluster analysis (HCA) was performed. HCA build a tree diagram, which
136 identify groups of similar observations in a dataset. These analysis were realized with R
137 statistical software (<http://www.r-project.org/>) using R Commander package (Rcmdr). For the
138 relationship between the two types of farming practices and soil fauna, we used Welch's t-test.
139 This test was carried out using R software.

140

141 **3. Results and discussion**

142 *3.1 Characterization of vegetables cropping systems*

143 *3.1.1 Description of 49 farms based on surveys*

144 The survey showed the diversity of agricultural practices in vegetables cropping systems in
145 Guadeloupe. In Basse-Terre and in Grande-Terre, we saw similar crops such as lettuce,
146 zucchini, tomatoes, melon, chili pepper, and eggplant. In addition, in Basse-Terre, we also
147 observed cucumber, pumpkin, cabbage, ochra and chives. We also observed various types of
148 cropping systems, from monoculture to polyculture, and a wide range of practices, from
149 conventional to agroecological.

150 Farming practices are mainly territorially anchored. Tillage is used to ameliorate soil
151 conditions in relation to the water balance and crop growth, to loose upper soil layers to
152 prevent soil compaction, to diminish weed growth and to prepare the seedbed ([27, 28, 29,
153 30]. Our results showed that in Grande-Terre, most farmers used deep tillage (76%) compared
154 to superficial tillage (24%). In this region, vertisols which are clay rich soils which are
155 extremely hard when they dry with cracks and polygonal structures [31] are dominant. Deep
156 tillage is thus used to prepare field for the next culture, by moving and mixing the topsoil with
157 crop residues, which are incorporated into the soil [28]. On the contrary, farmers from Basse-
158 Terre used superficial tillage (71%) rather than deep tillage (29%), due to the type of soils met
159 in this region. Ferralsols have loose and friable fragments [22]. Nitisols are very similar to
160 ferralsols but are an early stage. Finally, andosols are slightly sticky and friable to very friable
161 [32]. Tillage reduced soil organic matter availability by accelerating decomposition and by
162 increasing soil erosion and soil degradation [33]. Moreover, it has a detrimental effect on
163 environmental quality because of its impact on greenhouse gas emissions [34, 35]. Soil
164 disturbance such as tillage has a strong influence on soil fertility and water availability [36].
165 At the opposite, by minimizing mechanical disturbance of soil and macro-aggregate
166 destruction, reduced tillage strongly decrease soil erosion [37,38] and improve water use
167 efficiency [39]. Reduced tillage has thus positive effects on nutrient cycling and soil
168 biodiversity [40,41].

169 During the survey, we observed that use of synthetic pesticides was widely spread among the
170 different farms. In Guadeloupe, crop yield was subjected to pest damage and diseases mainly
171 during rainfall season. Farmers usually prevent economic loss due to pest by spreading heavy
172 pesticides treatments [42]. Additionally, the application rate of herbicides depended on the
173 area. Farmers from Grande-Terre combined herbicides and deep tillage. The mixture of those
174 two methods regulated the abundance of weed species in field [43]. In fact, Chauhan and
175 Johnson (2008) [44] showed that when seeds are deeply buried, the emergence rate was very
176 low.

177 33% of farmers in Grande-Terre and 11% of farmers in Basse-Terre applied mineral
178 fertilizers. Agricultural production has increased, since 1950s, due to the large input of
179 mineral fertilizers [45]. However, the intensive use of mineral fertilizers has a negative impact
180 on soil fertility (soil acidification) and yield production [46]. 25% of farmers applied organic
181 matter in Basse-Terre and 24% in Grande-Terre. Organic fertilizers are used as an alternative
182 to synthetic ones, in order to restore or ameliorate soil physical, chemical and biological
183 properties [47]. Organic matter is not only a source of plants nutrients in soils, but also, it has
184 an important role in preserving soil fertility, reducing soil erosion, nutrient cycling, water
185 retention and disease suppression [48, 49]. During the study, we noticed that in most cases,
186 farmers mixed organic matter and mineral fertilizer together, 54% in Basse-Terre compared to
187 33% in Grande-Terre. A meta-analysis, across sub-Saharan Africa, demonstrated that the use
188 of both input types leads to a greater crops production [50]. Other studies have reported that
189 organic input prevents the rapid leaching of nitrogen fertilizer by immobilizing the nitrogen
190 temporarily [51, 52, 53].

191 For the management of crops residues, most farmers left crops residues in the plot (68% and
192 52% for Basse-Terre and Grande-Terre respectively). Some farmers removed crop residues
193 from the field, 29% in Grande-Terre and 18% in Basse-Terre, or incorporated them into the
194 soil (19% and 14% for Grande-Terre and Basse-Terre respectively). Crop residues can serve
195 as a nutrient source for soil organisms [54]. Moreover, crop residues can ameliorate soil
196 structure, increase organic matter in soil and reduce evaporation [55]. At the same time, we
197 did an observation of soil biodiversity activity on soil surface (observation of ant nests and
198 earthworm casts), and most farms had an activity between high and medium. The presence of
199 ant nests and earthworm casts may be an indicator of soil health.

200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242

3.1.2. Typology of farms located on vertisols


We realized a typology on 18 farms located on vertisol in Grande-Terre based on PCA and AHC. The first two components of PCA explained nearly 43.85% of the total variation (Figure 1 a). Axis F1 has a positive correlation with organic matter, soil biodiversity and slash-and-burn. At the opposite, axis F1 has a negative correlation with herbicides and synthetic fertilizer. Axis F2 opposed plots with biological pesticides to plots using weed control.

Our results showed that farmers from type A are using conventional agricultural practices. These farms are the most numerous in Grande-Terre (Figure 1c) and are characterized by an intensive to medium application of synthetic fertilizers and herbicides. In this type, farmers do not use mulch and slash-and-burn methods. The observation on the soil activity showed low biodiversity (Figures 1a, 1b, 1c). At the opposite, farmers from type B are using alternative agroecological farming practices. In particular, these farms are characterized by application of organic matter, the use of biological pesticides or no pesticides, slash and burn and mulch. The residues are usually left on the field. The observation on soil activity showed a rich biodiversity (Figures 1a, 1b, 1c).


In our study, farmers from type A applied mineral fertilizer, which globally, improve crop yields and food security [56, 57]. Nevertheless, the overdose of mineral fertilizer contributed to soil deterioration, water pollution, and soil biodiversity through soil acidification [58, 59, 60]. Farmers from type A also applied high amount of herbicides which also had a negative effect on fauna, by reducing soil fauna abundance or fitness, due to the destruction of habitat and food resources [61]. On the contrary, in type B, the application of organic matter had a beneficial effect on diverse biological processes by being a food resource for various ecological groups in the community [62, 63]. In addition, farmers of type B applied slash-and-burn, an alternative method. By using this method, farmers can actually maintain carbon stock and increase biodiversity [64, 65, 66, 67]. Mulching also had a major impact on soil fauna abundance and diversity. Mulching is a form of cover crops that remains on the surface of the soil. It can be inorganic or organic material (plastic, straw, cover crop residues or live plant) and it is used to prevent soil erosion, increase water retention, pest control and weed control [10, 68, 69, 70]. However, a few number of surveyed farmers are using this method. Farmers, who were using cover crop method, had a positive feedback based on their crops production. Though, farmers, who used plastic, have trouble to recycle the plastic and they are planning to go to an ecological method.

Our results showed the impact of farming practices on soil biodiversity. In order to confirm this observation with quantitative data, we performed a soil macrofauna extraction on farms.

Figure 1 (a) Projection of the variables used to elaborate the farm typology with Principal Component Analysis (PCA), (b) Representation of farms classified by type based on the components of the PCA (● Type A and ■ Type B), and (c) Dendrogram chart obtained for the Agglomerative Hierarchical Clustering analysis (AHC) performed on components of the PCA, n represents the number of farms for each type. These analyses were carried out using R software.




243



244

REVIEW



245

246 3.1.3. Soil macrofauna on farms located on vertisols


247 Soil macrofauna were collected on selected farms. We found 171 ± 52 (mean \pm SE)
 248 individuals.m⁻² in type A, and 554 ± 239 individuals.m⁻² in type B. The abundance of soil
 249 fauna was slightly higher in type B (Figure 2b). However, there was no significant difference
 250 in soil fauna abundance between both types (t-test Welch; $P = .13$). In general, ecosystems
 251 engineers were more abundant than litter transformers and predators (Figure 2a). In type B,
 252 the amount of ecosystem engineers (432 ± 229 individuals.m⁻²) and predators (48.8 ± 16.88
 253 individuals.m⁻²) was slightly higher than in type A (ecosystem engineers: 116 ± 41
 254 individuals.m⁻², biological regulators: 24 ± 6 individuals.m⁻²). However, there was no
 255 significant difference between the quantity of ecosystems engineers and predators between
 256 type A and type B (t-test Welch; $P = .21$ and $P = 0.15$). On the other hand, the quantity of
 257 litter transformers was significantly different between the two types (t-test Welch; $P = .02$)
 258 (Figure 2a). The amount of litter transformers was higher in type B (72 ± 18 individuals. m⁻²)
 259 than in type A (30 ± 10 individuals. m⁻²). Also, in Figure 2c, the taxonomic richness was
 260 significantly higher in type B (11 ± 0.4 taxonomic richness) compared to type A ($6.5 \pm$
 261 0.61 taxonomic richness) (t-test Welch; $P < .001$).

262 Soil macrofauna may be used as bioindicators of soil health and contributed to ecosystems
 263 services [25]. Soil macrofauna play an important role in soil organic matter decomposition
 264 (litter transformers), regulations of pests (predators), formation of stable aggregates, water
 265 regulation and erosion control (ecosystems engineers) [71]. Our results showed that soil
 266 macrofauna may be directly or indirectly impacted by agricultural practices. In type A, we
 267 observed a number of conventional agriculture practices (deep tillage, application of high
 268 amounts of chemical pesticides, synthetic fertilizer, and herbicides), which are well known to
 269 have a negative impact on soil biodiversity [59]. Our study showed that litter transformers are
 270 strongly impacted by these conventional practices. They had an essential role in soil carbon
 271 sequestration [72]. As a consequence, by decreasing the quantity of litter transformers,
 272 conventional agriculture may have profound effects on climate change. On the contrary, by
 273 decreasing the input of synthetic fertilisants and herbicides, by reducing the rate of tillage and
 274 by increasing the application of organic matter, farmers in type B are stabilizing their soil.
 275 Moreover, type B applied mulching, which can have a positive effect on soil habitat.
 276 Mulching helps to preserve the ecosystem by reducing the rate of tillage. Sustainable


277 agriculture also had a beneficial impact on soil physical and chemical properties, such as,
278 aggregation and nitrogen content [73, 74], which indirectly impacted soil fauna abundance
279 and diversity. In order to overcome the impacts of conventional agriculture, sustainable
280 agriculture methods have been developed to minimize environmental footprints and sustain
281 natural environments and resources [75, 30]. Our study showed that in Guadeloupe, farmers
282 are looking for alternative agriculture practices in vegetables cropping systems. In order to
283 better understand the impact of those new agroecological practices, further physical, chemical
284 and soil fauna analyses should be realized.

285

286 **Figure 2** (a) Soil fauna abundance in 9 farms (type A and type B) in Guadeloupe (Grande-
287 Terre).(b) Soil fauna abundance in type A and B in Guadeloupe. (c) Taxonomic richness
288 abundance in type A and type B in Guadeloupe. Values with similar letters are not
289 significantly different (Welch t-test). These analyses were carried out using R software




290



291

c



292

293

Conclusion

294

295

296

297

298

299

300

301

302

303

304

305

306

References

307

308

1. Trewanas Antony. Malthus foiled again and again. *Nature*. 2002 ; 418 : 668-670.

309

310

2. Gliessman Stephen R. *Agroecology. The Ecology of sustainable food systems*. Second edition, CRC Press Taylor & Francis Group, Florida ; 2007.

311

312

3. Nair PKR. Grand Challenges in Agroecology and Land Use Systems. *Front Environ Sci*. 2014 ; 2 : 1-3.

313

314

4. Pimentel D. Environmental and economic costs of the application of pesticides, primarily in the United States. *Environ Dev Sustain*. 2005 ; 7 : 229-252.

315

316

317

5. Halweil B, Mastny L, Assadourian E, Flavin C, French H, Gardner G, Nierenberg D, Postel S, Renner M, Sarin R, Sawin J, Vickers A. *State of the World*. W.W. Norton & Company ; 2004.

318

319

320

6. Castillo LE, Martinez E, Ruepert C, Savage C, Gilek M, Pinnock M, Solis E. Water quality and macro invertebrate community response following pesticide applications in a banana plantation, Limon, Costa Rica. *Sci. Total Environ*. 2006 ; 367 : 418-432.

- 321 7. Kendall P, Petracco M. The current state and future of Caribbean agriculture. *J. Sustain.*
322 *Agric.* 2009 ; 33 : 780-797.
- 323 8. Chappell M, LaValle LA. Food security and biodiversity: can we have both? An
324 agroecological analysis. *Agric Human Values.* 2011 ; 28 : 3-26.
- 325 9. Bommarco R, Kleijn D, Potts SG. Ecological intensification: harnessing ecosystem services
326 for food security. *Trends Ecol Evol.* 2013; 28 : 230–238.
- 327 10. Altieri MA. The ecological role of biodiversity in agroecosystems. *Agric Ecosyst Environ.*
328 1999; 74 :19-31.
- 329 11. Andres C, Bhullar G. Sustainable Intensification of Tropical Agro-Ecosystems: Need and
330 Potentials. *Front Environ Sci.* 2016 ; 4 : 1-10.
- 331 12. Meynard JM. L'agroécologie, un nouveau rapport aux savoirs et à l'innovation. *Tropical*
332 *Issue.* 2017 ; 24.
- 333 13. Pretty JN. Agricultural sustainability: Concepts, principles and evidence. *Phil. Trans. R.*
334 *Soc. B.* 2008 ; 363 : 447-465.
- 335 14. The Statistics Portal. 2019. Accessed 10 March 2019. Available:
336 <https://www.statista.com/statistics/264065/global-production-of-vegetables-by-type/>.
- 337 15. Lester GE. Environmental regulation of human health nutrients (ascorbic acid, carotene,
338 and folic acid) in fruits and vegetables. *HortSciences.* 2006 ; 41 : 59-64.
- 339 16. Slavin JL, Llyord B. Health benefits of fruits and vegetables. *Advances in Nutrition.* 2012
340 ; 3(4) : 506-516.
- 341 17. AGRESTE. Enquête sur les pratiques phytosanitaires sur les légumes en 2013. 2013.
342 Accessed 23 March 2019. Available:
343 [http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/breve_1_-_pklegumes_2013_cle81ed6e-](http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/breve_1_-_pklegumes_2013_cle81ed6e-1.pdf)
344 [1.pdf](http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/breve_1_-_pklegumes_2013_cle81ed6e-1.pdf).
- 345 18. AGRESTE. Mémento de la statistique agricole. 2017. Accessed 25 March 2019. Available:
346 <http://daaf.guadeloupe.agriculture.gouv.fr/Memento-Agricole-Guadeloupe-2017>.
- 347 19. Clermont-Dauphin C, Cabidoche YM, Meynard JM. Effects of intensive monocropping of
348 bananas on properties of volcanic soils in the uplands of the French West Indies. *Soil Use*
349 *Manage.* 2004 ; 20 : 105-113.
- 350 20. Sierra J, Causeret F, Diman JL, Publicol M, Desfontaines L, Cavalier A, Chopin P.
351 Observed and predicted changes in soil carbon stocks under export and diversified agriculture
352 in the Caribbean. The case study of Guadeloupe. *Agric. Ecosyst. Environ.* 2015 ; 213 : 252-
353 264.
- 354 21. Bulletin de santé du végétale (BSV). Retrouver les bilans par filière de l'année 2016.
355 2016. Accessed 12 March 2019. Available:
356 [http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/BSV971_ToutesFilières_Bilan2016_cle4](http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/BSV971_ToutesFilières_Bilan2016_cle492392.pdf)
357 [92392.pdf](http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/BSV971_ToutesFilières_Bilan2016_cle492392.pdf); 2016.

- 360 22. Sierra J, Desfontaines L. Rapport : Les sols de la Guadeloupe. Genèse, distribution et
361 propriétés. 2018. Accessed 26 February 2019. Available:
362 <https://www6.antilles.inra.fr/astro/Ce-que-nous-savons-sur/No-6-Les-sols-de-la-Guadeloupe>
- 363 23. Anderson JM, Ingram JSI. Tropical Soil Biology and Fertility: A Handbook of Methods.
364 CAB International, Wallingford ; 1993.
- 365 24. Brussaard L. Soil fauna, guilds, functional groups and ecosystem processes. *Appl Soil*
366 *Ecol.* 1998 ; 9 : 123-135.
- 367 25. Turbé A, Toni DA, Benito P, Lavelle P., Lavelle P, Ruiz N, Wim HVdP, Labouze E,
368 Mudgal S. Soil biodiversity : functions, threats and tools for policy makers. Bio Intelligence
369 Service, IRD, and NIOO, Report for European Commission (DG Environment) ; 2010.
- 370 26. Jolliffe IT, Cadima J. Principal component analysis: a review and recent developments.
371 *Phil. Trans. R. Soc. A.* 2016 ; 374 (2065).
- 372 27. Wall HD, Bardgett RD, Behan-Pelletier V, Herrick JE, Jones H, Ritz K, Six J, Strong DR,
373 Van der Putten WH. Soil ecology and Ecosystem services. Oxford University Press, New
374 York ; 2012.
- 375 28. Welbaum GE. Vegetable Production and Practices. Virginia Tech University, USA ; 2015.
- 376 29. Gómez JA, Orgaz F, Gómez-Macpherson H, Villalobos FJ, Fereres E. Tillage, In:
377 Villalobos FJ, Fereres E, editors. Principles of Agronomy for Sustainable Agriculture, Cham:
378 Springer International Publishing ; 2016.
- 379 30. Stavi I, Bel G, Zaady E. Soil functions and ecosystem services in conventional,
380 conservation, and integrated agricultural systems. A review. *Agron Sustain Dev.* 2016 ; 36:
381 32.
- 382 31. Baize D, Girard, MC. Référentiel pédologique. Quæ Editions ; 2008.
- 383 32. Baize D, Girard MC. A Sound Reference Base for Soils: The "Référentiel Pédologique".
384 Quæ Éditions ; 1998.
- 385 33. Gliessman Stephen R. Agroecology. The Ecology of sustainable food systems. Third
386 edition, CRC Press Taylor & Francis Group, Florida ; 2015.
- 387 34. Schneider UA, Smith P. Energy intensities and greenhouse gas emission mitigation in
388 global agriculture. *Energy Effic.* 2009 ; 2 : 195-206.
- 389 35. Ji Q, Wang Y, Chen XN, Wang XD. Tillage effects on soil aggregation, organic carbon
390 fractions and grain yield in Eum-Orthic Anthrosol of a winter wheat-maize double-cropping
391 system, Northwest China. *Soil Use Manag.* 2015 ; 31 : 504-514.
- 392
393 36. Adimassu Z., Alemu G., Tamene L. Effects of tillage and crop residue management on
394 runoff, soil loss and crop yield in the Humid Highlands of Ethiopia. *Agric Syst.* 2019 ; 168
395 :11-18.
- 396
397 37. Six J, Bossuyt H, Degryze S, Denef K. A history of research on the link between (micro)
398 aggregates, soil biota, and soil organic matter dynamics. *Soil Tillage Res.* 2004 ; 79 : 7–31.

- 400 38. Liu, E., Teclerian SG, Yan C, Yu J, Gu R., Liu S, He W, Liu Q. Long-term effects of
401 no-tillage management practice on soil organic carbon and its fractions in the northern China.
402 *Geoderma*. 2014 ; 213 : 379-384.
- 403 39. Li L, Huang G, Zhang R, Bill B, Guangdi L, Kwong YC. Benefits of conservation
404 agriculture on soil and water conservation and its progress in China. *Agric. Sci. China*. 2011 ;
405 10 : 850-859.
- 406 40. Lal R. Restoring soil quality to mitigate soil degradation. *Sustainability*. 2015 ; 7 : 5875-
407 5895.
- 408 41. Pareja-Sánchez E, Plaza-Bonilla D, Ramos MC, Lampurlanés J, Álvaro-Fuentes
409 J, Cantero-Martínez C. Long-term no-till as a means to maintain soil surface structure in an
410 agroecosystem transformed into irrigation. *Soil Till Res*. 2017 ; 174 : 221-230.
- 411 42. National Research Council.. *Sustainable Agriculture and the Environment in the Humid*
412 *Tropics*. National Academy Press., Washington; 1993.
- 413 43. Mishra JS, Singh VP. Tillage and weed control effects on productivity of a dry seeded
414 rice-wheat system on a Vertisol in Central India. *Soil Till Res*. 2012 ; 123 : 11-20.
- 415 44. Chauhan BS, Johnson DE. Germination Ecology of Southern Crabgrass (*Digitaria ciliaris*)
416 and India Crabgrass (*Digitaria longiflora*): Two Important Weeds of Rice in Tropics. *Weed*
417 *Sci*. 2008 ; 56 : 722-728.
- 418 45. Cai A, Xu M, Wang B, Zhang W, Liang G, Hou E, Luo Y. Manure acts as a better
419 fertilizer for increasing crop yields than synthetic fertilizer does by improving soil fertility.
420 *Soil Till Res*. 2019 ; 189 : 168-175.
- 421 46. Zhu Q, Liu X, Hao T, Zeng M, Shen J, Zhang F, De Vries W. Modeling soil acidification
422 in typical Chinese cropping systems. *Sci Total Environ*. 2018 ; 613-614 : 1339-1348.
- 423 47. Celestina C, Hunt JR, Sale PWG, Franks AE. Attribution of crop yield responses to
424 application of organic amendments: A critical review. *Soil Till Res*. 2019 ; 186 : 135-145.
- 425 48. Loveland P, Webb J. Is there a critical level of organic matter in the agricultural soils of
426 temperate regions: a review. *Soil Till Res*. 2003 ; 70 : 1-18.
- 427 49. Murphy, BW. Impact of soil organic matter on soil properties—a review with emphasis on
428 Australian soils. *Soil Res*. 2015 ; 53 : 605-635.
- 429 50. Chivenge P, Vanlauwe B, Six J. Does the combined application of organic and mineral
430 nutrient sources influence maize productivity? A meta-analysis. *Plant Soil*. 2010 ; 342 : 1-30.
- 431 51. Vanlauwe B, Diels J, Aihou K, Iwuafor ENO, Lyasse O, Sanginga N, Merckx R. Direct
432 interactions between N fertilizer and organic matter: evidence from trials with ¹⁵N-labelled
433 fertilizer. In: Vanlauwe B, Diels J, Sanginga N, Merckx R, editors. *Integrated Plant Nutrient*
434 *Management in Sub-Saharan Africa: From Concept to Practice*. CAB International; 2002.

- 435 52. Bekunda MA, Nteranya S, Woomer PL. Restoring soil fertility in sub-sahara africa. *Adv*
436 *Agron.* 2010; 108 : 183–236.
- 437 53. Pincus L, Margenot A, Sixe J, Scowd K.. On-farm trial assessing combined organic and
438 mineral fertilizer amendments on vegetable yields in central Uganda. *Agric Ecosyst Environ.*
439 2016 ; 225 : 62-71.
- 440 54. Feng Y, Balkcom KS. Nutrient Cycling and Soil Biology in Row Crop Systems under
441 Intensive Tillage. In: Al-Kaisi MM, Lowery B, editors. *Soil Health and Intensification of*
442 *Agroecosystems.* Academic Press Elsevier Inc.; 2017.
- 443 55. Liang S, Li X, Wang J. *Advanced Remote Sensing.* Elsevier Inc ; 2012.
- 444 56. Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. Agricultural sustainability and
445 intensive production practices. *Nature.* 2002 ; 418 : 671-677.
- 446 57. Boli N, Mingzhu L, Shaoyu L, Lihua X, Yanfang W.. Environmentally friendly slow-
447 release nitrogen fertilizer. *J Agric Food Chem.* 2011 ; 59 : 10169-10175.
- 448 58. Tilman D, Fargione J, Wolff B, D'Antonio C, Dobson A, Howarth R, Schindler D,
449 Schlesinger WH, Simberloff D, Swackhamer D. Forecasting agriculturally driven global
450 environmental change. *Science.* 2001 ; 292 : 281-284.
- 451 59. Clermont-Dauphin C., Blanchart E, Loranger-Merciris G, Meynard JM. Cropping systems
452 for soil biodiversity and ecosystem services: prospects and research needs. *Sustainable*
453 *Agriculture Reviews .* 2014 ; 14 : 117-158.
- 454 60. Smith LED, Siciliano G. A comprehensive review of constraints to improved management
455 of fertilizers in China and mitigation of diffuse water pollution from agriculture. *Agriculture.*
456 *Ecosyst. Environ.* 2015 ; 209 : 15-25.
- 457 61. Prosser RS, Anderson JC, Hanson ML, Solomon KR, Sibley PK. Indirect effects of
458 herbicides on biota in terrestrial edge-of-field habitats: A critical review of the literature.
459 *Agric Ecosyst Environ.* 2016 ; 232 : 59-72.
- 460 62. Roger-Estrade J, Anger C, Bertrand M, Richard G. Tillage and soil ecology: Partners for
461 sustainable agriculture. *Soil Till Res.* 2010; 111 : 33-40.
- 462 63. Ayuke FO, Brussaard L, Vanlauwe B, Six J, Lelei DK, Kibunja CN, Pulleman MM. Soil
463 fertility management: Impacts on soil macrofauna, soil aggregation and soil organic matter
464 allocation. *Appl Soil Ecol.* 2011; 48 : 53-62.
- 465 64. Bruun TB, Neergaard A, Lawrence D, Ziegler AD. Environmental consequences of the
466 demise in swidden cultivation in southeast asia: carbon storage and soil quality. *Hum Ecol.*
467 2009 ; 37 : 375-388.
- 468 65. Padoch C, Pinedo-Vasquez M.. Saving slash and burn to save biodiversity. *Biotropica.*
469 2010 ; 42 : 550-552.
- 470 66. Ziegler AD, Fox JM, Webb EL, Padoch C, Leisz SJ, Cramb R, Mertz O, Bruun TB, Vien
471 TD. Recognizing contemporary roles of swidden agriculture in transforming landscapes of
472 southeast asia. *Conserv. Biol.* 2011 ; 25 : 846-848.

- 473 67. Lopes Thomaz E. Dynamics of aggregate stability in slash-and-burn system Relaxation
474 time, decay, and resilience. *Soil Till Res.* 2018 ; 178 : 50-54.
- 475 68. Saxton K, Chandler D, Stetler L, Lamb B, Clairborn C, Lee BH. Wind erosion and
476 fugitive dust fluxes on agricultural lands in the Pacific Northwest. *Trans. ASAE.* 2000 ; 43 :
477 623-630.
- 478 69. Gonzalez-Martin C, Teigell-Perez N, Valladares B, Griffin DW. The global dispersion of
479 pathogenic microorganisms by dust storms and its relevance to agriculture. *Adv. Agron.* 2014
480 ; 127 : 1-41.
- 481 70. Quintanilla-Tornel MA, Wang KH, Tavares J, Hooks CRR. Effects of mulching on above
482 and below ground pests and beneficials in a green onion agroecosystem. *Agric Ecosyst*
483 *Environ.* . 2016 ; 224: 75-85.
- 484 71. Lavelle P, Decaëns T, Aubert M, Barot S, Blouin M, Bureau F, Margerie P, Mora P, Rossi
485 JP. Soil invertebrates and ecosystem services. *Eur. J. Soil Biol.* 2006 ; 42 : 3-15.
- 486 72. Angst Š, Mueller CW, Cajthaml T, Angst G, Lhotáková Z, Bartuška M, Špaldoňová A,
487 Frouz J. Stabilization of soil organic matter by earthworms is connected with physical
488 protection rather than with chemical changes of organic matter. *Geoderma.* 2017 ; 289 : 29-35
- 489 73. Caviglia OP, Andrade FH. Sustainable intensification of agriculture in the Argentinean
490 Pampas: capture and use efficiency of environmental resources. *Am. J. Plant Sci. Biotechnol.*
491 2010 ; 3 : 1-8.
- 492 74. Lal R, Reicosky DC, Hanson JD. Evolution of the plow over 10,000 years and the
493 rationale for no-till farming. *Soil Tillage Res.* 2007 ; 93 : 1-12.
- 494 75. Liu G.B. Soil conservation and sustainable agriculture on the Loess Plateau: challenges
495 and prospects. *Ambio.* 1999 ; 28 : 663-668.
- 496 76. Altieri MA, Funes-Monzote FR, Peterson P. Agroecologically efficient agricultural
497 systems for smallholder farmers: contributions to food sovereignty. *Agron Sustain Dev.* 2012
498 ; 32 : 1-13.