

Spermatozoa characteristics, serum metabolites and testicular oxidative stress traits in guinea pigs (*Cavia porcellus*) fed on *Zanthoxylium leprieurii* fruit

10 **ABSTRACT**

Zanthoxylium leprieurii belong to aromatic plants. Its fruit is commonly used as spice in soups and as medicine in many African countries. Because of its phytochemical composition, it can also be used as antioxidant. In the present study, ~~we evaluated the effects of aqueous extract of *Z. leprieurii* fruit on reproductive function in male guinea pig (*Cavia porcellus*)~~ **was evaluated**. Fifty male guinea pigs with an average weight of 320.56 ± 30 g, aged 4 months were used. They were divided into 5 groups (G1, G2, G3, G4 and G5) of 10 animals each. During 60 days, animals of G1 were daily given distilled water orally, while G2 received 100 mg/kg body weight (bw) of vitamin C. In the other hand, G3, G4 and G5 received respectively by the same method 50, 100 and 200 mg/kg bw of aqueous extract of *Z. leprieurii*. At the end of the treatment, all animals were sacrificed for evaluating the genital organs weight, sperm characteristics, serum levels of reproductive hormones and stress biomarkers. Results revealed that the weight of testes, epididymis, vas deferens and accessory glands did not significantly affect ($p>0.05$) in cavies exposed to different treatments compared to control animals. There was a significant ($p<0.05$) increase in serum content of FSH at 100 mg/kg. bw (26.67 ± 3.51 ng ml⁻¹) and LH at 50 mg/kg. bw (10.71 ± 2.42 ng ml⁻¹) in animals exposed to aqueous extract of *Z. leprieurii* with reference to the control groups. In addition, there is a non-significant increase ($p>0.05$) of the level of testosterone in the treated cavies compared to controls. Aqueous extract induced significant ($p<0.05$) increase in sperm mobility and sperm count in treated cavies with respect to the controls. The testicular activities of superoxide dismutase, catalase and peroxidase increased significantly ($p<0.05$) in guinea pigs exposed to aqueous extract of *Z. leprieurii* compared with those of control (G1). The reverse effect was observed concerning the concentration of malondialdehyde. In conclusion, the aqueous extract of *Z. leprieurii* fruit efficiently improves male reproductive characteristics by ~~It is responsible for the increased level in reproductive hormones and an~~ **improvement** of anti-oxidative enzymes activities.

11
12 **Keywords:** Aqueous extract, *Z. leprieurii*, male guinea pig, reproduction.

13
14
15
16 **1. INTRODUCTION**

17
18 In Africa especially in tropical region, the use of plants and its extract for the treatment and management of diseases has been in existence since ancient times. Factors such as poverty and illiteracy still militate against availability and accessibility of conventional medical services. A larger number of these tropical plants and their extract possess diverse bioactive molecules such as phenols, terpenes, tannins, alkaloids rich in various properties: anti-oxidant, anti-inflammatory, anti-cancer, anti-microbial and aphrodisiac [1]. These properties can be in animal production to neutralize the effects of endogenous and exogenous in benefic of growth and productivity. Many studies ~~carry out~~ in the area of animal production have shown that the decrease in animal growth and productivity ~~is~~ **is** linked to poor quality of ~~feed~~ **feed** and water, climate changes, psychological and hormonal disorders and drug [2-5]. As a solution, the breeders previously used in animal diet, antibiotics as food additives to enhance their performances. Unfortunately, the use of that ~~feed~~ **feed** additive have been prohibited due to its side effects attached to microbial resistance, cell apoptosis and reprotoxicity [6]

28 nowadays, natural plants products w
 29 and often with minimal side effects,
 30 *Zanthoxylum leprieurii* due to their b
 31 animal reproduction characteristics [1
 32 200 and 300 mg/kg body weight for 7
 33 The seeds of *Zanthoxylum leprieur*
 34 west region. It belongs to the fami
 35 Ethnomedically, it is used in the trea
 36 rheumatism, neuralgia, fever, tooth
 37 compounds such as phenols, flavonoids, tannins, vitamins A, E and C.

Introduction
 Needs to very concisely written
 The introduction part may be briefed to the point
 What are the active ingredients in the plant and how these can improve the semen characteristic needs to be focused in the introduction

able all year round, cheap, accessible
 rmances. Many plants among which
 enes) have been reputed to improve
 tract of *Psidium guajava* at doses of
 bility and viability.¹⁵
 ameroon and especially those from
 et forming deciduous shrub or tree.
 ose vein, raynauld disease, arthritis,
 d that the fruit of this plant possess

38 Despite frequent and regular use, there is no reported work related to the effect of this plant on male reproductive
 39 function. The study was thus intended to bridge the gap in our continued efforts to establish the effects of local food
 40 spices and medications on male reproductive function.

41
 42
 43 **2. MATERIAL AND METHODS**
 44

45 **Selection of Animals and experimental feeding**

46 Fifty male guinea pigs (*Cavia porcellus*) reared at the Teaching and Research Farm of the University of Dschang were
 47 used. Their average body weight was 320.56±30 g at the beginning of the assay. They were identified individually using
 48 numbered earrings and housed in identical cages of 100 cm x 80 cm x 60 cm (length, width and height) under standard
 49 conditions with free access to water and feed (Table 1). All experiments were carried out in compliance with the
 50 recommendations Guide of the National Academy of Sciences on the care and use of laboratory Animals [17] and
 51 approved by the department of animal science.
 52
 53
 54
 55

Table 1: Composition and chemical characteristics of the experimental diet

Ingredients	Quantities (kg)	Chemical Characteristics
Corn	26.50	Metabolizable energy (kcal/kg) : 2600.00
Bran wheat	25.00	Crude protein (%) : 19.00
Rice bran	12.00	Crude cellulose (%) : 14.18
Soy beans cake	6.00	Calcium (%) : 10.05
Cotton cake	6.00	Phosphorus (%) : 0.68
Palm kernel cake	13.00	Sodium (%) : 0.28
Fish meal	2.00	Lysine (%) : 1.01
Sea shells	2.00	Methionine(%): 0.40
Palm oil	2.00	
Sal	0.50	
Premix10% *	5.00	

56 *Premix 5%: mixture of vitamins A, B complex, D, K, and E plus iron, Cu, Zn, Se, Mn, methionine, lysine
 57 principally and incorporated at 5% in diet.

58

59 **Plant material**

60 The seed of *Zanthoxylum* were bought from Dschang (Menoua division in Cameroon) local market. They were dried
 61 sheltered from the sun, and then grinded at the mill. The obtained powder was used for extractions, using 6 liters of
 62 distilled water for 1 kilogram of powder. The filtrate was dried in the oven at 45 °C to obtain a paste used to prepare the *Z.*
 63 *leprieurii* aqueous extract at different concentrations.

64

65 **Phytochemical screening of *Z. leprieurii***

66 The phytochemical screening of *Z. leprieurii* aqueous extract (Table 2) was done as described by Ngbede [18] and
 67 Ngbede [19].

68

69 **Table 2** : Phytochemical constituent of *Z. leprieurii* aqueous extract

Constituents	(+) present ; (-) absent
Alkaloïds	-
Phenols	+
Tannins	+
Saponins	+
Flavonoids	+
Steroids	-
Triterpenes	+

70

71 **Experimental design**

72 Animals were distributed into 5 groups (G1, G2, G3, G4 and G5) of 10 animals each, comparable in term of body weight.
 73 During 60 days, animals of G1 were daily given orally distilled water (1 ml/kg bw), G2 received 100 mg/kg bw of vitamin C
 74 diluted in distilled water, while G3, G4 and G5 received respectively 50, 100 and 200 mg/kg bw of aqueous extract of *Z.*
 75 *leprieurii*, dissolved in distilled water. The animal's body weight was recorded weekly and the doses of vitamin C and *Z.*
 76 *leprieurii* aqueous extract adjusted accordingly.

77 **Blood and organs collection**

78 Twenty-four hours after the last administration of the vitamin C and aqueous extract solutions, animals were anesthetized
 79 using ether vapor and blood samples were collected by cardiac puncture for the measurement of hormonal
 80 concentrations. After the sacrifice, testes, epididymis, vas deferens and sexual accessory glands were excised out and
 81 weighed.

82 The cauda epididymis of each animal were minced in 10 ml of 0.9% NaCl solution (37 °C) for sperm characteristics
 83 evaluation. For sperm mobility, a drop of the obtained solution was placed on a slide and observed at magnification 400x
 84 under the light microscope (Leica) and the mobility score was attributed according to the method described by Boiti.²⁰ The
 85 sperm count was done using the Thomas haemocytometer, while the integrity of the plasma membrane was evaluated
 86 using the hypo-osmotic test [21].

87 **Evaluation of oxidative stress indicators**

88 A 15% (weigh/volume) homogenate was prepared using left testis of each animal. Thus, the testis was crushed in cold
 89 0.9% NaCl solution followed by a centrifugation (3000 rpm, 30 min). The supernatant obtained was used to evaluate the
 90 testes content in oxidative stress indicators. The malondialdehyde concentration was determined using the thiobarbituric

acid metho, [22] the superoxide dismutase activity was evaluated according to the method describe by Dimo [23]. The catalase (CAT) activity was assessed using the chromic acetate method [24] and the glutathione peroxidases (GPx) activity was determined by the potassium iodate method.²⁵

Evaluation of reproductive hormones

The serum concentration in testosterone, FSH and LH were measured using the instructions from ELISA kit (Omega Diagnostics) (Scotland, United Kingdom).

Statistical analysis

Data analysis were performed using one-way ANOVA, followed by Duncan's test at 5 % significance. All analyses were done using SPSS IBM statistics software 20.0. The results are expressed as mean \pm standard deviation.

3. RESULTS AND DISCUSSION

Weight of genital organs in male guinea pigs

The effects of the aqueous extract of *Z. leprieurii* on the weight of sexual organs in guinea pig are presented in table 3. The weights of testes, epididymis, vas deferens and accessory glands were comparable ($p > 0.05$) among treatments. Although the values of these characteristics were statistically comparable, an increase was observed in the animals treated with aqueous extract with reference to the negative control (T0-).

Table 3: Effects of the aqueous extract of *Z. leprieurii* fruit on the relative weight of genital organs in cavy

Relative weight of genital organs (g/100 g bw)	Control		Doses of <i>Z. leprieurii</i> extract (mg/kg bw)			P-value
	T0 ⁻ (n=12)	T0 ⁺ (n=12)	50 (n=12)	100 (n=12)	200 (n=12)	
Testis	0.43 \pm 0.03 ^a	0.48 \pm 0.05 ^a	0.54 \pm 0.08 ^a	0.47 \pm 0.05 ^a	0.48 \pm 0.08 ^a	0.26
Epididymis	0.08 \pm 0.01 ^a	0.09 \pm 0.02 ^a	0.10 \pm 0.03 ^a	0.10 \pm 0.01 ^a	0.09 \pm 0.02 ^a	0.52
Vas deferens	0.42 \pm 0.09 ^a	0.40 \pm 0.08 ^a	0.43 \pm 0.08 ^a	0.43 \pm 0.04 ^a	0.46 \pm 0.08 ^a	0.89
Accessory glands	0.04 \pm 0.01 ^a	0.04 \pm 0.01 ^a	0.05 \pm 0.01 ^a	0.04 \pm 0.01 ^a	0.05 \pm 0.00 ^a	0.67

a: within the same raw, values with the same letters are not significantly different ($p > 0.05$) n: Number of cavy. Bw, body weight; T0-, group receiving 1 mL/kg bw of distilled water; T0+, group receiving 100 mg/kg bw of vitamin C; P, Probability.

Characteristics of spermatozoa in guinea pigs

Table 4 summarizes the effects of the aqueous extract of *Z. leprieurii* on caudal epididymal sperm characteristics in male guinea pigs. The sperm mobility, sperm count per cauda and per gram of epididymis and spermatozoa with entire plasma membrane increased significantly ($p < 0.05$) in animals treated with aqueous extract of *Z. leprieurii* compared to controls (T0- and T0+). However, the highest sperm mobility and the sperm count was observed in animals exposed to 200 mg/kg bw compared to the others.

Table 4: Effects of the aqueous extract of *Z. leprieurii* fruit on the caudal epididymal sperm characteristics in male guinea pigs

Sperm characteristics	Controls		Doses of <i>Z. leprieurii</i> extract (mg/kg bw)			P value
	T0 ⁻ (n=12)	T0 ⁺ (n=12)	50 (n=12)	100 (n=12)	200 (n=12)	
Mobility (%)	67.50 \pm 1.73	76.67 \pm 0.5	80.00 \pm 0.0	74.00 \pm 1.4	88.33 \pm 1.15 ^a	0.00

	e	8 ^c	0 ^b	1 ^d		
Number/cauda epididymis (10 ⁷)	06.50±0.17	04.67±0.6	07.81±0.3	10.50±0.4	10.42±0.38 ^a	0.00
	c	3 ^d	8 ^b	3 ^a		
Number/g of cauda epididymis (10 ⁷)	29.83±1.25	39.55±2.3	55.51±1.2	37.23±1.3	54.88±1.71 ^a	0.00
	c	7 ^b	3 ^a	8 ^b		
Spermatozoa with EPM (%)	60,0±0,50 ^c	52,5±0,58	75,0±0,82	72,67±0,5	75,33 ±0,58 ^a	0,00
		d	b	8 ^a		

120 a, b, c, d, e, within the same raw, values with the same letters are not significantly ($p>0.05$) different. N, Number
 121 of guinea pigs. Bw, body weight. T0-, group receiving 1 mL/kg bw of distilled water. T0+, group receiving 100
 122 mg/kg bw of vitamin C; EPM, entire plasma membrane.
 123

124 Effect of aqueous extract of *Z. leprieurii* on serum metabolites in male guinea pigs

125 As shown in Table 5, the total cholesterol decreased significantly ($p<0.05$) in *Z. leprieurii* aqueous extract treated guinea
 126 pigs irrespective of the dose as compared to the negative control. The oral administration of *Z. leprieurii* aqueous extract
 127 in guinea pigs did not significantly ($p>0.05$) affect the serum content in total proteins. With respect to the controls, the
 128 serum content in FSH increased in cavies exposed to *Z. leprieurii* aqueous extract, but that increase was significant only
 129 at dose of 100 mg/kg bw. The serum content in LH increased significantly ($p<0.05$) in cavies treated with 50 mg/kg bw of
 130 *Z. leprieurii* aqueous extract compared to control (T0-). The level of testosterone increased but not significantly ($p>0.05$) in
 131 guinea pigs treated with aqueous extract of *Z. leprieurii* at doses of 100 and 200 mg/kg bw compared to the controls.
 132

133 **Table 5:** Effects of aqueous extract of *Z. leprieurii* fruit on serum metabolites in male guinea pigs

Serum metabolites	Controls		Doses of <i>Z. leprieurii</i>			P-value
	extract (mg/kg bw)		50 (n=12)	100 (n=12)	200 (n=12)	
	T0- (n=12)	T0+ (n=12)				
Total cholesterol (mg/dl)	104.66±5.45 ^a	90.03±7.34 ^{bc}	85.36±10.74 ^c	90.03±7.72 ^{bc}	101.16±7.57 ^{ab}	0.02
Total proteins (g/dl)	3.76±0.65	4.16±0.94	3.63±0.6	4.04±0.18	3.78±0.42	0.81
FSH (mIU/ml)	14 ±3.67 ^b	12±3.74 ^b	16.5±4.81 ^b	26.67±3.51 ^a	18.25±4.72 ^b	0.00
LH (mIU/ml)	9.74±1.17 ^{ab}	7.92±1.71 ^{ab}	10.71±2.42 ^a	7.14±0.33 ^b	8.71±1.89 ^{ab}	0.04

Testosterone	1.36±0.38	1.43±0.31	1.21±0.12	1.53±0.46	1.57±0.32	0.54
(ng/ml)						

134 a, b, within the same line, values with the same letters are not significantly (P>0.05) different. n: Number of
 135 cavies. bw: body weight. T0-, group receiving 1 mL/kg bw of distilled water. T0+, group receiving 100 mg/kg
 136 bw of vitamin C; FSH, follicle stimulating hormone; LH, luteinizing hormone.

137 Effect of *Z. leprieurii* extract on some oxidative stress indicators in male guinea pigs

138 As shown in Table 6, the testicular activities of catalase (CAT), superoxide dismutase (SOD) and peroxidase glutathione
 139 (GPx) significantly increased (p<0.05) in cavies treated with aqueous extract of *Z. leprieurii* compared to the controls. The
 140 concentration of malondialdehyde (MDA) significantly decreased (p<0.05) in animals treated with the aqueous extract of
 141 *Z. leprieurii* irrespective of the dose compared to the controls.

142
 143 **Table 6:** Effects of aqueous extract of *Z. leprieurii* on oxidative stress indicators in male guinea pigs

Oxidative stress indicators	Controls		Doses of <i>Z. leprieurii</i>			P-value
	extract (mg/kg bw)		50	100	200	
	T0 ⁻ (n=12)	T0 ⁺ (n=12)	(n=12)	(n=12)	(n=12)	
MDA(μM/g of testis)	1.23±0.05 ^a	1.14±0.02 ^b	0.68±0.08 ^d	0.64±0.04 ^d	0.95±0.03 ^c	0.00
CAT(μM/min/g of testis)	6.68±0.33 ^c	6.21±0.04 ^d	7.89±0.0 ^g	7.72±0.05 ^b	12.25±0.02 ^a	0.00
SOD(U/min/mg of testicular protein)	0.32±0.01 ^c	0.47±0.03 ^c	0.81±0.16 ^b	0.87±0.14 ^{b^a}	1.04±0.07 ^a	0.00
GPx (μmol/min/g of testis)	26.99±0.68 ^c	25.84±0.95 ^d	30.05±0.44 ^b	32.83±0.55 ^a	29.44±0.3 ^b	0.00

144 Superscripts ^{a, b, c, d} within the same line differ (P<0.05), values with the same letters are not significantly
 145 (P>0.05) different. n, Number of cavies. bw:,body weight. T0-, group receiving 1 mL/kg bw of distilled water.
 146 T0+, group receiving 100 mg/kg bw of vitamin C; MDA, Malondialdehyde. CAT, Catalase; SOD, Superoxide
 147 dismutase; GPx, Glutathione peroxidase.

148 149 3.2. Discussion

150 The chemical screening of *Z. leprieurii* aqueous extract carried out in the present study revealed the presence of phenols,
 151 tannins, triterpenes, saponins flavonoids. These molecules have been reputed to have diverse activities (antioxidants,
 152 antibacterial, anti-inflammatory, antiseptic, antiparasitic, and immunomodulatory properties) susceptible to improve the
 153 animal reproductive characteristics [6, 5]

154 In this study, the effects of aqueous extract of *Z. lepreurii* fruit on reproductive parameters in male guinea pigs were
155 determined. The weights of the testis, epididymis, vas deferens and accessory sex glands increased dose-dependently in
156 animals submitted to aqueous extract of *Z. lepreurii* compared to the controls. These results are comparable to those
157 found in male rats treated with ethanol extract of the fruits of *Xylopiya aethiopica* at doses of 30, 100 and 300 mg/kg bw
158 [26]. In rats treated with aqueous extracts of *Psidium guajava* leaves (100, 200 and 300 mg/kg bw) [27, 28] and in male
159 guinea pigs treated with essential oil of *Psidium guajava* leaves (80, 100 and 120µl /kg bw), [29] the similar results were
160 also recorded. This observation might be explained by the action of antioxidant compounds such as phenols, flavonoids,
161 tannins, terpenoids and saponins revealed in this extract following the phytochemical tests [12]. These molecules could
162 have neutralized free radical attacks [30] or inhibiting enzymes responsible of their production and then protecting cells
163 against oxidative stress. The weight, size and secretory function of testes, epididymis and seminal vesicle are closely
164 regulated by androgens. [31]. In fact, Androgens, especially testosterone have anabolic properties which are
165 characterized by an increased synthesis of proteins and therefore muscle mass. Androgens then contribute to the
166 increased volume and weight of the testis and epididymis by stimulating protein synthesis [31, 32] The increase in the
167 serum content of testosterone observed in the present study would induce that of the sexual organ weights.

168 Sperm count, mobility and entire plasma membrane are considered to be important factors that affect fertility [33]. The
169 sperm with high density of spermatozoa could contained an important proportion of active spermatozoa capable to fertilize
170 an ovum [34]. It can also be diluted to have a high volume used to fertilize a large number of females in the case of
171 artificial insemination. In the other hand, spermatozoa with high mobility rate move rapidly in female genital track and has
172 an advantage to meet an ovum at the ampullary-isthmic junction and fuse with it [34]. In the present study, the *Z. lepreurii*
173 markedly increased the sperm count and the spermatozoa mobility in guinea pigs. This effect would be a result of *Z.*
174 *lepreurii* bioactive molecules with antioxidant properties. These molecules reduce the spermatozoa membrane and
175 nucleic acid attack by reactive oxygen species. This action subsequently decreases the death in the spermatozoa and
176 increases the spermatozoa mobility which is beneficial for reproduction.

177 The serum content of FSH in *Z. lepreurii* treated guinea pigs significantly increased at doses of 100 and 200 (mg/kg bw)
178 with respect to the controls. Some bioactive molecules of *Z. lepreurii* would have a possible action on pituitary gland cells
179 synthesizing the FSH. The elevated sperm count recorded in this study would result in FSH action. FSH has a key role in
180 the spermatogenesis. Together, FSH and testosterone support meiosis, exhibit an anti-apoptotic action on spermatocytes
181 and round spermatids. They act co-operatively to promote spermatid maturation and sperm release [35]. Testosterone is
182 the main male gonadal hormone produced by the interstitial cells of the Leydig in the testes in response to LH [36]. In this
183 study, both serum content in LH and testosterone tend to increase in *Z. lepreurii* treated guinea pigs. The *Z. lepreurii* rich
184 in bioactive molecules (phenols, alkaloids...) with antioxidant properties would protect the pituitary gland cells specialized
185 in the production of LH. The testosterone levels recorded would be a result of the relationship between LH and
186 testosterone.

187 Many environmental insults (poor quality of food and water intake, fluctuation of temperature, highly density in breeding...)
188 induce overproduction of reactive oxygen species responsible for animal cell membrane and nucleic acids impairment.
189 According to Tchoffo, [34] the substances with antioxidant properties inhibit the reactive oxygen species attacks and
190 subsequently improve the animal cell characteristics. In the present study, the malondialdehyde which is the major
191 resulting product from the membrane lipid peroxidation decreased significantly ($p<0.05$) in *Z. lepreurii* aqueous extract
192 treated male guinea pigs. Inversely, the activities of superoxide dismutase (SOD), catalase (CAT) and peroxidase
193 glutathione (GPx) increased significantly ($p<0.05$) in male guinea pigs exposed to aqueous extract of *Z. lepreurii*
194 compared to the controls. The phytochemical test on *Z. lepreurii* revealed that it contains bioactive molecules as phenols
195 and flavonoids. These molecules possess antioxidant properties responsible for *Z. lepreurii* aqueous extract effects. In
196 male animals, the spermatozoa membrane is characterized by a highly presence of polyunsaturated fatty acids rending
197 them susceptible to lipid peroxidation [34]. The *Zanthoxylum* due to their molecules with antioxidant properties could have
198 protect the spermatozoa from reactive oxygen species attacks and subsequently improve their characteristics as recorded
199 in the present study.

200 4. CONCLUSION

201 The results of this study revealed that the *Z. lepreurii* fruit aqueous extract efficiently improved reproductive
202 characteristics by increased level in reproductive hormones. It also induced an increase in testicular antioxidant enzymes
203 (SOD, catalase, peroxidase) activities and a decrease in MDA level. Based on these effects, the *Z. lepreurii* fruit aqueous
204 extract can be used in male animals to improve its reproductive performances.
205
206

207 ETHICAL APPROVAL

208 All experiments were carried out in compliance with the recommendations Guide of the National Academy of Sciences on
209 the care and use of laboratory Animals and approved by the department of animal science, FASA, University of Dschang
210

COMPETING INTERESTS

The authors declare that they have no conflict of interest.

REFERENCES

1. Raji Y, Fadare OO, Adisa RA, Salami SA. Comprehensive assessment of the effect of *Sphenocentrum jollyanum* root extract on male reproductive activity in albino rats. *Reprod Med Biol*. 2006; 23 (4):283-292; DOI: 10.1111/j.1447-0578.2006.00154.
2. Kandeel FR, Vivien KTK, Ronald SS. Male Sexual Function and its Disorder: Physiology, Pathophysiology, Clinical Investigation and Treatment. *Endocr rev*. 2001; 22 (3): 342–388.
3. Guay AT, Spark RF, Bansal S, Cunningham GR, Goodman NF, Nankin HR, Patak SM, Perez JB. American Association of Clinical Endocrinologist: Medical Guidelines for Clinical Practice for the evaluation and treatment of male sexual dysfunction. A Couple's Problem. *Endocrinol Pract*. 2003; 9(1):78–95.
4. Yakubu MT, Akanji MA, Oladiji AT. Male sexual dysfunction and methods used in Assessing Medicinal Plants with Aphrodisiac Potentials. *Pharmacogn Rev*. 2007; 1(1):49–56.
5. Tchoffo H, Kana JR, Ngoula F, Ngoumtsop VH, Ngouzeu MM, Tadondjou CD'A, Folack TLV. Effects of Ginger (*Zingiber officinale*, Roscoe) Essential Oil on Growth and Laying Performances, Serum Metabolites, and Egg Yolk Antioxidant and Cholesterol Status in Laying Japanese Quail. *J Vet Med*. 2019; Article ID 7857504, 8 pages.
6. Bertrand B, Collier JP. Bio markers exposure and effect, Editions Petiot from Terran. Thèse de doctorat publiée. Institut national polytechnique de Toulouse. Biotechnol, 2004 ; Taylor & Francis LTD.
7. Ufelle SA, Ukaejiofor EO, Neboh EE, Achukwu PU, Ghasi S, Ikekpeazu EJ, Maduka IC. The effect of crude methanolic leaf extract of *Bryophyllum pinnatum* on some haematological parameters in wistar rats. *Asian J Med Sci*. 2011; 3 (3): 121–124. DOI: [10.3923/rjpharm.2011.14.17](https://doi.org/10.3923/rjpharm.2011.14.17).
8. Yadav RNS, Agarwala M. Phytochemical analysis of some Medicinal Plants. *J Phytol*. 2011;3 (12):10–14.
9. Tillotson A. The One Earth Herbal Sourcebook: Everything You Need to Know About Chinese, Western, and Ayurvedic Herbal Treatments (Paperback); 2001 Edition Paperback 167. New York Twin stream/Kensington.
10. Baljinder S, Vikas A, Parveen B, Ranjit S, Dharmendra K. Pharmacological Potential of Plants used as Aphrodisiacs. *Inter J Pharm Sci Rev Res*. 2010; 5(1):104–113.
11. Yakubu MT, Akanji MA. Effect of aqueous extracts of *Massularia acuminata* stem on sexual behaviour of male Wistar rats. *Evid based complement Alternat Med*. 2011; Article ID 738103, 10 pages; dx.doi.org/10.1155/2011/738103.
12. Etuk EU and Muhammad AA. Fertility enhancing effects of aqueous stem bark extract of *Lophira lanceolata* in male Spargue dawley rats. *Inter J Plant Physiol Biochem*. 2009; 1(1): 001-004.
13. Sarieh S, Javad-Sharifi R, Farzaneh MR, Mohammad RS, Mohammad RA. Effects of Aqueous Root Extracts of *Anacyclus pyrethrum* on Gonadotropins and Testosterone Serum in Adult Male Rats. *Am J Phytomed Clin Therap*. 2014;2(6):767-772
14. Ngandjui A, Ngaha N M, Kenmogne H, Koloko BL, Massoma LD, Bend EF, Oundoum OP, Domkam J, Moundipa P, Dimo T. Evaluation of the fertility activity of the aqueous leaves extract of *Zanthoxylum macrophylla* (Rutaceae) on male rats. *J Phytopharmacol*. 2017; 6(5): 277-281.
15. Ekaluo UB, Erem FA, Omeje IS, Ikpeme EV, Ibiang YB, Ekanem EB. Aqueous leaf extract of guava: A non-toxic male fertility booster. *IOSR J. Environ. Sci. Toxicol. Food Technol*. 2013; 3: 33-35
16. Abdou BA. Contribution to the study of a functional food based on spices from Cameroon: physico-chemical and functional characterization. Thèse de Doctorat PhD, Université de Ngaoundéré. 2009, 117P.
17. NAS. Guide for the care and use of laboratory animals, Eight Ed. NW, Washington, DC: *Natl Acad Pres*. 2013, p11-132.
18. Banso A. and Ngbede JE. Phytochemical screening and in vitro antifungal properties of Fagaraz anthoxyloides. *J Food, Agri Environ*. 2006; 4(3-4): 8-9.
19. Ngbede J, Yakubu RA, Nyam DA. Phytochemical screening for active compounds in *Canarium scheinfurthii* leaves from Jos North, plateau state, Nigeria. *Res J Biol Sci*. 2008; 3(9) 1076–1078.
20. Boiti C, Castellini C, Theau-Clément M, Besenfelder U, Liguori, L, Renieri T, Pizzi F. Guidelines for the handling of rabbit bucks and semen. *World Rabbit Sci*. 2005; 13:71– 9.
21. Kumi-Diaka J. Subjecting canine semen to the hypo-osmotic test. *Theriogenology*. 1993; 39(6):1279-1289.
22. Nilsson UA, Olsson LI, Carlin G. Inhibition of lipid peroxidation by spin labels. *J Biol chem*. 1989; 264:11131-11135.
23. Dimo T, Tsala DE, Dzeufiet DPD. Effects of *Alafia multiflora* stapf on lipid peroxidation and antioxidant enzyme status in carbon tetrachloride-treated rats. *Pharmacol online*. 2006; 2:76-89.
24. Sinha AK. Colorimetric assay of catalase. *Analytical biochemistry*. 1972 ; 47: 389-394.

- 270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
25. Kodjio N, Atsafack SS, Njateng GSS, Sokoudjou JB, Kuate J-R, Gatsing G. The typhoid fever induced in Wistar rats model. *J Adv Medical Pharm Sci.* 2016; 7(3):1-13. DOI: 10.9734/JAMPS/2016/24949.
 26. Woode E, Alhassan A, Chrissie S, Abaidoo. Effect of ethanolic fruit extract of *Xylopia aethiopica* on reproductive function of male rats. *Int J Pharm Biomed Res.* 2011; 2:161-165.
 27. Ekaluo UB, Erem FA, Omeje IS, Ikpeme EV, Ibiang YB. Aqueous leaf extract of guava: a non-toxic male fertility booster. *J Environ Sci, Toxicol Food Technol.* 2013a; 3:21-23.
 28. Ekaluo UB, Erem FA, Omeje IS, Ikpeme EV, Ibiang YB. Is aqueous leaf extract of guava spermatotoxic in rat? *J Environ Sci, Toxicol Food Technol.* 2013b; 3:21-23.
 29. Ngoula F, Nouboudem SC, Kenfack A, Tadondjou CD, Kana JR, Kouam J-M, Tsafack B, Ngoumtsop H, Nguemdjo TM, Akassa H, Ousmane IAD, Kamtchouing P, Galeotti M, Tchoumboue J. Effect of Guava (*Psidium guajava*) Leaves Essential Oil on Some Reproductive Parameters in Male Guinea Pig (*Cavia porcellus*). *Biol Syst.* 2014; 3:125; DOI:10.4172/2329-6577.1000125.
 30. Hodek P, Trefil P, Stiborova M. Flavonoids-potent and versatile biologically active compounds interacting with cytochromes P450. *Chem-Biol Interact.* 2002; 139 :1-2. DOI: 10.1016/S0009-2797(01)00285-X.
 31. Hazard J, Perlemuter L, Abramovici Y, Bourgeon M. *Endocrinologie.* Masson, Paris, 2000; 484p.
 32. Gayraud V. *Mammalian reproductive physiology.* Ecole Nationale Vétérinaire. Toulouse. 2007 ; 198p.
 33. Kobeasy MI, El-Naggar AY, Abdallah AA. A novel method for protective role against reproductive toxicity of carbofurem in male rats using palm pollen grains and vanadyl (II) folate as a new compound. *J Chem Pharm Res.* 2015; 7(4):1142-1148.
 34. Tchoffo H, Kana JR, Ngoula F, Folack, TLV, Adoum G, Moussa MO, Ngouzeu MWM. Growth performance, serum biochemical profile, oxidative status and fertility traits in male Japanese quail fed on ginger (*Zingiber officinale*, Roscoe) essential oil. *Vet Med Inter.* 2018; Article ID 7682060, 8p, <https://doi.org/10.1155/2018/7682060>.
 35. Chauhan NS, Dixit VK. Effects of *Bryonia laciniosa* seeds on sexual behavior of male rats. *Int J Impot Res.* 2010; 22:190-195.
 36. Chauhan NS, Rao CV, Dixit VK. Effect of *Curculigo orchioides* rhizomes on sexual behavior of male rats. *Fitoterapia.* 2007; 78:530-534.