

## Original Research Article

# Growth and Yield Response of Bush Bean (*Phaseolus vulgaris* L.) as Influenced by Different Levels of Nitrogen and Phosphorus Application

### ABSTRACT

The experiment was conducted at Horticulture Farm at Sher-e-Bangla Agricultural University, Bangladesh during the period December, 2014 to March 2015 to evaluate the effect of different levels of nitrogen and phosphorous on the growth and yield of bush bean. The two factor experiment was laid out in Randomized Complete Block Design with three replications. The treatment was comprised of two factors- Factor A: levels of nitrogen i) 0 kg/ha, ii) 20 kg/ha, iii) 40 kg/ha and Factor B: levels of phosphorous – i) 0 kg/ha, ii) 50 kg/ha, iii) 75 kg/ha, iv) 100 kg/ha. The results revealed that most of the growth and yield contributing parameters were significantly influenced by the different levels of nitrogen and phosphorous application. The maximum promotive effect on growth and yield of bush bean was associated with 40 kg N/ha and 75 kg P<sub>2</sub>O<sub>5</sub>/ha. Again their combined application enhanced maximum vegetative growth and with higher pod yield and seed yield. Therefore, application of 40 kg N/ha with 75 kg P<sub>2</sub>O<sub>5</sub>/ha can be conducive for bush bean cultivation in Bangladesh with higher yield.

Comment [H1]: P or P2O5?

*Keywords: growth, yield, bush bean, Nitrogen, Phosphorous*

### 1. INTRODUCTION

Bush bean (*Phaseolus vulgaris* L.) is a versatile short duration legume crop with high grain yielding potential and can be used both as pulses and vegetable. Although the crop has originated in South America it has gained popularity all over the world for its high yielding potential and diversified use. Immature pods are marketed fresh, frozen and canned. Its dry seeds are used in preparations with fish, meat and other vegetables. Foliage of the plant may also provide hay, silage and green manures. After harvest, plants can be used as fodder to feed the cattle, sheep and horses. Its edible pods supply huge protein, carbohydrate, fat, fiber, thianin, riboflavin, Ca and Fe [1]. Due to its sublime nutritional quality and diversified use, it has gained huge popularity in Bangladesh over the recent years. Besides these, the crop has eminent export potential which has drawn the attention of Agriculture industry and exporters. So there is tremendous scope for the cultivation of this important vegetable crop in our country with the export potential. However, to ensure proper growth and yield, the crop needs optimum nutrient management. Nitrogen is one of the key elements for growth and development of a crop plants [2]. Meanwhile, it has been reported

35 that, unlike other leguminous crops, it does not nodulate with the native rhizobia [3].  
36 | Therefore, nitrogen input during production is **very** crucial for this crop. Nitrogen deficiency  
37 constraints leaf area expansion, enhances leaf senescence, inhibits photosynthetic rate in  
38 most of the crops and consequently reduces the crop productivity [4, 5]. In addition to this,  
39 deficiency of phosphorous is now considered as one of the major constraints to successful  
40 production of legumes and upland crops in Bangladesh [6]. The most obvious effect of  
41 phosphorus is on the root development particularly of the lateral and fibrous rootlets that are  
42 essential to fix the atmospheric nitrogen in legume crops. Phosphorous also make its  
43 contribution through seed formation. In case of application of various fertilizer doses, there  
44 | were significant differences in pod number per plant in bush bean [7]. **In experiment, t**The  
45 plant height, number of branches, length of pod per plant and seed yield per pod increase  
46 with successive increase in the dose of nitrogen as well as phosphorous [8]. Therefore,  
47 optimum combination of nitrogen and phosphorous may bring about considerable increase in  
48 the yield of Bush bean due to their complementary effects. A detailed systematic study is  
49 needed to find out the requirements and effect of nitrogen and phosphorous for maximizing  
50 the yield of bush bean in Bangladesh. Considering the above, the present investigation was  
51 undertaken to study the growth and yield response of bush bean under different levels of  
52 nitrogen and phosphorous to find out the best combination of nitrogen and phosphorous for  
53 | maximizing the production of **B**bush bean.

## 54 **2. MATERIAL AND METHODS**

### 55 **2.1 Experimental site**

56  
57  
58  
59 The experiment was conducted at the Horticultural farm of Sher-e-Bangla Agricultural  
60 University, Dhaka, Bangladesh from December 2014 to March 2015. The location of the  
61 experimental site was 23074/N latitude and 90035/E longitude and at an elevation of 8.2 m  
62 from sea level. The climate of experimental site was under the subtropical climate,  
63 characterized by three distinct seasons, the winter season from November to February and  
64 the pre-monsoon or hot season from March to April and the monsoon period from May to  
65 October. **The soil of the experimental area belongs to the Modhupur Tract (AEZ No 28).** It  
66 had shallow red brown terrace soil. The selected plot was medium high land and the soil  
67 series was Tejgaon.

### 68 **2.2 Planting material**

69  
70  
71 The variety of bush bean used in the experiment was BARI jharsheem. The seeds were  
72 collected from Horticulture Research Centre, Bangladesh Agriculture Research Institute  
73 (BARI), Gazipur, Bangladesh.

### 74 **2.3 Experimental design and treatments**

75  
76  
77 The two factor experiment was laid out in Randomized Complete Block Design with three  
78 replications. The experimental plot was first divided into three blocks. Each block consisted  
79 of 12 unit plots. Thus the total number of plots was 36. The treatment was comprised of two  
80 factors- Factor A: levels of nitrogen i) 0 kg/ha, ii) 20 kg/ha, iii) 40kg/ha and Factor B: levels of  
81 | **phosphorous  $P_2O_5$  - i) 0 kg/ha, ii) 50 kg/ha, iii) 75 kg/ha, iv) 100 kg/ha.** Different  
82 combinations of nitrogen and phosphorous were assigned randomly to each block as per  
83 design of the experiment. The size of the unit plot was 3m×4. A distance of 0.75m between  
84 the plots and 1m between the blocks were kept.

### 85 **2.4 Growth condition of Bush bean and measurement of parameters**

Comment [H2]: Only the soil?

Formatted: Subscript

Formatted: Subscript

88 Seedlings were grown following proper methods and all of the cultural practices were done  
89 properly. Application of manure and fertilizers were applied as per treatment. Healthy and  
90 uniform sized seedlings were transplanted in the main field. Intercultural practices were done  
91 as per requirements. For controlling leaf caterpillars *Nogos* @ 1 ml/L water were applied two  
92 times at an interval of 10 days starting soon after the appearance of infestation. Immature  
93 green pods were harvested at tender stage, suitable for use as vegetable. First harvest was  
94 done at 55 days after sowing (DAS) and was weighted to estimate the fresh pod yield. Again  
95 the rest of the pods were harvested at mature stage when the pods become yellow and fully  
96 dry. These seeds were collected from the pods and sun dried seeds were weighted to  
97 estimate the seed yield.

## 98 2.5 Data collection and statistical analysis

99  
100  
101 Different yield contributing data have been recorded from the mean of five harvested plants  
102 which was selected at random of each unit plot of every harvesting stage. The plants in the  
103 outer rows and the extreme end of the middle rows were excluded from the random  
104 selection to avoid the border effect. The height of the plants was measured from the ground  
105 level to the tip of the highest leaves. Green pods were harvested from each unit plot at 4  
106 days interval and their total weight was recorded. Harvesting was done for four times and  
107 their weight was recorded in each unit plot and expressed in kilogram (Kg). The green pod  
108 yield per plant was finally converted to yield per hectare and expressed in ton (t). The data  
109 obtained for different parameters were statistically analyzed to find out the significance  
110 difference of variety and different fertilizer application on yield and yield contributing  
111 characters of *cabbage*. The mean values of all the characters were calculated and analysis  
112 of variance was performed by the 'F' (variance ratio) test. The significance of the difference  
113 among the treatment combinations means was estimated by the Duncan's Multiple Range  
114 Test (DMRT) at 5% level of probability.

Comment [H3]: Really?

## 115 3. RESULTS AND DISCUSSION

### 116 3.1 Plant Height (cm)

117  
118  
119  
120 Different levels of nitrogen and phosphorous exhibited significant variation in respect of plant  
121 height of bush bean **at different days** after sowing (Table 1). In all the dates of observation  
122 plant height gradually increased with the increasing nitrogen level. At final harvest the  
123 maximum plant height (43.82cm) was obtained from the plant grown with 40kg N/ha and the  
124 minimum (37.85cm) was recorded from control treatment. Nitrogen is **the most essential**  
125 nutrient for plant growth which might have facilitated the highest plant height N<sub>2</sub>. Similar  
126 result was supported by [9] who reported that plant height increased with increasing rate of  
127 N doses up to 120 kg/ha.

Comment [H4]: Only for this?

Comment [H5]: So, the results were not similar.

128  
129 Plant height was also significantly influenced by the application of different levels of  
130 phosphorous at different growth stages (Table 2). Plant height was gradually increased with  
131 the passage of time up to the final harvest. At harvest, the tallest plant (41.98 cm) was  
132 produced **on soil with application of from the plant growth with** 100 kg P<sub>2</sub>O<sub>5</sub>/ha whereas  
133 the shortest plant was recorded in the controlled treatment. Similar result was reported by  
134 [10,11].

Comment [H6]: Why?

135  
136 In case of combined effect of **Nitrogen nitrogen** and **Phosphorousphosphorous**, the  
137 maximum plant height (44.60 cm) was observed from the treatment combination N<sub>2</sub>P<sub>2</sub>  
138 whereas the minimum plant height was observed from the control treatment. The present  
139 result is in agreement with the findings of [9].  
140

141 **3.2 Number of pod/plant**

142

143 The number of pod per plant differed significantly due to the application of different levels of  
144 nitrogen (Table 1). The maximum number of pods per plant (13.26) was produced from the  
145 40kg N/ha. On the contrary, minimum number of pod per plant was recorded from the control  
146 treatment. It might be due to the adequate supply of nitrogen which facilitated the increased  
147 number of pod bearing branches.

Comment [H7]: in which nitrogen contributes to the formation of pods?

148

149 The number of pod per plant was also showed significant effect on increasing the number of  
150 pod per plant (Table 2). The maximum number of pods per plant (13.62) was recorded from  
151 50 kg P<sub>2</sub>O<sub>5</sub>/ha meanwhile the minimum number of pod per plant was recorded from the  
152 control treatment. The result is in conformity with the [11,12, 13].

153

154 Combined effect of different levels of nitrogen and phosphorous on the number of pod per  
155 plant showed significant variation (Table 3). The maximum number of pods per plant (13.90)  
156 was recorded from the combined treatment of 40kg N/ha and 50 kg P<sub>2</sub>O<sub>5</sub>/ha. On the contrary  
157 minimum number of pods per plant was found from the N<sub>2</sub>P<sub>3</sub>. It has been reported that P  
158 uptake reduced above 60 kg P<sub>2</sub>O<sub>5</sub>/ha.

Comment [H8]: Why?

159

160 **3.3 Pod length (cm)**

161

162 Nitrogen had significant influence in respect to pod length (Table 1). Pod length gradually  
163 increased with the increasing dose of nitrogen fertilizer. The highest length of green pod of  
164 bush bean (17.51 cm) was found in the crop grown with the higher dose of nitrogen (40 kg  
165 N/ha) and lowest was observed from the control treatment. Length of the green pod was also  
166 influenced significantly by the different doses of phosphorous (Table 2). The maximum  
167 length of green pod (16.99 cm) was obtained from 100 kg P<sub>2</sub>O<sub>5</sub>/ha. On the other hand  
168 minimum pod length was recorded from 50 kg P<sub>2</sub>O<sub>5</sub>/ha. It was observed that the interaction  
169 effect of different doses of nitrogen and phosphorous on green pod length was significant  
170 (Table 3). The highest pod length (18.05 cm) was recorded from the treatment combination  
171 40 kg N/ha and 75 kg P<sub>2</sub>O<sub>5</sub>/ha and the lowest was recorded from the control treatment.

Comment [H9]: There are no explications.

172

173 **3.4 Pod weight/plant**

174

175 A significant variation was observed in respect of pod weight per plant ad different level of  
176 nitrogen (Table 1). The highest pod weight per plant (85.26 g) was obtained from 40 kg N/ha  
177 and the lowest pod weight (48.57 g) per plant was recorded from the control. The level of  
178 phosphorous was also significantly influenced the pod weight per plant of bush bean in the  
179 similar way. The highest pod weight (72.27 g) was obtained from the plant which was treated  
180 with 75 kg P<sub>2</sub>O<sub>5</sub>/ha and the lowest (62.10 g) from the control treatment. The combined effect  
181 of different levels of nitrogen and phosphorous on pod weight per plant was found  
182 statistically significant (Table 3). The maximum pod weight per plant (89.16 g) was obtained  
183 from the treatment N<sub>2</sub>P<sub>1</sub> whereas the minimum was obtained from the control treatment.

Comment [H10]: There are no explications.

184

185 **3.5 Pod yield**

186

187 The result indicated that nitrogen had significant effect on pod yield per hectare (Table  
188 1).The maximum pod yield (16.56 t/ha) was obtained from 40 kg N/ha and the minimum pod  
189 yield (10.04 t/ha) in control. Phosphorous also influenced significantly the pod yield per  
190 hectare (Table 2).The combined effect of different level of nitrogen and phosphorous on pod  
191 yield per hectare was found to be statistically significant (Table 3). The highest pod yield per  
192 hectare (18.61 t/ha) was achieved from the treatment combination 40 kg nitrogen per

193 hectare with 75 kg P<sub>2</sub>O<sub>5</sub>/ha whereas the lowest value per hectare (9.22 t/ha) was found from  
194 the control combination. Similar kind of result was reported by [14,15].

195  
196  
197

### 3.6 No of seeds/pod

Comment [H11]: Number?

198 The number of mature seeds per pod was significantly influenced by the application of  
199 nitrogen (Table 1). The highest seed per pod (5.85) was obtained from the plant receiving 40  
200 kg N/ha. The lowest number of mature seed per pod (5.31) was recorded from control  
201 treatment. This result might be due to the better growth and development and larger pod  
202 formation with higher rate of N application. Similar kind of result was reported by [1].  
203 Different doses of phosphorous also significantly influenced the number of seed per pod  
204 (Table 2). Number of seed per pod increased with increasing rates of phosphorous fertilizer.  
205 It was maximum (5.63) at 100 kg P<sub>2</sub>O<sub>5</sub>/ha which was statistically similar to 75 kg P<sub>2</sub>O<sub>5</sub>/ha  
206 whereas the minimum number of seed (5.13) was obtained with zero phosphorous. There  
207 was also significant interaction between nitrogen and phosphorous in respect of seed per  
208 pod (Table 3). The maximum number of seeds per pods (6.16) was recorded from the  
209 treatment combination 40 kg N/ha with 50 kg P<sub>2</sub>O<sub>5</sub>/ha whereas the minimum number of seed  
210 per pod was recorded in the control treatment.

Comment [H12]: ????

Comment [H13]: Why, when phosphorus is used alone, the best value is one and when in interaction with nitrogen, the result is another?

211  
212

### 3.7 Thousand Seed weight

213 A significant variation was observed in respect of 1000 seed weight due to different nitrogen  
214 levels (Table 1). Maximum weight of 1000 seed (230.75 g) was recorded from application of  
215 40 kg N/ha. Meanwhile minimum was obtained from the control treatment. The level of  
216 phosphorous were also significantly influenced the 1000 seed weight (Table 2). The highest  
217 1000 seed weight (236.00 g) was recorded at 75 kg P<sub>2</sub>O<sub>5</sub>/ha while the lowest was obtained  
218 from the control treatment. The combined effect of nitrogen and phosphorous was found to  
219 be significant (Table 3). The highest weight of 1000 seed (255.00 g) was obtained from the  
220 treatment combination of 40 kg N/ha with 75 kg P<sub>2</sub>O<sub>5</sub>/ha. The lowest 1000 seed weight  
221 (211.00 g) was recorded from the control treatment.

222  
223  
224

### 3.8 Seed yield/plot

225 The result indicated that nitrogen had significant effect on seed yields per plot (Table 1).  
226 Seed yields per plot were varied from 2.65 kg to 3.3 kg. The maximum seed yield per plot  
227 was produced from the 40 kg N/ha and minimum seed yield per plot was obtained from the  
228 control.

229 Phosphorous also influenced significantly the seed yield per plot (Table 2). Maximum seed  
230 yield per plot (3.08 kg) was obtained with the 75 kg P<sub>2</sub>O<sub>5</sub>/ha while the minimum was  
231 recorded in the control treatment.

232 The combined effect of different levels of nitrogen and phosphorous on seed yield per plot  
233 was found to be statistically significant (Table 3). The highest yield per plot (3.53 kg) was  
234 obtained from the treatment combination of 40 kg N/ha with 50 kg P<sub>2</sub>O<sub>5</sub>/ha whereas the  
235 lowest value was found from the control treatment.

236  
237

### 3.9 Seed Yield (t/ha)

238 The result indicated that nitrogen had significant effect on seed yield per hectare (Table 1).  
239 The maximum seed yield (2.71 t/ha) was obtained from 40 kg N/ha and the minimum (2.23  
240 t/ha) in control treatment.  
241

242 Phosphorous also influenced significantly the seed yield per hectare (Table 2). The maximum seed yield per hectare (2.58) was obtained  
 243 with the 75 kg P<sub>2</sub>O<sub>5</sub>/ha whereas the minimum was obtained from the control treatment. The combined effect of different levels of nitrogen  
 244 and phosphorous on seed yield per hectare was found to be statistically significant (Table 3). The highest yield per hectare (2.96 t) was  
 245 obtained from the treatment combination of 40 kg N/ha with 50 kg P<sub>2</sub>O<sub>5</sub>/ha whereas the lowest value was found from the control treatment.  
 246 The result is in agreement with [ 16, 17].  
 247

248 **Table 1: Effect of Nitrogen on the growth and yield contributing attributes of bush bean**

Nitrogen Level	Plant height (cm)	Number of pod plant <sup>-1</sup>	Pod length (cm)	Pod weight plant <sup>-1</sup>	Pod yield (t ha <sup>-1</sup> )	No of seeds pod <sup>-1</sup>	1000 seed weight (g)	Seed yield plot <sup>-1</sup> (kg)	Seed yield (tha <sup>-1</sup> )
N <sub>0</sub>	37.85 c	13.22 b	14.85 b	48.57 c	10.04 c	5.13 c	225.00 b	2.65 c	2.23 c
N <sub>1</sub>	41.34 ab	13.24 ab	16.58 ab	66.93 b	13.62 ab	5.35 b	228.50 ab	2.94 ab	2.42 b
N <sub>2</sub>	43.83 a	13.26 a	17.51 a	85.26 a	16.56 a	5.85 a	230.75 a	3.23 a	2.71 a
LSD ( 0.05)	0.988	0.065	0.488	0.065	0.100	0.088	1.955	0.037	0.046
CV	4.25	3.20	4.16	6.40	2.21	1.63	7.56	3.54	4.12

Formatted: Highlight

249 *Means in a same column followed by different letter (s) are significantly different at P<0.05*

250  
251

252 **Table 2: Effect of Phosphorous on the growth and yield contributing attributes of bush bean**

Phosphorus Level	Plant height (cm)	Number of pod plant <sup>-1</sup>	Pod length (cm)	Pod weight plant <sup>-1</sup>	Pod yield (tonha <sup>-1</sup> )	No of seeds pod <sup>-1</sup>	1000 seed weight (g)	Seed yield plot <sup>-1</sup> (kg)	Seed yield (tonha <sup>-1</sup> )
P <sub>0</sub>	39.83 c	13.20 b	16.29 b	62.10 c	10.76 c	5.13 b	223.33 c	2.65b	2.17 b
P <sub>1</sub>	40.66 b	13.62 a	15.93 c	68.25 b	13.51 b	5.44 b	227.00 bc	2.94 b	2.48 b
P <sub>2</sub>	41.55 ab	13.12 b	16.03 bc	72.27 a	15.35 a	5.56 ab	236.00 b	3.08 a	2.58 a
P <sub>3</sub>	41.98 a	12.99 c	16.99 a	65.08 bc	14.55 ab	5.63 a	226.00 bc	3.06 a	2.58 a
LSD ( 0.05)	1.142	0.075	0.564	0.415	0.115	0.10	2.258a	0.043	0.053
CV	4.55	1.19	4.65	5.87	4.12	3.07	8.67	2.43	4.11

Formatted: Highlight

253 *Means in a same column followed by different letter (s) are significantly different at P<0.0*

Comment [H14]: ????

254 | **Table 3: Combined effect of Nitrogen and Phosphorous on the growth and yield contributing attributes of bush bean**

Nitrogen Level	Plant height (cm)	Number of pod plant <sup>-1</sup>	Pod length (cm)	Pod weight plant <sup>-1</sup>	Pod yield (tonha <sup>-1</sup> )	No of seeds pod <sup>-1</sup>	1000 seed weight (g)	Seed yield plot <sup>-1</sup> (kg)	Seed yield (tonha <sup>-1</sup> )
N <sub>0</sub> P <sub>0</sub>	35.27d	13.20 b	15.04 c	40.66 e	9.22 d	4.73 d	211.00 d	2.21 d	1.89 d
N <sub>0</sub> P <sub>1</sub>	36.33 cd	13.46 b	13.56 d	42.66 de	9.80 cd	4.97 cd	235.00 b	2.50 cd	2.13 c
N <sub>0</sub> P <sub>2</sub>	39.82 c	12.83 bc	15.16 c	57.71 c	11.81 c	5.31 c	225.00 bc	2.87 c	2.41 bc
N <sub>0</sub> P <sub>3</sub>	39.97 c	13.33 b	15.65 c	53.23 cd	10.93 c	5.54 ab	231.00 bc	2.91 bc	2.49 bc
N <sub>1</sub> P <sub>0</sub>	40.46 bc	12.96 bc	16.09 bc	63.72 bc	10.15 c	5.22 c	219.00 cd	2.77 c	2.19 c
N <sub>1</sub> P <sub>1</sub>	41.03 bc	13.51 b	16.17 bc	72.95 b	13.58 bc	5.22 c	218.00 cd	2.80 c	2.35 bc
N <sub>1</sub> P <sub>2</sub>	41.30 b	13.53 b	16.22 bc	71.44 b	15.63 b	5.36 ab	227.00 bc	3.16 b	2.62 bc
N <sub>1</sub> P <sub>3</sub>	42.57 b	13.45 b	17.82 ab	59.60 c	15.15 b	5.58 ab	236.00 b	3.07 bc	2.52 bc
N <sub>2</sub> P <sub>0</sub>	43.77 b	13.50 b	17.76 ab	81.93 ab	12.91 bc	5.46 ab	228.00 c	2.95 c	2.42 bc
N <sub>2</sub> P <sub>1</sub>	43.53 b	13.90 a	18.05 a	89.16 a	17.14 ab	6.16 a	229.00 c	3.53 a	2.96 a
N <sub>2</sub> P <sub>2</sub>	44.60 a	13.00 b	16.71 bc	87.67 ab	18.61 a	6.01 a	255.00 a	3.22 b	2.73 b
N <sub>2</sub> P <sub>3</sub>	43.40 b	12.20 c	17.51 ab	82.82 ab	17.57 ab	5.75 ab	223.00 bc	3.19 b	2.73 b
LSD(0.05)	1.978	0.131	0.977	0.720	0.200	0.177	3.910	0.075	0.09
CV (%)	2.85	2.59	3.54	3.44	4.87	1.91	3.01	1.52	2.19

255 | Means in a same column followed by different letter (s) are significantly different at P<0.05; Factor A: levels of nitrogen i) 0 kg/ha, ii) 20 kg/ha, iii) 40 kg/ha  
 256 | and Factor B: levels of **phosphorous P<sub>2</sub>O<sub>5</sub>** – i) 0 kg/ha, ii) 50 kg/ha, iii) 75 kg/ha, iv) 100 kg/ha.

Formatted: Highlight

Formatted: Subscript

Formatted: Subscript

257  
258  
259  
260  
261  
262  
263  
264

#### 4. CONCLUSION

The result of the present study revealed that both nitrogen and phosphorous significantly influenced on the green pod and seed yield of bush bean. Application of 40 kg N/ha with 75 kg P<sub>2</sub>O<sub>5</sub>/ha has emerged as the best treatment for increasing growth and yield contributing attributes of bush bean and therefore can be conductive for bush bean cultivation in Bangladesh with higher yield.

265  
266

#### REFERENCES

267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308

1. Chaudhari CS, Mendhe SN, Pawar WS, Lngole AS and Nikam RR. Nutrient Management in French bean. J. of Soil and Crops.2001; 11(1): 137-139.
2. Rana NS and Singh R. Effect of nitrogen and phosphorous on growth and yield of French bean. Indian J.Agron.1998; 43(2): 367-370.
3. Singh Ak, and Singh SS. Effect of planting dates, nitrogen and phosphorus level on yield contributing factors in French bean. Legume Research.2000; 23(1): 33-36.
4. Tewari JK and Singh SS. Effect of nitrogen and phosphorous on growth and seed yield of French bean (*Phaseolous vulgarais*).Vegetable Sci. 2000; 27(2): 172-175.
5. Reddy M, Malla, Padmaja B, Reddy and Ram R. Response of French bean to irrigation schedules and nitrogen levels in Talangana region of Andhra Pradesh. J. Food legumes. 2008; 23(1): 38-40.
6. Mozumder SN, Moniruzzaman M, Islam MR and Alam SN. Effect of planting time and spacing on the yield performance of bush bean (*Phaseolus vulgaris* L.) in the eastern hilly area of Bangladesh. Legume Res .2003 26(4): 242- 247.
7. Kaisar MO, Zaman S, AHoque AKMS,. Rahman SML and Zaman MM. Effect of nitrogen and phosphorus on growth, yield and profitability of French bean. Bangadesh J. Pro. Sci. & Tech. 2007; 5(2): 373-376.
8. Dhanjal R, Prakash O and Ahlawat IPS. Physiological variation in French bean (*Phaseolus vulgaris*) cultivars as affected by plant density and nitrogen. Indian J. Plant Physiol. 2003; 8: 34-37.
9. Ferdous AKM, Khaliq QA, Hoque MM, Sirajul Karim AJM and Bhuyian MSA. Effect of nitrogen fertilizer on growth, nitrogen and phosphorus uptake and yield in edible podded pea. Bangladesh Agron. J. 2004; 10 (1 & 2): 133-140.
10. Kumar M, Sinha KK and Roy Sharma RP. Effect of organic manure, NPK and boron application on the productivity of French bean in sandy loam soil of north Bihar. Indian J. Pulses Res. 2004; 17:42-44.
11. Zaman-Allah M, Sifi BL, Taief B, Aouni E, Drevon JJ. Symbiotic response to low phosphorus supply in two common bean (*Phaseolus vulgaris* L.) genotypes. Symbiosis. 2007; 44:109–113.

- 309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335
12. Bargaz A, Ghoulam C, Faghire, Aslan A, Drevon JJ. The nodule conductance to the O<sub>2</sub> diffusion increases with high phosphorus content in the Phaseolus vulgaris-rhizobia symbiosis. Symbiosis.2011; 53:157–164.
  13. Hafez M, Magda M, Asmaa R, Mahmoud M. Response of snap bean (Phaseolus vulgaris L.) to nitrogen fertilizer source. Ann. Agric. Sci., Mashtohor. 2004; 42(1):261–270.
  14. Voisin AS, Bourion V, Duc G, Salon C. Using an ecophysiological analysis dissect genetic variability and to propose an ideotype for nitrogen nutrition in pea. Ann Bot.2007; 100:589–598.
  15. Saxena K, Arun-Srivastava K, Singh RB. Response of French bean to nutrient application (NPK) in relation to physiological traits and their consequent effect on yield. Farm Sci J. 2003; 12(2):150–152.
  16. Piha MI, Munns DN. Nitrogen fixation potential of beans (Phaseolus vulgaris L.) compared with other grain legumes under controlled conditions. Plant Soil. 1987; 98(2):75–82.
  17. Abdel-Mawgoud AMR, EL-Desuki M, Salman SR, Abou-Hussein SD. Performance of some snap bean varieties as affected by different levels of mineral fertilizers. Agron. J. 2005; 4(3):242–247.

Formatted: Portuguese (Brazil)