

Original Research Article

Long term effects of integrated plant nutrition system on rice yield, nitrogen dynamics and biochemical properties in soil of rice-rice cropping system

Comment [H1]: It needs to explain, better, some results, besides just presenting the results obtained.

ABSTRACT

Long term fertilization from manure and fertilizer influences crop yield as well as soil nutrient cycling. Therefore, a field experiment was conducted from 2010 to 2014 in order to observe the long term effects of integrated application of manure and fertilizers on rice yield, soil nitrogen dynamics and soil biochemical properties. The field experiment was carried out in a rice-fallow-rice cropping pattern under wetland condition with four fertilizer management practices: NPKS fertilizer as soil test based (STB), Cow dung (CD) @ 3 t ha⁻¹ + NPKS as integrated plant nutrition system (IPNS) [CD+IPNS], Poultry manure (PM) @ 2 t ha⁻¹ + NPKS as IPNS [PM+IPNS] and N control. The rice grain yield, soil N dynamics and other the biochemical properties assessed in our study significantly varied with the organic sources used in IPNS system. The annual grain yield of double cropped rice with PM + IPNS was at par with the STB treatment over the five cropping years while CD+IPNS responded from the third year. After five years of rice cropping PM+IPNS showed better impact on soil nitrogen dynamics and biochemical properties compared to STB fertilizer management. Considering the soil health, our study suggests that PM+IPNS could be a good practice for sustainable rice production in long run reducing the use of chemical fertilizer.

Keywords: rice yield, mineral nitrogen, nitrogen balance, urease enzyme activity, free living nitrogen fixing bacteria, soil depth

1. INTRODUCTION

Rice-fallow-rice is the dominant cropping system in Bangladesh which occupied about 27% of the net cropped area [1]. Among the major nutrient elements that limit rice crop growth and yield, nitrogen (N) considered as the most important. Farmers generally meet the crop N demand by applying chemical N fertilizer because of its effectiveness in improving yield [2]. However, the intensive application of chemical N fertilizer in the cropping system has raised the concerns of sustainable crop production. Moreover, the injudicious use of chemical N fertilizer coupled with modern high yielding varieties and little or no use of organic matter lead to soil organic matter deficiency, soil acidity and poor impoverishment of the soil physical properties [3, 4, 5]. On the other hand, manure alone cannot substitute inorganic fertilizer to maintain the desired level of yield of the high yielding varieties [6]. Therefore, the combined application of manure and fertilizer has emerged as an effective approach for sustainable crop production. Many studies have reported significant improvement in soil physical, chemical and biological properties in organic matter application system [7,8,9].

33 Inappropriate fertilizer management and excessive use of N fertilizer causes N losses
34 through ammonia volatilization and leaching [10]. It is a major challenge in N management to
35 minimize the use of N fertilizer in crop production avoiding N deficiency in soil. Combined
36 application of fertilizer and manure is a recommended measure for reducing N loss and
37 increasing N use efficiency [11]. The combined application of organic and inorganic fertilizers
38 enhances N balance and minimizes losses by converting inorganic N to organic forms [12]
39 resulting sustainable productivity [13, 14]. Crop cultivation with integrated approach of
40 organic and inorganic fertilizers ensures plant nutrients in readily available form, good soil
41 health and sustainable yield [15].

42 The term soil health refers to the chemical, physical and biological properties that enable soil
43 to perform a wide range of function. The N fertilization from chemical and organic sources
44 influences N cycling, soil C storage and mineralization rates [16], and also effects temporal
45 and spatial distribution of inorganic soil N [17]. Soil biological properties respond quickly
46 under different soil managements than chemical or physical properties [18, 19]. Soil enzyme
47 activity is recognized as a good indicator of soil health as it reflects the effects of cultivation,
48 soil properties and pedological amendments [18, 19]. Soil microbes also play an important
49 role in nutrient cycling while the diversity of microbial population is largely affected by
50 fertilizer management practices. The abundance of microbial community is directly or
51 indirectly influenced by the fertilization practices [20].

52 Among the organic sources that are available to farmers in Bangladesh, cow dung and
53 poultry manure are good source of organic matter and can improve soil fertility providing
54 essential plant **nutrients elements** [21, 22]. The application of fertilizers and manures of
55 different properties differentially affects the physical, chemical and biological properties of
56 soil [23]. N mineralization in wet land rice soil is significantly influenced by the quality and
57 quantity of the organic matter applied [24]. However, many studies reported that the effects
58 of organic matter application can be observed after 3 to 5 years [25, 26]. Therefore, it is
59 important to study the long-term effects of different organic fertilizers in combination with
60 chemical N fertilizer on yield and soil properties. In Bangladesh several studies have been
61 conducted on integrated application of manure and fertilizers on rice yield which are mainly
62 short term study [27, 28] while there is lack of long term studies. Accordingly, this study was
63 carried out to determine the long term effect of combined application of manure and fertilizer
64 on soil N dynamics and biochemical properties from a five-year field experiment. Among the
65 biochemical properties we assessed free living nitrogen fixing bacteria population, urease
66 enzyme activity and soil organic carbon (**OC**) responses to applied treatment at different soil
67 depths.

69 2. MATERIALS AND METHODS

71 2.1 Site description

73 The experiment was conducted at experimental farm of Bangladesh Rice Research Institute
74 (23.59° N, 90.24° E, 8.4 m elevation) from 2010 to 2014. The climate of the experimental site
75 is subtropical in nature and experiences periodic south western monsoon with an average
76 annual rainfall of 2000 mm. The 80% of the rainfall occurs from mid-June to end of
77 September. The lowest mean temperature (15°C) prevails in January and highest (30°C) in
78 May. The soil of the experimental site belongs to the order Inceptisols in USDA soil
79 classification having soil texture of silty clay loam. The initial surface (0-20 cm) soil sample
80 had a bulk density of 1.40 g cm⁻³ and contained 12.21 g kg⁻¹ **organic carbon OC**, 1.1 g kg⁻¹
81 total nitrogen (**TN**), 19 mg kg⁻¹ available **phosphorus P**, 0.14 cmol_c **potassium K** in kg⁻¹ soil
82 and 28 mg kg⁻¹ available sulphur.

84 2.2 Experimental design and crop management

85

Formatted: Subscript

86 The experiment was consisted of four treatments arranged in a randomized complete block
 87 design with three replications. The treatments were: Soil Test Based dose of NPKS fertilizer
 88 (STB), Cow dung (CD) @ 3 t ha⁻¹ + NPKS as integrated plant nutrition system (IPNS)
 89 [CD+IPNS], Poultry manure (PM) @ 2 t ha⁻¹ + NPKS as IPNS [PM+IPNS] and N control. The
 90 STB dose of NPKS in Boro (dry season) was 138-10-80-5 kg ha⁻¹ and in T. Aman (wet
 91 season) it was 100-10-80-5 kg ha⁻¹. The quantity of N nutrient supplied from per ton cow
 92 dung was 5 kg N, 1.5 kg P and 5 kg K and per ton poultry manure supplied 19 kg N, 5 kg P
 93 and 7.5 kg K. The N control treatment received the STB dose of chemical fertilizer except N.
 94 The STB dose was calculated using the BARC fertilizer recommendation guide [29]. In
 95 CD+IPNS and PM+IPNS treatments fertilizer dose was calculated by subtracting the nutrient
 96 supplied with manures from the STB dose [29]. The treatment details of the experiment are
 97 presented in Table 1. Urea, tripple super phosphate, muriate of potash and gypsum were
 98 used as fertilizer source for N, P, K and S nutrients. Manures and all the fertilizers except
 99 urea were applied at the time of final land preparation and incorporated into the soil. Urea
 100 was applied in three equal splits. The first split was applied at the time of final land
 101 preparation and the remaining two splits were applied at maximum tillering and 5-10 days
 102 before panicle initiation stage of the crop. The unit plot size was 6 m x 7 m. Rice crops were
 103 grown following Boro-Fallow-T. Aman cropping pattern. In Boro season rice variety (BRRI
 104 dhan28) was transplanted in the first week of January and was harvested in May. In T.
 105 Aman season rice variety (BRRI dhan49) was transplanted in the first week of August and
 106 harvested in last week of November.

107 **Table 1: Nutrient added from different sources in each cropping season**

Treatment	Source	Nutrient added								Annual N rate
		Boro				T. Aman				
		N	P	K	S	N	P	K	S	
STB	Fertilizer	138	10	80	5	100	10	80	5	238
CD+IPNS	Fertilizer	123	5.5	65	5	85	5.5	65	5	238
	Cow dung	15	4.5	15	0	15	4.5	15	0	
	Total	138	10	80	5	100	10	80	5	
PM+IPNS	Fertilizer	100	0	65	0	62	0	65	5	238
	Poultry manure	38	10	15	0	38	11.2	15	0	
	Total	138	10	80	0	100	11.2	80	5	
N control	Fertilizer	0	10	80	5	0	10	80	5	0

109 2.3 Plant and soil sample collection

110 Plant samples were collected at maturity stage of the crop growth for calculating grain yield
 111 and nutrient analysis. Grain yield was calculated from a harvesting area of 5 m² and
 112 adjusted to 14% moisture content. Total nitrogen content of plant tissue was estimated by
 113 Micro Kjeldhal method [30] and the crop N uptake was calculated from dry biomass (grain +
 114 straw) weight and N concentrations [28].

115 For this study soil samples were collected from four depths: 0 to 5, 5 to 10, 10 to 15 and 15
 116 to 20 cm from each plot after the harvest of T. Aman crop in 2014. The soil samples were
 117 air-dried, grinded and passed through 2-mm sieve and stored in polythene bags at room
 118 temperature for total N, incubated ammonium and organic carbon (OC) determination. For
 119 determining NH₄⁺-N, NO₃⁻-N, free living N fixing bacteria population and urease enzyme
 120 activity fresh soils were collected and refrigerated at 4° C.

121 2.4 Net N balance

122 After ten cropping seasons the N balance was estimated as follows:
 123
 124
 125
 126

127 Net N balance = (Total N removal from ten cropping seasons + change in soil total N after 10
128 cropping seasons) – (N addition through manures and fertilizer)
129 In calculating net N balance, N inputs from other sources like rainfall, irrigation, crop
130 residues, biological N fixation etc. and N outputs from various N loss mechanisms were not
131 considered in this study.
132

133 **2.5 Soil ammonium N, nitrate N, mineralized N, total N and organic carbon** 134 **determination**

135
136 Soil NH_4^+ -N and NO_3^- -N was determined from moist soil extracting with 2 M KCl [31] followed
137 by distillation with MgO and Devarda alloy [32]. The mineralized N in soils received different
138 fertilizers treatments was determined using a soil incubation study described by Sahrawat,
139 1983 [33]. Total N was determined by Kjeldhal digestion, distillation and titration [34]. Total
140 organic carbon (TOC) was determined following modified Walkley-Black method [35]. All the
141 measurements were expressed on dry (105°C) soil basis.
142

143 **2.6 Free living nitrogen fixing bacteria (NFB) population and urease enzyme** 144 **activity**

145
146 The free living NFB population was counted as colony forming unit culturing on N free media
147 [36] using serial dilution method [37].
148 Urease enzyme activity was determined by measuring the amount of ammonium released
149 from the hydrolysis of urea and expressed as mg NH_4^+ released g^{-1} soil h^{-1} described by
150 Tabatabai and Bremner, 1979 [38].
151

152 **2.7 Statistical analysis**

153
154 Data related to rice yield, soil ammonium N, nitrate N, total N, mineralizable N, organic
155 carbon, free living nitrogen fixing bacteria population and urease enzyme activity in
156 different treatments and soil depths were analyzed using ANOVA at 5% level of significance.
157 Mean comparison was done using Least **significance Significance test Test (LSDLST)**.
158 Results are presented as mean with standard error of three replicates. Statistical analysis
159 was performed with STAR (Statistical Tool for Agricultural research) version 2.0.1.
160


161 **3. RESULTS AND DISCUSSION**

162 **3.1 Grain yield**

163
164 The different fertilizer management practices showed significant effects on rice grain yield
165 and the yield scenario of five years rice cropping with different fertilization is shown in Figure
166 1. During the five years cropping, the annual rice grain yield in PM+IPNS was at par with the STB
167 treatment. The annual grain yield in CD+IPNS was significantly lower compared to STB
168 treatment in the first two cropping years. However, from the third year it resulted in similar
169 annual rice grain yield to STB. The omission of N fertilizer drastically reduced rice grain yield
170 compared to N fertilized treatments from the third year (Figure 1). Our findings indicate that
171 PM performed better than CD in obtaining the rice grain yield at par with STB treatment. The
172 variation in crops yield with the types of organic matter used and their combination with
173 inorganic fertilizers has been confirmed by previous studies [39, 40, 41, 42]. The better
174 performance of PM on rice grain yield over CD might be due to higher nutrient content and
175 low to optimum C:N ratio in PM than CD. Therefore, nutrient release from PM is faster
176 resulting higher nutrient uptake and thereby higher yield. On the other hand manure like CD
177 having high C:N ratio initially favors nutrient immobilization which eventually resulted in lower
178 grain yield. These findings were supported by Rahman et al., 2016 [43]. Moreover, it is well
179

Comment [H2]: ?????

180 documented that integrated application of manure and fertilizers in balanced doses increase
 181 crop yield by improving soil physical, chemical and biological properties [44, 45].
 182



183 **Figure 1: Fertilization effects on rice grain yield in Boro and T. Aman seasons during**
 184 **five years of cropping.**

185 *Error bars represent LSD value*

188 3.2 Net nitrogen balance

189 The **net N nitrogen** balance in the 0-20 cm soil depth after five years of rice-rice cropping
 190 system as affected by different fertilizer management practices was shown in Table 2. All the
 191 fertilized treatments received an equal N input of 1380 kg ha⁻¹ from fertilizer and manures.
 192 However, the highest (991 kg ha⁻¹) N removed by the rice crop was from STB treated soil
 193 resulting in a decrease (406 kg ha⁻¹) of soil N with a net N balance of -795 kg ha⁻¹. Among
 194 the manure treated soils, PM+IPNS showed an increase (910 kg ha⁻¹) in soil N with a net
 195 positive N balance of 435 kg ha⁻¹ while CD+IPNS resulted in a negative net N balance of 148
 196 kg ha⁻¹. Our study suggests that combined application of manure and fertilizer for long term
 197 can build up soil N while sole application of inorganic fertilizer or no nitrogen fertilizer
 198 resulted in a higher negative N balance. Kumar and Mukhopadhyay, 2017 [46]; Tadesse et
 199 al., 2013 [47] reported a higher positive N balance with combined application of manure and
 200 fertilizer in rice based cropping system. Tiwari et al., 2010 [48] reported a negative **nitrogen**
 201 **N** balance with inorganic fertilizers only and positive with manures. The positive balance in
 202 PM treated soil might be attributed due to the variation in N addition from different sources,
 203 other than fertilizer and manure and variation in crop N removal. Because N inputs were
 204 equal in all fertilized treatments and N removal (except N losses) were considered, the
 205 positive N balance in the PM treated soil might be attributed to higher biological N fixation by
 206 free living N fixing bacteria which population was recorded higher in PM amended soil
 207 (Figure 4.a). Ladha et al., 2000 [49] also reported similar findings. The higher negative N
 208 balance in STB treatment where only chemical fertilizer was applied may be due to higher N
 209 loss through ammonia volatilization which was reported by several studies [50, 51, 52].
 210

211
 212
 213
 214

215
216

Table 2: Fertilization effects on net N balance after five years of rice cropping

Treatment	N Added		Initial soil N	N removed	Soil N after 10 th crop	Net N balance
	Fertilizer	Manure				
	(kg ha ⁻¹)					
STB	1380	0	3080	991.085	2674	-794.915
STB+CD	1230	150	3080	902.537	3409	-148.463
STB+PM	1000	380	3080	905.444	3990	435.444
N control	0	0	3080	456.3195	2156	-467.681

Formatted Table

217
218
219

3.3 Soil ammonium N

After ten cropping season soil NH₄⁺-N was significantly varied with fertilization practices and soil depths. The variation in NH₄⁺-N concentration due to different fertilizer management practices was much higher at the top 0-5 cm soil depth where significantly highest NH₄⁺-N was found with PM+IPNS treatment (25.49 mg kg⁻¹) followed by CD+IPNS treatment (19.83 mg kg⁻¹). At 5-10 cm soil depth NH₄⁺-N concentration was statistically similar with PM+IPNS (15.90 mg kg⁻¹) and STB (15.01 mg kg⁻¹) fertilizer management practices. In 10-15 and 15-20 cm soil depth the NH₄⁺-N concentration was identical between PM+IPNS and CD+IPNS treatments. In STB, CD+IPNS and PM+IPNS treatments NH₄⁺-N concentration was higher at top soil layer and decreased with the increase of soil depths. However, in N control treatment the NH₄⁺-N concentration was lower at 0-5 soil depth then it increased at 5-10 cm soil depth and then it gradually decreased up to 20 cm soil depth (Figure 2.a).

NH₄⁺-N is the most reduced and preferred forms of N in flooded soil [53, 54]. NH₄⁺-N builds up in wetland soils particularly in soils high in organic matter or when easily decomposable organic matter is added in high amounts [53; 55, 56]. This might be the reason of higher NH₄⁺-N in manure treated soil especially in PM treated soils which is easily decomposable than CD. The lower concentration of NH₄⁺-N in deeper soil layer might be attributed to its adsorbance on soil colloids which leads to lower percolation loss [57]. Lee and Choi, 2017 [5]; Mi et al., 2019 [58] also reported decreased NH₄⁺-N with the increase of soil depth. As the mineralization of N from chemical N fertilizer is high the NH₄⁺ released from STB treatment is subjected to instant crop uptake [5] and volatilization loss which resulted in lower concentration of NH₄⁺ in the post harvested soil [52].


241
242
243

3.4 Soil nitrate N

Soil NO₃⁻-N was significantly affected by fertilizer management practices along the soil depths. At all the four soil depths the NO₃⁻-N concentration was significantly higher with PM+IPNS treatment and lowest with N control treatment. In CD+IPNS, PM+IPNS and N control treatments the NO₃⁻-N concentration was significantly higher at 0-5 cm soil depth and then it sharply decreased and remained static at 5-20 cm soil depths. However, in STB fertilizer treatment the NO₃⁻-N concentration was significantly higher at 5-10 cm soil depth and was identical at 0-5, 10-15 and 15-20 cm soil depths (Figure 2.b).

In this study the higher NO₃⁻ concentration in the PM+IPNS treated soil might be due to higher NH₄⁺ concentration in this soil as the NH₄⁺ adsorbed onto clay particles is gradually converted to NO₃⁻ via nitrification [5]. In an incubation study Murugan and Swarnam [59] found higher NO₃⁻ in manure treated soils compared to inorganic N treated soils. The lower NO₃⁻-N concentration in the deeper soil may be due to less aerobic conditions in flooded soil [60]. Mi et al., 2019 [58] also reported lower NO₃-N with increasing soil depth.

257



Comment [H3]: The legend is in different directions relative to the other graphics.

258

259 **Figure 2: Fertilization effects on soil ammonium N and nitrate N after five years of rice**
 260 **cropping.**

261 *Error bars represent standard error (n=3) of the means of ammonium N and nitrate N.*

262

263 3.5 Total N

264

265 The soil total N content was significantly affected with the different fertilization practices
 266 within a soil profile of 20 cm. At 0-5 cm soil depth the total N content was significantly higher
 267 with PM+IPNS followed by CD+IPNS while at 5-10 cm soil depth the trend was reverse. At
 268 10-15 cm soil depth the effect of CD+IPNS and PM+IPNS treatments on soil total N was
 269 statistically similar. Significantly lowest soil total N was observed in N control treatment at all
 270 four soil depths. Regardless of the fertilizer treatments an overall decreasing trend of soil
 271 total N was observed with the increase of soil depths (Figure 3.a).

272 Previous studies revealed that combined application of different organic matter and chemical
 273 fertilizer for long term significantly increased soil total N compared to only chemical fertilizer
 274 [61, 62, 63]. Regardless of the fertilizer treatment total N decreased with the increase in soil
 275 depth which coincides with the results of Lee and Choi, 2017 [5].

276

277 3.6 Mineralizable N

278

279 Mineralizable N was determined from the anaerobic incubation of dried soils after harvesting
 280 of 10th rice crop. Different fertilization practices showed significant effect on soil
 281 mineralizable N at different soil depths after 14 days of incubation. PM+IPNS treated soils
 282 resulted significantly highest mineralizable N at all four soil depths (0-5, 5-10, 0-15 and 15-
 283 20 cm) compared to CD+IPNS and STB treated soils. In the PM+IPNS treated soil
 284 mineralizable N ranged from 384.50-220.00 mg kg⁻¹. The N controlled soils resulted in
 285 lowest mineralizable N at all soil depths. Irrespective of fertilizer treatments, mineralizable N
 286 was highest at upper 0-5 cm soil layer and was lowest at 15-20 cm soil depth although in N

Comment [H4]: Why?

Comment [H5]: But, why?

287 control treatment after 0-5 cm the variation in mineralizable N was statistically similar (Figure
288 3.b).

289 The higher soil residual mineral N levels found in our study with the PM+IPNS treatment
290 might be due to slow release of the N during the crop growth period. Mineralizable N is
291 affected by the quantity and quality of organic matter applied in the soil [64] which might be
292 cause of variation in mineralizable N in PM and CD treated soils. Sahrawat, 1983 [33]
293 reported that mineralizable N in anaerobic incubation was significantly and positively
294 correlated with soil OC and total N. This might be the reason of higher mineralizable N in
295 manure treated soils in our study which contained higher OC and total N compared to
296 chemical fertilizer treated soils. Pal et al., 2015 [65] found highest mineralizable N with the
297 highest rate of chemical fertilizer combined with farm yard manure while Myint et al., 2010
298 [66] reported highest mineralized N with sole application of mineral N fertilizer than sole
299 application of manure.

300


301 3.7 Total organic carbon

302

303 Soil total organic carbon (TOC) significantly varied with fertilizer management practices
304 and soil depth. Application of PM @ 2 t ha⁻¹ for five consecutive cropping seasons resulted
305 in significantly highest accumulation of TOC at all the four studied soil depths followed by
306 application of CD @ 3 t ha⁻¹. In PM+IPNS treated soil TOC varied from 16.87-8.45 g kg⁻¹.
307 The lowest TOC was found in N controlled soil. Regardless of the fertilizer treatments, soil
308 TOC decreased with the increase of soil depth (Figure 3.c).

309 Previous studies showed that combined application of manure and fertilizer increased soil
310 OC [67, 68, 61]. Similar result was also found in our study where higher soil OC content was
311 found with PM and CD amended soil. However, the accumulation of OC was higher in PM
312 than CD. Rahman et al. 2016 [43] also reported increased OC content in PM treated soil
313 over CD after 2.5 years of rice cultivation. The increased soil OC content in the top soil can
314 be explained by the higher accumulation of manure and crop residues in the top arable
315 layer soil.

Comment [H6]: In the real, lower CN relation.



Comment [H7]: Idem.

316

317 Figure 3: Fertilization effects on soil total N, mineralizable N and organic carbon after
318 five years of rice cropping.

319 Error bars represent standard error (n=3) of the means of total N, mineralizable N and
320 organic carbon.

321

322

323

324

3.8 Free living nitrogen fixing bacteria population

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

3.9 Urease enzyme activity

344

345

346

347

348

349

350

351

352

353

354


355

356

357

358

Urease activity is a good indicator of changes in soil quality for soil management [72]. In this study soil urease activity was significantly affected by fertilizer management practices and soil depths. Regardless of the soil depth urease activity was significantly higher in PM+IPNS followed by CD+IPNS treated soils compared to STB fertilized soil. In PM+IPNS treated soil the urease activity ranged between 51.12-23.36 $\mu\text{g NH}_4^+ \text{g}^{-1}\text{h}^{-1}$ and in CD+IPNS treated soil it was 42.54-18.43 $\mu\text{g NH}_4^+ \text{g}^{-1}\text{h}^{-1}$. The lowest urease activity was found in N control treated soil. Urease activity was higher at upper soil layer and decreased with the increase of soil depth in all applied treatments (Figure 4.b). The higher urease activity in the organic matter amended soil and upper soil layer is due to high OC and microbial activity which was reported by several studies [73, 74, 75]. The OC provides energy for microbial activity resulting in higher enzyme activity. The constituents of the organic matter also influence the soil urease activity and it is positively correlated with OC and total N [76]. Application of PM performed better than CD which might be due to high N content in PM.



Comment [H8]: Idem.
What is NFB (log 10)?

359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390

Figure 4: Fertilization effects on free living N fixing bacteria (NFB) and urease enzyme activity at different soil depths after five years of rice cropping.

Error bars represent standard error (n=3) of the means of free living N fixing bacteria and urease enzyme activity.

4. CONCLUSION

From this study it can be concluded that long term (5 years) application of organic and inorganic fertilizers following IPNS approach in the rice-fallow-rice cropping system has profound influence on rice yields, N forms and biochemical properties. Application of PM @ 2 t ha⁻¹ in each cropping season resulted in annual grain yield at per the STB treatment during the five years of rice cropping. After five years of rice cropping PM+IPNS resulted in a positive soil N balance while it was negative in CD+IPNS and STB treatments. The consecutive use of PM+IPNS also showed significant effects on N forms (ammonium N, nitrate N, mineralizable N and total N), organic carbon and biochemical properties (free living N fixing bacteria population and urease enzyme activity). Considering the soil health, our study suggests that PM+IPNS could be an effective fertilization practice for sustainable rice production in long run reducing the use of chemical fertilizer.

COMPETING INTEREST

The authors have declared that no competing interests exist.

REFERENCES

- Nasim M, Shahidullah SM, Saha A, Mutaleb MA, Aditya TL, Ali MA, Kabir MS. Distribution of Crops and Cropping Patterns in Bangladesh. Bangladesh Rice Journal. 2017;21(2):1-55.

- 391 2. Zhu ZL, Chen DL. Nitrogen fertilizer use in China—Contributions to food production,
392 impacts on the environment and best management strategies. *Nutrient Cycling in*
393 *Agroecosystems*. 2002;63(2-3):117-27.
- 394 3. Celik I, Gunal H, Budak M, Akpınar C. Effects of long-term organic and mineral fertilizers
395 on bulk density and penetration resistance in semi-arid Mediterranean soil conditions.
396 *Geoderma*. 2010;160(2):236-43. doi:10.1016/j.geoderma.2010.09.028
- 397 4. Liu E, Yan C, Mei X, He W, Bing SH, Ding L, Liu Q, Liu S, Fan T. Long-term effect of
398 chemical fertilizer, straw, and manure on soil chemical and biological properties in
399 northwest China. *Geoderma*. 2010;158(3-4):173-180.
400 doi:10.1016/j.geoderma.2010.04.029
- 401 5. Lee J, Choi HL. The dynamics of nitrogen derived from a chemical nitrogen fertilizer with
402 treated swine slurry in paddy soil-plant systems. *PloS one*. 2017;12(3):e0174747.
403 doi:10.1371/journal.pone.0174747
- 404 6. Efthimiadou A, Bilalis D, Karkanis A, Froud-Williams B. Combined organic/inorganic
405 fertilization enhance soil quality and increased yield, photosynthesis and sustainability of
406 sweet maize crop. *Australian Journal of Crop Science*. 2010;4(9):722.
- 407 7. Chang EH, Chung RS, Wang FN. Effect of different types of organic fertilizers on the
408 chemical properties and enzymatic activities of an Oxisol under intensive cultivation of
409 vegetables for 4 years. *Soil science and plant nutrition*. 2008;54(4):587-99.
410 doi:10.1111/j.17470765.2008.00264.x
- 411 8. Surekha K, Latha PC, Rao KV, Kumar RM. Grain yield, yield components, soil fertility,
412 and biological activity under organic and conventional rice production systems.
413 *Communications in soil science and plant analysis*. 2010;41(19):2279-92.
414 doi:10.1080/00103624.2010.507834
- 415 9. Subehia SK, Sepelhy S, Rana SS, Negi SC, Sharma SK. Long-term effect of organic
416 and inorganic fertilizers on rice (*Oryza sativa* L.)–wheat (*Triticum aestivum* L.) yield, and
417 chemical properties of an acidic soil in the western Himalayas. *Experimental agriculture*.
418 2013;49(3):382-94. doi:10.1017/S0014479713000173
- 419 10. Moe K, Mg KW, Win KK, Yamakawa T. Effect of combined application of inorganic
420 fertilizer and organic manures on nitrogen use and recovery efficiencies of hybrid rice
421 (Palethwe-1). *American Journal of Plant Science*. 2017;8:1043-1064.
422 doi:10.4236/ajps.2017.85069
- 423 11. Wen Z, Shen J, Blackwell M, Li H, Zhao B, Yuan H. Combined applications of nitrogen
424 and phosphorus fertilizers with manure increase maize yield and nutrient uptake via
425 stimulating root growth in a long-term experiment. *Pedosphere*. 2016;26(1):62-73.
426 doi:10.1016/S1002-0160(15)60023-6
- 427 12. Kramer AW, Doane TA, Horwath WR, Van Kessel C. Combining fertilizer and organic
428 inputs to synchronize N supply in alternative cropping systems in California. *Agriculture,*
429 *Ecosystems & Environment*. 2002;91(1-3):233-43. doi: 10.1016/S0167-8809(01)00226-2
- 430 13. Satyanarayana V, Vara Prasad PV, Murthy VR, Boote KJ. Influence of integrated use of
431 farmyard manure and inorganic fertilizers on yield and yield components of irrigated
432 lowland rice. *Journal of plant nutrition*. 2002;25(10):2081-90. doi:10.1081/PLN-
433 120014062
- 434 14. Jobe B. Integrated nutrient management for increased rice production in the inland
435 valleys of the Gambia. In: Sanyang S, Ajayi A, Sy AA, editors. *Proceedings of the*
436 *Second Biennial Regional Rice Research Review*; 2003.
- 437 15. Lakshmi CS, Rao PC, Sreelatha T, Madahvi M, Padmaja G, Rao PV, Sireesha A.
438 Nitrogen use efficiency and production efficiency of rice under rice-pulse cropping
439 system with integrated nutrient management. *Journal of Rice Research*. 2012;5(1):2.
- 440 16. Sardans J, Peñuelas J, Estiarte M. Changes in soil enzymes related to C and N cycle
441 and in soil C and N content under prolonged warming and drought in a Mediterranean
442 shrubland. *Applied Soil Ecology*. 2008;39(2):223-35. doi: 10.1016/j.apsoil.2007.12.011

- 443 17. Shi Z, Li D, Jing Q, Cai J, Jiang D, Cao W, Dai T. Effects of nitrogen applications on soil
444 nitrogen balance and nitrogen utilization of winter wheat in a rice-wheat rotation. *Field*
445 *Crops Research*. 2012;127:241-7. doi: 10.1016/j.fcr.2011.11.025
- 446 18. Nannipieri P, Kandeler E, Ruggiero P. Enzyme activities and microbiological and
447 biochemical processes in soil. In: Burns RG, Dick RP, editors. *Enzymes in the*
448 *Environment*. New York: Marcel Dekker, 2002.
- 449 19. Paz-Ferreiro J, Gascó G, Gutiérrez B, Méndez A. Soil biochemical activities and the
450 geometric mean of enzyme activities after application of sewage sludge and sewage
451 sludge biochar to soil. *Biology and fertility of soils*. 2012;48(5):511-7.
452 doi: 10.1007/s00374-011-0644-3
- 453 20. Gu Y, Wang Y, Lu SE, Xiang Q, Yu X, Zhao K, Zou L, Chen Q, Tu S, Zhang X. Long-
454 term fertilization structures bacterial and archaeal communities along soil depth gradient
455 in a paddy soil. *Frontiers in microbiology*. 2017;8:1516. doi:10.3389/fmicb.2017.01516
- 456 21. Khanam M, Rahman MM, Islam MR, Islam MR. Effect of manures and fertilizers on the
457 growth and yield of BRRI Dhan30. *Pakistan Journal of Biological Sciences*. 2001; 4:
458 172-174. doi: 10.3923/pjbs.2001.172.174
- 459 22. Mitchell CC, Tu S. Nutrient accumulation and movement from poultry litter. *Soil Science*
460 *Society of America Journal*. 2006; 70(6): 2146-2153. doi:10.2136/sssaj2004.0234
- 461 23. Hossain MA, Shamsuddoha AT, Paul AK, Bhuiyan MS, Zobaer AS. Efficacy of different
462 organic manures and inorganic fertilizer on the yield and yield attributes of boro rice. *The*
463 *Agriculturists*. 2011;9(1&2):117-25.
- 464 24. Sahrawat KL. Organic matter and mineralizable nitrogen relationships in wetland rice
465 soils. *Communications in Soil Science and Plant Analysis*. 2006;37(05-06):787-96.
466 doi:10.1080/00103620600564034
- 467 25. Ohyama N, Katano M, Hasegawa T. Effects of long term application of organic materials
468 to the paddy field originated from Aso volcanic ash [Japan] on the soil fertility and rice
469 [*Oryza sativa*] growth, 1: Effects on the rice growth and nutrient uptake for the initial
470 three years. *Proceedings of Faculty of Agriculture-Kyushu Tokai University (Japan)*.
471 1998.
- 472 26. Uenosono S, Nagatomo M. Effect of application of only manure on yield and quality of
473 paddy rice. *Kyushu Agricultural Research (Japan)*. 1998.
- 474 27. Ali ME, Islam MR, Jahiruddin M. Effect of integrated use of organic manures with
475 chemical fertilizers in the rice-rice cropping system and its impact on soil health.
476 *Bangladesh Journal of Agricultural Research*. 2009;34(1):81-90.
- 477 28. Sarkar MU, Rahman MM, Rahman GK, Naher UA, Ahmed MN. Soil test based inorganic
478 fertilizer and integrated plant nutrition system for rice (*Oryza sativa* L.) cultivation in
479 inceptisols of Bangladesh. *The Agriculturists*. 2016;14(1):33-42. doi:
480 10.3329/agric.v14i1.29098
- 481 29. BARC (Bangladesh Agricultural Research Council). *Fertilizer Recommendation Guide*
482 2012. Dhaka, Bangladesh: BARC; 2012.
- 483 30. Yoshida S, Forno DA, Cock JH. *Laboratory manual for physiological studies of*
484 *rice*. Philippines: Los Banos IRRI; 1971.
- 485 31. Maynard DG, Kalra YP. Nitrate and exchangeable ammonium nitrogen. In: Carter MR,
486 editor. *Soil sampling and methods of analysis*. Boca Raton: Lewis Publishers; 1993.
- 487 32. Bremner JM, Keeney DR. Steam distillation methods for determination of ammonium,
488 nitrate and nitrite. *Analytica Chimica Acta*. 1965; 32: 485-495.
- 489 33. Sahrawat KL. Mineralization of soil organic nitrogen under waterlogged conditions in
490 relation to other properties of tropical rice soils. *Soil Research*. 1983; 21:133-138.
- 491 34. Bremner JM. Determination of nitrogen in soil by the Kjeldahl method. *The Journal of*
492 *Agricultural Science*. 1966; 55:11-33.
- 493 35. Page AL, Miller RH, Keeney DR. *Methods of soil analysis- Part 2*. Madison: American
494 *Society of Agronomy*; 1982.

- 495 36. Dobereiner J, Day JM. Associative symbioses in tropical grasses: characterization of
496 microorganisms and dinitrogen-fixing sites. In: Newton WE, Nyman CJ, editors.
497 Proceedings of the 1st international symposium on nitrogen fixation. Pullman:
498 Washington State University Press; 1976
- 499 37. Dhingra OD, Sinclair JB. Basic plant pathology methods. London: CEC press; 2000.
- 500 38. Tabatabai MA, Bermner JM. Use of P-nitrophenyl phosphate for assay of soil
501 phosphatase activity. *Soil Biology and Biochemistry*. 1969; 1:301-307.
- 502 39. Gupta V, Sharma RS, Vishwakarma SK. Long-term effect of integrated nutrient
503 management on yield sustainability and soil fertility of rice (*Oryza sativa*)–wheat
504 (*Triticum aestivum*) cropping system. *Indian Journal of Agronomy*. 2006;51(3):160-4.
- 505 40. Saha PK, Ishaque M, Saleque MA, Miah MA, Panaullah GM, Bhuiyan NI. Long-Term
506 Integrated Nutrient Management for Rice-Based Cropping Pattern: Effect on Growth,
507 Yield, Nutrient Uptake, Nutrient Balance Sheet, and Soil Fertility. *Communications in
508 Soil Science and Plant Analysis*. 2007;38(5-6):579-610. doi:
509 10.1080/00103620701215718
- 510 41. Baishya LK, Rathore, SS, Singh D, Sarkar D, Deka BC. Effect of integrated nutrient
511 management on rice productivity, profitability and soil fertility. *Annals of Plant and Soil
512 Research*. 2015;17, 86-90.
- 513 42. Gawde N. Long term effect of integrated nutrient management on soil nutrient status
514 under rice-wheat cropping system in Inceptisols. *International Journal of Chemical
515 Studies*. 2017; 5:1050-1057.
- 516 43. Rahman F, Rahman MM, Rahman GM, Saleque MA, Hossain AS, Miah MG. Effect of
517 organic and inorganic fertilizers and rice straw on carbon sequestration and soil fertility
518 under a rice–rice cropping pattern. *Carbon Management*. 2016;7(1-2):41-53. doi:
519 10.1080/17583004.2016.1166425
- 520 44. Rahman MM. Potential supplies and use efficiencies of nutrients from different organic
521 wastes under tomato cultivation. *Annals Bangladesh Agriculture*. 2012;16: 25-39.
- 522 45. Rahman MM. Nutrient-use and carbon-sequestration efficiencies in soils from different
523 organic wastes in rice and tomato cultivation. *Communications in Soil Science and Plant
524 Analysis*. 2013;44(9):1457-1471. doi: 10.1080/00103624.2012.760575
- 525 46. Kumar B, Mukhopadhyay SK. Effect of integrated nutrient management on system
526 productivity, nutrient uptake, nitrogen balance, soil structural properties and nitrogen use
527 efficiency under wheat-rice cropping system. *Journal of Pharmacognosy and
528 Phytochemistry*. 2017;SP1:1030-1033.
- 529 47. Tadesse T, Dechassa N, Bayu W, Gebeyehu S. Effects of farmyard manure and
530 inorganic fertilizer application on soil physico-chemical properties and nutrient balance in
531 rain-fed lowland rice ecosystem. *American Journal of Plant Sciences*. 2013;4(02):309.
532 doi: 10.4236/ajps.2013.42041
- 533 48. Tiwari KR, Sitaula BK, Bajracharya RM, Børresen T. Effects of soil and crop
534 management practices on yields, income and nutrients losses from upland farming
535 systems in the Middle Mountains region of Nepal. *Nutrient Cycling in Agroecosystems*.
536 2010;86(2):241-53. doi:10.1007/s10705-009-9289-0.
- 537 49. Ladha JK, Dawe D, Ventura TS, Singh U, Ventura W, Watanabe I. Long-term effects of
538 urea and green manure on rice yields and nitrogen balance. *Soil Science Society of
539 America Journal*. 2000; 64: 1993–2001. doi:10.2136/sssaj2000.6461993x
- 540 50. Huang S, Lv W, Bloszies S, Shi Q, Pan X, Zeng Y. Effects of fertilizer management
541 practices on yield-scaled ammonia emissions from croplands in China: a meta-analysis.
542 *Field Crops Research*. 2016;192:118–125. doi: 10.1016/j.fcr.2016.04.023
- 543 51. Ke J, Xing X, Li G, Ding Y, Dou F, Wang S, Liu Z, Tang S, Ding C, Chen L. Effects of
544 different controlled-release nitrogen fertilisers on ammonia volatilisation, nitrogen use
545 efficiency and yield of blanket-seedling machine-transplanted rice. *Field Crops
546 Research*. 2017;205,147–156. doi: 10.1016/j.fcr.2016.12.027

- 547 52. Liu T, Huang J, Chai K, Cao C, Li C. Effects of N fertilizer sources and tillage practices
548 on NH₃ volatilization, grain yield, and N use efficiency of rice fields in Central
549 China. *Frontiers in Plant Science* . 2018;9:385. doi: 10.3389/fpls.2018.00385
550 53. Ponnampertuma FN. Effects of flooding on soils. In: Kozłowski T, edotor. *Flooding and*
551 *plant growth*. New York: Academic Press; 1984.
552 54. George T, Ladha JK, Buresh RJ, GarrityDP. Managing native and legume fixed nitrogen
553 in lowland rice-based cropping systems. *Plant and Soil*. 1992;141: 69–91.
554 55. Narteh LT, Sahrawat KL. Influence of flooding on electrochemical and chemical
555 properties of West African soils. *Geoderma*. 1999;87(3-4): 179–207.
556 56. Narteh LT, Sahrawat KL (2000) Ammonium in solution of flooded West African soils.
557 *Geoderma*. 2000;95(3-4): 205–214.
558 57. Cassman KG, Peng S, Olk DC, Ladha JK, Reichardt W, Dobermann A, Singh U (1998)
559 Cassman KG, Peng S, Olk DC, Ladha JK, Reichardt W, Dobermann A, Singh U.
560 Opportunities for increased nitrogen-use efficiency from improved resource
561 management in irrigated rice systems. *Field crops research*. 1998;56(1-2):7-39. doi:
562 10.1016/S0378-4290(97)00140-8
563 58. Mi W, Gao Q, Xia S, Zhao H, Wu L, Mao W, Hu Z, Liu Y. Medium-term effects of
564 different types of N fertilizer on yield, apparent N recovery, and soil chemical properties
565 of a double rice cropping system. *Field Crops Research*. 2019;234: 87-94.
566 doi.org/10.1016/j.fcr.2019.02.012
567 59. Murugan AV, Swarnam TP. Nitrogen release pattern from organic manures applied to an
568 acid soil. *Journal of Agricultural Science*. 2013;5(6):174..doi:10.5539/jas.v5n6p174
569 60. Roldán A, Salinas-García JR, Alguacil MM, Díaz E and Caravaca F. Soil enzyme
570 activities suggest advantages of conservation tillage practices in sorghum cultivation
571 under subtropical conditions. *Geoderma*. 2005;129(3-4):178–185. doi:
572 10.1016/j.geoderma.2004.12.042
573 61. Li W, Chen H, Cao C, Zhao Z, Qiao Y, Du S. Effects of long-term fertilization on organic
574 carbon and nitrogen dynamics in a Vertisol in Eastern China. *Open Journal of Soil*
575 *Science*. 2018; 8:99-117. doi: 10.4236/ojss.2018.83008
576 62. Chang EH, Wang CH, Chen CL, Chung RS (2014) Effects of long-term treatments of
577 different organic fertilizers complemented with chemical N fertilizer on the chemical and
578 biological properties of soils. *Soil Science and Plant Nutrition*. 2014;60(4):499-511. doi:
579 10.1080/00380768.2014.917333
580 63. Gao X, Han X, Zhan X, Sun Z, Jiang L, Chen H. Effect of long-term fertilization on total
581 nitrogen storage in a brown soil. *Plant Nutrition and Fertilizer Science*. 2009;15(3):567-
582 72.
583 64. Sahrawat KL. Nitrogen mineralization in lowland rice soils: the role of organic matter
584 quantity and quality. *Archives of Agronomy and Soil Science*. 2010;56(3):337-53.
585 65. Pal B, Pati S, Badole S, MalothuV, Patra PK. Effect of integrated nutrient management
586 on nitrogen dynamics in soil of rice-potato based cropping sequence. *Journal of Applied*
587 *and Natural Science*. 2015;7(2): 652-655. doi: 10.31018/jans.v7i2.661
588 66. Myint AK, Yamakawa T, Kajihara Y, Zenmyo T. Application of different organic and
589 mineral fertilizers on the growth, yield and nutrient accumulation of rice in a Japanese
590 ordinary paddy field. *Science World Journal*. 2010; 5(2): 47-54.
591 doi:10.4314/swj.v5i2.61513
592 67. Stark C, Condron LM, Stewart A, Di HJ, O'Callaghan M. Influence of organic and mineral
593 amendments on microbial soil properties and processes. *Applied Soil Ecology*.
594 2007;35(1):79-93. doi: 10.1016/j.apsoil.2006.05.001
595 68. Kuntal MH, Anand S, Mishra B, Manna MC, WanjariRH, Mandal KG, Misra AK. Impact of
596 long-term application of fertilizer, manure and lime under intensive cropping on physical
597 properties and organic carbon content of an Alfisol. *Geoderma*. 2008;148:173–179. doi:
598 10.1016/j.geoderma.2008.09.015

- 599 69. Mujiyati M, Supriyadi S. Effect of manure and NPK to increase soil bacterial population
600 of *Azotobacter* and *Azospirillum* in chili (*Capsicum annuum*) cultivation. *BioScience*.
601 2009;1(2):59-64. doi: 10.13057/nusbiosci/n010202
- 602 70. Shrestha RK, Ladha JK (1996) Genotypic variation in promotion of rice dinitrogen
603 fixation as determined by nitrogen-15 dilution. *Soil Science Society of America Journal*.
604 1996;60(6):1815-1821. doi: 10.2136/sssaj1996.03615995006000060029x
- 605 71. Ayuni N, Radziah O, Naher UAA, Panhwar QA, Halimi MS. Effect of nitrogen on
606 nitrogenase activity of diazotrophs and total bacterial population in rice soil. *Journal of*
607 *Animal and Plant Sciences*. 2015; 25(5):1358-1364.
- 608 72. Diaz-Marcote I, Polo A (1995) MSW compost for the restoration of degraded soil. In
609 *World Congress on Waste Management. Proc. of the 25th Anniversary of the*
610 *International Solid Waste Association. 15th-20th October, 1995 ISWA, Vienna (CD*
611 *Rom)*.
- 612 73. Chakrabarti K, Sarkar B, Chakraborty A, Banik P, Bagchi DK. Organic recycling for soil
613 quality conservation in a sub-tropical plateau region. *Journal of Agronomy and crop*
614 *Science*. 2000 Apr;184(2):137-42. doi: 10.1046/j.1439-037x.2000.00352.x
- 615 74. Chang EH, Chung RS, Tsai YH. Effect of different application rates of organic fertilizer
616 on soil enzyme activity and microbial population. *Soil Science and Plant Nutrition*. 2007
617 Apr 1;53(2):132-40. doi: 10.1111/j.1747-0765.2007.00122.x
- 618 75. Goutami N, Rani PP, Pathy RL, Babu PR. Soil properties and biological activity as
619 influenced by nutrient management in rice-fallow sorghum. *International Journal of*
620 *Agricultural Research, Innovation and Technology*. 2015; 5:10-14. doi:
621 10.3329/ijarit.v5i1.24581
- 622 76. Dharmakeerthi RS, Thenabadu MW. Urease activity in soils: a review. *Journal of the*
623 *National Science Foundation of Sri Lanka*. 1996;24:159-195. doi:
624 10.4038/jnsf.v24i3.5548