

Original Research Article

SOIL FERTILITY DELINEATION TECHNIQUES AND RICE PRODUCTION

Comment [H1]: Suggestion: Soil fertility levels in Bangladesh for rice cultivation, geo-referenced.

ABSTRACT

Determination of soil fertility with minimum data set for crop zoning and devising fertilizer recommendations. Soil fertility evaluation method based on pH, cation exchange capacity (CEC), soil organic carbon (SOC), available phosphorus (P), sulphur (S), zinc (Zn), boron (B) and exchangeable potassium (K). The data were collected from existing literatures and scoring was done on 0–100 scale. The lowest score was assigned for the minimum value of tested attributes and then gradually higher scoring values. Arithmetic, weighted, geometric and most minimum of mean scores were calculated and their performances were compared with grain yield of dry season irrigated (Boro) rice. Soil fertility in 10-12 and 39-52 percent % areas of the country are very low and low, respectively. Medium fertile and fertile soils are distributed in 17-41% and in about 8% areas of the country. About 55% soils scored 70–95 (medium to high SOC) and the rest belongs to inferior quality. In some areas P build up has taken place (25% areas), but widespread K mining. Sulphur and Zn status in about 40 areas are low to very low (scored <35 and <40). Soils of the major areas of the country are with low pH (5.0-6.0) and CEC in the range of 15-25 cmol_c kg⁻¹. Weighted mean score and most minimum of eight attributes score showed good relationships with dry season irrigated rice yields than other tested methods indicating that this technique can be used for soil fertility rating in tropical countries.

Comment [H2]: As it is written, it seems that one sentence has no relation to the other. Make a connection between sentences.

Comment [H3]: What country?

Comment [H4]: %?

Key words: Soil attributes, Score, Weighted mean, Most minimum mean, Maps

26 **1. INTRODUCTION**

27 Global population is increasing and so does the demand for food production, which has
28 already created tremendous pressure on soil, a finite resource for mankind. It is our obligation
29 to keep soil healthy and productive through appropriate amendments and crop management
30 practices [1]. Indigenous nutrient supplying capacity and fertilizer management may make a
31 soil fertile for one type of crop but could be deficient for the others. So, determination of soil
32 fertility range would be important not only for producing healthy crops economically but also
33 for maintaining its productivity for future generations. Soils in Bangladesh are exposed to
34 high temperatures mostly; plenty of rainfall and greater pressure from growing two or more
35 crops in a year with or without balanced fertilizations [2] and thus nutrients mining are
36 widespread. New nutrient deficiencies are emerging [3], and there might be potential hidden
37 hunger for many others that need to be identified for efficient crop production.

38 Soil fertility varies among regions indicating that variable amounts of fertilizers need to be
39 applied for different types of crop production. Inadequate dose will impair crop yield, while
40 overdose can cause not only economic losses but also could be responsible for environmental
41 pollutions [4]. So, a broad knowledge on soil fertility can provide a better perception on
42 current nutrient status, distribution patterns and trends [5] that can be obtained through geo-
43 statistical and geospatial analyses [6,7]. Such analyses help in decision making processes for
44 precision agriculture and thus for improvement of crop productivity [8,9].

45
46 Soil fertility can be determined in different ways [10,11] by using soil pH, SOC, P, K,
47 exchangeable calcium (Ca), magnesium (Mg) and aluminium (Al), S, etc [12,13]. Mbogoni et
48 al [14] evaluated soil fertility by using average weighted data on SOC, soil pH, total N,
49 electrical conductivity, C/N ration, available P, exchangeable Ca, Mg and texture for rice
50 based system productivity improvement. Khaki et al [15] utilized square-root method as
51 parametric approach and Joint Fuzzy Membership functions to compute soil fertility index

52 (SFI). They found both the system suitable for soil fertility mapping and showed good
53 relations with rice yield. Desavathu et al [7] used soil pH, EC, N, P and for soil fertility
54 evaluation through inverse distance weightage interpolation. Thus it is found that researchers
55 had taken initiative for making soil fertility maps for specific locations or regions but a
56 simple method for a the country is still lacking. Therefore, the objective of this study was to
57 use geo-referenced data on selected soil attributes for preparation of soil fertility maps using
58 average, weighted mean, geometric mean and most minimum value techniques for
59 Bangladesh and to establish their relationships with rice yields.

60

61 **2. MATERIALS AND METHODS**

62 Data on **soil organic carbon (SOC)**, available **phosphorus (P)**, **sulphur (S)**, **zinc (Zn)**, and
63 **boron (B)**, **cation exchange capacity (CEC)**, soil pH and exchangeable K were collected
64 from Bangladesh Agricultural Research Council website, Soil Resource Development
65 Institute and existing available literatures. Average Boro clean rice (dry season irrigated crop,
66 hereafter as Boro rice) yields from 2007 to 2013 were collected from different volumes of
67 Bangladesh Bureau of Statistics and its relationships were established with soil fertility
68 scores. Although crop yields vary depending on inherent soil fertility, some other factors like
69 electrical conductivity, water quality (such as salinity) and its availability, agronomic
70 management practices, other biotic and abiotic factors also greatly influences crop
71 productivity. Nonetheless, inclusion of all those factors that influence soil fertility is beyond
72 the scope of the present investigation.

73 **2.1. Scoring criteria and map preparation**

74 Soil nutrient status in Bangladesh has been classified as very low, low medium, optimum and
75 high based on different ranges (Table 1). This classification system was considered for
76 assigning scoring values (Table 2) against each selected soil attribute. The scoring scale, as

77 considered in the present investigation, was 0–100. Attribute-wise soil fertility ratings over
78 different locations of Bangladesh were made by using MS-Excel Macros and IDRISI3.2.

79
80 Soil fertility scores, as determined by arithmetic mean (AM), geometric mean (GM),
81 weighted mean and (WM) and most minimum attribute (MAtrib_{score}) techniques, were used to
82 find out their relationships with Boro rice yields (64 districts of Bangladesh from 2007 to
83 2013) through regression analyses. Considering higher R² values, final soil fertility rating
84 maps were prepared based on weighted mean scores (Equa. I) and scores of the most
85 minimum of eight parameters for each district (Equa. II). Among soil attributes the most
86 limiting factors dictate crop yield, so we have provided weight to such factors in determining
87 WM as follows:

88
89
$$WM = ([SOC_{score}] * [P_{score}] * [K_{score}] * [CEC_{score}] * [pH_{score}])^{(1/5)} * 0.5 + [S_{score}] * 0.25 +$$

90
$$([Zn_{score}] * [B_{score}])^{(1/2)} * 0.25 \dots\dots\dots (I)$$

91 where, SOC_{score} is the soil organic carbon, P_{score}, K_{score}, CEC_{score}, pH_{score}, S_{score}, Zn_{score} and
92 B_{score} stand for **phosphorus**, **Kpotassium**, **CECcation exchange capacity**, soil pH,
93 **sulphur**, **Znzinc** and **Bboron** scores, respectively.

94
95 MAtrib_{score} for selected eight soil parameters were determined as follows:

96
$$MAtrib_{score} = \text{Geomean}(\text{Small}(\text{Atrib1:Atrib8,1}), \text{Small}(\text{Atrib1:Atrib8,2}), \dots\dots\dots,$$

97
$$\text{Small}(\text{Atrib1:Atrib8,8})) \dots\dots\dots (II)$$

98 where, Atrib1, 2, 3, ..., 8 are the soil parameters considered first, second, etc.

99
100 GM score was calculated as follows:

101
$$GM = ([B_{score}] * [K_{score}] * [P_{score}] * [CE_{score}] * [pH_{score}] * [SOC_{score}] * [S_{score}] * [Zn_{score}])^{(1/8)} \dots (III)$$

102
103 AM score was computed as follows:

104 $AM = ([B_{score}] + [K_{score}] + [P_{score}] + [CEC_{score}] + [pH_{score}] + [SOC_{score}] + [S_{score}] + [Zn_{score}]) / 8 \dots\dots (IV)$

105

106 Scores for most minimum of 1, 2, 3, 4, 5, 6 and 7 soil attributes were also found out in similar
107 fashion of Equa. II. The maps of tested attributes were prepared by using IDRISI3.2. Soil
108 fertility rating maps on the basis of WM and most minimum of eight attributes were used for
109 soil fertility delineation in Bangladesh. The other maps prepared based on different
110 techniques were used as supplementary figures.

111 3. RESULTS

112 Soil organic carbon, a vital component of fertility index showed >95 score for about 25%
113 areas in Bangladesh (Fig. 1a). About 55% soils had 70–95 score (medium to high SOC) and
114 the rest belongs to inferior quality. The scores for soil P varied from <10 to >75 in which
115 very low (<7 ppm), low (7-15 ppm), optimum (15-30 ppm) and high (>30 ppm) P levels
116 covered about 22.64, 47.74, 12.98 and 16.64 percent areas in the country (Fig. 1b). About
117 25% soils are with optimum/high (>80 score) K fertility. Majority areas (~43%) bear low K
118 (0.091-0.18 meq 100 g⁻¹ soil) and the rest belong to very low (<10 score) and medium (40-80
119 score) K categories y (Fig. 2a). The least score (<10) for S indicated that about 15.62% soils
120 are very poor (<7.5 ppm); 26.04% low and 14.54% medium and 43.79% areas are with
121 optimum/high S fertility status (Fig. 2b). In about 37.61% areas (score >75), soil Zn contents
122 are optimum to high (>1.351 ppm), 20.77% areas (40-75 score) are with medium Zn
123 containing soils and 41.61% soils scored <10 to 40 indicating (Fig. 3a) that Zn application is
124 a must practice for Bangladesh. **The content of BBoron fertility** is very low to low in about
125 50% soils (score <10 to 40) and rest of the soils had medium B content (Fig. 3b).

126

127 Soil pH score varied from <25 to >85 depending on locations and soil types in the country.
128 Maximum area coverage was 44.59% followed by 32.25% in the pH range of 5.0-6.0 and 6.5-
129 7.5, respectively (Fig. 4a). Soil pH below 5.0 and above 7.5 covers about 7.29% areas of the

130 country. The rest of the soils (15.22% areas) are with pH range of 6.0-6.5. The CEC scores
131 ranged from <25 to >85 depending on location in the country (Fig. 4b). The CEC of major
132 soils (47.46%) are 15-25 $\text{cmol}_c \text{kg}^{-1}$ followed by less than 15 $\text{cmol}_c \text{kg}^{-1}$ in 37.72% areas of
133 the country. Higher CEC (>25 $\text{cmol}_c \text{kg}^{-1}$) was found in 14.81% areas only.

134
135

136 **3.1. Soil fertility relationships with rice yield**

137 Soil fertility score based on different techniques and their relationships with clean rice yields
138 are shown in Fig. 5. About 49% yield variabilities are explained by the WM and most
139 minimum of eight tested soil attributes score (MAttrib-8). The performances of AM and GM
140 techniques in explaining yield variabilities were the least compared to others. Most minimum
141 1-7 soil attributes score explained Boro rice yield variabilities by about 23-42 percent.

142

143 **3.2. Soil fertility status**

144 Soil fertility scores varied from <35 to >60 with WM score technique and it was <35 to >55 with the
145 MAttrib-8 (Fig. 6). In the lowest soil fertility score (<35), area coverages are 10-12% of the country
146 based on above stated two techniques. The largest areas (28-30%) fall within the score of 40-45 under
147 both the techniques. Areas covered by higher scores (>55) were only about 16% of the country. Soil
148 fertility scores of 35-40 represented 9.41% and 24.08% areas under WM and MAttrib-8 techniques,
149 respectively. Similarly, 45-50 and 50-55 scores under WM and MAttrib-8 represented about 20% and
150 11-16% areas, respectively of the country. Based on GM, AM and MAttrib-1 to MAttrib-7 soil
151 fertility score varied greatly and represented different areas of the country, but major areas showed
152 low fertility score (data not shown). There were variations in the highest and the lowest scores
153 because of method employed (Table 3). The standard deviations were ± 8.52 , ± 7.19 , 7.73 and ± 8.52
154 for GM, AM, WM and MAttrib-8 means score, respectively having corresponding co-efficient of
155 variations of 19.62%, 14.22%, 16.58% and 19.62%.

156

157

158

159

160 4. DISCUSSION

161 In about 29% areas of the country, the SOC was at medium category; although there are high
162 and very high SOC in certain areas, especially with peat soils. In general, SOC was higher in
163 low lying areas, the single cropped zones, which remain 5–6 months under water in a year.
164 This level of SOC specifically in about 18% areas of the country is still inadequate for
165 satisfactory crop production [16]. As population pressure is increasing, farmers are using
166 such lands to increase total production through cropping intensification resulting in depletion
167 of SOC along with other **essential** plant nutrients. The decrease rate of SOC is comparatively
168 faster with arable cropping over time [17] with or without addition of organic manures. So,
169 we have found lower SOC rating in intensely cropping zones of Bangladesh. Partial
170 productivity of applied fertilizers is also decreasing indicating that nutrients from organic
171 matter (OM) need to be added that has been observed in our experiments at BRRI. Most soils
172 showed good response when OM was incorporated either from poultry litter, cow dung,
173 vermicompost [18,19] or green manuring because SOC influences soil pH, buffering
174 capacity, nutrient supplies and soil biological activity [20].

175

176 Although available P in the category of very low and low cover a larger area (about 70%), in
177 some areas its **build up** has taken place (Fig. 1b) because of cropping patterns followed, **soil**
178 **acidity**, fertilizer management options and inherent characteristics of parent materials [16,21].
179 As a greater area suffers from available P, corrective measures have to be taken for profitable
180 production [22]. This scenario is also true for global perspective in which P is depleting by
181 5.1 kg ha⁻¹ yr⁻¹ [23]. However, majority of the farmers in Bangladesh prefer to add N
182 fertilizer because of its immediate visible effects [2] and thus nutrient imbalance impose
183 negative impact on soil properties and crop production as a whole.

Formatted: Highlight

Formatted: Highlight

184
185 Potassium levels in major areas were very low to low (Fig. 2a) indicating that K mining was
186 taking place because of its substandard dose used by the farmers. Since farmers generally use
187 more N fertilizer and minimum K rate, the later is depleting rapidly in many areas of
188 Bangladesh [21, 22, 24]. In the global perspective, K is also depleting by $38.8 \text{ kg ha}^{-1} \text{ yr}^{-1}$
189 [23]; although its build up is not either uncommon in some areas because of excessive use
190 | with certain crops [16].

191 | Though S and Zn deficiencies are widespread in the country, its wet and dry depositions are
192 | also taking place because of industrial development [25]; but S fertilizer application still
193 | improves rice yields in many areas of the country.

194 | The scenario of B fertility is not healthy because in some areas it has depleted severely over
195 | time [16]. Yields of wheat, mustard and papaya reduce greatly in many parts of the country
196 | without B application. The depletion of soil fertility in areas with high cropping intensities
197 | [26] indicated that replenishment of removed nutrients were not taking place or it is beyond
198 | the capacity of the soils to supply major nutrients for growing high yielding crop varieties.
199 | There are evidences that Zn and B contents have been depleted severely from 1991 to 2012 in
200 | some selected areas of Bangladesh and thus crop productivity is declining [16].

201
202 | Lower soil pH covers quite larger areas in the north and north-east part and higher pH in the
203 | southern part of the country where **essential** plant nutrients availability is a limiting factor for
204 | satisfactory crop production without proper amendment. In some cases soil pH is increasing,
205 | especially in northern part of the country and thus playing a negative role on nutrient
206 | availability. It was reported that nutrient availability from applied fertilizers may be
207 | unavailable by more than 33-75% if soil pH ranges from 4.5 to 5.5 [27]. Generally, major
208 | nutrients are available for plants when soil pH varies from 6.5 to 7.5 [28]. Among others, soil
209 | P and many micronutrients become unavailable when pH exceeds 7.5, but molybdenum (**Mo**)

210 availability increases in alkaline pH. Moreover, CEC also depends on soil pH in a neutral soil
211 will have higher CEC than acidic soils [29]. Low CEC indicates light textured soils having
212 tendency of K and Mg deficiencies and faster decrease in soil pH [29, 30]. In such situations
213 frequent liming is needed for sandy type soils than clay category for profitable crop
214 cultivation.

Comment [H5]: What is light texture?

215
216 We have seen good relationships ($R^2 = 0.49$) of WM and MAttrib-8 scores with rice yields,
217 which is similar to the findings of Vasu et al. [31]. In Bangladesh, no grouping of soils has
218 been made based on combine scores or combine effects of different soil attributes; but
219 component-wise soil fertility delineations are available [22,26, 32]. So, our efforts are to
220 group soil fertility status combining all tested attributes as score <35 (very low fertility), 35-
221 45 (low fertility), 45-55 (medium fertile) and >55 (fertile). Accordingly, 10-12 and 39-52
222 percent areas of the country represented very low and low soil fertility, respectively (Fig. 6).
223 Medium fertile and fertile soils are distributed in 17-41% and in about 8% areas of the
224 country. These findings clearly indicate that special cares are needed for efficient and
225 economic crop production in major areas of Bangladesh. However, crop yields not only
226 depend on soil fertility, but also on other factors like water availability, temperature, and so
227 on. Moreover, soil fertility scores alone cannot explain yield variability of a crop rather it can
228 provide an indication for fertilizer rate determination and crop zoning for profitable farming.

229
230 Population pressure is increasing in Bangladesh, while soil fertility is decreasing indicating
231 that we are manipulating our soils beyond its bearing capacity. In general, nutrient mining is
232 taking place in Bangladesh at about $100 \text{ kg ha}^{-1} \text{ yr}^{-1}$ [22, 33] also reported low to very low
233 soil fertility for most of the studied soils in Bangladesh. This scenario is also true in terms of
234 global scale where soil fertility problems are associated with human-induced nutrient
235 depletion [23]. Besides, soil nutrient availability is limiting in cultivated lands of tropical

236 countries because of low inherent soil fertility [34]. Calcium deficiencies are emerging in
237 some agro-ecological zones (AEZ-3 and 21) of Bangladesh [3] and there might be hidden
238 hunger for micronutrients and thus reducing soil fertility and ultimately crop yield, but not
239 considered in the present investigation because of unavailability of data for the whole
240 country. In time series analyses for nutrient depletion, it was found that the contents of
241 exchangeable K, Ca and Mg have declined in all physiographic units except Old Himalayan
242 Piedmont and Madhupur Tract after 27 years of crop cultivation [35]. In one of our study, it
243 was also found that soil nutrient ratios have been changed in many places of Bangladesh
244 because of over exploitation of inherent soil fertility and thus Ca:P and N:Zn were playing
245 significant negative role with wet season rice yields under unfavourable ecosystems of
246 Bangladesh [36]. Similarly P: K ratio was acting antagonistically in agricultural ecological
247 zone 3, 18 and 26 of Bangladesh. All these factors indicate that we have to know our soils
248 before its use for crop production. Determination of soil fertility status by combining
249 important but minimum attributes can help in this regard for profitable farming and to
250 recuperate soil fertility through crop and fertilizer management.

251

252 **5. CONCLUSION**

253 A simple method of soil fertility evaluation for a country with minimum data sets is very
254 much desirable for proper crop zoning and delineating agronomic management options for
255 satisfactory crop production. We have determined soil fertility scores using pH, CEC, SOC,
256 available P, S, Zn, B and exchangeable K and following geometric, arithmetic, weighted and
257 mean approaches along with most minimum of tested attributes score. Weighted mean and
258 most minimum of soil attribute scoring methods showed better relationships with dry season
259 irrigated rice yields in Bangladesh indicating that this technique can be employed for soil
260 fertility assessment and its subsequent use for crop zoning and for determination of fertilizer
261 rates in similar environments around the globe.

262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288

REFERENCES

1. MacCarthy DS, Agyare WA, Vlek PLG, Adiku SGK. Spatial variability of some soil chemical and physical properties on agricultural landscape. *West Afr. J. Appl. Ecol.* 2013; 21: 47-61.
2. Biswas JC, Maniruzzaman M, Sattar MA, Neogi MG. Improvement of rice yield through fertilizer and cultural management at farmer's field. *Bangladesh Rice J.* 2008; 13: 9-14.
3. Saha PK, Islam S, Islam M N, Biswas JC, Haque MM. Soil plant nutrient status under intensive rice-farming systems in unfavourable eco-system of Bangladesh. *Intl. J. Biol. Pharma. Sci.* 2016; 2: 1-11.
4. Yang Y, Zhang S. Approach of developing spatial distribution maps of soil nutrients. In L. Daoliang (ed.) *Proc IFIPTC on Computer and Computing Tech Agric.* 2008; 1:565-571.
5. Dafonte JD, Ulloa GM, Jorge P, Glécio S, Vázquez M, Vidal E. Mapping of soil micronutrients in an European Atlantic agricultural landscape using ordinary kriging and indicator approach. *Bragantia* 2010; 69: 175-186.
6. Behera SK, Shukla AK. Spatial distribution of surface soil acidity, electrical conductivity, soil organic carbon content and exchangeable potassium, calcium and magnesium in some cropped acid soils of India. *Land Degrad. Dev.* 2015; 26: 71-79.
7. Desavathu RN, Nadipena AR, Peddada JR. Assessment of soil fertility status in Paderu mandal, Vaisakhapatnam district of Sndra Pradesh through Geopsatial techniques. *The Egyptian J. Remote Sensing Space Sci.* 2018; 21: 73-81.

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- 289 **8.** Markoski M, Arsov S, Mitkova T, Stamenkovska J. The benefit of GIS technologies
290 and precision agriculture principles in soil nutrient management for agricultural crop
291 production. *Bulg. J. Agric. Sci.* 2015; 21: 554-559.
- 292 **9.** Rahman AMAE, Natarajan A, Hedge R. Assessment of land suitability and capability
293 by integrating remote sensing and GIS for agriculture in Chamarajanagar district,
294 Karnataka, India. *Egypt. J. Remote Sensing Space Sci.* 2016; 19: 125-141.
- 295 **10.** Khalid R, Mahmood T, Bibi R, Siddique MT, Alvi S, Naz SY. Distribution and
296 indexation of plant available nutrients of rainfed calcareous soils of Pakistan. *Soil*
297 *Environ.* 2012; 31: 146-151.
- 298 **11.** Salehi N, Sepanlow MG, Gorzin BJ. An evaluation of soil fertility using soil organic carbon,
299 potassium, phosphorus and salinity factors for rice cultivation by fuzzy logic and AHP
300 technique. *Int. J. Agri. Crop Sci.* 2013; 5: 2233-2241.
- 301 **12.** Moran EF, Brondizio ES, Tucker JM, da SilvaForsberg MC, McCracken S, Falesi I.
302 Effects of soil fertility and land use on forest succession in Amazonia. *For. Ecol.*
303 *Manag.* 2010; 139: 93-108.
- 304 **13.** Panwar P, Pal S, Reza SK, Shama B. Soil fertility index, soil evaluation factor, and
305 microbial indices under different land uses in acidic soil of humid subtropical India.
306 *Commun. Soil Sci. Plan Anal.* 2011; 42: 2724-2737.
- 307 **14.** Mbogoni JDJ, Kiwambo BJ, Urassa GJ, Assenga SV. Soil fertility appraisal for
308 enhancing productivity in rice-based system of Ruvu basin, Agricultural Research
309 Institute Mlingano, Tanga, Tanzania; 2011.
- 310 **15.** Khaki BD, Honarjoo N, Davatgar N, Jalalian A, Golsselfidi HT. Assessment of two
311 soil fertility indexes to evaluate paddy fields for rice cultivation. *Sustainability* 2017;
312 9: 1299b (doi:10.3390/su9081299).
- 313 **16.** Siddique MMNEA, Halim MA, Kararuzzaman M, Karim D, Sultana J. Comparative insights
314 for investigation of soil fertility degradation in a piedmont area which cover the Anjamkhor

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- 315 union of Baliadangi upazila, Thakurgaon, Bangladesh. IOSR J. Environ. Sci. Toxico Food
316 Tech. 2014; 8: 82-87.
- 317 **17.** Zahid AM, Hossain MB, Halim MA, Hossain MA, Shahreen F. Organic matter and
318 plant nutrient depletion in major soil series in the high Ganges river floodplain. Intl. J.
319 Sustain. Agric. Tech. 2011; 7: 30-37.
- 320 **18.** Saleque MA, Abedin MJ, Bhuiyan NI, Zaman SK, Panaullah GM. Long-term effects of
321 inorganic and organic fertilizer sources on yield and nutrient accumulation of lowland rice.
322 Field Crop Res. 2004; 86: 53-65.
- 323 **19.** Haque MM, Akter M, Biswas JC. Performance of vermicompost and poultry manure
324 on rice yield and soil health. Annual Research Review Workshop 2016-17. Soil
325 Science Division, Bangladesh Rice Research Institute, Gazipur. 75pp; 2017.
- 326 **20.** Karlen DL, Andrews SS, Wienhold BJ, Zobeck TM. Soil quality assessment: past,
327 present and future. J. Integrated Biosci. 2008; 6: 3-14.
- 328 **21.** FRG (Fertilizer Recommendation Guide). Bangladesh Agricultural Research Council,
329 Farmgate Dhaka, 1215. In press; 2018.
- 330 **22.** Shil NC, Saleque MA, Islam M R, Jahiruddin M. Soil fertility status of some of the intensive
331 crop growing areas under major agro-ecological zones of Bangladesh. Bangladesh J. Agril.
332 Res. 2016; 41: 735-757.
- 333 **23.** Tan ZX, Lal R, Wiebe KD. Global soil nutrient depletion and yield reduction. J.
334 Sustainable Agric. 2005; 26: 123-146.
- 335 **24.** Rahman MH, Islam MR, Jahiruddin M, Rafii MY, Ismail MR, Malek MA.
336 Fertilization for increased crop production and nutrient balance in the maize-legume-
337 rice cropping pattern. J. Food Agric. Environ. 2013; 11: 653-656.
- 338 **25.** Biswas JC, Haque MM, Akter M, Hossain ATMS, Khan FH, Baki MZI, Sarker ABS,
339 Islam MR. Element Composition of the Atmospheric Depositions in Bangladesh. J.
340 Environ. Protec. 2018; 9:948-956.

Formatted: Portuguese (Brazil)

- 341 **26.** Jahiruddin M, Satter MA. Land and soil resource management report, in: Research
342 Priority in Agriculture and Development Vision Document-2030 and Beyond; 2010.
- 343 **27.** CU (Cornell University). How dose soil pH affect fertilizers and fertility. Cornell
344 University Cooperative Extension of Suffolk County. Extension Education Center,
345 423 Griffing Avenue, NY ([https://s3.amazonaws.com/assets.cce.cornell.edu/](https://s3.amazonaws.com/assets.cce.cornell.edu/attachments/3271/ow_does_soil_ph_affect_fertilizers_and_fertility.pdf?1413386058)
346 [attachments/ 3271/ow_does_soil_ph_affect_fertilizers_and_fertility.pdf?1413386058](https://s3.amazonaws.com/assets.cce.cornell.edu/attachments/3271/ow_does_soil_ph_affect_fertilizers_and_fertility.pdf?1413386058)
347 (Access on 6-2-2019).
- 348 **28.** IPNI (International Plant Nutrition Institute). Soil pH and the availability of plant
349 nutrients. IPNI, Georgia 30092-2806 USA (www.ipni.net); 2010.
- 350 **29.** CU (Cornell University). Cation exchange capacity (CEC). Agronomy fact sheet
351 series, Fact sheet number 22. Cornell University Cooperative Extension, College of
352 Agriculture and Life Sciences (<http://nmsp.css.cornell.edu>); 2007.
- 353 **30.** Noble AD, Gillman GP, Ruaysoongnern S. A cation exchange index for assessing
354 degradation of acid soil by further acidification under permanent agriculture in the
355 tropics. European J. Soil Sci. 2007; 51: 233-243.
- 356 **31.** Vasu D, Singh SK, Ray SK, Duraisami VP, Tiwary P, Chandran P, Nimkar AM,
357 Anantwar SG. Soil quality index (SQI) as a tool to evaluate crop productivity
358 in semi-arid Deccan plateau, India. *Geoderma*, [http:// dx. doi. org/ 10. 1016 / j.](http://dx.doi.org/10.1016/j.geoderma.2016.07.010)
359 [geoderma. 2016.07.010](http://dx.doi.org/10.1016/j.geoderma.2016.07.010) 0016-7061/© 2016 Elsevier B.V.
- 360 **32.** Rijmpa J, Jahiruddin M. Final; report on national strategy and p[lan for use of soil
361 nutrient balance in Bangladesh. SFFP consultancy report, DAE, Khamarbari, Dhaka;
362 2004.
- 363 **33.** Hossain M, Bayes A Islam SMF. A Diagnostic Study on Bangladesh Agriculture.
364 RED Working Paper Series, BRAC; 2017.

Formatted: Portuguese (Brazil)

Field Code Changed

Formatted: Portuguese (Brazil)

Formatted: No underline, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- 365 **34.** FAO (Food and Agriculture Organization). Status and trends in land and water
366 resources. Pages 19-60 in The State of the World's land and Water Resources for
367 Food and Agriculture: Managing Systems At Risk. FAO, UN and Earthscan; 2011.
- 368 **35.** Ali MM, Shaheed SM, Kubota D, Masunaga T, Wakatsuki T. Soil degradation during
369 the period 1967 – 1997 in Bangladesh. II. Selected chemical characters. Soil Sci.
370 Plant Nut. 1997; 43: 879-890.
- 371 **36.** Biswas JC, Haque MM, Saha PK. Rice yield potential under unfavorable soil
372 ecosystems in Bangladesh. Asian J Soil Sci. Plant Nutr. 2017;1: 1-10.

373

374 **Figure titles**

375 Fig. 1. Status of (a) soil organic carbon and (b) phosphorus in Bangladesh

376 Fig. 2. Status of (a) soil potassium and (b) sulphur in Bangladesh

377 Fig. 3. Status of (a) soil zinc and (b) boron in Bangladesh

378 Fig. 4. Distribution patterns of (a) soil pH (b) CEC in different parts of Bangladesh

379 Fig. 5. Relationships of clean Boro rice yields with scores of different soil attributes,
380 Bangladesh

381 Fig. 6. Soil fertility variations in Bangladesh as per (a) weighted mean and (b) most minimum
382 of eight soil-attributes scores

383 **Supple. Fig. 1. Soil fertility for Bangladesh according to (a) arithmetic mean, (b) geometric**
384 **mean, (c) one most minimum, (d) two most minimum, (e) three most minimum, (f) four most**
385 **minimum and (g) five most minimum, (h) six most minimum and (i) seven most minimum**
386 **attributes**

387

388

389

390

Comment [H6]: ???

Formatted: No underline, Highlight

Formatted: No underline, Highlight




Fig. 1 a & b

391
392

393

394

395

396




Fig. 2 a & b

397

398

399

400

401




Fig. 3 a & b


402

403

404

405

406



407

408


409

Fig. 4 a & b

410

411


412



413


414

415



416
417
418
419
420
421
422
423
424

Fig. 5



425
426
427
428
429
430

Fig. 6 a & b

Table 1. Soil nutrient status and its classifications in Bangladesh

	Critical limit	Very low	Low	Medium	Optimum	High
SOC (%)	-	<0.336	0.337-0.574	0.575-1.148	1.489-2.308*	>2.308**
Olsen P (ppmmg)	10	<7.50	7.51-15.00	15.1-22.5	22.51-30.00	30.1-37.50

dm⁻³

Formatted: No underline, Superscript

Bray P (mg dm⁻¹

³ppm)

7

<5.25

5.25-10.50

10.51-15.75

15.76-21.00

21.10-26.25

S (mg dm⁻³ppm)

10

<7.50

7.51-15.00

15.1-22.5

22.51-30.00

30.1-37.50

K (cmol_c dm⁻³meq

100 g⁻¹ soil)

0.12

<0.09

0.091-0.18

0.181-0.27

0.271-0.36

0.361-0.45

Formatted: No underline, Subscript

Ca (cmol_c dm⁻³meq

100 g⁻¹ soil)

2

<1.50

1.51-3.00

3.1-4.50

4.51-6.00

6.1-7.50

Formatted: No underline, Subscript

Mg (cmol_c dm⁻¹

³meq100 g⁻¹ soil)

0.5

<0.0375

0.376-0.75

0.751-1.25

1.16-1.50

1.51-1.875

Formatted: No underline, Subscript

Formatted: No underline, Superscript

Cu (mg dm⁻³ppm)

0.6

<0.15

0.151-0.30

0.31-0.45

0.451-0.60

0.61-0.75

Zn (mg dm⁻³ppm)

0.2

<0.45

0.451-0.90

0.91-1.35

1.351-1.81

1.81-2.25

Fe (mg dm⁻³ppm)

4

<3.00

3.10-6.00

6.1-9.00

9.1-12.00

12.1-15.00

Mn (mg dm⁻³ppm)

1

<0.75

0.756-1.50

1.51-2.25

2.56-3.00

3.1-3.75

B (mg dm⁻³ppm)

0.2

<0.15

0.151-0.30

0.31-0.45

0.451-0.60

0.61-0.75

Mo (mg dm⁻³ppm)

0.1

<0.075

0.076-0.15

0.151-0.225

0.226-0.30

0.31-0.375

431 FRG, 2012; *High and **Very high

432

433

434

435

436 **Table 2. Scoring criteria for different nutrient levels**

Soil nutrients		Soil pH		SOC		CEC	
Status	Score	Range	Score	Range	Score	Range	Score
Very low	5	<5.0	25	<0.336	40	<5	25
Low	30	5.0-5.5	45	0.337-0.574	70	5-10	40
Medium	70	5.5-6.0	65	0.575-1.148	85	10-20	65
Optimum	100	6.0-6.5	75	1.489-2.308	95	20-30	75
High	100	6.5-7.0	85	>2.308	100	30-40	80
		7.0-7.5	95			40-50	85
		>7.5	25			>50	100

437

438

439

440

441 **Table 3. Soil fertility scoring variations due to methods**

	Geometric	Arithmetic	Weighted	MAttrib-8
	mean	mean	mean	
Maximum	60.19	63.38	61.55	60.19
Minimum	24.20	31.20	30.10	24.20
Mean	43.42	50.57	46.60	43.42
Sd (\pm)	8.52	7.19	7.73	8.52
CV(%)	19.62	14.22	16.58	19.62

442

443

444

445

446

UNDER PEER REVIEW