

**CORRELATION ANALYSIS OF TOXIC METALS DISTRIBUTION AND POLLUTION INDICES IN SOIL, BEANS AND MAIZE SAMPLES OF KANO STATE, NIGERIA**

**ABSTRACT**

Correlational study and evaluation of pollution indices of toxic metals distribution in soil and crops of a population is imperative for accessing risk of chronic diseases associated with these metals. Correlational analysis for the distribution of; lead (Pb), cadmium (Cd), chromium (Cr) and mercury (Hg) in soil (S), bean (B) and maize (M) was conducted around Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and Wudil (WDL) as sampling zones around Kano State, Nigeria. The samples were collected from farm harvests in each of the sampling zones under the study and designed as; BKRS, BKRB, BKRM, DBTS, DBTB DBTM, GRZS, GRZB GRZM, UGGS, UGGB, UGGM, WDLs, WDLB, WDLM. The metal Concentration was determined using atomic absorption spectrometry (AAS). Results in mg/kg across the local governments indicates respective ranges for Hg, Pb, Cd, and Cr of; 0.33 - 3.13, 0.14 - 0.84, 0.02 - 0.05 and 0.01- 0.49 in soil, 0.04-4.23, 0.06-0.23, 0.02-0.04 and 0.00-0.10 in maize and 0.20-4.23, 0.16-0.19, 0.03-0.04 and 0.00-0.03 in beans. Although with exception of mercury, the ranges of the toxic metals are within the tolerable ranges set by International Standard Tolerable Limits and European Regulatory Standard. Potential hazard may be speculated because the detected levels are on higher tolerable ranges. A higher level of mercury in almost all the samples indicates potential hazards associated with human activities in those areas. Strong positive correlation between soils samples with respect to level of some of the toxic metal may suggest a common nature of the soil, while the negative correlation may be due variation in agrochemicals in-use. For the pollution load index, Wudil had the highest soil pollution load index for Hg ( $3.13 \pm 0.16$ ), Cd ( $1.6 \times 10^{-2} \pm 0.01$ ) and Cr ( $4.9 \times 10^{-3} \pm 0.01$ ), while Ungogo had the highest pollution load for Pb. Also, all grains within the study zones exhibited a positive transfer factor, except Cr in Bunkure, Danbatta and Gwarzo. It may be concluded that crops grown in those areas may bioaccumulates some of these toxic metals, thereby incorporating them into food chain, hence potential health risk.

**Keywords:** *toxic metals, Soil, Farm produce, Correlation, Distribution, pollution index Kano-*

**LGAs**

**1.0 INTRODUCTION**

Toxic metals are confined to those metals having higher atomic number (Norman, 1981), associated with little or no biochemical functions. However; they have many industrial and agricultural applications (Alloway, 1995) and are relatively high density metallic elements with

39 potential toxicity at low concentration (Holding, 2004). According to WHO (2014) toxic metals  
40 occur in trace amount and are at least five times denser than water, They are stable and persistent  
41 in environmental samples, hence bioaccumulates to pass up via food chain to humans. According  
42 to Alhassan *et al* (2012) sources of heavy metals are varied and include natural sources and all  
43 human activities, with possibility of polluting the environment. In another perspective,  
44 environmental toxic metals are due to direct deposition from mining activities, industrial  
45 processes as well as waste water from domestic processes, in addition to agricultural practices  
46 (Madyiwa, 2006). Heavy metals contamination to agricultural soils could be through a variety of  
47 sources among which is land application of biosolids, fertilizers, livestock manure,  
48 agrochemicals (Pesticides), irrigation activities and atmospheric droplet. Worthy consideration is  
49 accumulation of these metals in agricultural soils and possible negative impacts on soil fertility  
50 and large potential accumulation in crops for possible incorporation to human food chains (Gray  
51 *et al.*, 2003). Unlike the inorganic, the organic (alkylated) forms are readily taken up by body  
52 tissues and can be retained for a considerable length of time (Garrett *et al.*, 1992) this is due to  
53 their lipid solubility. The tissue penetrating power of organometals crown them more toxic to  
54 humans than the inorganic form (Carpenter, 2001).

55 Human exposure to soil accumulated toxic metal may be chronic via food chain transfer or acute  
56 by direct ingestion or dermal contact. The former may be associated with; mental lapses, kidney,  
57 liver, lung and gastrointestinal tract abnormalities, central nervous insufficiency, lower energy  
58 levels, damage to blood compositions, kidneys, and other vital organs (Gray *et al.*, 2003). Long  
59 term exposure to low dose may result in slowly progressing physical, muscular, and neurological  
60 degenerative processes that mimic Alzheimer`s disease, Parkinson`s disease, muscular  
61 dystrophy, and multiple Sclerosis (Srivastava and Goyal, 2010). Allergies are not uncommon and

62 repeated long-term contact with some metals or their compounds may even cause cancer. The  
63 letter, depends on the degree to which a system, tissue, organ or cell is affected by a heavy metal  
64 type and the individuals degree of exposure (Vallero and Letcher, 2013).

65 Mercury combines with other elements to form organic and inorganic mercury compounds.  
66 Metallic mercury used to produce chlorine gas, caustic soda, in thermometers and dental fillings,  
67 switches, light bulbs, and batteries, including coal-burning power plants are the largest human-  
68 caused source of mercury emissions to the air in the United States. Mercury in soil and water is  
69 converted by microorganisms to methyl mercury, a bio accumulating toxin (Sabine and Wendy,  
70 2009). The EPA reported that mercuric chloride and methyl mercury are possible human  
71 carcinogens and nervous system is very sensitive to all forms of mercury. Exposure to high  
72 levels can permanently damage the brain, kidneys, and developing fetuses. Effects on brain  
73 functioning may result in irritability, shyness, tremors, changes in vision or hearing, and memory  
74 problems. Short-term exposure to high levels of metallic mercury may cause lung damage,  
75 nausea, vomiting, diarrhea, increases in blood pressure or heart rate, skin rashes, and eye  
76 irritation (Davidson *et al.*, 2004).

77 Exposure to high lead levels can severely damage the brain and kidneys and ultimately cause  
78 death. In pregnant women, high levels of exposure to lead may cause miscarriage. High level  
79 exposure in men can damage the organs responsible for sperm production (Emsley, 2011).

80 Chromium, chromium (III) compounds and chromium metal are not considered a health hazard,  
81 while the toxicity and carcinogenic properties of chromium (VI) have been known since late 19th  
82 century, while Cadmium exposure is a phenomenon of the early 20th century, and onwards. In  
83 Japan in 1910, the Mitsui Mining and Smelting Company discharging cadmium into the  
84 Jinzugawa River, as a byproduct of mining operations. Residents in the surrounding area

85 subsequently consumed rice grown in cadmium-contaminated irrigation water, experienced  
86 softening of the bones and kidney failure (Gray *et al.*, 2003).

87

## 88 **2.0 STUDY AREA**

89 Kano State was formally established on April 1, 1968. It is situated in a semi-arid region, located  
90 between latitudes 10.30°N to 13°N and longitude 7.40°E and 10.39°E. Kano city is at 472.45  
91 meters above sea level. The state is bordered by Jigawa State in the north-east, Katsina State in  
92 the north-west and Kaduna State is on the southern boundary. According to 2006 census, the  
93 state has a population of 9.5 million with projection of over 13 million in 2018. It has a total land  
94 area of 20,760 square kilometres with 1, 754, 200 hectares for agricultural and 75,000 hectares  
95 forest vegetation and grazing land. The state is noted for its fairly stable climate with relatively  
96 minor changes in temperature and humidity (Stilwell, 2000). Agriculture is the mainstay of the  
97 economy involving at least 75% of the rural population. Important crops produced in the State  
98 include maize, beans, rice, corn and varieties of vegetables.

99

## 100 **3.0 MATERIALS AND METHODS**

### 101 **3.1 Sample Collection and Preparation**

102 The samples were dehusked into grains washed with de-ionized water and dried to constant  
103 weight in hot air oven at 105°C for 6hrs, ground to powder and stored in clean air tied  
104 containers.

105

106

107

108

109 **3.2 ANALYSIS**

110 Five 5.0g of the prepared samples of subsection 3.1 were air dried, ground in an agate mortar  
111 and was placed in quartz crucibles for dry ashing and mineralized thermally in a muffle furnace  
112 at 450°C for 12hrs. The mineralized samples was then acid digested according to Miller (1998),  
113 by dissolving in 20ml of 1:1 (v/v) concentrated HNO<sub>3</sub> and HCl acids in 100ml volumetric flask.  
114 The flask was then heated in an electro thermal heater with gentle swirling till digestion  
115 completed by evolution of white fumes. The cooled digests was filtered through what man No 1  
116 filter paper into 50ml volumetric flask and was diluted to the mark with de-ionized water. The  
117 heavy metals content of the samples were determined using the atomic absorption  
118 spectrophotometer (AAS) and the concentration was calculated using the relation  $y = mx + c$   
119 from calibration of each metal standard (AOAC, 2005).

120

121 **3.3 Determination of Pollution Load Index (PLI), Transfer Factor (TR), Daily Intake of**  
122 **Metals (DIM) and Health Risk Index (HRI)**

123

124 **3.3.1 Pollution Load Index (PLI)**

125 The following modified equation was used to determine the pollution load index (PLI) level in  
126 soils (Liu *et al.*, 2005).

$$PLI = \frac{C_s}{C_r}$$

127 Where  $C_{soil}$ = Concentration of metal in soil.

128  $C_{reference}$ = Reference standard value of metal in soil.

129

130

131

132 **3.3.2 Transfer Factor (TF)**

133 Transfer factor (TF) from soil to grains was calculated as described by Liu *et al.* (2005).

134

$$TF = \frac{C_g}{C_s}$$

135 Where  $C_g$  = Concentration of metal in grains

136  $C_s$  = Concentration of metal in soil.

137

### 138 3.3.3 Daily Intake of Metals (DIM)

139 The Daily intake of metals (DIM) was determined as described by Wang *et al.* (2005).

$$DIM = \frac{C_g \times C_f \times D_{fi}}{B_{abw}}$$

140 Where:  $C_g$  = concentration of heavy metals in grains (mg/kg)

141  $C_f = 0.085$

142  $D_{fi}$  (daily food intake) = 0.345 kg per day for adult

143  $B_{abw}$  (average body weight) = 55.90 kg for adult (Wang *et al.*, 2005).

144

### 145 3.3.4 Health Risk Index (HRI)

146 Health risk index was calculated as described by US-EPA (2002).

$$HRI = \frac{DIM}{RFD}$$

147 Where  $RFD$  = reference oral dose (USEPA, 2002).

148  $DIM$  = Daily intake of metals

149

150

151

### 152 Statistical analysis:

153 The data were expressed as mean  $\pm$  standard deviation, two-way ANOVA was conducted to

154 examine the interaction between the source (soil, beans and maize) and location (Local

155 Government Areas) on the level of heavy metals (Hg, Pb, Cd and Cr). SPSS version 20.0 was  
156 used for statistical analyses and  $P < 0.05$  was set as level significant.


157

## 158 4.0 RESULT AND DISCUSSION

### 159 4.1 RESULTS

160 Figure 1 shows the imperfect linear distribution of mean Hg concentration in samples of Bunkure  
161 (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and Wudil (WDL) local government  
162 areas. The near zero linear coefficient (r) correlation is suggesting the distribution of Hg across  
163 the local governments is not perfectly correlated. This is further supported by the square of  
164 coefficient of multiple variations ( $R^2$ ), which gave a measure of the total variation in the  
165 dependent variable (soil), explained by variations in the explanatory variable (farm produce).

166




167

168 *Figure 1: Non-linear distribution of mean Hg levels in sample of Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG)*  
169 *and Wudil (WDL) local government of Kano state Nigeria.*

170


171 Figure 2 displays weak linear correlations for the distribution of mean Pb levels in samples  
 172 around Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and Wudil (WDL)  
 173 local government of Kano state Nigeria, as indicated by the value of linear coefficient (r). The  
 174 measure of the total variation in the dependent variable ( $R^2$ ) further justifies that.



176  
 177 *Figure 2: Non-linear distribution of mean Pb concentration in sample of Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo*  
 178 *(UGG) and Wudil (WDL) local government of Kano state Nigeria.*  
 179

180 Strong non – linear correlation ( $r = 0.0015$ ) for the distribution of mean Cd levels in samples  
 181 around Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and Wudil (WDL)  
 182 local government of Kano state Nigeria is shown on Figure 3 with a reasonable measure of the  
 183 total variation in the dependent variable ( $R^2$ ).





184


185 *Figure 3: imperfect linear distribution of mean Cd value in sample of Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and*  
 186 *Wudil (WDL) local government of Kano state Nigeria.*

187

188 Figure 4 shows an imperfect linear distribution of Cr in samples of Bunkure (BKR), Danbatta  
 189 (DBT), Gwarzo (GRZ), Ungogo (UGG) and Wudil (WDL) local government of Kano state  
 190 Nigeria. The near zero linear coefficient (r) correlation is suggesting the distribution of the Cr  
 191 across the local governments is not perfectly correlated. This is further substantiated by the value  
 192 of the square of coefficient of multiple variations ( $R^2$ ), which gave a measure of the total  
 193 variation in the dependent variable.

194

195



196

197 *Figure 4: imperfect linear distribution of mean Cr value in sample of Bunkure (BKR), Danbatta (DBT), Gwarzo (GRZ), Ungogo (UGG) and*  
 198 *Wudil (WDL) local government of Kano state Nigeria.*

199

200

UNDER PEER REVIEW

Table 1 shows the coefficient correlation matrix for Mercury (Hg) distribution in soil, beans and maize of the five local government areas of the study. It show a significant perfect correlation ( $P < 0.01$ ) between; WDLS vs WDLM, DBTS vs GRZB, UGGS vs BKRB, WDLS vs BKRM, UGGM vs GRZM, UGGM vs WDLB and imperfects positive correlation between GRZS vs GRZB.

**Table 1: coefficient correlation of Mercury (Hg) distribution in soil, beans and maize of five local government areas of Kano state Nigeria**

	Hg DBTS	Hg DBTB	Hg DBTM	Hg BKRS	Hg BKRB	Hg BKRM	Hg GRZS	Hg GRZB	Hg GRZM	Hg UGGS	Hg UGGB	Hg UGGM	Hg WDLS	Hg WDLB	Hg WDLM
HgDBTS	1.000														
HgDBTB	.000	1.000													
HgDBTM	.000	-1.000**	1.000												
HgBKRS	.816	.500	-.500	1.000											
HgBKRB	-1.000**	.000	.000	-.816	1.000										
HgBKRM	.333	-.816	.816	.000	-.333	1.000									
HgGRZS	.816	-.500	.500	.500	-.816	.816	1.000								
HgGRZB	1.000**	.000	.000	.816	-1.000**	.333	.816	1.000							
HgGRZM	.333	.816	-.816	.816	-.333	-.333	.000	.333	1.000						
HgUGGS	-1.000**	.000	.000	-.816	1.000**	-.333	-.816	-1.000**	-.333	1.000					
HgUGGB	.000	1.000**	-1.000**	.500	.000	-.816	-.500	.000	.816	.000	1.000				
HgUGGM	.333	.816	-.816	.816	-.333	-.333	.000	.333	1.000**	-.333	.816	1.000			
HgWDLS	.333	-.816	.816	.000	-.333	1.000**	.816	.333	-.333	-.333	-.816	-.333	1.000		
HgWDLB	.333	.816	-.816	.816	-.333	-.333	.000	.333	1.000**	-.333	.816	1.000**	-.333	1.000	
HgWDLM	.333	-.816	.816	.000	-.333	1.000**	.816	.333	-.333	-.333	-.816	-.333	1.000**	-.333	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed).

Table 2 shows coefficient correlation of Lead (Pb) distribution in soil, and farm produce (beans and maize) in five local government areas of the study. The lead distributions show significant negative correlation ( $P < 0.01$ ) between soil in DBT and the farm produce, maize in BKR and beans in WDL. In some instances it shows significant positive correlation ( $P < 0.01$ ) between; DBTS vs UGGS and UGGB, BKRB vs WDLM, GRZB vs GRZM and WDLM and UGGM vs WDL. Imperfect positive correlation was shown between soil sample of each local government and respective farm produce.

Table 2: coefficient correlation of Lead (Pb) distribution in soil, beans and maize of five local government areas of Kano state Nigeria

	Pb DBTS	Pb DBTB	Pb DBTM	Pb BKRS	Pb BKRB	Pb BKRM	Pb GRZS	Pb GRZB	Pb GRZM	Pb UGGS	Pb UGGB	Pb UGGM	Pb WDL	Pb WDLB	Pb WDLM
<b>PbDBTS</b>	1.000														
<b>PbDBTB</b>	-1.000**	1.000													
<b>PbDBTM</b>	-1.000**	1.000**	1.000												
<b>PbBKRS</b>	-1.000**	1.000**	1.000**	1.000											
<b>PbBKRB</b>	-.333	.333	.333	.333	1.000										
<b>PbBKRM</b>	1.000**	-1.000**	-1.000**	-1.000**	-.333	1.000									
<b>PbGRZS</b>	.333	-.333	-.333	-.333	.333	.333	1.000								
<b>PbGRZB</b>	-.333	.333	.333	.333	1.000**	-.333	.333	1.000							
<b>PbGRZM</b>	-.333	.333	.333	.333	1.000**	-.333	.333	1.000**	1.000						
<b>PbUGGS</b>	1.000**	-1.000**	-1.000**	-1.000**	-.333	1.000**	.333	-.333	-.333	1.000					
<b>PbUGGB</b>	1.000**	-1.000**	-1.000**	-1.000**	-.333	1.000**	.333	-.333	-.333	1.000**	1.000				
<b>PbUGGM</b>	.333	-.333	-.333	-.333	.333	.333	1.000**	.333	.333	.333	.333	1.000			
<b>PbWDL</b>	.333	-.333	-.333	-.333	.333	.333	1.000**	.333	.333	.333	.333	1.000**	1.000		
<b>PbWDLB</b>	-.333	.333	.333	.333	-.333	-.333	-1.000**	-.333	-.333	-.333	-.333	-1.000**	-1.000**	1.000	
<b>PbWDLM</b>	-.333	.333	.333	.333	1.000**	-.333	.333	1.000**	1.000**	-.333	-.333	.333	.333	-.333	1.000

\*\*Correlation is significant at the 0.01 level (2-tailed).

Table 3 show the coefficient correlation for the distribution of Cd in the local government area under study. It show significant positive correlation ( $P < 0.01$ ) between; DBTB vs DBTS, BKRS vs DBTM, GRZS vs DBTS and DBTB, GRZM vs BKRB, UGGB vs BKRM, UGGM vs BKRB, WDLs vs DBTS, DBTB and GRZS, and WDLM, DBTM and BKRS. The distribution projects significant negative correlation between BKRM vs DBTM and BKRS, UGGB vs DBTM and BKRS, WDLM vs BKRM and UGGB. Cd distribution in soil of each local government under the study show negative correlation with that of maize except for UGG that shows significant positive correlation ( $P < 0.01$ ). The distribution show positive correlation with that of beans except in UGG and WDL.

**Table 3: coefficient correlation of Cadmium (Cd) distribution in soil, beans and maize of five local government areas of Kano state Nigeria**

	Cd DBTS	Cd DBTB	Cd DBTM	Cd BKRS	Cd BKRB	Cd BKRM	Cd GRZS	Cd GRZB	Cd GRZM	Cd UGGS	Cd UGGB	Cd UGGM	Cd WDLs	Cd WDLB	Cd WDLM
<b>CdDBTS</b>	1.000														
<b>CdDBTB</b>	1.000**	1.000													
<b>CdDBTM</b>	-.333	-.333	1.000												
<b>CdBKRS</b>	-.333	-.333	1.000**	1.000											
<b>CdBKRB</b>	.333	.333	.333	.333	1.000										
<b>CdBKRM</b>	.333	.333	-1.000**	-1.000**	-.333	1.000									
<b>CdGRZS</b>	1.000**	1.000**	-.333	-.333	.333	.333	1.000								
<b>CdGRZB</b>	.816	.816	.000	.000	.816	.000	.816	1.000							
<b>CdGRZM</b>	.333	.333	.333	.333	1.000**	-.333	.333	.816	1.000						
<b>CdUGGS</b>	.333	.333	.333	.333	1.000**	-.333	.333	.816	1.000**	1.000					
<b>CdUGGB</b>	.333	.333	-1.000**	-1.000**	-.333	1.000**	.333	.000	-.333	-.333	1.000				
<b>CdUGGM</b>	.333	.333	.333	.333	1.000**	-.333	.333	.816	1.000**	1.000**	-.333	1.000			
<b>CdWDLs</b>	1.000**	1.000**	-.333	-.333	.333	.333	1.000**	.816	.333	.333	.333	.333	1.000		
<b>CdWDLB</b>	.000	.000	-.816	-.816	-.816	.816	.000	-.500	-.816	-.816	.816	-.816	.000	1.000	
<b>CdWDLM</b>	-.333	-.333	1.000**	1.000**	.333	-1.000**	-.333	.000	.333	.333	-1.000**	.333	-.333	-.816	1.000

\*\* . Correlation is significant at the 0.01 level (2-tailed).

The coefficient correlation of Chromium (Cr) distribution in the local government areas under study is shown on Table 4. It shows significant correlation ( $P < 0.01$ ) between; DBTS vs DBTB, DBTM, BKRB and GRZM, DBTB vs BKRB and GRZM, DBTM vs BKRB, GRZM, BKRS vs GRZB, GRZS vs WDLB, UGGS vs UGGB, UGGM vs WDLM. The lead distributions show significant negative correlation ( $P < 0.01$ ) in some instances Table 4. Positive correlation exists in each local government between soil and farm produce except in BKR and beans in GRZ and WDL that shows negative correlation.

**Table 4: coefficient correlation of Chromium (Cr) distribution in soil, beans and maize of five local government areas of Kano state Nigeria**

	Cr DBTS	Cr DBTB	Cr DBTM	Cr BKRS	Cr BKRB	Cr BKRM	Cr GRZS	Cr GRZB	Cr GRZM	Cr UGGS	Cr UGGB	Cr UGGM	Cr WDLB	Cr WDLM	
<b>CrDBTS</b>	1.000														
<b>CrDBTB</b>	1.000**	1.000													
<b>CrDBTM</b>	1.000**	1.000**	1.000												
<b>CrBKRS</b>	-.500	-.500	-.500	1.000											
<b>CrBKRB</b>	1.000**	1.000**	1.000**	-.500	1.000										
<b>CrBKRM</b>	-.500	-.500	-.500	-.500	-.500	1.000									
<b>CrGRZS</b>	-.816	-.816	-.816	.000	-.816	.816	1.000								
<b>CrGRZB</b>	-.500	-.500	-.500	1.000**	-.500	-.500	.000	1.000							
<b>CrGRZM</b>	1.000**	1.000**	1.000**	-.500	1.000**	-.500	-.816	-.500	1.000						
<b>CrUGGS</b>	.816	.816	.816	-.816	.816	.000	-.333	-.816	.816	1.000					
<b>CrUGGB</b>	.816	.816	.816	-.816	.816	.000	-.333	-.816	.816	1.000**	1.000				
<b>CrUGGM</b>	.816	.816	.816	.000	.816	-.816	-1.000**	.000	.816	.333	.333	1.000			
<b>CrWDLB</b>	-.816	-.816	-.816	.816	-.816	.000	.333	.816	-.816	-1.000**	-1.000**	-.333	1.000		
<b>CrWDLM</b>	-.816	-.816	-.816	.000	-.816	.816	1.000**	.000	-.816	-.333	-.333	-1.000**	.333	1.000	
	.816	.816	.816	.000	.816	-.816	-1.000**	.000	.816	.333	.333	1.000**	-.333	-1.000**	1.000

\*\* . Correlation is significant at the 0.01 level (2-tailed).

The pollution load index of toxic metals in five Local Government Areas, Kano State, Nigeria is shown on Table 5. The result show that the highest soil pollution load (PL) of Mercury was in Wudil ( $3.13 \pm 0.16$ ) followed by Ungogo soil with Hg PLI of  $2.72 \pm 0.48$ , Gwarzo ( $2.51 \pm 0.46$ ) Danbatta ( $2.24 \pm 1.30$ ) and Bunkure with the lowest mercury PLI of  $0.33 \pm 0.44$ . Ungogo shows the highest soil PL of Lead ( $5.6 \times 10^{-3} \pm 0.04$ ) while the lowest soil Lead PL was in Bunkure ( $0.93 \times 10^{-3} \pm 0.08$ ). Also, Wudil show the highest PL of Cadmium ( $1.6 \times 10^{-2} \pm 0.01$ ) and PL of Chromium ( $4.9 \times 10^{-3} \pm 0.01$ ).

**Table 5: Toxic metals pollution load index in soil of five Local Government Areas, Kano State, Nigeria**

Study area	Hg	Pb	Cd	Cr
<b>Bunkure</b>	$0.33 \pm 0.44$	$0.93 \times 10^{-3} \pm 0.08$	$0.67 \times 10^{-2} \pm 0.00$	$0.1 \times 10^{-3} \pm 0.02$
<b>Danbatta</b>	$2.24 \pm 1.30$	$1.66 \times 10^{-3} \pm 0.13$	$1.0 \times 10^{-2} \pm 0.00$	$0.2 \times 10^{-3} \pm 0.02$
<b>Gwarzo</b>	$2.51 \pm 0.46$	$1.9 \times 10^{-3} \pm 0.05$	$1.3 \times 10^{-2} \pm 0.00$	$1.5 \times 10^{-3} \pm 0.03$
<b>Ungogo</b>	$2.72 \pm 0.48$	$5.6 \times 10^{-3} \pm 0.04$	$1.3 \times 10^{-2} \pm 0.00$	$4.4 \times 10^{-3} \pm 0.04$
<b>Wudil</b>	$3.13 \pm 0.16$	$3.3 \times 10^{-3} \pm 0.01$	$1.6 \times 10^{-2} \pm 0.01$	$4.9 \times 10^{-3} \pm 0.01$

Table 6 shows the transfer factor of toxic metals from soil to grains in the five Local Government Areas under study. From the result, all the grains within the study zones exhibited a positive transfer factor, with the exception of Chromium in Bunkure, Danbatta and Gwarzo, where the transfer factor was zero.

**Table 6: The transfer factor of toxic metals from soil to grains from five Local Government Areas, Kano State, Nigeria.**

Study area	Hg		Pb		Cd		Cr	
	Beans	Maize	Beans	Maize	Beans	Maize	Beans	Maize
<b>Bunkure</b>	0.60	0.12	1.21	0.42	1.5	1	0	0
<b>Danbatta</b>	0.70	0.54	0.64	0.84	1	1	0	0
<b>Gwarzo</b>	0.67	0.87	0.55	0.62	0.75	0.75	0	0
<b>Ungogo</b>	1.12	1.23	0.21	0.27	1	1	0.06	0.22
<b>Wudil</b>	1.35	1.35	0.38	0.46	0.80	0.80	0.06	0.12

Table 7 show the daily intake of toxic metals in the five Local Government Areas under study. From the results, beans and maize from Wudil and Ungogo and maize from Gwarzo provided the highest daily intake of Mercury. This was followed by Gwarzo beans, beans and maize from Danbatta and Bunkure beans. The daily intake of Chromium from Bunkure beans and maize, Danbatta beans and maize and Gwarzo beans and maize was zero.

**Table 7: Daily intake of toxic metals in beans and maize across five Local Government Areas under study**

Study area	Hg		Pb		Cd		Cr	
	Beans	Maize	Beans	Maize	Beans	Maize	Beans	Maize
<b>Bunkure</b>	$1.0 \times 10^{-4}$	$2.0 \times 10^{-5}$	$8.9 \times 10^{-5}$	$3.1 \times 10^{-5}$	$1.5 \times 10^{-5}$	$1.0 \times 10^{-5}$	0	0
<b>Danbatta</b>	$8.2 \times 10^{-4}$	$6.4 \times 10^{-4}$	$8.3 \times 10^{-5}$	$1.0 \times 10^{-5}$	$1.5 \times 10^{-5}$	$1.5 \times 10^{-5}$	0	0
<b>Gwarzo</b>	$8.8 \times 10^{-4}$	$1.1 \times 10^{-3}$	$8.3 \times 10^{-5}$	$9.4 \times 10^{-5}$	$1.5 \times 10^{-5}$	$1.5 \times 10^{-5}$	0	0
<b>Ungogo</b>	$1.6 \times 10^{-3}$	$1.7 \times 10^{-3}$	$9.4 \times 10^{-5}$	$1.2 \times 10^{-4}$	$2.0 \times 10^{-5}$	$2.0 \times 10^{-5}$	$1.5 \times 10^{-5}$	$5.2 \times 10^{-5}$
<b>Wudil</b>	$2.2 \times 10^{-3}$	$2.2 \times 10^{-3}$	$9.9 \times 10^{-5}$	$1.2 \times 10^{-4}$	$2.0 \times 10^{-5}$	$2.0 \times 10^{-5}$	$1.0 \times 10^{-5}$	$3.1 \times 10^{-5}$

The health risk index of toxic metals across the five Local Government Areas, Kano State, Nigeria is presented on Table 8. From the results beans and maize from Wudil had highest health risk index due high Mercury content, followed by Ungogo, Gwarzo, Danbatta and Bunkure with the lowest health risk index of Mercury. All the remaining toxic metals showed a health risk index of less than one.

**Table 8: Health risk index in beans and maize across five zones in Kano State, Nigeria**

Study area	Hg		Pb		Cd		Cr	
	Beans	Maize	Beans	Maize	Beans	Maize	Beans	Maize
<b>Bunkure</b>	0.33	0.10	0.025	0.008	0.015	0.01	0	0
<b>Danbatta</b>	2.7	2.1	0.023	0.003	0.015	0.015	0	0
<b>Gwarzo</b>	2.9	3.6	0.023	0.026	0.015	0.015	0	0
<b>Ungogo</b>	5.3	5.6	0.026	0.035	0.02	0.02	0	0
<b>Wudil</b>	7.3	7.3	0.028	0.035	0.02	0.02	0	0



## 4.2 DISCUSSION

Current study evaluated the levels of toxic metals (mercury, lead, cadmium and chromium) in soil and farm produce (maize and beans) of five local governments of Kano state. Mean Pb, Cd and Cr concentration of the soil samples are within the respective limits of 150mg/Kg, 3mg/kg and 100 mg/Kg set by International Standard tolerable limits (2001). However, with exception of BKRS, all soil samples have mean mercury (Hg) concentration above 1mg/Kg, the limits set by international standard tolerable limits (2001). Mean Pb, Cd and Cr levels in beans and maize of the study area were all within the respective limits of 0.3mg/kg, 0.2 mg/kg and 0.2mg/kg set by International Standard tolerable limits (2001). While mean Hg concentrations in beans and maize are above 0.05mg/kg tolerable limit set by international standard tolerable limits (2001). Although, the mean levels of Pb, Cr, Cd are within tolerable limit, however, attention should be focused on low level chronic exposure to these toxic metals, mercury inclusive.

The study shows that the spatial soil distribution of the toxic metals in each of the study areas is significantly ( $P < 0.01$ ) correlated with levels in the farm produce. The significant ( $P < 0.01$ ) negative correlation may indicates uptake/removal of the toxic metals from soil into the crops, while the instances of positive significant ( $P < 0.01$ ) correlation may indicates additional point sources of these metals, possible from the application of agrochemicals (fertilizers, pesticides and herbicides) and/or from non-point sources such as run off from environmental waste deposits of organic manure of animal origin, this is in line with the findings of Alhassan *et al* (2012) who established higher than reference value of some of these toxic metals in road site dust.

The established levels of these toxic metals in the farm produce relative to soil concentration could be alarming and may indicate their buildup to food chain. It may therefore serve as means of exposing human population to these toxic metals. This may be convinced by considering the outstanding characteristics of elements “non-destroyable and bioaccumulating”. The finding of

the study established alarming concentration of Hg in most of the sample analyzed, despite the non or less toxic effect due to exposure of low levels of metallic mercury speculated by some researchers (Davidson *et al.*, 2004), biotransformation of the inorganic Hg to organic Hg such as methyl mercury may occur, that potentiate the toxic effects of the mercury which includes lung damage, nausea, vomiting, diarrhea, increase in blood pressure or heart rate, skin rashes, eye irritation, permanently damage to brain, kidneys, and developing fetuses (Sabine and wendy, 2009 and Davidson *et al.*, 2004). Alhassan *et al* (2012) pointed out that human exposure to lead may result in many biological effects depending on the level and duration of exposure. Lead resembles calcium as divalent ion, it is therefore handled by the body as if it were  $\text{Ca}^{2+}$ , hence replacing the calcium in many biological systems and protein matrix with concomitant loss in biological activity. In addition other biochemical toxicity of lead could be associated to its affinity to cell membrane and mitochondria, where it interferes with oxidative phosphorylation, it also interferes with Na, K and Ca ATP-ase pumps (Schumacher *et al.*, 1991). Chronic lead exposure to blood level  $> 1.4 \mu\text{g/l}$  in children is believed to cause mental retardation, selective deficits in language, cognitive function, balance, behavioural and school performance. Across all age chronic exposure to lead is associated kidney damage and interstitial nephritis, affect sperm and ova, anaemia, CNS defects, peripheral neuropathies, and reduced birth weight. Susceptibility to lead toxicity is governed by genetic factors, as established in Caucasians population with 15% variant of gene code for aminolevulinic acid dehydrogenase, hence become more susceptible to lead exposure (Pirzada *et al.*, 2009).

Cadmium is a very toxic metal, found in all soils, including coal and mineral fertilizers. Cadmium has many uses, including batteries, pigments, metal coatings, and plastics. It is used extensively in electroplating (Peter, 2005). Cadmium and cadmium compounds are known

human carcinogens. Smokers get exposed to significantly higher cadmium levels than non-smokers. Severe damage to the lungs may occur through breathing high levels of cadmium common to cigarette smokers. High levels ingestion induces severe stomach irritation that induces vomiting and diarrhea, while Long-term exposure to lower levels leads to a buildup in the kidneys and possible kidney disease, lung damage, and fragile bones (Somers, 1983).

Chromium is found in rocks, animals, plants, and soil, its compounds bind to soil and are not likely to migrate to ground water but, they are very persistent in sediments in water. Chromium is used in metal alloys such as stainless steel; protective coatings on metal (electroplating); magnetic tapes; and pigments for paints, cement, paper, rubber, composition floor covering and other materials. Its soluble forms are used in wood preservatives (Lynch, 2005). Chromium (VI) compounds are toxins and known human carcinogens, whereas Chromium (III) is an essential nutrient. Breathing high levels can cause irritation to the lining of the nose; nose ulcers; runny nose; and breathing problems, such as asthma, cough, shortness of breath, or wheezing (Alabdulaaly *et al.*, 2009).

The level of Cadmium and Chromium in the soil, beans and maize samples from all the local government are not considered a health hazard. However, chronic exposure to such a low levels may lead to the toxicity and carcinogenic properties of chromium, softening of bones and kidney failure (Akintola, 2008). It is worth noting that most of the above mentioned sign and symptoms of heavy metals toxicity could be very common clinical cases in most hospitals nowadays and may not be unconnected to buildup of these toxic metals in foods and ultimately in human tissues.

Based on the findings of this study, mercury was found to possess a health risk index in four of the five Local Government Areas studied. The calculated health risk index of mercury in the

studied areas was greater than one, and which shows obvious risk of exposure by the population in the zones (Zhan *et al.*, 2012). According to the world health organization (2007), the majority of mercury in the environment results from coal-fired power station, residential heating systems and waste incinerators. Mercury is also released into the environment in the process of mining gold and other metals. WHO (2007), also established that consumption of contaminated fish, shellfish and marine mammals is the main source of methyl mercury exposure, especially for people who rely on predatory fish as their main source of protein. A recently conducted study by Zhang *et al.* (2010) found an elevated concentration of methyl mercury in rice (9.3ug/kg) throughout the Guizhou province of China. According to the Al-saleh and Shinwari, (2001), Mercury continues to be found in rice even in regions where mercury based pesticides are no longer in use. A study conducted in Saudi Arabia found that the concentration in rice was (3.18 ug/kg). In addition to past or ongoing fungicide use, mining and other industrial activities introduce another route of mercury exposure into the food chain through grain consumption in some regions of the world.

## **CONCLUSION**

Heavy metals contamination constitutes a major factor of environmental pollution. However, from this study, it appears that all the toxic metals of the samples (soils, beans and maize) were below the maximum tolerable levels and might appear harmless. However, it is imperative to note that chronic exposure to these even at low concentrations should be avoided.

## **REFERENCES**

- Alloway B.J (1995). Heavy metals in soils 2<sup>nd</sup> edition, Blackie academic and professional, Glasgow. pp. 500.
- Alhassan, A.J. Sule, M.S. Atiku, M.K., Wudil, A.M., Dangambo, M.A., Mashi, J.A. And Ibrahim, N.A. (2012). Study of Correlation between Heavy Metal Concentration, Street

Dust and Level of Traffic in Major Roads of Kano Metropolis, Nigeria.

- Al-Saleh I and Shinwari N. (2001). Report on the levels of cadmium, lead and mercury in imported rice grain samples. *Biol Trace Elements Health Research*, 83 (1):91-96.
- AOAC. (2005). Official Methods of Analysis, association of analytical Chemists. Washington DC, 15 edition. Pp 11-14.
- Alabdulaaly, A. I., & Khan, M. A. (2009). Heavy metals in cooler waters in Riyadh, Saudi Arabia. *Environmental Monitoring and Assessment*. 157, 23-28.
- Carpenter D.O. (2001). Effects of metals on the nervous system of Humans and metals. *International Journal of Occupational medicine and Environmental Health*. 14(3); 209-218.
- Davidson PW, Myers GJ, Weiss B (2004). "Mercury exposure and child development outcomes". *Pediatrics* **113**: 1023–9.
- Emsley, J (2011). Nature's Building Blocks. Oxford University Press. ISBN 9780199605637. Pp 34.
- Garrett N.E, Garrett R.J.B, Archdeacon J.W. (1992). Placental transmission of mercury to foetal rat. *Toxicological Application pharmacological*. 22; 649-654.
- Gray C.W, McLaren R.C, Roberts A.H.C. (2003). Atmospheric accessions of heavy metals to some New Zealand pastoral soils. *The science of the total Environment*. 305; 105-115.
- Holding BV (2004). Heavy Metals ([http: www. Lenntech. Com/heavy metals.htm](http://www.Lenntech.Com/heavy%20metals.htm)).
- International Standard Tolerable Limits (2001). Maximum levels of heavy metals in soil and foods. ISBN 9780123970268.
- Liu, W. H., Zhao, J. Z., Soderlund, L., and Liu, G. H. (2005). Impacts of sewage irrigation on heavy metal distribution and contamination in Beijing, China. *Environment international journal*. 31: 313-812.
- Lynch, E., and Braithwaite, R. (2005). A Review Of The Clinical And Toxicological Aspects of "Traditional" (Herbal) Medicines Adulterated With Heavy Metals. Sand Well and West Birmingham Nhs. In: UK Trust Regional Laboratory for Toxicology City Hospital Birmingham UK, pp. 769-778.
- Madyiwa S. (2006). Modeling lead and cadmium uptake by star Grass under irrigation with treated waste water. Published philosophiae. Doctor Thesis, University of Pretoria, South Africa. pp. 44-51
- Peter, B. W. (2005). Municipal Solid Waste Compositing: Potential Effect of Heavy Metals in

Municipal Solid Waste Composts on Plants and Environment. Boyce Thompson Institute for Plant Research at Cornell University: 1-5.

- Pirzada, H., Ahmad, S. S., Rashid, A., & Shah, T. (2009). Multivariate analysis of selected roadside plants (*Dalbergia sissoo* and *Cannabis sativa*) for lead pollution monitoring. *Pakistan Journal Bot*, 41, 1729-1736.
- Sabine Martin and Wendy Griswold. (2009). Centre for hazardous substances Research. Kansas University. U.S.
- Schumacher, M., Bosgue, M. A., Domingo, J. L., & Corbella, J. (1991). Dietary Intake of Lead and Calcium from Foods in Tarragona Province, Spain. *Bull Environ Contaminants and Toxicology*, 46, 320-328.
- Somers, E. (1983). The Toxic Potential of Trace Metals in foods. A Review. *Journal of Food Science*. 39, 215-217.
- Srivastava, S; Goyal, P (2010). Novel Biomaterials: Decontamination of Toxic Metals from Wastewater. Springer-Verlag. ISBN 978-3-642-11329-1.
- Stilwell, Sean (2000). "Power, Honour and Shame: The Ideology of Royal Slavery in the Sokoto Caliphate". *Africa: Journal of the International African Institute* (Edinburgh University Press) 70 (3): 394–421.
- USEPA, IRIS. (2002). United State, Environmental Protection Agency, Integrated Risk Information System. <http://www.epa.gov/iris/subs>. Accessed 25 February 2010.
- Vallero, DA; Letcher, TM (2013). Unravelling environmental disasters. Elsevier.
- Wang, S. T. (2005). Study on predicting effects of the soil polluted with heavy metals on vegetables and method of assessing quality of the soil in vegetable plot irrigated with a sewage-a case study on Baoding suburb. Agricultural University of Hebei, China. Pp 1-43.
- World Health Organization (WHO). (2014) Joint FAO/WHO Food Standards Programme. Codex Committee on Food Additives and Contaminants. Thirty-second session. Draft Maximum levels of lead. Prepared by Denmark pp 389.
- World Health Organization (WHO), (2007). *Exposure to mercury: A major public health concern*. Geneva, Switzerland, World Health Organization (<http://www.who.int/phe/news/Mercury-flyer.pdf>, accessed 10 May 2010).
- Zhan-Jun Xue, Shu-Qing Liu, Yan-Ling Liu and Yong-Lu Yan (2012). Health risk assessment of heavy metals for edible parts of vegetables grown in sewage-irrigated soils in suburbs of Baoding city, china. *Environmental Monitoring Assess* 184:3503-3513.