

Original Research Article

Dried Pig Faeces: Impacts on Growth, Haematology and Histology of *Clarias gariepinus*

ABSTRACT

Aims: Improper utilization and disposal of pig faeces in pig farms have been a great concern due to their hazards and risks they constitute to the environment. This situation calls for conversion of animal waste to wealth through effective utilization of pig faeces for fish feed formulation. This project was therefore designed to provide information on safe level of Dried Pig Faeces (DPF) on growth, histology and haematology of *Clarias gariepinus*.

Study design: A total number of one hundred and fifty samples of *C. gariepinus* were collected for the feeding trial experiment. The fish meal was partially and completely replaced by different graded inclusion levels of DPF at 0%, 25%, 50%, 75% and 100% respectively and was coded T1, T2, T3, T4 and T5 respectively.

Materials and Methods: Growth was determined by calculating growth parameters according to standard formulae. Histology of vital organs and haematology of the fish samples were determined according to standard methods. Data obtained were subjected to statistical analysis using ANOVA and Duncan multiple range test was used to separate the means.

Results: The Mean Weight Gain (MWG) revealed a highest value in T3 (69.40g) which was significant among the treatments, while 100% inclusion of DPF exhibited lowest MWG (21.07%). The Packed Cell Volume, Haemoglobin and Red Blood Cell values obtained decreased with increase in DPF inclusions, while an increase in these parameters were observed in T4 (26.50%). White Blood Cell count was significant in T3 ($16,150 \pm 2.07g/l$) and decreased in T4 ($15,800 \pm 1.01g/l$) and T5 ($12,650 \pm 0.85g/l$) respectively. Histological results confirmed the absence of lesions in the heart, gills and the intestine of fish samples fed DPF in all the treatments. Lesions was severed in the liver in T4 and T5.

Conclusions: The findings of this study revealed that 50% inclusion of DPF is the safe level for growth and survival of *C. gariepinus*. Since the histological results also revealed no lesions in all the vital organs, except the liver that had severe lesions, DPF can be recommended as an alternative feed ingredient that could substitute fish meal in the fish diet.

Comment [U1]: Abstract too long, summarize it, it should not be more than 300 wards.

Comment [U2]: Summaries this area

Keywords: *Clarias gariepinus*, Dried Pig Faeces, Haematology, Histology, Nutrient Utilization.

Comment [U3]: It should be between four to eight words

1. INTRODUCTION

Integrated fish farming has been practiced globally to maximize usage of land and space acquired for agricultural purposes in order to reduce cost of aquaculture production. Fish are sometimes raised with pigs by constructing pig's pen on the pond embankment and the **faecesfeces** are allowed to drain directly into the ponds. This integrated system has been universally practiced for pork and manure production. The excreta from the pens of pigs fertilizes the pond and enhances massive production of zooplankton and phytoplankton which serves as natural food for the fish. Fish feed directly on digestive food materials that are not completely digested in the digestive system of pigs. This farming system has been successfully practiced with fish farming with the aim of using the **faecesfeces** as an alternative source of feed to the fish. In part of Asia where improper utilization and disposal of pig **faecesfeces** in pig farms have been a great concern due to their hazards to the environments, research efforts are on viable integrated farming approach that involves livestock such as poultry, swine, duck and crops or vegetables [1,2,3,4]. In Teagasc, Ireland, economic analysis showed that land-spreading of pig manure for its fertilizer value is the most economic use for pig manure than other technological feasible alternative uses of pig manure [5]. In Nigeria, pigs are reared mainly to provide pork and manure for improving soil fertility, while studies on integrated farming involving pig and fish **has have** received little attention. The effective utilization of this obvious wastes from pig production will further enhance fish production. This study was therefore designed to provide information on growth, nutrient utilization, haematobiochemical profile and tissue pathology of *Clarias gariepinus* fed diet with graded inclusion levels of DPF.

2. MATERIALS AND METHODS

2.1 Study Site

The feeding trial experiment was conducted at the Teaching and Research Farm, Osun State University, Ejigbo Campus.

2.2 Collection of Fish Samples

One hundred and fifty samples of *C. gariepinus* of mean weight 23.51g were procured from Turning Point Fish Farm, Ibadan. The fish samples were conditioned and transported from the farm to the laboratory inside fifty (50) liters container. Fish samples were acclimatized for fourteen (14) days before the commencement of the experiment.

2.3 Collection of Pig Manure

Fresh pig **faeces** were collected from pigs' pen at the Piggery Unit, Department of Animal Science, Osun State University, Ejigbo Campus. In order to prevent growth of molds which could be injurious to fish, pig **faeces** were distributed inside stainless bowls and sun dried for three weeks. After drying, DPF were stored inside a zip lock nylon at room temperature.

2.4 Experimental Procedure

Fish samples were stocked at 30 fish per treatment and replicated thrice inside experimental bowls containing fifty liters of water. Fish meal in the fish diets were replaced with 0%, 25%, and 50%, 75% and 100% DPF and were coded T1, T2, T3, T4 and T5 respectively. The feeding trial experiment was conducted for a period of twelve weeks. Fish samples were fed with formulated diets at 5% body weight. Fish samples were weighed fortnightly and new weight was used to adjust the quantity of feed fed to the fish samples.

2.5 Formulation of Diet with Varying Levels of DPF

Fish diets (Table 1) were formulated by grinding DPF with blender and mixing the grounded DPF with other feed ingredients like fish meal, maize, soya bean meal, groundnut cake, wheat offal, cassava powder, bone meal, salt and fish premix. Mixed DPF and other feed ingredients were moistened with warm water and pelletized with locally fabricated pelleting machine. The wet pellets were sundried and stored inside zip lock nylon at room temperature.

Table1. Formulated Diet with Varying Levels of DPF

Ingredients	(T1)	(T2)	(T3)	(T4)	(T5)
Maize	29.56	29.56	29.56	29.56	29.56
FM	46.10	34.58	23.05	11.53	-

DPD	-	11.52	23.05	34.57	46.10
Wheat offal	4.89	4.89	4.89	4.89	4.89
Cassava powder	4.03	4.03	4.03	4.03	4.03
SBC	7.72	7.72	7.72	7.72	7.72
GNC	5.85	5.85	5.85	5.85	5.85
BM	1.00	1.00	1.00	1.00	1.00
Salt	0.25	0.25	0.25	0.25	0.25
Fish premix	0.60	0.60	0.60	0.60	0.60
TOTAL	100	100	100	100	100

KEYS:

- FM = Fish meal
- DPD = Dried pig droppings
- SBC = Soya bean cake
- GNC = Groundnut cake
- BM = Bone meal

2.6 Collection and Determination of Haematological Profile

After the feeding trial experiment, blood samples of the control and test fish were collected into heparinised bottles by inserting 2 ml syringe needle into the posterior caudal vein. Packed Cell Volume (PCV) and haemoglobin (Hb) concentration were analysed immediately after collection. Red blood cells (RBC) and white blood cells (WBC) were counted with Neubauer's improved haemocytometer using Hyem's and Turk's solution as a diluting fluid respectively. Mean Corpuscular Haemoglobin (MCH) and Mean Cell Volume (MCV) were calculated respectively using standard procedure [6,7].

2.7 Histological Analysis of Vital Organs

Fish samples were anaesthetized with Benzocaine. Vital organs such as gills, liver, intestine, heart and kidneys were removed after dissection of the fish samples. Histological analysis and examinations of the organs were determined according to standard procedures [8]. The organs were fixed inside 4% formaldehyde (Lach-Ner, Czech Republic). Tissues of the organs were stained with Hematoxylin and

Eosin. Microphotographs of the slides of the tissues of the organs were taken using a Leica DM LS microscope (Leica, Austria) with a light microscope (U-MDOB, Olympus optical Co. Ltd., Japan).

2.8 Analysis of Proximate Composition

Proximate composition (crude protein, crude fat, crude fibre, moisture contents and ash) of DPF, fish carcass and fish diets (Tables 2 and 3) were determined according to the procedures described by [9].

2.9 Microbial Load Analysis

The microbial load of DPF and fish diets were carried out according to procedures described by [10,11].

2.10 Statistical Analysis

The data collected were analyzed using Statistical Package for Social Sciences (SPSS), Version 11 (2001) and Statistical Analysis Software (SAS), Version 8 (2001). Duncan's Multiple Range Test was used to compare the differences among the means. The significant level was set at 5%.

3. RESULTS

Analysis of proximate composition of DPF as shown in Table 2 revealed that crude protein, crude fibre, fat and ash had 22.16%, 7.45%, 1.70% and 4.24% respectively. Similarly, the proximate composition of fish carcass as shown in Table 3 revealed 27.75% (crude protein), 0.01% (crude fibre), 18.85% (fat) and 0.12% (ash).

Table 2: Proximate Composition Of Dpf

S/N	Parameters	Values (%)
1	Crude Protein	22.16
2	Crude Fibre	7.45
3	Fat	1.70
4	Ash	4.24

Table 3: Proximate Composition of Fish Carcass

S/N	Parameters	Values (%)
1	Crude Protein	27.75
2	Crude Fibre	0.01
3	Fat	18.85
4	Ash	0.12
5	Moisture content	28.56
6	Carbohydrate	53.28

The results obtained after the microbial count analysis of DPF is shown in Table 4. The values recorded for TBC, TCC and TFC are 2.2×10^5 cfu/g, 1.6×10^5 cfu/g and 0.6×10^5 cfu/g respectively. The microbial count of feed formulated with DPF is presented in Table 5. TBC, TCC and TFC showed that the counts increased across the table as the inclusion increased. TBC showed a significant count $p > 0.05$ in T5 among the treatments. Similarly, TCC was observed to have the highest significant value $p > 0.05$ in T5, whereas the lowest count (0.34×10^5 cfu/g) was recorded in 50% (T1) inclusion. TFC in T5 was significant different $p > 0.05$ among the treatments.

Table 4: Microbial Counts of DPF

	Parameters (%)	Values
1	Total Bacterial Count (TBC)	2.2×10^5 cfu/g

2	Total Coliform Count (TCC)	1.6 x10 ⁵ cfu/g
3	Total Fungi Count (TFC)	0.6 x10 ⁵ cfu/g

Table 5: Microbial Load of Feeds

Parameter	T1	T2	T3	T4	T5
Total Bacterial Count.(TBC)	1.43 ^d x10 ⁵ cfu/g	2.22 ^d x10 ⁵ cfu/g	5.34 ^c x10 ⁵ cfu/g	14.1 ^b x10 ⁵ cfu/g	15.9 ^a x10 ⁵ cfu/g
Total Coliform Count.(TCC)	0.34 ^c x10 ⁵ cfu/g	0.45 ^b x10 ⁵ cfu/g	0.41 ^b x10 ⁵ cfu/g	1.73 ^a x10 ⁵ cfu/g	1.82 ^a x10 ⁵ cfu/g
Total fungi count(TFC)	0.10 ^b x10 ⁵ cfu/g	0.10 ^b x10 ⁵ cfu/g	0.11 ^b x10 ⁵ cfu/g	0.43 ^a x10 ⁵ cfu/g	0.49 ^a x10 ⁵ cfu/g

Mean with the same superscript are not significantly different (P>0.05)

Haematological parameters of the fish samples fed DPF are presented in Tables 6a and b. The PCV, Hb and RBC values obtained in *C. gariepinus* fed with compounded feed of DPF at varying concentrations decreased with increase in concentration while an increase in these parameters were observed in T4. White Blood Cell count increased in T3(16,150±2.07) and decreased in T4(15,800±1.01) and T5(12,650±0.85). Platelet concentration in treatment T2(210,000±0.51) first increased and then followed a downward trend pattern up to T5(159,000±0.07) when an increase was experienced. A non-significant increase from T1 (control, 65±0.62) was recorded for lymphocyte across the varying concentrations the test fishes were exposed to. For Neutrophils, a non-significant decrease was observed in treatment T2(26.5±0.09) and T3(23.5±1.12) in relation to the control. A steady decrease in Monocyte value was recorded from T1(control, 5.0±0.03) to T4(2.5±0.06) while a marginal increase was observed for T5(3.0±0.02). A significant increase was recorded for Eosinophils in treatment T3(4.5±0.02) and T4(4.5±0.04) which decreased significantly for T5(2.0±0.01). For MCV and MCH, a significant increase was recorded for the test fishes in T5 while no significant increase was observed for MCHC across the treatments.

Table 6a: Haematological parameters of the test and control fish fed with DPF

Treatment	PCV(%)	Hb (g/dl)	RBC (10 ⁶ /L)	WBC (10 ⁹ /L)	Platelet (10 ⁷ /L)	Lymphocyte (10 ³ /mm ³)
T1	36.24±0.50 ^a	11.7±0.07 ^a	3.27±0.04 ^a	14,000±1.01 ^{ab}	198,000±0.23 ^a	65±0.62 ^{ab}

T2	28.50±0.23 ^b	9.5 ±0.02 ^{ab}	3.26±0.10 ^a	16,200±0.90 ^a	210,000±0.51 ^a	67±0.91 ^a
T3	22.50±1.05 ^c	7.2 ±2.01 ^b	2.18±0.09 ^b	16,150±2.07 ^a	186,000±0.49 ^{ab}	69±0.98 ^a
T4	26.50±0.87 ^b	8.3 ±0.65 ^b	2.76±0.42 ^{ab}	15,800±1.01 ^a	112,500±0.93 ^d	68±0.52 ^a
T5	21.00±0.41 ^c	6.7 ±0.12 ^c	1.42±0.04 ^c	12,650±0.85 ^c	159,000±0.07 ^c	70±0.61 ^a

Mean with the same superscript are not significantly different (P>0.05)

Table 6b: Haematological parameters of the test and control fish fed with DPF

Treatment	Neutrophil (%)	Monocyte (%)	Eosinophil (%)	Basophil (%)	MCV (fl)	MCH (pg)	MCHC (g/l)
T1	28.0±0.10 ^a	5.0±0.03 ^a	2.0±0.01 ^b	0±0.02 ^b	110.08 ^{ab}	35.78 ^{ab}	32.28 ^a
T2	26.5±0.09 ^a	4.0±0.01 ^a	2.0±0.06 ^b	0.5±0.01 ^a	87.42 ^a	29.14 ^a	33.33 ^a
T3	23.5±1.12 ^c	3.0±0.01 ^{ab}	4.5±0.02 ^a	0±0.00 ^b	103.21 ^{ab}	33.03 ^{ab}	32.00 ^a
T4	25.5±1.05 ^{ab}	2.5±0.06 ^{bc}	4.5±0.04 ^a	0.5±0.01 ^a	96.01 ^a	30.07 ^a	31.32 ^a
T5	25.0±0.07 ^{ab}	3.0±0.02 ^{ab}	2.0±0.01 ^b	0±0.00 ^b	147.89 ^c	47.18 ^c	31.90 ^a

Mean with the same superscript are not significantly different (P>0.05)

The growth parameters of fish samples fed DPF are shown in Table 7. The MWG revealed a highest value in T3 (69.40g) which was significant among the treatments, while 100% inclusion of DPF exhibited lowest MWG (21.07%). In the same trend, SGR of T3 varied significantly p>0.05 compared to other treatments. FCR revealed a significant value p>0.05 in T3 which showed the best conversion rate (2.71). PER showed no significant variation among the treatments fed DPF. The survival rate of all the treatments showed no significant variation during the period of the experiment.

TABLE 7: Growth Parameters of *Clarias gariepinus* fed with Varying Levels of DPF

Treatment	T1	T2	T3	T4	T5
MIW(g)	23.83 ^a ±0.43	22.73 ^a ±0.55	23.30 ^a ±0.87	23.57 ^a ±0.2	24.13 ^a ±0.27

MFW(g)	86.67 ^a ±6.34	84.15 ^a ±7.19	92.70 ^a ±17.9	68.93 ^{ab} ±1.4	45 ^b ±4.65
MWG(g)	62.83 ^a ±5.9	61.42 ^a ±7.72	69.40 ^a ±7.1	45.37 ^{ab} ±1.6	21.07 ^b ±4.67
SGR	1.55 ^{ab} ±0.70	1.57 ^{ab} ±0.13	1.62 ^a ±0.18	1.29 ^b ±0.03	0.74 ^c ±0.87
FCR	2.88 ^{bc} ±0.01	2.88 ^{bc} ±0.03	2.71 ^c ±0.02	3.21 ^a ±0.03	3.19 ^{ab} ±0.19
PER	0.12 ^a ±0.002	0.13 ^a ±0.004	0.13 ^a ±0.005	0.14 ^a ±0.00	0.13 ^a ±0.008

Means with same superscript were not significantly different (P>0.05)

The histological examinations of the heart muscles, gills, intestines, livers and kidneys were examined in this experiment. The results revealed that the heart muscles, gills and intestines of all the treatments had no observable lesions. Histological examinations of the kidneys are shown in Figures 1a, b, c, d and e. The results revealed that T1, T2 had no observable lesions, though T3 had no observable lesions, there were vacuolar degeneration of tubular epithelium observed in the kidney. Conspicuous vacuoles in tubular epithelial cells of the kidney were observed in T4 and T5. Cross sections of Figures 2a and c revealed that there were diffuse vacuolation of hepatocytes in T1 and T3, while Figure 2b revealed that T2 had vasculitis in portal areas of the liver in addition to diffuse vacuolation of hepatocytes. Figures 2d and e had severe diffuse vacuolation of hepatocytes in the liver.




Fig.1a. Cross section of kidney
of fish fed 0% inclusion of DPF




Fig. 1b. Cross section of kidney fish fed 25%
inclusion of DPF




Fig. 1c. Cross section of kidney of fish fed 50% inclusions of DPF.




Fig. 1d. Cross section of kidney of fish fed 75% inclusions of DPF.




Fig. 1e. Cross section of kidney of fish fed 100% inclusions of DPF




Fig. 2a. Cross section of liver of fish fed 0% Inclusion of DPF.




Fig. 2b. Cross section of liver of fish fed 25% inclusions of DPF.




Fig. 2c. Cross section of liver of fish fed 50% Inclusions of DPF.




Fig. 2d. Cross section of liver of fish fed 75% inclusions of DPF.




Fig. 2e. Cross section of liver of fish fed 100% inclusions of DPF

4. DISCUSSION

This study evaluates the growth, nutrient utilization, haematobiochemical profile and tissue pathology of *C. gariepinus* fed diet with graded inclusion levels of DPF. In this study, the optimal growth was achieved at 50% inclusion level of DPF while 75% and 100% inclusions caused poor growth. In a similar experiment, [12] recorded a different value of 25% inclusion of dried poultry droppings fed *C. gariepinus*, while 100% inclusion caused poor growth as recorded in this experiment. The poor growth at the highest inclusions could be attributed with the high levels of microbe counts recorded in the feed formulated with DPF. Histological results confirmed the absence of lesions in the heart, gills and the intestine of fish samples fed DPF. Hepatic lesions in the liver were severe due to detoxification of high loads of DPF. A readily available and fast means of assessing clinical and nutritional health status of animals on feeding trials may be the use of blood analysis, because ingestion of dietary components have measurable effects on blood composition [13,14] and may be considered as appropriate measure of long term nutritional status [15]. The decline observed in PCV and RBC of the fish exposed to varying concentration of DPF may be due to the harmful effects of high dietary content of toxicant, possibly ammonia. This low level is not too different from 30% PCV, Hb content of 10.30g/dl and RBC of 7.10 x106/ml as reported by [16]. The reduction in the values of haemoglobin when compared with the control could be as a result of the destruction or inhibition of erythrocyte production. This is in consonance with

Comment [U4]: Explain how your result is connected to your citation. Use a connective sentences to connect your result to other finding. Connective statement like; in line with, agree with etc.

the report of [17]. The increase in WBC count in T2 and T3 could be attributed to the increase in leucocytes synthesis as a defence mechanism against possible microbes in the feed.

Comment [U5]: This area is Ok. Do similar example above.

Comment [U6]: Are you referring to this statement as being report by [17]? If no, please state his/her finding. Such as; Who reported that..... Or revealed that.....etc.

5. CONCLUSIONS

It was apparent in the study that the use of some feed ingredients introduce feed microbes into fish feeds. In order to ensure food safety, there is urgent need for autoclaving alternative feed ingredients before compounding them as fish feeds for fish use.

REFERENCES

1. Korikantimath VS, Manjunath BL. Integrated farming systems for sustainability in Agricultural production. Proceedings of National Symposium on “New Paradigms in Agronomic Research”. Indian Society of Agronomy. Navsari Agriculture University Gujarat. 2008;279-281.
2. Shoko AP, Getabu A, Mwayuli G, Mgaya YD. Growth performance, yields and economic benefits of Nile tilapia *Oreochromis niloticus* and Kales Brassica oleracea cultured under vegetable-fish culture integration. Tanzania Journal of Science. 2011; 37: 37-48.
3. Ogello EO, Obiero K, Munguti JM. Lake Victoria and the Common Property Debate: Is the Tragedy of the Commons a Threat to its Future? Lakes, reservoirs and ponds. 2013; 7(2): 101-126.
4. Sahoo UK, Singh SL. Integrated Fish-Pig and Fish-Poultry Farming in East Kalcho, Saiha District of Mizoram, North-East India: An Economic Analysis. International Journal of Agriculture and Forestry. 2015; 5(5): 281-286.
5. The Pig Site. Latest Cost of Pig Production Report. 5m Publishing, Benchmark House, 8 Smithy Wood Drive, Sheffield, S35 1QN, England. 2014.
6. Dacie JV, Lewis SM. Practical haematology (6th Ed). ELBS and Churchill, Livingston. 1984
7. Joshi PK, Bose M, Harish D. Haematological changes in the blood of *Clarias batrachus* exposed to mercuric chloride. Eco toxicological Environmental Monitoring. 2002; 12(2):119-122.

8. Delahoy MJ, Wodnik B, McAliley L, Penakalapati G, Swarthout J, Freeman MC, Levy k. Pathogens transmitted in animal feces in low- and middle-income countries. *International Journal of Hygiene and Environmental Health*. 2018; 221: 661–676.
9. AOAC. International official methods of analysis (18th Ed). AOAC International, Gaithersburg, MD. 2005.
10. Cruickshank R, Duguid JP, Marmion BR, Swain RHA. (1975): *Medical Microbiology*, 12th Ed., Living stone, London, New York. 1975; 812-825.
11. Ayandiran TA, Ayandele AA, Dahunsi SO, Ajala OO. Microbial Assessment and Prevalence of Antibiotic Resistance in Polluted Oluwa River, Nigeria. *The Egyptian Journal of Aquatic Research*. 2014; 40(3):291-299.
12. Ajiboye AO, Owoseni AA, and Ajani F. Effects of Varying Levels of Dried Poultry Droppings on Haematology, Microbial Properties and Nutrient Utilization of *Clarias gariepinus*. *Not. Sci. Biol*. 2017; 9(4):460-465.
13. Church JP, Judd JT, Young CW, Kebay JL, Kim WW. Relationships among diet constituents and specific serum clinical components of subjects eating self-selected diets. *Afric. J. Clin. Nutr*. 1984; 40:1338-1344.
14. Maxwell MH, Robertson GW, Spences and McCongrouodala CC. Composition of haematological values in restricted and ad libitum feed domesticated fowls. *RBC Characteristics*. *British Poultry*. 1990; 60:14741484.
15. Olabanji RO, Farinu GO, Akinlade JA, Ojebiyi OO. Growth performance and haematological characteristics of weaner rabbits fed different levels of wild sunflower (*Tithonia diversifolia* Hems L A. Gray) leaf blood meal mixture. *Proc. of 32nd Animal Conf. of Nig. Soc. for Anim. Prod*. 2007; 207-209.
16. Bawala TO, Akpan U, Ogunnowo AO, Fasae OA, Sogunle OM. Influence of Mg supplementation on the haematological indices of young West African Dwarf (WAD) goats. *Proc. of the 32nd Annual Conf. of NSAP*. 2007; 76-78.
17. Agbon O, Omoniyi II, Teko AA. “Acute toxicity of tobacco (*Nicotiana tobaccum*) leaf dust on *Oreochromis niloticus* and haematological changes resulting from sublethal exposure,” *Journal of Aquatic Science*. 2002; 17(1): 5–8.

UNDER PEER REVIEW

UNDER PEER REVIEW