

Original Research Article

Comment [m1]: It is very well-marked that this study is acceptable with minor revision and useful for publish in this journal.

NOTE: There is no discussion in the article. Please, Discuss with literature data.

In addition to, please add some pape and you can use them in manuscript.

9. . Sevindik M, Akgul H, Pehlivan M, Selamoglu Z. Determination of therapeutic potential of *Mentha longifolia* ssp. *longifolia*. *Fresen Environ Bull*, 2017; 26: 4757-4763.

Phenolic and mineral characteristics of seed coats and kernels

ABSTRACT

Aims: The objective of the present work is the investigation of the physico-chemical characteristics of seed coats and kernels from 24 species with medicinal and food applications.

Methodology: Seeds from 24 species (2 herbs, 11 vines and 11 trees), belonging to 13 families, were sampled in Raipur (India) in 2017. The collected seeds were dried and weighted, after which seed coats were manually peeled and weighed separately. Phenolic and mineral contents in the seed coats and kernels were analyzed by spectrophotometric and X-ray fluorescence (XRF) techniques, respectively.

Results: The seed coat fraction represented from 12% to 95% of the seed mass, depending on the species. The concentrations of total polyphenols, flavonoids and minerals in the seed coats varied from 1800 to 32300 mg/kg, from 1200 to 26900 mg/kg, and from 5876 to 36499 mg/kg, respectively. Where as in the seed kernels, TPh, Fla and minerals ranged from 780 to 31760 mg/kg, from 300 to 31300 mg/kg, and from 12595 to 40810 mg/kg, respectively. Elements: P, S, K, Mg, Ca and Fe were found to be the main macro- and micro-elements. Seed coats from Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae families featured the highest total polyphenol contents, and those from Lauraceae and Rutaceae families showed the highest flavonoid concentrations. The highest total mineral contents corresponded to seed coats from Lauraceae, Rutaceae and Euphorbiaceae families.

23 **Conclusion:** Indian-laurel and curry tree stand out as promising phytochemical and nutrient
24 sources.

25 **Keywords:** Seed coat, Seed Kernel, Total polyphenol, Flavonoid, Mineral.

26 INTRODUCTION

27 The seed coat protects the internal parts of the seed from fungi, bacteria and insects, and prevents
28 water loss. It is composed of cellulose, fiber, polyphenols, starch, wax, etc. Its outer layer, called
29 testa, is generally hard and thick, while its inner layer, known as the tegmen, is softer [1].
30 Enrichment of various compounds (viz. minerals, cellulose, fiber, polyphenols, starch, wax, etc.)
31 in seed coats has been reported in the literature [2-7]. Among these phytochemicals: polyphenols
32 have become the subject of increasing research efforts owing to their potential beneficial effects
33 on human health [8,9].

34 Among the plants found in Raipur area, Black Siris (*Albizia odoratissima* (L.f.) Benth.), Malabar
35 spinach (*Basella rubra* L., syn. *Basella alba* L.), wax gourd (*Benincasa hispida* (Thunb.) Cogn.),
36 squash (*Cucurbita maxima* (Duchesne) Duchesne ex Poir.), watermelon (*Citrullus lanatus*
37 (Thunb.) var. *lanatus*), Persian melon (*Cucumis melo* var. *cantalupo* Ser.), Liane Cacorne
38 (*Entada gigas* (L.) Fawc. & Rendle), tree cotton (*Gossypium arboreum* L.), physic nut (*Jatropha*
39 *curcas* L.), Persian walnut (*Juglans regia* L.), hyacinth bean (*Lablab purpureus* (L.) Sweet),
40 calabash (*Lagenaria siceraria* Standl.), Chinese-okra (*Luffa acutangula* Roxb.), sponge gourd
41 (*Luffa aegyptiaca* Mill.), Indian-laurel (*Litsea glutinosa* (Lour.) C.B.Rob.), Indian-lilac (*Melia*
42 *azadirachta* L., syn. *Azadirachta indica* A.Juss.), bitter melon (*Momordica charantia* L.), curry
43 tree (*Murraya koenigii* Spreng.), emblic (*Phyllanthus emblica* L.), East Indian kino (*Pterocarpus*
44 *marsupium* Roxb.), Indian sandalwood (*Santalum album* L.), Ceylon-oak (*Schleichera oleosa*

(Lour.) Oken), clearing-nut-tree (*Strychnos potatorum* L.f.), and Indian tuliptree (*Thespesia populnea* Sol. ex Corrêa) are widely used as medicine, food and fodder for animals [10-23].

Accumulation of the nutrients and polyphenols in some seed coats were reported [6, 24-28]. In this work, the physical and chemical characteristics of the seed coats and kernels from these 24 species (2 herbs, 11 vines and 11 trees) are analyzed, with emphasis on their polyphenol contents.

METHODS AND MATERIALS

Sample collection and handling

Seeds from aforementioned twenty-four species were collected in Raipur area (21.25°N 81.63°E), Chhattisgarh, India, during their maturation period in 2017. The seeds were manually separated and sundried in a glass room for one week, after which they were further dried in a hot air oven at 50 °C for 24 h. The mass of the seeds was measured using an AG245 (Mettler Toledo, Columbus, OH, USA) electronic balance. The seed coats were then carefully peeled with the aid of a surgical blade and their mass was measured. The separated seed coats and kernels were crushed into fine powder, and particles of mesh size ≤ 0.1 mm were sieved out. The samples were preserved in a deep freezer at -4 °C till the analyses were conducted.

Analyses

Sigma-Aldrich AR grade reagents were used for analysis of polyphenols. 0.1 g of powdered seed coat were extracted with an acetone:water mixture (7:3, v/v), as recommended by Bertaud et al. [29]. An appropriate fraction was allowed to react with Folin-Ciocalteu reagent for color development, and absorbance was measured at $\lambda=740$ nm with a UV-1800 (Shimadzu, Kyoto, Japan) UV-Vis spectrophotometer [30]. Three replicates for each solvent extract were performed

to determine the total phenolic content (TPh), which was expressed in terms of tannic acid equivalents by using a standard calibration curve. For flavonoid (Fla) analysis, a fraction of the extract was reacted with an aluminum chloride solution to develop a yellow colored complex, measuring the absorbance at $\lambda=410$ nm [31]. The Fla concentration was determined with the aid of standard quercetin calibration curve and indicated in terms of quercetin equivalents. Three replicates for each solvent extract were performed, and results are presented as average values across the three replicates.

A III Tracer-SD portable XRF (Bruker, Billerica, MA, USA) spectrophotometer was used for the quantification of 15 elements: K, Rb, Mg, Ca, Sr, Al, P, S, Cl, Ti, Mn, Fe, Cu, Zn and Pb. Standard brown and white cowpea (*Vigna unguiculata* (L.) Walp.) seeds were used as reference material to standardize the analyte concentration [32].

Statistical analyses

Cluster analysis was used to assess similarities in the micro- and macro-elements content in the seed coats. IBM (Armonk, NY, USA) SPSS v.25 software was used.

RESULTS AND DISCUSSION

Physical characteristics

The physical characteristics of the seeds and seed coats under study (show in **Fig. 1**) are summarized in **Table 1**. Large differences in seed mass were found, with average values ranging from 25 to 23623 mg per seed, with the highest weights for *Entada gigas* (23623 mg), followed by *Juglans regia* (12200 mg). The seed coat mass represented from 12 to 95% of the total seed weight.

Fig. 1. Seed samples from: (1) *P. embilica*, (2) *P. marsupium*, (3) *L. aegyptiaca*, (4) *B. hispida*, (5) *C. lanatus*, (6) *C. melo*, (7) *C. maxima*, (8) *L. acutangula*, (9) *L. siceraria*, (10) *M. charantia*, (11) *L.*

Deleted: *Phyllanthus*

Deleted: *Pterocarpus*

Deleted: *Luffa*

Deleted: *Benincasa*

Deleted: *Citrullus*

Deleted: *Cucumis*

Deleted: *Cucurbita*


Deleted: *Luffa*

Deleted: *Lagenaria*

Deleted: *Momordica*

Deleted: *Litsea*

101 | *glutinosa*, (12) *G. arboreum*, (13) *T. populnea*, (14) *S. album*, (15) *M. koenigii*, (16) *S. oleosa*, (17) *M.*
 102 | *azadirachta*, (18) *S. potatorum*, (19) *A. odoratissima*, (20) *B. rubra*, (21) *L. purpureus*, (22) *J. curcas*,
 103 | (23) *J. regia*, and (24) *E. gigas*.
 104



Deleted: *Gossypium*

Deleted: *Thespesia*

Deleted: *Santalum*

Deleted: *Murraya*

Deleted: *Schleichera*

Deleted: *Melia*

Deleted: *Strychnos*

Deleted: *Ibizzia*

Deleted: *Basella*

Deleted: *Lablab*

Deleted: *Jatropha*

Deleted: *Juglans*

Deleted: *Entada*

105

Table 1. Physico-chemical characteristics of seeds and seed coats. Total phenolic contents and flavonoid contents correspond to the seed coat and kernel samples.

Species	Family	Type	Seed mass (mg)	Color	Seed coat (%)	Seed coat, mg/kg		Kernel, mg/kg	
						TPh	Fla	TPh	Fla
<i>B. rubra</i>	Basellaceae	V	38	BrB	47±2	11400	10500	3457	1650
<i>C. maxima</i>	Cucurbitaceae	V	132	YeW	18±1	3100	1900	4931	1100
<i>L. siceraria</i>	Cucurbitaceae	V	216	WhBr	42±2	14400	8900	1956	1400
<i>C. lanatus</i>	Cucurbitaceae	V	38	ReBr	49±2	18500	13100	2278	1280
<i>L. aegyptiaca</i>	Cucurbitaceae	V	105	B	43±2	3100	2500	780	620

Deleted: *Basella*

Deleted: *Cucurbita*

Deleted: *Lagenaria*

Deleted: *Citrullus*

Deleted: *Luffa*

<i>C. melo</i>	Cucurbitaceae	V	25	LY	28±1	2900	2100	965	300	Deleted: <i>Cucumis</i>
<i>L. acutangula</i>	Cucurbitaceae	V	122	B	47±2	8300	7200	2144	1380	Deleted: <i>Luffa</i>
<i>B. hispida</i>	Cucurbitaceae	V	64	YeW	47±	2900	2600	4074	2280	Deleted: <i>Benincasa</i>
<i>M. charantia</i>	Cucurbitaceae	V	189	YeBr	35±1	30847	1700	1769	1180	Deleted: <i>Momordica</i>
<i>J. curcas</i>	Euphorbiaceae	H	758	Br	47±2	14700	4000	1501	4260	Deleted: <i>Jatropha</i>
<i>L. purpureus</i>	Fabaceae	V	293	DBr	34±1	22000	3700	1260	2550	Deleted: <i>Lablab</i>
<i>A. odoratissima</i>	Fabaceae	T	159	LBr	42±	27000	4100	2492	4300	Deleted: <i>Albizia</i>
<i>E. gigas</i>	Fabaceae	V	23623	DBr	40±1	26900	3900	18840	2650	Deleted: <i>Entada</i>
<i>P. marsupium</i>	Fabaceae	T	933	LY	93±3	25800	3800	31760	5800	Deleted: <i>Pterocarpus</i>
<i>J. regia</i>	Juglandaceae	T	12200	PY	32±1	9600	1900	1045	1520	Deleted: <i>Juglans</i>
<i>L. glutinosa</i>	Lauraceae	T	248	DBr	43±2	29200	26900	4931	3880	Deleted: <i>Litsea</i>
<i>S. potatorum</i>	Loganiaceae	T	280	B	24±1	26000	15000	2707	1640	Deleted: <i>Strychnos</i>
<i>G. arboreum</i>	Malvaceae	H	82	Br	48±2	4000	3500	7263	7160	Deleted: <i>Gossypium</i>
<i>T. populnea</i>	Malvaceae	T	162	LBr	47±2	16800	8000	15839	12020	Deleted: <i>Thespesia</i>
<i>M. azadirachta</i>	Meliaceae	T	972	DBr	65±2	1800	1200	1822	1300	Deleted: <i>Melia</i>
<i>P. embilica</i>	Phyllanthaceae	T	920	PW	95±3	27000	3500	4476	3750	Deleted: <i>Phyllanthus</i>
<i>M. koenigii</i>	Rutaceae	T	155	B	12±1	32300	25300	3457	3650	Deleted: <i>Murraya</i>
<i>S. album</i>	Santalaceae	T	180	DBr	40±2	10900	6200	7075	2750	Deleted: <i>Santalum</i>
<i>S. oleosa</i>	Sapindaceae	T	352	DBr	49±2	8500	5400	2198	1950	Deleted: <i>Schleichera</i>

V = Vien, H = Herb, T = Tree, BrB = Brownish black, YeW = Yellowish white, WhBr = Whitish brown, ReBr Reddish brown, B = Black, YeBr = Yellowish brown, DBr = Dark brown, LuB = Luster black, LY = Light Yellow, PY = Pale yellow, DB = Dark black, PW = Pale white

124 Polyphenol contents

125 The concentration of TPh and Fla in the seed coats and kernels varied from 1800 to 32300
 126 mg/kg, from 780 to 31760 mg/kg, from 1200 to 26900 mg/kg and from 300 to 12020 mg/kg,
 127 respectively with a mean value of 15748, 5376, 6954 and 2932 mg/kg (**Table 1**). The [TPh]/[Fla]
 128 ratio in the studied seed coats and kernels ranged from 1.1 to 18.1 and 0.4 to 7.1 with a mean

value of 3.5 and 1.9, respectively. Higher contents of TPh and Fla in the seed coats than the seed kernels was marked.

Large variations in the polyphenol content were observed from one species to another, with noticeably higher TPh and Fla values in seed coats and kernels from tree species (**Fig. 2**).

Similarly, remarkable differences in the polyphenol content of the seed coat and kernel samples were detected as a function of the family (**Fig. 3**). In the case of seed coats, four families showed the highest TPh contents: Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae. Whereas in the case of kernels, three families exhibited the highest TPh contents: Fabaceae, Malvaceae and Santalaceae. As regards Fla contents, the highest concentrations corresponded to seed coats from Lauraceae and Rutaceae families.

Mineral contents

The mineral contents of 15 elements (viz. K, Rb, Mg, Ca, Sr, Al, P, S, Cl, Ti, Mn, Fe, Cu, Zn and Pb) in the seed coats are summarized in **Table 2**. The total concentrations (Σ_{M15}) ranged from 5876 to 36499 mg/kg, with the highest values for seed coats from *Jatropha curcas*. Remarkably high mineral contents were observed in the seed coats from three families: Lauraceae, Rutaceae and Euphorbiaceae (**Fig. 4**).

P and K nutrients were abundant in the seed coats, ranging from 99 to 4983 mg/kg and from 1714 to 21982 mg/kg, respectively. The highest P contents was observed in seed coats from Cucurbitaceae family, while the highest K contents (>15000 mg/kg) were detected in seed coats from *P. marsupium*, *L. glutinosa*, *T. populnea* and *M. koenigii*.

Deleted: *Pterocarpus*

Deleted: *Litsea*

Deleted: *Thespesia*

Deleted: *Murraya*




Fig. 2. Polyphenol concentration variation in seed coats with respect to plant types. (A): Seed coats and (B): Seed kernels.




Fig. 3. Polyphenol concentration variation of seed coats with respect to family. (A) seed coat, (B) seed kernel

Table 2. Mineral contents in the seed coats from the 24 species under study, expressed in mg/kg.

Species	Mg	Al	P	S	Cl	K	Ca	Rb	Sr	Ti	Mn	Fe	Cu	Zn	Pb
<i>B. rubra</i>	1012	78	99	194	71	6305	3409	12	4	39	71	389	4	33	3

<i>C. maxima</i>	1762	67	4220	1307	1178	10591	992	13	3	7	85	583	2	4	2
<i>L. siceraria</i>	2020	55	2799	966	88	11324	3833	9	2	9	47	295	2	5	1
<i>C. lanatus</i>	1913	41	3474	1865	78	3247	2965	10	32	8	57	524	11	10	1
<i>L. aegyptiaca</i>	3344	67	2273	909	55	6791	4080	9	1	11	31	142	3	9	1
<i>C. melo</i>	1638	44	4983	1719	101	3913	338	21	1	12	37	364	8	49	1
<i>L. acutangula</i>	1754	81	3486	1302	142	7134	845	9	3	7	20	125	12	10	1
<i>B. hispida</i>	561	98	1878	1063	131	5859	1604	14	3	8	70	313	1	13	2
<i>M. charantia</i>	2642	432	2441	1444	88	5470	2763	8	2	9	77	308	1	3	1
<i>J. curcas</i>	4002	47	1991	1264	91	14636	14210	15	27	11	40	147	13	4	1
<i>L. purpureus</i>	1382	38	2344	1156	48	8541	3176	4	3	7	32	84	6	9	1
<i>A. odoratissima</i>	1738	46	1745	3140	71	8049	6256	15	9	8	132	125	3	5	1
<i>E. gigas</i>	1096	35	104	195	65	6278	3405	16	12	11	58	65	17	27	1
<i>P. marsupium</i>	2098	61	985	1897	59	15236	6685	18	29	38	79	491	914	3	3
<i>J. regia</i>	105	55	254	116	66	7297	2292	3	9	12	17	68	4	4	1
<i>L. glutinosa</i>	542	43	1559	1424	121	21982	3403	23	12	27	94	181	35	27	1
<i>S. potatorum</i>	5808	68	375	975	105	1714	7734	5	2	7	127	166	6	4	1
<i>G. arboreum</i>	2067	77	4001	1399	91	9312	1387	1	5	9	14	137	5.5	40	1
<i>T. populnea</i>	662	87	1631	1062	4859	16894	3256	17	4	11	33	419	402	1	1
<i>M. azadirachta</i>	323	844	826	725	132	7515	2205	9	3	30	28	685	5.5	12	1
<i>P. embilica</i>	175	59	465	243	111	3405	1008	6	20	9	18	348	4	4	1
<i>M. koenigii</i>	1395	49	1311	480	81	20233	6401	7	13	11	21	271	5	11	1
<i>S. album</i>	911	54	886	1100	77	12405	5280	26	29	23	70	327	7	16	1
<i>S. oleosa</i>	1635	65	1814	1227	68	2568	4982	3	13	12	34	334	14	6	1

219

220 S and Cl concentrations in seed coats were in the 116–3140 mg/kg and in the 48–4859 mg/kg

221 range, respectively. The highest values for S and Cl corresponded to *A. odoratissima* and *T.*

222 *populnea*, respectively.

Deleted: Albizzia

Deleted: Thespesia

Mg and Ca elements, probably present as silicates, ranged from 105 to 5808 mg/kg and from 338 to 14210 mg/kg, respectively. In this case, the highest concentrations of Mg and Ca were observed in seed coats from *S. potatorum* and *J. curcas*.

The concentrations of other elements in the seed coats, expressed in mg/kg, were in the following ranges: 1–26 (Rb), 1–32 (Sr), 35–844 (Al), 7–39 (Ti), 14–132 (Mn), 65–685 (Fe), 1–914 (Cu), 1–49 (Zn) and 1–3 (Pb). The highest concentrations of Rb, Sr, Al, Ti, Pb, Mn, Fe, Cu and Zn were found in the seed coats from *S. album*, *C. lantus*, *M. azadirachta*, *B. rubra*, *P. marsupium*, *A. odoratissima*, *M. azadirachta*, *P. marsupium* and *C. melo*, respectively. It is worth noting that seed coats from *T. populnea* featured high contents of Cl, K, Fe and Cu.




Fig. 4. Total mineral content familywise.

Noticeable differences in the mineral contents were also found depending on plant type. Total mineral contents were at least 29 and 55% higher in seed coats from trees and herbs, respectively, than in vine samples (**Fig. 5A**). Higher concentrations of major elements (P, S, Mg, Ca and Al) were observed in the herb samples (**Fig. 5B**), while those of Cl, K, Mn, Cu, Ti and Sr

Deleted: *Strychnos*

Deleted: *Jatropha*

Deleted: *Santalum*

Deleted: *itrullus*

Deleted: *Melia*

Deleted: *Basella*

Deleted: *Pterocarpus*

Deleted: *Albizia*

Deleted: *Melia*

Deleted: *Pterocarpus*

Deleted: *Cucumis*

Deleted: *Thespesia*

Fig. 4. Total mineral contents in seed coats as a function of the family to which the plant species belongs.

were higher in the tree samples (**Fig 5C**). As regards samples from vines, high contents of Rb, Fe and Zn were detected (**Fig 5D**).

On the basis of their mineral contents, the seed coats from the 24 species under study were categorized into two groups by using cluster analysis (**Fig. 6**). Group-I consisted of 19 species, and the other 5 species were included in group-II, in such a way that the mean concentration value of Σ_{M15} in the seed coats that belonged to group-II was at least twice that of group-I ones.

Correlation coefficients

The correlation coefficients (r) for the seed coat samples from species belonging to the Fabaceae family are shown in **Table 3**. Good correlations were found among K, Mg, Ca, Al, Sr, Ti, Fe, Cu and Pb, suggesting similarities in their bioaccumulation. Between TPh and Fla contents and Cl, Rb and Mn were well correlated, which would point to their accumulation via bond formations with phenolic groups. In addition, good correlations were observed among S, Mg, Ca and Mn, which may be ascribed to the accumulation of the latter three as sulfur compounds.

Distribution of minerals in seed coats and seed kernels

The content of the minerals in the seed coat and the kernels of the prominent seeds is shown in **Tables 4**. The nutrients are well abundant in the seed coats than the kernels. Fifteen elements were detected in the coats. However, in the kernels, only ten elements were identified. Generally, higher concentration of nutrients in the seed coats than the kernel was marked, **Fig. 6**.


(A)

(B)

Ca	0.50	0.46	0.92	0.88	0.01	0.80	0.49	0.68	1.00								
Rb	0.91	0.54	0.39	0.52	-0.75	0.19	0.75	0.33	0.65	1.00							
Sr	0.41	-0.07	0.70	0.85	-0.50	0.11	0.17	0.84	0.68	0.75	1.00						
Ti	0.18	-0.26	0.74	0.89	-0.32	0.10	-0.06	0.93	0.63	0.57	0.97	1.00					
Mn	0.71	0.90	0.50	0.40	0.00	0.83	0.84	0.07	0.78	0.59	0.20	0.06	1.00				
Fe	0.13	-0.21	0.86	0.96	-0.14	0.28	-0.06	0.98	0.73	0.51	0.93	0.98	0.16	1.00			
Cu	0.11	-0.30	0.79	0.91	-0.22	0.15	-0.12	0.97	0.65	0.51	0.94	0.99	0.05	0.99	1.00		
Zn	0.23	0.00	-0.87	-0.73	-0.67	-0.81	0.14	-0.67	-0.71	0.07	-0.27	-0.39	-0.40	-0.58	-0.48	1.00	
Pb	0.11	-0.29	0.80	0.92	-0.21	0.16	-0.12	0.97	0.65	0.50	0.94	0.99	0.06	0.99	1.00	-0.49	1.00

TPh = Total polyphenol content, Fla = Flavonoid content

274



275

Fig. 6. Distribution of total elements in seed coats and kernels.

276

Table 4. Distribution of elements in the seed kernel, mg/kg.

Seed	Mg	P	S	K	Ca	Mn	Fe	Cu	Zn	Rb	Sr
<u>C. maxima</u>	1130	5505	1614	3920	102	63	157	14	78	11	2
<u>J. curcas</u>	1905	5513	1596	8032	2509	26	96	10	42	16	2
<u>P. marsupium</u>	5555	6743	5993	17119	5095	41	145	39	50	20	11

Deleted: Cucurbita

Deleted: Jatropha

Deleted: Pterocarpus

M. koenigii	551	1436	525	13130	2891	10	110	8	15	4	5
-----------------------------	-----	------	-----	-------	------	----	-----	---	----	---	---

Deleted: *Murraya*

280

281 **CONCLUSIONS**

282 Seed coats are major sources of polyphenols and minerals. At least two folds higher nutrients in
 283 seed coats than the kernels are available. Remarkably high total polyphenol contents (of up to
 284 32300 mg/kg) were detected in the seed coats from tree species of the Loganiaceae,
 285 Phyllanthaceae, Lauraceae and Rutaceae families, while the highest flavonoid concentrations (of
 286 up to 26900 mg/kg) corresponded to seed coats from the latter two families. As regards mineral
 287 contents, the highest total values were observed in the seed coats from three families: Lauraceae,
 288 Rutaceae and Euphorbiaceae. The highest concentrations of major elements (P, S, Mg, Ca and
 289 Al) were observed in seed coats from herb species, while those of Cl, K, Mn, Cu, Ti and Sr were
 290 higher in the tree samples. In turn, samples from vines featured high contents of Rb, Fe and Zn.
 291 Seed coats from Indian-laurel and curry tree stand out as particularly promising phytochemical
 292 and nutrient sources.

293 **CONSENT**

294 Not applicable.

295 **ETHICS APPROVAL**

296 Not applicable.

297 **CONFLICT OF INTEREST**

298 The authors declare no conflict of interest.

299

301 **REFERENCES**

- 302 1. Moïse JA, Han S, Gudynaitė-Savitch L, Johnson DA, Miki BLA. Seed coats: Structure,
 303 development, composition, and biotechnology. In *Vitro Cellular & Developmental Biology*
 304 - Plant. 2005;41(5):620-644. <https://doi.org/10.1079/IVP2005686>
- 305 2. Attree R, Du B, Xu B. Distribution of phenolic compounds in seed coat and cotyledon, and
 306 their contribution to antioxidant capacities of red and black seed coat peanuts (*Arachis*
 307 *hypogaea* L.). *Industrial Crops and Products*. 2015;67:448-456.
 308 <https://doi.org/10.1016/j.indcrop.2015.01.080>
- 309 3. Zhang RF, Zhang FX, Zhang MW, Wei ZC, Yang CY, Zhang Y, Tang XJ, Deng YY,
 310 Chi JW. Phenolic composition and antioxidant activity in seed coats of 60 Chinese black
 311 soybean (*Glycine max* L. Merr.) varieties. *Journal of Agricultural and Food Chemistry*.
 312 2011;59(11):5935-5944. <https://doi.org/10.1021/jf201593n>
- 313 4. Abutheraa R, Hettiarachchy N, Phillips GK, Horax R, Chen P, Morawicki R, Kwon YM.
 314 Antimicrobial activities of phenolic extracts derived from seed coats of selected soybean
 315 varieties. *Journal of Food Science*. 2017;82(3):731-737. [https://doi.org/10.1111/1750-](https://doi.org/10.1111/1750-3841.13644)
 316 [3841.13644](https://doi.org/10.1111/1750-3841.13644)
- 317 5. Phommalath S, Teraishi M, Yoshikawa T, Saito H, Tsukiyama T, Nakazaki T, Tanisaka T,
 318 Okumoto Y. Wide genetic variation in phenolic compound content of seed coats among
 319 black soybean cultivars. *Breeding Science*. 2014;64(4):409-415. doi:10.1270/jsbbs.64.409
- 320 6. Ribeiro ND, Maziero SM, Prigol M, Nogueira CW, Rosa DP, Possobom MTD. Mineral
 321 concentrations in the embryo and seed coat of common bean cultivars. *Journal of Food*
 322 *Composition and Analysis*. 2012;26(1-2):89-95. <https://doi.org/10.1016/j.jfca.2012.03.003>

- 323 7. Wu XJ, James R, Anderson AK, Mineral contents in seed coat and canning quality of
 324 selected cultivars of dark red kidney beans (*Phaseolus vulgaris* L.). Journal of Food
 325 Processing and Preservation. 2005;29(1):63-74. [https://doi.org/10.1111/j.1745-](https://doi.org/10.1111/j.1745-4549.2005.00013.x)
 326 4549.2005.00013.x
- 327 8. Ganesan K, Xu B. A critical review on polyphenols and health benefits of black soybeans.
 328 Nutrients. 2017;9(5):455. doi:10.3390/nu9050455
- 329 9. Sevindik M, Akgul H, Pehlivan M, Selamoglu Z. Determination of therapeutic potential of
 330 *Mentha longifolia* ssp. *longifolia*. Fresen Environ Bull, 2017; 26: 4757-4763.
- 331 10. Sivasankar V, Moorthi A, Sarathi Kannan D, Suganya devi P. Anthocyanin, and its
 332 antioxidant properties in selected fruits. Journal of Pharmacy Research. 2011;4(3):800-806.
- 333 11. Prasad DMR, Izam A, Khan MMR. *Jatropha curcas*: Plant of medical benefits. Journal of
 334 Medicinal Plants Research. 2012;6(14):2691-2699. <https://doi.org/10.5897/JMPR10.977>
- 335 12. Saboo SS, Thorat PK, Tapadiya GG, Khadabadi SS. Ancient and recent medicinal uses
 336 of cucurbitaceae family. International Journal of Therapeutic Applications. 2013;9:11-19.
- 337 13. Hassan GA, Bilal T, Ahmad BT, Sameena W, Irshad AN. Economic and ethno-medicinal
 338 uses of *Juglans regia* L. in Kashmir Himalaya. Unique Journal of Ayurvedic and Herbal
 339 Medicines. 2013;1(3):64-67.
- 340 14. Ramana KV, Raju AJS. Traditional and commercial uses of *Litsea glutinosa* (Lour.) C.B.
 341 Robinson (Lauraceae). Journal of Medicinal Plants Studies. 2017;5(3):89-91.
- 342 15. Yadav KN, Kadam PV, Patel JA, Patil MJ. *Strychnos potatorum*: Phytochemical and
 343 pharmacological review. Pharmacognsy Review. 2014;8(15):61-66. doi: 10.4103/0973-
 344 7847.125533

- 345 16. Singh E, Sharma S, Pareek A, Dwivedi J, Yadav S, Sharma S. Phytochemistry, traditional
 346 uses and cancer chemopreventive activity of Amla (*Phyllanthus emblica*): The Sustainer.
 347 Journal of Applied Pharmaceutical Science, 2011;2(1):176-183.
- 348 17. Kumar SR, Loveleena D, Godwin S. Medicinal property of *Murraya Koenigii*- A review.
 349 International Research Journal of Biological Sciences. 2013;2(9):80-83.
- 350 18. Kumar R, Anjum N, Tripathi YC. Phytochemistry and pharmacology of *Santalum album* L.:
 351 A review. World Journal of Pharmaceutical Research. 2015;4(10):1842-1876.
- 352 19. Bhatia H, Kaur J, Nandi S, Gurnani V, Chowdhury A, Reddy PH, Vashishtha A, Rathi B. A
 353 review on *Schleichera oleosa*: Pharmacological and environmental aspects. Journal of
 354 Pharmacy Research. 2013;6(1):224-229. <https://doi.org/10.1016/j.jopr.2012.11.003>
- 355 20. Sathyanarayana T, Sarita T, Balaji M, Ramesh A, Boini MK. Antihyperglycemic and
 356 hypoglycemic effect of *Thespesia populnea* fruit in normal and alloxan-induced diabetes in
 357 rabbits. Saudi Pharmaceutical Journal. 2004;12(2):107-111.
- 358 21. Al-Snafi AE. The pharmacology and medical importance of *dolichos lablab* (*Lablab*
 359 *purpureus*)-A review. IOSR Journal of Pharmacy. 2017;7(2):22-30. DOI: 10.9790/3013-
 360 0702012230
- 361 22. Van Andel TR, Croft S, Van Loon EE, Quiroz D, Towns AM, Raes N. Prioritizing West
 362 African medicinal plants for conservation and sustainable extraction studies based on market
 363 surveys and species distribution models. Biological Conservation. 2015;181:173-181.
 364 <https://doi.org/10.1016/j.biocon.2014.11.015>
- 365 23. Moraghan JT, Grafton K. Genetic diversity and mineral composition of common bean seed.
 366 Journal of the Science of Food and Agriculture. 2001;81(4):404-408.
 367 [https://doi.org/10.1002/1097-0010\(200103\)81:4<404](https://doi.org/10.1002/1097-0010(200103)81:4<404)

- 368 24. Moraghan JT, Grafton K. Distribution of selected elements between the seed coat and
369 embryo of two black bean cultivars. *Journal of Plant Nutrition*. 2002;25(1):169-176.
370 <https://doi.org/10.1081/PLN-100108788>
- 371 25. Tajoddin M, Shinde M, Lalitha J. Polyphenols of mung bean (*Phaseolus aureus* L.) cultivars
372 differing in seed coat color: Effect of dehulling. *Journal of New Seeds*. 2010;11(4): 369-379.
373 <https://doi.org/10.1080/1522886X.2010.520146>
- 374 26. Yang QQ, Gan RY, Ge YY, Zhang D, Corke H. Polyphenols in common beans (*Phaseolus*
375 *vulgaris* L.): Chemistry, analysis, and factors affecting composition. *Comprehensive*
376 *Reviews in Food Science and Food Safety*. 2018;17(6):1518-1539.
377 <https://doi.org/10.1111/1541-4337.12391>
- 378 27. Marles MA, Vandenberg A, Bett KE. Polyphenol oxidase activity and differential
379 accumulation of polyphenolics in seed coats of pinto bean (*Phaseolus vulgaris* L.)
380 characterize postharvest color changes. *Journal of Agricultural and Food Chemistry*.
381 2008;56(16):7049-7056. doi: 10.1021/jf8004367.
- 382 28. Zhong L, Wu G, Fang Z, Wahlqvist ML, Hodgson JM, Clarke MW, Junaldi E, Johnson SK.
383 Characterization of polyphenols in Australian sweet lupin (*Lupinus angustifolius*) seed coat
384 by HPLC-DAD-ESI-MS/MS. *Food Research International*. 2019;116:1153-1162.
385 <https://doi.org/10.1016/j.foodres.2018.09.061>
- 386 29. Bertaud F, Tapin-Lingua S, Pizzi A, Navarrete P, Petit-Conil M. Characterisation of
387 industrial barks for their tannin contents for further green-wood based adhesives
388 applications. *InTech Fibre*. COST FP0901-Hamburg. 2010.

- 389 30. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other
390 oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in*
391 *Enzymology*. 1999;299:152-178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)
- 392 31. Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis
393 by two complementary colorimetric methods. *Journal of Food and Drug Analysis*.
394 2002;10(3):178-182.
- 395 32. Towett, E. K., Shepherd, K. D., Drake, B. L. (2016) Plant elemental composition and
396 portable X-ray fluorescence (pXRF) spectroscopy: quantification under different analytical
397 parameters, *X-Ray Spectrom*. 45, 117–124. DOI 10.1002/xrs.2678