

Original Research Article

THE EFFECTS OF ORGANIC AND CHEMICAL INORGANIC FERTILIZER APPLICATIONS ON YIELD AND PLANT VEGETATIVE GROWTH OF EGGPLANT (*Solanum melongena* L.)

Abstract

A field experiment was conducted to evaluate the effects of organic and chemical fertilizers on yield and plant vegetative growth of eggplant (*Solanum melongena* L.) between 22.05.2018 and 12.09. 2018 under field conditions in Bandırma, Turkey. Treatments were control with no fertilizer (CONT), developed organic fertilizer (DOF), organic fertilizer (OF) chemical fertilizer (CHF), each treatment has 3 replicates with ?? plants. Developed oOrganic fertilizer and OFrganic fertilizer were applied in (2) ton/da, chemical fertilizer (15N, 15P, 15K, 20 SO₃) was applied in 40 kg/da. Yield and plant vegetative growth weight were determined. Organic and chemical fertilizer applications had a significant effect ($P<0.05$) on yield and plant vegetative growth. The eggplant yield at CONT, DOF, OF, CHF were 3922, 4593, 4375, 4579 kg.da⁻¹ respectively. The plant vegetative growth at CONT, DOF, OF, CHF were 3548, 4018, 3818, 3882 kg.da⁻¹ respectively. The difference between fertilizers (DOF, OF, and CHF) was not very high so that organic fertilizers are competitive and may be a suitable replacement for chemical fertilizer.

KEYWORDS: Organic fertilizer, vegetative growth, fruit yield, eggplant

INTRODUCTION

Eggplant (*Solanum melongena* L.), also known as aubergine, brinjal or guinea squash is one of the non-tuberos species of the night shade family Solanaceae (Kantharajah et al., 2004). The varieties of *Solanum melongena* L. show a wide range of fruit shapes and colors, ranging from oval or egg-shaped to long club shaped; and from white, yellow, green through degrees of purple pigmentation to almost black. It is an economically important crop in Asia, Africa and the sub-tropics (India, Central America) and it is also cultivated in some warm temperate regions of the Mediterranean and South America (Sihachkr et al., 1993). Eggplant fruits are known for being low in calories and having a mineral composition beneficial for human health. They are also a rich source of potassium, magnesium, calcium and iron (Zenia and Halina, 2008). According to the reports released in 2013, in 162 countries, 1.8 million organic certified producers, 37.2 million ha in the field of organic farming (Willer et al., 2013). According to the data obtained from TURKSTAT (2014) as of 2013 (213) kinds of products in Turkey, at 769014 hectares, 60797 Manufacture, 1.620.466 tons of organic herbal production. [The amount of organic vegetable production in Turkey around 30,000 tons. The major share respectively tomatoes, cucumber, beans, pepper, lettuce and eggplant (Hekimoğlu and Altındöğür, 2010). Fertilizers are used to improve soil fertility but intensive inorganic fertilizer usage in agriculture causes so many health problems and unrecoverable environmental pollution. Thus, to reduce and eliminate the adverse effects of sSynthetic fertilizers on human health and environment, now-a-days a new agricultural practice have been developed called as organic agriculture, sustainable agriculture or ecological agriculture (Aksoy, 2001; Chowdhury, 2004). Inorganic fertilizers are known for their high cost and their negative environmental effects if managed poorly (Morris et al., 2007). The Ddirect effects are due to readily available nutrients from applied compost while indirect effects are through the improvement of soil structure and nutrient and

Formatted: English (United States)

Comment [P1]: Title should be modified according to context

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Comment [P2]: The units must be the same style throughout the entire manuscript ton/da or ton da⁻¹. Check the entire manuscript.

Comment [P3]: Give units % or kg/da

Comment [P4]: Plant organs fresh weight or dry weight???

Comment [P5]: Are the differences significant or not???

Formatted: Font: Italic

Comment [P6]: Sentence is incomplete

Comment [P7]: It is not given in the list

Comment [P8]: Sentence is incomplete

Formatted: English (United States)

Comment [P9]: Italic or regular. Check others in the text

47 water retention favorable to plant growth. In a particular experiment, the authors found that application of
48 organic amendments hasve potential to increase the growth and chemical composition of two cultivars of
49 safflower and therefore, might be a good alternative to chemical fertilizers (Naderi and Bijanzadeh,
50 2014). There is an effect of different organic manures and inorganic fertilizer levels in soil were observed
51 on growth performances and yield of eggplant (Vijaya, 2011). Louisa and Taguiling, 2016 reported that
52 compost stimulates plant growth in terms of total height, number of leaves, fresh weight of plant biomass,
53 and initial yield of green pepper (*Capsicum annuum*), eggplant (*Solanum melongena*), and Okra
54 (*Abelmoscus esculentus*).
55 This study aims to determine the effect of organic and developed organic fertilizers as compared to
56 chemical inorganic fertilizers on plant growth and yield of eggplant.

Comment [a10]: A pre-sentence should be written, which justifies this sentence.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

57 2. MATERIALS AND METHODS

58 2.1. MATERIALS

59 **Research Area:** The experiment carried out in a field near Akçapınar village which located at 40 ° 16
60 '44.4252' North and 28 ° 4' 18.9552 " Eastern latitudes and longitudes. The height altitude from the sea
61 level is 41 m). Akçapınar is bound to the of Bandırma district, Balıkesir governorate, Bandırma is
62 located to the south of the Marmara Sea. Karacabey (Bursa) to the east, Gönen to the west, Erdek
63 and Kapıdağ to the north, Manyas and Kuşgözü to the south.

Comment [a11]: This information is not necessary because meridians and parallels are given.

64 **Climate:** Bandırma is under the climate of the Mediterranean Sea and the Black Sea. In addition, due to
65 the fact that is located on the transition area of the terrestrial climate of the Balkans, various climate
66 features are observed in the district. In addition, due to the fact that the terrestrial climate from the
67 Balkans is located on the transition area, various climate features are observed in the district, the
68 lowest temperature according to 52-year climate data is -14.6 °C (January 15, 1954), while the high
69 temperature was recorded as 42.4 °C (July 9, 2000). The average annual temperature is 14 °C. The
70 dominant wind direction is North-Northeast. The average wind speed is 15 km/h. The average annual
71 rainfall in the district is 703.3 mm. The annual relative humidity average is 73%.

72 **Properties of the Experimental Area Soil:** From the soil profile opened in the experiment area,
73 disturbed and undisturbed soil samples were taken from 0-30, 30-60 and 60-90 cm soil layers and
74 physical and chemical analyses were performed on these samples. The pH of the test soil was pH 8-8.1,
75 salt content 0.76-0.84 ds/m, organic matter % 0.68-1.2, total nitrogen 9.4-9.7 kg.da⁻¹, phosphorus 1.44 -
76 1.69 kg.da⁻¹, potassium 59.8-83.8 kg.da⁻¹, calcium varied from 1971 to 2230 kg.da⁻¹, and volume weight
77 (As) ranged from 1.28 to 1.31 g.cm⁻³. In addition, a 0-30 cm layer of the soil profile; 30-60 cm sandy
78 clay Tin, 60-90 cm layer has clayey texture.

Comment [a12]: Modify sentence

79 **Properties of Irrigation Water:** Irrigation water was taken from a deep well near the experiment field.
80 The water from the deep well is was pumped into a pool. A sample was taken from irrigation water and
81 analyzed. Irrigation water is was in C3S1 class. The water is was 3 in salinity and 1 in sodium.

Comment [a13]: What area 3 and 1?

82 **Class A pan Evaporation:** Class A pan evaporation used in the experiment is was 121 cm in diameter
83 and 25 cm in height and was made of 2 mm thick galvanized steel plate. The Wooden base was put at
84 the bottom of the pan allowing air flow 10 cm height. Changes in the water level at the surface are
85 measured. It was covered on a wire mesh to prevent the water intake by the insects and animals.

86 **Fertilizers:** Three different fertilizers were used in the study. 1-Chemical Inorganic fertilizer (CHIF), 2-
87 Organic fertilizer (OF), and 3- Developed organic fertilizer (DOF). CHIF; N, P, K, SO₃ (15-15-15 + 20),
88 OF; DOF; 75% cow manure and 25% poultry manure (The difference between OF and DOF is that
89 developed organic fertilizer is mixed with small particles from the organic fertilizer it's about 50

Formatted: Subscript

microns in volume). In both organic fertilizers, the organic matter content is 33%, the humic + volvic content **is was** 18.4% and the pH is 7.4. The organic fertilizers both were prepared in an aerobic fermentation method (windrow), whereas the level of humidity, oxygen and temperature were monitored. The humidity at the beginning of fermentation was 60% and the process of flipping and aeration were performed when the temperature reached **at** about 60 degrees Celsius and when the oxygen rate reduced above 5% for 45 days. This process called composting which is a naturally occurring process in which bacteria, fungi, and other microorganisms convert organic material into a stabilized product termed as compost. **This means that microorganisms do the composting work for you. Your role in managing the composting process is to make sure that the microorganisms have the environment they need in order to do their work quickly and effectively.**

2.2. Method

Fertilizer Treatment : Three fertilizer treatments were applied; the first treatment is organic fertilizer (OF) applied at 2 **f**Ton/da, the second fertilizer is developed **edment** organic fertilizer (DOF) applied at 2 **f**Ton/da and the third treatment is **chemical inorganic** fertilizer (**CHFIF**) applied at 40 kg/da. Each treatment has 3 replicates.

Soil Preparation and Planting First, the experimental area was tilled and then parcels were created. Then the fertilizers applied **a**According to the experiment plan. After that, the drip irrigation system **has been applied.was established**. The eggplant **seedlingsapling** were planted according to the experiment plan on 02/06/2017, the spacing between **the lines as planting linesrow is was** 70 cm, the spacing **between the plant and the otherwithin rows** is 40 cm, each parcel contains 6 **linesrows**, in each **line row** (20) plants and each **experiment pieceplot** contains (120) plants, the **plot** dimensions **arewere**: 8 m*4.2 m=33.6 m². **There are 9 parcels, with an area of 33.6 m² for each parcel and the total area of the experiment pieces is 302.4 m², the space between the parcel and the other is 2 m,The parcels were separated from each others 2 m space and** the total area of the experiment **is was** 659.2 m², the total number of **eEggplant (Solanum mMelongena) saplingsseedling was is** (1080).

Formatted: Superscript

Formatted: Superscript

Calculation of Irrigation Water Amount The amount of irrigation water was calculated according to the method given by Gençoğlu and Ark., (2006), **taking e**Evaporation quantities were measured at 09:00 a.m. in class A **evaporation** pan **evaporation**. These measured values **are were** used to calculate the amount of irrigation water. The amount of irrigation water **is was** calculated by the equation given below.

$$V = E(\text{pan}) * A * P * IR$$

Formatted: English (United States)

V: water volume / L

E: A Pan-Evaporation / mm

A: parcel area / m²

Formatted: Superscript

P: percentage of coverage

IR: Irrigation Applied rate (100%). Drip irrigation method was used, irrigation was applied as 100% of evaporation from Class A pan evaporating, and the calculated irrigation water amounts were applied to the parcels by using water meters. **The E**experiment was irrigated 12 times. The total amount of irrigation water applied to the experimental **pieces area** was 305 mm/da, in addition to 143 mm rainfall.

Evaluation of Data: Statistical analyzes were performed on the parcels (randomized blocks) of the obtained data according to the experimental designThe experiment was established according to

randomized blocks design. Each treatment was replicated times with plant ANOVA one way statistical analyses were performed by using SPSS (24.0v) Program.

Result and Discussion

Eggplant plant began to fruiting one month after planting (22.06.2018). It was harvested 5 times, the first harvest was in (19.07.2018), and the last harvest was in (12.09.2018). According to the results of variance analysis conducted in order to determine the effects of the kind of fertilizer on the yield, it was found significant at 5% level ($p < 0.05$). (Urkurkar, 2010 and Yolcu, 2011) also reported significant differences in yield with the application of different fertilizer sources. According to the LSD test there are 2 groups, DOF, OF and CHF treatment, in A group, the control in B group. The heights yield at DOF (4593 kg.da⁻¹) and the lowest yield was at control (3922 kg.da⁻¹). There were no significant differences between the three fertilizers treatments (DOF, OF, CHF) (table 1). These results can be explained that adding organic fertilizer to the soil make mineral more available in the soil in addition to what organic fertilizer contain from mineral.

Table 1. Total Yield Results (kg.da⁻¹)

Replicates	DOF	OF	CHF	Control
R1	4630	4827	4539	3886
R2	4713	3987	4561	3828
R3	4437	4311	4638	4051
Average	4593 ^a	4375 ^a	4579 ^a	3922 ^b




Figure 1. Effect of three different fertilizers on yield

Plant Vegetative Growth: According to the results of variance analysis conducted in order to determine the effects of the kind of fertilizer on the plant vegetative growth of eggplant it was found significant at 5% level ($p < 0.05$). (Lanyasunya; 2007, Efthimiadou; 2009, Fateh, et al, 2009 and Urkurkar et al., 2010) also reported significant differences in vegetative growth with the application of different fertilizer sources. According to the LSD test there are 2 groups, DOF, OF, CHF treatment, in A group and the control in B group. The heights plant vegetative growth at DOF (4018 kg.da⁻¹), the lowest plant vegetative growth was at control (3548 kg.da⁻¹). There were no significant differences between the three fertilizers treatments (DOF, OF, CHF). (Table 2). These result can be explained that organic fertilizer contains Azote which is important to vegetative growth.

Comment [P15]: Add replication and plant number for each replication

Formatted: English (United States)

Comment [P16]: Add significance level and comperison methods. LSD, Tukey ???

Formatted: English (United States)

Comment [P17]: Et al., in the list

Comment [P18]: The citations must be done according to common wrting formats.

Formatted: Font: (Default) Times New Roman, Font color: Black, English (United States)

Formatted: Font: (Default) Times New Roman, Font color: Black, English (United States)

Formatted: Font: (Default) Times New Roman, Font color: Black, English (United States)

Comment [P19]: Writing character should be same in all manuscript

Formatted ... [1]

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted ... [2]

Formatted ... [3]

Formatted ... [4]

Formatted ... [5]

Formatted ... [6]

Formatted ... [7]

Formatted ... [8]

Formatted ... [9]

Formatted ... [10]

Formatted ... [11]

Formatted ... [12]

Formatted ... [13]

Formatted ... [14]

Comment [P20]: No information ws given as to waht R1, R2 and R3 (Table 1 and 2) represent

Formatted: English (United States)

Formatted ... [15]

Formatted: English (United States)

Formatted: English (United States)

Formatted ... [16]

Formatted ... [17]

Table 2. Total vegetative growth Results (kg.da⁻¹)

Replicates	DOF	OF	CHF	Control
R1	3991	3869	3957	3592
R2	3987	3895	4083	3571
R3	4077	3690	3606	3483
Average	4018 ^a	3818 ^a	3882 ^a	3548 ^b




Figure 2. Effect of three different fertilizers on Plant Vegetative Growth

4. CONCLUSIONS

According to Bandırma Akçapınar conditions, during the growing period of the eggplant it was found as fertilizers application there were significant differences between the control treatment and the other treatments ($P < 0.05$), in yield and plant vegetative growth. No significant differences were found between the three fertilizers treatments (DOF, OF, CHF) in yield and plant vegetative growth ($P > 0.05$). The highest fruit yield of the experiment (4593 kg.da⁻¹) was obtained at DOF treatment; the lowest yield in the experiment (3562 kg.da⁻¹) was obtained at control. The heights plant vegetative growth at DOF treatment (4018 kg.da⁻¹), and the lowest vegetative growth was at control (3548 kg.da⁻¹).

5. REFERENCES

- Aksoy, U., (2001). Ecological Farming. II. Ecological Farming Symposium in Turkey. 14-16 December. Antalya.
- Anonymous, [http://www.bandirma.gov.tr/climate and geographic Web site.](http://www.bandirma.gov.tr/climate%20and%20geographic%20Web%20site/)
- Chowdhury, R., (2004). Effects of chemical fertilizers on the surrounding environment and the alternative to the chemical fertilizers IES- Envis Newsletter. (3): pp 4-5.
- Efthimiadou, A., D. Bilalis, A. Karkanis, B. Froud-Williams, I. Eleftherochorinos, (2009). Effects of Cultural System (Organic and Conventional) on Growth, Photosynthesis and Yield Components of Sweet Corn (Zea mays L.) under Semi-Arid Environment. Not Bot Hort Agrobot Cluj. 37, 104-111.
- FAO Irrigation and Drainage Paper No. 56 (2006) FAO Irrigation Paper No. 33

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States), Highlight

Formatted: Font: 11 pt, Highlight

Formatted: English (United States), Highlight

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Comment [P21]: The letter characters must be same throughout the entire manuscript.

Formatted: English (United States)

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: 11 pt, Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Font: 11 pt, Highlight

Comment [P22]: Each references in the list must be written in the same format according to instruction for authors.

Comment [P23]: It was not cited in the text

Comment [P24]: It was not cited in the text

179 Fateh, E., Chaichi, M.R., Sharifi Ashorabadi, E., Mazaheri, D., Jafari, A.A., Rengel, Z., (2009). Effects of
180 organic and chemical fertilizers on forage yield and quality of globe artichoke (*Cynara scolymus* L.).
181 Asian Journal of Crop Science, 1: 40-48.

Formatted: Font: 11 pt, Italic

182 Gençoğlu, C., Altunbey, H., Gençoğlu, S., (2006). Response of Green Bean (*P. Vulgaris* L.) to
183 Subsurface Drip Irrigation and Partial Rootzone-Drying Irrigation. Agricultural water management,
184 84(3): 274-280.

Formatted: Font: 11 pt, Italic

185 Gul, I., Tan, M., Dumlu, Z., (2009). The effects of chemical fertilizer, farmyard manure and some soil
186 amendments on seed yield of common vetch. I. GAP Organic Agriculture Congress, 17-20 Kasım
187 Şanlıurfa.

Comment [P25]: English

Comment [P26]: It was not cited in the text

188 Hekimoğlu B, Altındeğer, M., (2010). Samsun governorate vegetable production sector.
189 <http://samsuntarim.gov.tr/ya-publications/technical information /liflets/garden/ vegetable sector.pdf>.

190 Kantharajah A.S., Golegaonkar P.G., (2004). Somatic Embryogenesis in Eggplant Review. J. Sci. Hortic.
191 99:107-117.

192 Lanyasunya, T.P., Wang, H., Mukisira, E. A., Lukibisi, F.B., Kuria, D.M., Kibitok, N.K., (2007). Effect
193 of manure and fertilizer application on yield of sorghum alnum harvested at different maturity stages.
194 J Anim Vet Adv. 6: 879-882.

195 Morris M, VA Kelly, RJ Kopicki and D Byerlee, D., (2007). Fertilizer Use in African Agriculture:
196 Lessons Learned and Good Practice Guidelines. Washington, DC: The World Bank. The Rain Forest
197 Area of Nigeria. Applied Tropical Agriculture 5: pp 20-23.

198 Naderi, R. and Bijanzadeh, E. (2014). Organic amendments and nitrogen effects on growth and chemical
199 composition of two cultivars of safflower (*Carthamus tinctorius* L.) Australian Journal of Crop
200 science. 8(4):577-581.

201 Louisa, MA. and. Taguiling G., 2016. Response of Some Vegetable Plants to Green Biomass-Enriched
202 Compost. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)* e-ISSN: 2319-2380,
203 p-ISSN: 2319-2372. Volume 9, Issue 5 Ver. II (May. 2016), PP 67-74 www.iosrjournals.org

Comment [P27]: () ??

204 Sihachkr D., Chaput M.H., Serraf L., Ducreux G., 1993. Regeneration of plants from protoplasts of
205 eggplant (*Solanum melongena* L.). In: Bajaj, Y.P.S. (Ed.), Biotechnology in Agriculture and Forestry,
206 Plant Protoplasts and Genetic Engineering. Springer, Berlin. pp. 108-122.

Comment [P28]: () ??

207 Vijaya. K.S., Seethalakshmi, S., (2011). Response of Eggplant (*Solanum melongena* L.) To Integrated
208 Nutrient Management Amended Soil. International Journal of Scientific & Engineering Research
209 Volume 2, Issue 8, August-2011 1 ISSN 2229-5518.

Formatted: Font: 11 pt, Not Italic

210 Urkurkar, J.S., Tiwari, A., Chitale, S., Bajpai, R.K., (2010). Influence of long- term use of inorganic and
211 organic manures on soil fertility and sustainable productivity of rice (*Oryza sativa*) and wheat
212 (*Triticum aestivum*) in inceptisols. Indian J. Agr Sci., 80: 208-212.

Formatted: Font: 11 pt, Italic

213 Yolcu, H., (2011). The Effects of Some Organic and Chemical Fertilizer Application on Yield,
214 Morphology, Quality and Mineral Content of Common Vetch (*Vicia sativa* L.) Turkish Journal of
215 Field Crops, 2011, 16, 197-202.

Formatted: Font: 11 pt, Italic

216 Willer H, Lernoud J, Kilcher L. (Eds.), 2013. The World of Organic Agriculture. Statistics and Emerging
217 Trends 2009. IFOAM, Bonn, FiBL, Frick, ITC, Ge-neva, pp. 340.

218 Zenia M., Halina B., (2008). Content of Microelements in Eggplant Fruits Depending on Nitrogen
219 Fertilization and Plant Training Method. J. Elementol. 13(2):269-274.▲

Comment [P29]: () ??

Formatted: English (United States)

UNDER PEER REVIEW

Page 4: [1] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [1] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [1] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [1] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [2] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [2] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [3] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [3] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [4] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [4] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [5] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [5] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [6] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [6] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		

Page 4: [7] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [7] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [8] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [8] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [9] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [9] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [10] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [10] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [11] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [11] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [12] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [12] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [13] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		
Page 4: [13] Formatted	PC	02-Jul-19 9:05:00 AM
English (United States)		

