

Original Research Article

EFFECT ON ~~SOME~~ PROXIMATE, CHEMICAL AND FUNCTIONAL CHARACTERISTICS DURING THE TRANSFORMATION OF CHHATTISGARH LOCAL PADDY VARIETY TO FLAKED RICE (POHA)

ABSTRACT

The present study was undertaken to study the effect of ~~rice flaking process~~ on some proximate, functional and chemical characteristics ~~during the transformation~~ of Chhattisgarh local paddy variety ~~to flaked rice (Poha)~~. In the proximate analysis, change in ~~crude f~~ Fat content was found to ~~be significant (P>0.05)~~ and it ~~was~~ ranged from paddy 3.20% to 0.96% for thick and 1.00% for thin flaked rice, and the change in ~~crude f~~ Fiber content from paddy to flaked rice was found to be 2.00% to 1.85% for thick and 1.79 for thin flaked rice. Similarly, change in ~~crude p~~ Protein content from paddy to flaked rice was found to be 4.23% to 4.04% for thick and 3.76 for thin flaked rice and change in ~~a~~ Ash content from paddy to flaked rice was found to be 1.00% to 1.35% for thick and 1.24% for thin flaked rice. ~~During the investigation of~~ Chemical properties, it was found that the change in ~~showed that a~~ Amylose content from paddy to flaked rice ~~was found to be significant (P>0.05)~~ and it was ranged from paddy 19.27% to 18.19% for thick and 18.26% for thin flaked rice. Similarly, change in ~~s~~ Starch content from paddy to flaked rice was found to be 74.15% to 73.89% for thick and 73.47% for thin flaked rice. While the investigation on functional properties, it was found to be significant (P>0.05) result in Water absorbing (WAI) and Water solubility index (WSI). Change in Water absorbing index (WAI) from paddy to flaked rice was found to be 2.19 g/g to 5.18 g/g for thick and 5.05 g/g for thin flaked rice. ~~S~~ Similarly, change in Water solubility index (WSI) from paddy to flaked rice was found to be 3.25% to 4.86 for thick and 5.09% for thin flaked rice

Keywords – Paddy, ~~r~~ Raw rice, chemical and functional properties of flaked rice.

1. INTRODUCTION

Comment [h1]: This word 'some' should not be part of title. There is nothing like 'some proximate'.

Comment [h2]: Check your Table and report based on what you did. You did not subject your data to ANOVA, so delete the highlighted words 'be significant (P>0.05)'.

Comment [h3]: When you are talking about range, it is from least to highest value i.e. between 2 values. Please correct and do same for others.

Comment [h4]: Kindly rewrite these sentences bearing in mind the corrections above.

The grain, called rice (*Oryza sativa* L.) for more than 8000 years, has been the companion of humankind. It is the most important food commodity in Asia, particularly in South and South-East Asia, where more than 90% of rice is produced and consumed. Paddy (*Oryza sativa* L.) is a major food grain in India. It is grown under wide agro-climatic conditions. Several varieties of paddy are being grown in the world. India produces varieties of rice depending upon the climate, cultivation situation, rainfall and socio-economic factors and such as numerous varieties of paddy are produced in India. However, there are over 7,000 varieties of rice around the world. Before the rice grain is consumed, paddy undergoes several post-harvest operations. The maximum recovery of head rice, yield and the quality of rice depend mainly on the variety of paddy as well as the parboiling characteristics of paddy. (Dahare *et al.*, 2019).

Rice is a regular component of the African diet, usually consumed as a whole grain; which contributes more to the total calorie intake. In the major rice consuming countries, rice quality dictates the market value of the commodity and plays an important role in the development and adoption of new varieties (Juliano and Villarreal, 1993; Fitzgerald *et al.*, 2009). A significant variation in physical, milling and cooking quality has been shown among rice varieties produced in different parts of the world due to diverse genetic and environmental factors (Singh *et al.*, 2005; Izawa, 2008).

Paddy (*Oryza sativa* L.) is one of the most important staple food crops which is a major source of nutrients in many parts of the world. Paddy is second largest major cereal crop a member of the grass family (Graminaceae), which produces starchy seeds. Rice is used as an important staple food by the people in many parts of the world after wheat. Rice is used as a source of nourishment for more than half of the world's population (Dahare *et al.*, 2017).

The paddy grain is made up of hull or husk (18 - 28%) and the caryopsis or the brown rice (72 - 83%). The brown rice consists of a brownish outer layer (pericarp, tegmen and aleurone layers) called the bran (5 - 8%), the germ or embryo (2 - 3%) connected on the ventral side of the grain, and the edible portion endosperm, (89 - 94%) (Ray Lantin, 1999). Apart from production, the success of rice industries depends on the milling quality of rice. According to the qualities of rice, it is used for different industrial purpose. Chalky, medium, bold rice is more preferred by "Poha" Industries, than translucent for rice grain varieties

Comment [h5]: Italicize and check the spelling.

Comment [h6]: Is it a compound name? If no, please use only last name.

having translucent character fine, slender, with better Head Rice Recovery (HRR) are preferred. HRR is an important trait of rice makes the variety important for industrial purpose. If in any variety HRR is more it has better economic importance, but this trait varied within the varieties if grown in different seasons.

The whole rice is a good source of carbohydrate as well as other ingredients such as vitamins, minerals, and oryzanol as compared to with rice (Kumar *et al.*, 2016; Bhattacharya, 2011). It also contains valuable nutrients including dietary fiber and phytochemicals, which have been linked to minimizing the risk of various diseases (Maisont and Narkruga, 2009). It is utilized mostly at the household level where it is consumed as boiled, fried or ground rice with stew or soup (Osaretin and Abosede, 2007). Consumer demand for good quality rice is high due to their patronage for imported rice. Since rice production is the major occupation of most farmers In India, and to ensure that locally processed rice varieties remain vital and relevant to the rural economy and agricultural production, there is the need to evaluate their quality so as to compare them with their imported counterparts. Therefore, the objective of the present study was to evaluation “effect on some proximate, chemical and functional characteristics during the transformation of Chhattisgarh local paddy variety to flaked rice

Comment [h7]: Kindly rewrite as ‘....evaluate the effect of processing Chhattisgarh local paddy variety to flaked rice on proximate’

2. MATERIALS AND METHODS

2.1 Proximate analysis

2.1.1 Moisture content

The moisture content of the sample was determined by standard air-oven method (Ranganna, 1995). A test sample of 5 g was kept for 24 hours in hot air electric oven maintained at 105°C after 24 hours the sample drawn from the oven and placed in desiccators for cooling to ambient temperature. After cooling, the weight of the sample was taken precisely. The loss in weight was determined and moisture content was calculated by using the following equation (1) expression:

Comment [h8]: This should be AOAC and not Ranganna. Check your manual and correct others.

Comment [h9]: What do you mean? Kept or dried

Comment [h10]: Specify the model, company and country of origin. Please do same for other equipment used.

$$MC_{wt}(\%) = \frac{W_1 - W_2}{W_1} \times 100 \quad \dots 1$$

Comment [h11]: Let 1 and 2 be subscripts i.e. W_1 , W_2 and indicate what they stand for.

2.1.2 Crude pProtein content

Nitrogen (N₂ %) of brown rice samples was estimated by using auto Kjeldahl equipment (Kel plus, pelican system, India) according to the method described in AOAC (Year). Digestion of brown rice (0.5 g sample) was carried out in the auto Kjeldahl equipment at 420°C for 2.30 hours. The digested sample obtained was distilled with 40% NaOH (sodium hydroxide) and 4% boric acid. The vapor of ammonia obtained after distillation was collected in boric acid (distillation time approximately 7 min.) and then titrated against 0.1 N HCL (hydrochloric acid). The percentage of N₂ of brown rice sample was calculated by using the following equation (Ranganna, 1995).

$$\text{Nitrogen (\%)} = \frac{14.01 \times (SR - BR) \times 0.1 \times 100}{1000 \times W_s} \quad \dots 2$$

$$\text{Protein (\%)} = N \times 6.25 \quad \dots 3$$

2.1.3 Crude Fat content

Crude fat was determined by using the Soxhlet apparatus (AOAC, 2000). Oven dry beaker and sample at 100°C for half hours. Keep them in desiccators to avoid moisture content gain from the atmospheres. Weight the beakers and note the reading as initial weight. Carefully weight 5 gm of sample flour and keep in cellulose thimble. The thimble was then placed in a beaker and beaker is filled with petroleum ether (boiling point 40-60°C) about 80 ml of the beaker. Then beaker is now placed in soxhlet apparatus with a thimble for 2 h at 90°C. the petroleum ether was then removed by evaporation and the beaker with residue in an oven at 105°C for 30 min., cooled in desiccators and weight. The percentage of oil was calculated by using the following equation

$$\text{Fat content (\%)} = \frac{W_2 - W_1}{W} \times 100 \quad \dots 4$$

2.1.4 Ash content

Ash content was determined according to (AOAC, 2000) procedure. One (1) g of the sample was taken in a silica crucible and weighed. It was made to ash in a muffle furnace

Comment [h12]: You did not analyse only brown rice, so delete brown.

Comment [h13]: Repetition.

Comment [h14]: Please indicate the meaning of SR, BR and Ws.

Comment [h15]: 30 min is better.

Comment [h16]: Please rewrite and use past tenses, weighed' not 'weight', 'g' not 'gm'

Comment [h17]: Please change to '% Crude fat'. Also check equation (1) for more information and use it to correct equations (5) and (9).

at 600°C for 4 hours. The crucible was cooled in the desiccators and weighted, and the value of ash content was calculated by using the following equation.

$$\text{Ash content (\%)} = \frac{W_2 - W_1}{W} \times 100 \quad \dots\dots 5$$

2.1.5 Crude Fiber content

The crude fiber was determined by using the fibra plus apparatus (Sadasivam and Manickam, 2005). Oven dry crucible and sample at 100°C for 30 min ~~half hours~~. Keep them in desiccators to avoid moisture content gain from the atmospheres. Weight the crucibles and note the reading as initial weight (W_1). Carefully weight 2 gm of sample grind flour and keep in the crucible. The crucible was then placed in a fibra plus apparatus. And after that, the 1.25 % sulfuric acid (H_2SO_4) is filled from the top of the apparatus up to the 150 ml of the crucible. And the sample is boiled in apparatus at 400°C for 40 ~~minutes~~. After completion, the acid wash drains the acid and wash the sample twice and thrice with distilled water. During drainage ensure that the knob is in vacuum mode. After acid wash, a similar process is done. The 1.25 % NaOH (sodium hydroxide) is filled from the top of the apparatus up to the 150 ml of the crucible. And the sample is boiled in apparatus at again 400°C for 40 ~~minutes~~. After completion, the acid wash drains the acid and wash the sample twice and thrice with distilled water. During drainage ensure that the knob is in vacuum mode. After alkali wash takes out crucibles and dries them in a hot air oven until the crucible is free from moisture. ~~C~~ooled in desiccators. Weight the crucible and record the reading as (W_2). Place all the crucibles in a muffle furnace at 600°C for ashing. Cool down the hot crucible after ashing to room temperature using a desiccator. Now weight the crucible and record the reading as (W_3). The fiber content of the sample is calculated using the following equation.

$$\text{Fiber content (\%)} = \frac{W_1 - W_3}{W} \times 100 \quad \dots\dots 6$$

2.2 Chemical properties

2.2.1 Starch content determination

Starch is an important polysaccharide. Starch, which is composed of several glucose molecules, is a mixture of two types of component namely Amylose and Amylopectin. Starch is hydrolyzed into simple sugar by dilute acid and the quantity of sample sugar is measured colorimetrically.

A 500 mg sample of grind powder of each grain variety was weighted and kept in a centrifuge tube and homogenized the sample in hot 80% ethanol to remove sugar. Centrifuged at 600 rpm for 5 minutes at 25°C and retained residue. (Sadasivam and Manickam, 2005). Washing the residue repeatedly with hot 80% ethanol until the washing did not give the colour with anthrone reagent. In dried sample of residue 5.0 ml of distilled water and 6.5 ml of perchloric acid were added and centrifuged at 6000 rpm for 20 minutes at 0°C and supernatant was saved, the centrifuge of the sample was repeated with perchloric acid and distilled water and supernatant were saved and diluted sample with volume makeup with 100 ml distilled water. After that, pipette out 0.2 ml of supernatant and make up the volume to 1 ml with distilled water in each tube 4 ml of anthrone reagent was added carefully and also 4 ml anthrone reagent was added in standard solution prepared by taking 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 and 1.0 ml of working standard of glucose with makeup 1 ml distilled water solution. Heated up 8 minute in boiling water bath and cooled down rapidly and read the intensity of green to dark color at 630 nm.




Fig. 2.2: Standard graph of glucose solution using anthrone reagent

Comment [h18]: Please delete, it is irrelevant here.

Comment [h19]: Be consistent, 'g' is most acceptable. You can convert mg to g

Comment [h20]: Why did you separate this reference?

Comment [h21]: Check the temperature

Comment [h22]: 'saved' or 'sieved'. Please correct.

Comment [h23]: Where is Fig. 2.1?

2.2.2 Amylose content determination

Starch is composed of two compounds, namely Amylose and Amylopectin. Amylose is a linear or non branched polymer of glucose. The glucose units are joined by α -1-4 glucosidic linkage.

Comment [h24]: Delete

A 100 mg of grind powdered sample of each grain variety was weighed and put into a conical flask and 1 ml of distilled ethanol and 10 ml of 1N NaOH were added. (Sadasivam and Manickam, 2005). Then the sample was heated in a boiling water bath at 100°C for 10 minutes. 100 ml volume was made up by adding distilled water. 2.5 ml of the extract was taken then 20 ml distilled water and 3 drops of 0.1 % phenolphthalein were added. 0.1 N HCl solution was added drop by drop until the pink colour was disappeared. After that 1 ml of iodine reagent (KI solution) was added and 50 ml volume was made up by adding distilled water then the colour of the solution was read at 590 nm. The blank solution was prepared by diluting 1 ml iodine reagent into 50 ml distilled water. A standard graph of Amylose content was developed by taking the color of standard amylose solution at 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 and 1 ml concentration and read the intensity of color at 590 nm.

Comment [h25]: Reference should come immediately after 'added' before full stop.




Fig. 2.3: Standard graph of amylose using iodine reagent

Comment [h26]: Change to Fig. 2.2

2.3 Determination of Functional properties

2.3.1 Water absorbing index (WAI) and water solubility index (WSI)

One gram sample taken in centrifuge tube was added with 10 ml distilled water and agitated for 30 min followed by its centrifugation at 3000 rpm for 25 min. The decanted

centrifuged tube with settled gel at the base was weighed and used in the calculation of WAI (Eq. (8)). The supernatant obtained during WAI estimation was used to determine WSI by decanting it into a pre-weighed evaporating dish whose final weight after oven drying at 103°C was recorded and used in the calculation of WSI by using Eq. (9) (Stojceska *et al.*, 2008; Kumar *et al.*, 2016).

Comment [h27]: 'et' remove the full stop

$$\text{WAI (g/g)} = \frac{\text{Weight of gel}}{\text{The dry weight of the sample}} \quad \dots 7$$

$$\text{WSI (\%)} = \frac{\text{Weight of dry solid in the supernatant}}{\text{The dry weight of the sample}} \quad \dots 8$$

2.3.2 Swelling power

The swelling power of rice flour sample was determined by measuring the water uptake of the sample (Duangrutai Thumrongchote, 2012). The 500 mg of rice flour was weighed into a centrifuge tube and 15 ml of distilled water was added. The suspension was heated in a water bath at 80°C for 30 min. and then centrifuged at 4000 rpm for 20 min. the supernatant was carefully poured into the aluminum dish (of known weight) before drying at 105°C to constant weight ~~and weighing~~. The sediment was collected and weighed. The swelling power was calculated by using the following equation

Comment [h28]: Use only last name.

$$\text{Swelling Power (g/g)} = \frac{W_{ws}}{W_f - W_t} \quad \dots 9$$

3. RESULT AND DISCUSSION

3.1 Proximate composition analysis

At the time of the experiment, The result of proximate composition of..... is presented in Table 1. The initial moisture content of the Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, were was found to be 11.31%, 13.89 % and 13.79%, (wet basis) respectively. The moisture content of the rice varieties differed significantly ($p > 0.05$). Moisture content was higher in thick and thin flaked rice as compared

Comment [h29]: Use another word. There are no superscripts.

to brown rice, Due to the soaking process, the paddy kernel absorbs water resulting in its increased moisture content. Their values were below 14% optimal values for bag storage of grains (Juliano and Villarreal, 1993). Low moisture content is known to enhance keeping the quality of rice under storage (Table 1).

The value of Crude fat content of for Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 3.20%, 0.96%, and 1.00%, respectively. The crude fat content was nearly same for thick and thin flaked rice of selected varieties and not shown a big difference, but the fat content of raw rice is higher than their value-added product and showing the significant result ($p > 0.05$). Since fat is more on the bran layer and the more this layer is removed during milling the less the fat content of the milled rice (Okaka, 2005). Higher the fat contents expose the grains to spoilage during storage due to oxidation (Table 1).

Comment [h30]: delete

The crude fiber content of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was were found to be 2.00%, 1.85%, and 1.79%, respectively. The crude protein content of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 4.23%, 4.04%, and 3.76% respectively. The nutritional value of rice depends on the total quantity and quality of protein. On the basis of crude protein content, all the varieties contained sufficient amount which was is slightly lower than the reported values of 7% (Dipti *et al.*, 2002; Dutta *et al.*, 1998; Dipti *et al.*, 2002). Crude Fiber content and pProtein contents of brown rice, thick and thin flaked rice were almost similar and did not differ significantly ($p < 0.05$) (Table 1).

Comment [h31]: delete

The ash content of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 1.00%, 1.35%, and 1.24% respectively. Ash residual is generally taken to be a measure of the mineral content of materials. High ash content in milled rice is an indication of a good quality of minerals in the rice sample (Dipti *et al.*, 2003). All the tested varieties showed high ash content which ranged from 1.05 to 1.39%. The ash content of all the tested varieties did not differ significantly ($p < 0.05$) (Table 1).

Comment [h32]: Which varieties? You mentioned one variety. Please correct the sentence.

Comment [h33]: Check your Table 1, these values are not correct.

3.2 Chemical properties of rice

The starch content of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 74.15%, 73.89%, and 73.47%

respectively. Similarly, the values of Amylose content of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 19.27%, 18.91%, and 18.26% respectively. Amylose content in brown rice was significantly ($P>0.05$) higher than thick flaked rice. Processing of paddy into flaked rice at high-temperature results in starch gelatinization with a portion of amylose getting converted into resistant starch yielding flaked rice with higher resistant starch content in the process of rice parboiling (Chitra *et al.*, 2010; Mahanta and Bhattacharya, 2010; Ibukun, 2008). The Amylopectin content in Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 79.78%, 78.80%, and 80.76% respectively (Table 2).

Table 1: Proximate composition of different varieties of brown and flaked rice.

Parameter	Chhattisgarh local paddy variety (Dokra-Dokri)		
	Brown rice	Thick flaked rice	Thin flaked rice
Moisture content (%)	11.31 \pm 0.38	13.89 \pm 0.17	13.79 \pm 0.15
Fat content (%)	3.20 \pm 0.12	0.96 \pm 0.08	1.00 \pm 0.06
Fiber content (%)	2.00 \pm 0.49	1.85 \pm 1.53	1.79 \pm 1.00
Protein content (%)	4.23 \pm 0.52	4.04 \pm 0.38	3.76 \pm 0.46
Ash content (%)	1.00 \pm 0.10	1.35 \pm 0.35	1.24 \pm 0.15

Mean \pm Standard deviation values

3.3 Functional properties of rice

The WAI and WSI of brown, thick and thin flaked rice are shown in Table 2. The water absorbing index (WAI) of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 2.19 g/g, 5.18 g/g and 5.05 g/g, respectively. The water solubility index (WSI) of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 3.25%, 4.86, and 5.09%, respectively. The WAI and WSI of flaked rice increased significantly ($p > 0.05$) during the processing of paddy. WAI of thick and thin flaked rice was found to be greater than the brown rice. Processing of paddy to flaked rice resulted in physicochemical changes of grains. Roasting at high temperature resulted in decreasing moisture content of grains that led to the dry heat gelatinization. Both roasting and flaking resulted in the damage of some starch granules leading to their enhanced water absorption

capacity. (Kumar *et al.* 2016). The swelling power (SP) of Chhattisgarh local paddy variety (Dokra-Dokri) and their value-added product thick and thin flaked rice, was found to be 6.11%, 6.51% and 6.29% respectively (Table 2).

Table 2: Chemicals and functional properties of different varieties of paddy

Parameter	Chhattisgarh local paddy variety (Dokra-Dokri)		
	Brown rice	Thick flaked rice	Thin flaked rice
Starch content (%)	74.15 ± 1.20	73.89 ± 0.96	73.47 ± 1.37
Amylose content (%)	19.27 ± 0.22	18.91 ± 0.16	18.26 ± 0.25
Amylopectin (%)	79.78 ± 0.20	78.80 ± 0.16	80.76 ± 0.25
Water absorbing index (g/g)	2.19 ± 0.24	5.18 ± 0.29	5.05 ± 0.15
Water solubility index (%)	3.25 ± 1.03	4.86 ± 1.11	5.09 ± 0.75
Swelling power (%)	6.11 ± 0.07	6.51 ± 0.04	6.29 ± 0.8

Mean ± Standard deviation values

CONCLUSION

Their proximate chemical and functional were all within acceptable levels. Therefore, the Chhattisgarh local paddy variety (Dokra-Dokri) are of good quality. This investigation into the properties of grains gives rise to a number of conclusions. This study concludes with information on proximate, chemical and functional properties of Chhattisgarh local paddy variety (Dokra-Dokri) which may be useful for developing much more better quality of flaked rice (*poha*). For developing of flaked rice it has been necessary that the flaked rice show higher or good result is some quality parameter like fat content, fiber content, protein content, starch content, water absorbing/solubility index and swelling power, etc. In this investigation we found in the case of proximate analysis, the Chhattisgarh local paddy variety (Dokra-Dokri) gives good or significant ($p > 0.05\%$) result in terms of fat content and moisture content. Similarly, investigation of chemical and functional properties we found that Chhattisgarh local paddy variety (Dokra-Dokri) gives good or significant ($p > 0.05\%$) result in terms of starch content and Amylose content. Significant ($p > 0.05\%$) difference was observed in functional properties of flaked rice with increase in water absorption index (WAI) and water solubility index (WSI).

Comment [h34]: Remove the full stop.

Comment [h35]: delete

Comment [h36]: Kindly rewrite the conclusion, base it on your findings and avoid irrelevant information.

REFERENCES

- AOAC, 2000. Official Methods of Analysis of the Association of Official Analytical Chemists; Association of Official Analytical Chemists, Inc: Washington.
- Bhattacharya, K.R., 2011. Rice quality: A guide to rice properties and analysis. Woodhead Publishing Limited, Cambridge.
- Chitra, M., Singh, V., Ali, S.Z., 2010. Effect of processing paddy on digestibility of rice starch by in vitro studies. *J. Food Sci. Technol.* 47 (4), 414–419.
- Dahare, R., Tankesh Kumar Nishad and Bhupendra Sahu. 2019. Evaluation of Some Physical and Engineering Properties of Chhattisgarh Popular Paddy Varieties for Suitability of Flaked Rice (POHA). *International Journal of Current Microbiology and Applied Sciences*. 8(03): 1506-1513.
- Dahare, R., Pisalkar, P.S., Mishra, N.K., Patel, S. And Sahu, B. 2017. Studies on physical, gravimetric and frictional properties of paddy. *Journal of agricultural issues*. 22(2): 31-35.
- Dipti, S.S, Hossain, S.T, Bari, M.N. and Kabir, K.A. 2002. Physicochemical and Cooking Properties of some fine rice varieties. *Pakistan Journal of Nutrition*. 1(4): 188-190.
- Dipti, S.S., Bari, M.N. and Kabir, K.A. 2003. Grain Quality Characteristics of some Beruin Rice Varieties of Bangladesh. *Pakistan Journal of Nutrition*. 2(4): 242-245.
- Dutta, R.K, Lihiri, B.P, and Basetmia, M.A. 1998. Characterization of Some Aromatic and Fine Rice Cultivars in Relation to their Physicochemical Quality of Grain. *Indian Journal of Plant Physiology*. 3: 61-64.
- Duangrutai Thumrongchote, Toru Suzuki, Kalaya Laohasongkram and Saiwarun Chaiwanichsiri. 2012. Properties of Non-glutinous Thai Rice Flour: Effect of rice variety. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 3(1): 153.
- Fitzgerald, M. A., McCouch, S. R., and Hall, R. D. 2009. Not just a grain of rice: the quest for quality. *Trends in Plant Science*. 14(3): 133–139.

- Juliano, B.O. and Villarreal, C.P. 1993. Grain Quality Evaluation of World Rice. P.148. International Rice Research Institute, Philippines.
- Ibukun, E.O., 2008. Effect of prolonged parboiling duration on proximate composition of rice. *Sci. Res. Essay*. 3, 323–325.
- Izawa, T. 2008. The process of rice domestication: a new model based on recent data. *Rice*. 1(2): 127–134.
- Kumar, S., Haq, R., and Prasad, K., 2016. Studies on physicochemical, functional, pasting and morphological characteristics of developed extra thin flaked rice. *Journal of Saudi Society of Agricultural Sciences*.
- Mahanta, C.L., and Bhattacharya, K.R. 2010. Relationship of starch changes to puffing expansion of parboiled rice. *J. Food Science and Technology*. 47(2): 182–187.
- Maisont, S. and Narkruga, W. 2009. Effects of Some Physicochemical Properties of Paddy Rice Varieties on Puffing Qualities by Microwave “ORIGINAL”. *Kasetsart Journal of Natural Sciences*. 43: 566 – 575.
- Okaka, J.C. 2005. Handling, Storage and Processing of Plant Foods. OCJ Publishers Enugu. pp3-5.
- Osaretin, A.T.E and Abosede, C.O. 2007. Effect of cooking and soaking on physical characteristics and sensory evaluation of indigenous and foreign rice varieties in Nigeria. *African Journal of Biotechnology*. 6(8) :1016-1020.
- Rangana, S. 1995. Hand Book of Analysis and Quality Control for Fruit and Vegetable Products. Tata McGraw-Hills Publishing Company Limited, New Delhi,
- Ray Lantin, 1999. Rice: Postharvest operations. IRRI, Philippines. 4 - 5.
- Sadasivam, S. and Manickam, A. 2005. Biochemical methods. 2nd Edition, New age International, New Delhi.
- Singh, N., Kaur, L., Sodhi, N. S., and Sekhon, K. S. 2005. Physicochemical, cooking and textural properties of milled rice from different Indian rice cultivars. *Food Chemistry*. 89(2): 253-259.

Stojceska, V., Ainsworth, P., Plunkett, A., Ibanoglu, E., Ibanoglu, S. 2008. Cauliflower by-products as a new source of dietary fiber, antioxidants, and proteins in cereal-based ready-to-eat expanded snacks. *Journal of Food Engineering*. 87: 554–563.