

## Original Research Article

# Natural Regeneration and Ecological Succession in an Urban Fragment of the Atlantic Forest in Pernambuco, Brazil

### ABSTRACT

The development of studies on natural regeneration in fragments of the Atlantic Forest **contributes assist** to evaluate the regenerative power of forests against natural and anthropic disturbances. So, the objective of the work was to analyze the structure of the regenerative component and ecological succession of arboreal species in an urban fragment of Atlantic Forest. The study was **realized undertaken** in Parque Estadual Dois Irmãos (PEDI), in Recife, Pernambuco, in which 10 subplots of 1 m x 25 m (25 m<sup>2</sup>) each were installed. As inclusion criterion, the regenerating individuals of arboreal species should have **a** height equal to or greater than one meter and circumference at the height of the chest (CAP 1.30 m) inferior to 15 cm. The individuals were classified as the ecological groups origin (native or exotic species) and were calculated **via** phytosociological parameters **such as**, heights classes for regenerating individuals and diversity index. For **data the analysis of the data the software** Mata Nativa version 4.05 was used. The families that presented the highest number of species were: Myrtaceae and Fabaceae. It was observed that 15% of the species belong to the **pioneer group of the pioneers, while 48% of the species** belonged to the **group of the** initial secondary **group**, and 22% to the **group of the** late secondary **groupones**. The distribution of individuals of the species *Hirtella racemosa* and *Chamaecrista ensiformis* in the different size and relative density classes allows us to affirm that these species act directly in the process of ecological succession. The fragment is in the initial secondary stage of succession.

**Keywords:** Dense Ombrophylous Forest; Secondary succession; Phytosociological parameters, **regeneration**.

## 1. INTRODUCTION

The Brazilian Atlantic Forest is a rainforest, considered one of the major hotspots for being one of the richest biomes in **biodiversity in the worldglobally** [1, 2], presenting approximately 14,000 different **plant** species **of plants**, [3]. However, over the past 50 years **(up to 2015)**, the original coverage of this biome has been reduced to 8% [4], **causedue byto** the advanced stage of fragmentation caused by **the** anthropogenic **activities such as ... (cite source)**.

The development of studies on natural regeneration in forest fragments is of great importance **for in** understanding the ecological functioning of these ecosystems, because they contribute to **the assessment of** the regenerative power of forests in the face of natural and anthropogenic disturbances [5] and to understand the development of forests [5, 6]. In addition, it is possible to identify and quantify the species present, as **well as also** to evaluate and monitor their distribution [7]. However, understanding the pattern of regeneration of tree species is a complex activity, because it depends on the relationship between intrinsic and extrinsic factors, linked to the physiological and ecological characteristics of the species and the **ever changing** environmental conditions [8]. The **environmental** areas that are in the

31 | process of regeneration, function as a habitat for several native species, while also acting **mainly on** carbon  
32 | sequestration agents [9]. The history of land use, present fauna in the regeneration area, physical, chemical and biological  
33 | characteristics of the soil, seed bank and proximity to native forests are the main factors that influence the rate of  
34 | regeneration [10]. Thus **are considered as** regenerating individuals are considered, those plants with their height equal to  
35 | or greater than one meter, justifying that in this phase the individuals have already adapted to the environment,  
36 | minimizing their mortality rate and facilitating the morphological characterization for the subsequent species  
37 | classification [11,12]. The number of species present in natural regeneration is directly influenced by the species that  
38 | occupy the upper strata in the forest, by the propagation of propagules, the quantity and quality of light, the type of  
39 | substrate and other growth factors [13]. Therefore, through the characterization and evaluation of natural regeneration it is  
40 | possible to predict the regenerative potential of ecosystems, acting as an important subsidy in management decisions  
41 | [14]. Cannot have all these small paragraphs focussing on regeneration – all info related to regeneration must be  
42 | contained in one paragraph only.


43 |  
44 | The weighting (amount of information supplied) between regeneration and succession is not equal – please correct.

45 |  
46 | Nothing is stated for fragmentation - cause and effect

47 |  
48 | The context, it can be said that the process of ecological succession is influenced by the morphological and physiological  
49 | characteristics of the plant species, with their interaction with other species (plants and / or animals) and through their  
50 | interaction with **the** abiotic components [15]. These factors have a strong influence on the structure, abundance of  
51 | species and community diversity [16]. Therefore, to know the forest stock and its distribution in the plant community, a  
52 | qualitative-quantitative study of natural regeneration is necessary [17]. So, knowledge of the development and temporal  
53 | dynamics of the vegetation cover is a valuable tool for landscape planning and for decisions about conservation strategies  
54 | and restoration of forest resources. This study aimed to analyze the structure of the regenerative component and  
55 | ecological succession of tree species in an urban fragment of Atlantic Forest in Recife, Pernambuco, Brazil

## 56 | 57 | 2. MATERIAL AND METHODS

58 |  
59 | The indicate year/season of study study was carried out in Parque Estadual Dois Irmãos (PEDI), located in the  
60 | metropolitan region of Recife, state of Pernambuco, **between the neighborhoods of Dois Irmãos, Apipucos, Sítio dos**  
61 | **Pintos, Macaxeira and Córrego Jenipapo**, located at the geographic coordinates 07° 59' 30" and 08° 01' 00" S and 34°  
62 | 56' 30" and 34° 57' 30" W. The forest fragment had is 384.4 hectares, going to 1,157.72 hectares unclear - how big is this  
63 | fragment?, through state decree n. 40,547 of March 28, 2014 [18]. Of the total area, 14 hectares are occupied by the zoo  
64 | (indicate name of Zoo).



67 |  
68 |  
69 | **Figure 1. Location of the Atlantic Forest fragment in Recife, Pernambuco, Brazil, with emphasis on the study**  
70 | **area.**

71 | Figure 1 – not displaying!

72 | The climate of the region is As' tropical humid coastal, with average monthly temperatures over 25.5 ° C [19]. The annual  
73 | rainfall is greater than 1,600 mm and relative **air** humidity is around 80% [20]. The vegetation cover present in the area is

a fragment of Ombrophilous Lowland Forest and the soils are constituted by Yellow Latosols, Yellow Argisols and Gleysols, whose texture varies from sandy to sandy-clay, with acid pH of 4 to 5 [21, 22, 23].

The study was carried out in the new area [define new area] of PEDI, in which 10 plots of 10 mx 25 m (250 m<sup>2</sup>) each were installed randomly. For the study of natural regeneration, a subplot of 1 m x 25 m (25 m<sup>2</sup>) was installed on the left side of each plot, identifying all the regenerating individuals inside.

As inclusion criterion, individuals of tree species with height equal to or greater than 1 meter and circumference at breast height (CAP 1.30 m) of less than 15 cm were considered regenerating. The individuals were identified, whenever possible at family and species level and, measured the heights and circumference of the height of the base (CAB 0.30 m).

When it was not possible to identify plants in the field, the reproductive and / or vegetative botanical material was collected for later taxonomic identification in the Herbário Sérgio Tavares from to de Ciência Florestal da Universidade Federal Rural de Pernambuco (DCFL/UFRPE). The species were classified into families according to the Angiosperm Phylogeny Group IV system [24]. For a description of the botanical nomenclature and their respective authors, the List of Brazilian Flora Species available in the virtual environment <http://floradobrasil.jbrj.gov.br/>.

Individuals were classified according to successional groups and origin (exotic or native species). Three height classes (H) were used for species classification [11], as presented in Table 1.

The phytosociological parameters (density, frequency and relative dominance and importance value) were calculated, the diversity was estimated according to the diversity indexes of Shannon - Weaver (H'), Pielou Equability (J) and Simpson's (C) dominance index. For the analysis of the data the software Mata Nativa version 4.05 was used.

**Table 1. Height classes for regenerating individuals.**

Classes	Height of individuals
1	1.0 ≤ H ≤ 2.0 m
2	2.0 < H ≤ 3.0 m
3	H > 3.0 e CAP < 15.0 cm

Source: [11].

### 3. RESULTS AND DISCUSSION

In the survey of the natural regeneration of the tree species [in the methodology it MUST be stated that only trees were part of the study, not shrubs] in the implanted portion [define this portion], 339 individuals belonging to 46 species and 26 families were sampled (Table 2). The number of species found in the surveys are consequences the result of a set of variables, such as site topography, soil characteristics, pluviometric indexes, degradation status and successional stage [25].

The most representative families were Myrtaceae, Fabaceae and Erythroxylaceae with 80, 39 and 30 individuals, respectively, together accounting for 43.95% of the richness. The most representative families were Myrtaceae, Fabaceae and Erythroxylaceae with 80, 39 and 30 individuals, respectively, together accounting for 43.95% of the wealth. [Duplication of the previous sentence] Two of these families of higher representativity were also found in a study of natural regeneration of native tree species in sub-forest of *Eucalyptus saligna* Smith [26]. [Discuss reason(s) for the abundance of these families] What conclusions can be drawn, what recommendations can you make?

**Table 2. Phytosociological parameters of the natural regeneration in an urban fragment of the Atlantic Forest in Recife, Pernambuco, Brazil (where: NI: Number of Individuals; RD: Relative density (%); RF: Relative frequency (%); RDo: Relative dominance (%); VI: Value of importance (%); O: Origin; N: Native; E: Exotic).**

Family/Species	NI	RD	RF	RDo	VI (%)	O
<b>Anacardiaceae</b>						
<i>Anacardium occidentale</i> L.	1	0.29	0.83	1.06	0.73	N
<i>Tapirira guianensis</i> Aubl.	2	0.59	1.65	0.78	1.01	N
<i>Thyrsodium spruceanum</i> Benth.	14	4.13	4.13	6.15	4.81	N
<i>Thyrsodium</i> sp.	1	0.29	0.83	0.04	0.39	N

**Annonaceae**

<i>Xylopia frutescens</i> Aubl.	28	8.26	6.61	8.46	7.78	N
---------------------------------	----	------	------	------	------	---

**Apocynaceae**

<i>Himatanthus bracteatus</i> (A. DC.) Woodson	7	2.06	4.13	2.59	2.93	N
--	---	------	------	------	------	---

<i>Rauwolfia</i> sp.	1	0.29	0.83	0.08	0.4	-
----------------------	---	------	------	------	-----	---

**Araliaceae**

<i>Schefflera morototoni</i> (Aubl.) Maguire et al.	1	0.29	0.83	0.16	0.43	N
---	---	------	------	------	------	---

**Burseraceae**

<i>Protium heptaphyllum</i> (Aubl.) Marchand	1	0.29	0.83	0.09	0.4	N
--	---	------	------	------	-----	---

**Celastraceae**

<i>Maytenus distichophylla</i> Mart. Ex Reissek	11	3.24	4.13	2.92	3.43	N
---	----	------	------	------	------	---

<i>Maytenus guianensis</i> Klotzsch ex Reissek	3	0.88	1.65	0.30	0.94	N
--	---	------	------	------	------	---

**Chrysobalanaceae**

<i>Hirtella racemosa</i> Lam.	41	12.09	5.79	7.38	8.42	N
-------------------------------	----	-------	------	------	------	---

<i>Licania kunthiana</i> Hook.f.	1	0.29	0.83	0.11	0.41	N
----------------------------------	---	------	------	------	------	---

**Clusiaceae**

<i>Clusia nemorosa</i> G.Mey.	8	2.36	3.31	2.89	2.85	N
-------------------------------	---	------	------	------	------	---

**Erythroxylaceae**

<i>Erythroxylum citrifolium</i> A.St.-Hil	28	8.26	5.79	4.75	6.26	N
---	----	------	------	------	------	---

<i>Erythroxylum</i> sp.	1	0.29	0.83	1.37	0.83	-
-------------------------	---	------	------	------	------	---

<i>Erythroxylum squamatum</i> Sw.	1	0.29	0.83	0.04	0.39	N
-----------------------------------	---	------	------	------	------	---

**Fabaceae**

<i>Abarema cochliacarpus</i> (Gomes) Barneby & J.W.Grimes	2	0.59	0.83	0.53	0.65	N
---	---	------	------	------	------	---

<i>Abarema filamentosa</i> (Benth.) Pittier	4	1.18	2.48	1.87	1.84	N
---	---	------	------	------	------	---

<i>Chamaecrista ensiformis</i> (Vell.) H.S.Irwin & Barneby	31	9.14	4.13	16.66	9.98	N
--	----	------	------	-------	------	---

<i>Plathymenia foliolosa</i> Benth.	1	0.29	0.83	0.60	0.57	N
-------------------------------------	---	------	------	------	------	---

<i>Pterocarpus</i> sp.	1	0.29	0.83	0.25	0.46	-
------------------------	---	------	------	------	------	---

**Hypericaceae**

<i>Vismia guianensis</i> (Aubl.) Choisy.	3	0.88	2.48	1.35	1.57	N
--	---	------	------	------	------	---

**Lacistemataceae**

<i>Lacistema robustum</i> Schnizl.	2	0.59	0.83	0.72	0.71	N
------------------------------------	---	------	------	------	------	---

**Lauraceae**

<i>Ocotea glomerata</i> (Nees) Mez	1	0.29	0.83	0.03	0.38	N
------------------------------------	---	------	------	------	------	---

**Lecythidaceae**

<i>Lecythis pisonis</i> Cambess.	2	0.59	0.83	1.07	0.83	N
----------------------------------	---	------	------	------	------	---

**Melastomataceae**

<i>Miconia albicans</i> (Sw.) Triana	1	0.29	0.83	0.18	0.44	N
--------------------------------------	---	------	------	------	------	---

<i>Miconia ciliata</i> (Rich.) DC.	7	2.06	3.31	1.29	2.22	N
------------------------------------	---	------	------	------	------	---

<i>Miconia guianensis</i> (Aubl.) Cogn	1	0.29	0.83	0.06	0.39	N
--	---	------	------	------	------	---

<i>Miconia prasina</i> (Sw.) DC.	5	1.47	2.48	1.98	1.98	N
----------------------------------	---	------	------	------	------	---

**Meliaceae**

<i>Trichillia lepidota</i> Mart.	1	0.29	0.83	0.47	0.53	N
----------------------------------	---	------	------	------	------	---

**Moraceae**

<i>Sorocea hilari</i> Gaudich.	2	0.59	0.83	0.26	0.56	N
--------------------------------	---	------	------	------	------	---

**Myrtaceae**

<i>Myrcia guianensis</i> (Aubl.) DC.	55	16.22	5.79	11.46	11.16	N
--------------------------------------	----	-------	------	-------	-------	---

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not All caps

Formatted: Font: 10 pt, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not Bold, Not All caps

Formatted: Font: 10 pt, Not All caps

<i>Myrcia splendens</i> (Sw.) DC.	10	2.95	4.13	1.18	2.75	N
<i>Myrcia sylvatica</i> (G.Mey.) DC.	8	2.36	4.13	3.19	3.23	N
<i>Myrcia tomentosa</i> (Aubl.) DC	4	1.18	2.48	1.38	1.68	N
<i>Myrciaria ferruginea</i> O.Berg.	3	0.88	2.48	0.39	1.25	N
<b>Nyctaginaceae</b>						
<i>Guapira laxa</i> (Netto) Furlan	5	1.47	0.83	1.41	1.24	N
<b>Ochnaceae</b>						
<i>Ouratea polygyna</i> Engl.	1	0.29	0.83	0.47	0.53	N
<b>Peraceae</b>						
<i>Pera ferruginea</i> (Schott) Müll. Arg.	1	0.29	0.83	0.08	0.4	N
<i>Pogonophora schomburgkiana</i> Miers ex Benth.	20	5.9	2.48	8.68	5.69	N
<b>Polygonaceae</b>						
<i>Coccoloba</i> sp.	8	2.36	3.31	1.64	2.43	-
<b>Rubiaceae</b>						
<i>Salzmannia</i> sp.	1	0.29	0.83	0.04	0.39	-
<b>Salicaceae</b>						
<i>Casearia javitensis</i> Kunth	5	1.47	2.48	2.48	2.14	N
<b>Sapindaceae</b>						
<i>Allophylus edulis</i> (A.St.-Hil. Et al.) Hieron. Ex Niederl	2	0.59	1.65	0.28	0.84	N
<i>Cupania racemosa</i> (Vell.) Radlk	2	0.59	0.83	0.85	0.76	N
<b>Total</b>	<b>339</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>-</b>

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150


151

In relation to the phytosociological structure, the seven species of greatest Importance Value were *Myrcia guianensis*, *Chamaecrista ensiformis*, *Hirtella racemosa*, *Xylopia frutescens*, *Erythroxylum citrifolium*, *Pogonophora schomburgkiana* e *Thyrsodium spruceanum*, which together made up 54.1% of the total **Value of Importance (VI)** (Table 2). **[Discuss reason(s) for the abundance of these species]**. The representativeness of these species was also highlighted in other regeneration works, such as that of **Santiago et al.** [27] in a secondary forest fragment of the Botanic Garden of UFJF, **[insert author name]**[28] in a fragment of the Ombrophilous Dense Forest of Terras Baixas and **[insert author name]**[29] in a fragment of Atlantic forest in the city of Igarassu - PE. **Thus, the results of this work have demonstrated** that most of these species are able **tofor colonization and development in** of the PEDI fragment. **What characteristic makes them to be successful? What conclusions can be drawn in terms of veld management, what management recommendations can be make?**

The families that presented the highest number of species, **orderly** in descending order were: Myrtaceae and Fabaceae with five species, Anacardiaceae and Melastomataceae with four species, Erythroxylaceae with three species, Apocynaceae, Peraceae, Celastraceae and Chrysobalanaceae with two species each, together they represented 78.47% of advanced **(secondary?)** natural regeneration. These families are among the most important **found** in fragments of Atlantic Forests and corroborate **with** the results of other works carried out in the Atlantic Forest [1, 30, 31]. The importance of the Fabaceae family is emphasized, since it ensures productivity in most terrestrial ecosystems, due to their performance in nitrogen fixation. **What about the other families, what makes them to be important?**

The three species that have the highest VI stand out due to the high values of density and dominance in the area, *Myrcia guianensis* was the one with the highest values for all the estimated parameters. **[Possible reasons?]** The ecophysiological and environmental characteristics in which the species are inserted, some may not reach large diameters [32]. **[Explain this statement – why not?]**


Among the species raised in the study, the nine species that presented the number of individuals greater or equal to ten were: *Myrcia guianensis*, *Hirtella racemosa*, *Chamaecrista ensiformis*, *Xylopia frutescens*, *Erythroxylum citrifolium*, *Pogonophora schomburgkiana*, *Thyrsodium spruceanum*, *Maytenus distichophylla* and *Myrcia splendens* (Figure 2).



152  
153  
154 **Figure 2. Regenerating species with numbers of individuals equal to or greater than 10 in an urban fragment of**  
155 **Atlantic Forest in Recife, Pernambuco, Brazil.**

156  
157 All are native species of the Atlantic Forest biome, and among them five species are initial secondary. The nine species  
158 together represent 67.55% of all regenerating individuals in the area and are the most homogeneous species, since they  
159 were sampled in almost all plots. What makes these species to be successful in the study area?

160  
161 Regarding the successional classification of species, it is observed that 47.8348% belong to the group of the initial  
162 secondary, 21.7422% to the late secondary group and 15.22% to the pioneer group, equivalent to 22, 10 and 7 species,  
163 respectively (Figure 3). Of the total sampled, 7 species were not classified [give reason for not classifying them],  
164 representing 13%. The successional stage of a forest fragment is indicated by the successional group that presents a  
165 percentage greater than 50% of individuals [33].  
166  
167




168  
169 **Figure 3. Percentage of successional groups observed in an urban fragment of Atlantic Forest in Recife,**  
170 **Pernambuco, Brazil (being: PI: Pioneers, IS: Initial secondary, LS: Late secondary, NC: No classification).**

171  
172 The greater number of initial secondary species may be related, since the area does not present clearings, hindering the  
173 permanence of the pioneer species, seeing that they need the solar incidence to survive. Combine these 4 small (1  
174 sentence each) paragraphs into one larger paragraph – a paragraph cannot consist of one sentence only.

175  
176 The greatest cause of precedence begins with the fact that the forest fragment has several years of natural regeneration,  
177 may not be able to progress to other successional phases, or this process is occurring slowly [34].  
178

179 The increase of the initial secondary species in a forest fragment may indicate what occurred previously in that site **some**  
180 **as either** fragmentation, distribution and / or disturbance [35].  
181

182 In relation to regeneration by height class, the percentages were: 35.10%, 30.38% and 34.51%, for classes 1, 2 and 3,  
 183 respectively. Of the 46 species sampled, 15 were found in the three classes, 10 species in two classes and 21 species  
 184 only in one height class (Figure 4). DISCUSS THIS RESULT – WHAT ARE THE IMPLICATIONS FOR SUCCESSION?



187  
 188  
 189 **Figure 4. Relation of the 10 species with the highest values of total natural regeneration of the sampled**  
 190 **population, expressed as a percentage, sampled in an urban fragment of Mata Atlântica in Recife, Pernambuco,**  
 191 **Brazil (being: NRC1: Natural regeneration in Class 1; NRC2: Natural regeneration in Class 2; NRC3: Natural**  
 192 **regeneration in Class 3; TNR: Total natural regeneration).**

194 Among the species collected in this study, the ones with the highest percentages of Natural Regeneration in Class 1 were:  
 195 *Myrcia guianensis* (5.90%), *Erythroxylum citrifolium* (5.60%), *Hirtella racemosa* (4.72%) and *Xylopia frutescens*  
 196 (2.36%). In Class 2 of height were: *Myrcia guianensis* (6.49%), *Hirtella racemosa* (5.31%), *Chamaecrista ensiformis*  
 197 (3.83%) and *Xylopia frutescens* (2.36%). While in Class 3 height, the following stand out: *Chamaecrista ensiformis*  
 198 (4.13%), *Myrcia guianensis* (3.83%), *Xylopia frutescens* (3.54%) and *Pogonophora schomburgkiana* (3.24%).  
 199 DISCUSS THIS RESULT – WHAT ARE THE IMPLICATIONS FOR SUCCESSION?

200 Of the total sampled species, 27 presented total natural regeneration values lower than 1.0, with the passage of time  
 201 these species may present greater difficulty to regenerate or may occur the establishment of late species in these  
 202 ecosystems [36]. The occurrence of low regeneration of these species may be related to environmental conditions, mainly  
 203 by the closure of the canopy that will provide better conditions to the species that are tolerant to shading [37], for these  
 204 reasons it is necessary to monitor these species in the long term [5].

205 Regarding diversity, the Shannon-Weaver index for regenerating individuals was 3.04 nats.ind-1 exhibiting high diversity  
 206 in the area, a value similar to that found by [38] with 3.01 in an area of restoration with planting in high diversity. The  
 207 greater diversity of these individuals indicates that the process of restoration of that area is occurring properly, because it  
 208 is expected that in the initial phases an increase in diversity occurs with the occurrence of the establishment of new  
 209 species [39]. With this, it can be seen that a community that presents high diversity is directly related to its richness [40].  
 210 DISCUSS THIS RESULT – WHAT ARE THE IMPLICATIONS FOR SUCCESSION? The Pielou equability was 0.79  
 211 indicating that the species found are nicely distributed. The value was lower than that found by [41] with 0.81 in the  
 212 Atlantic Forest fragment and similar to that found by [27] in the Semi-deciduous Seasonal Forest area with 0.74. These  
 213 results are common for fragments of Atlantic forests in the state of Pernambuco [5]. The Simpson dominance index was  
 214 0.93 indicating that this area has a high diversity. The diversity indexes are considered high for the secondary formations  
 215 of Atlantic Forest, showing that these formations are well conserved [42].

216  
 217  
 218  
 219 **4. CONCLUSION SHORT COMMUNICATION PAPERS DO NOT AS A RULE HAVE CONCLUSION**  
 220 **SECTION – RATHER INCORPORATE THESE INSIGHTS INTO THE ABOVE TEXT.**

221 **The most representative families were: Myrtaceae, Fabaceae and Anacardiaceae.**  
 222  
 223 **The distribution of the individuals of *Hirtella racemosa* and *Chamaecrista ensiformis* in the different size and**  
 224 **relative density classes, allows to affirm that these species act directly in the process of ecological succession.**  
 225

226  
227 **From the obtained results, it can be inferred that the fragment is in the initial secondary stage of succession.**

228  
229 **Species that present low natural regeneration may, over time, be replaced by other species as the changes in the**  
230 **forest composition of that site occur. Therefore, it is necessary to have a continuous monitoring of these species**  
231 **in the new area of Parque Estadual Dois Irmãos.**

## 232 233 **COMPETING INTERESTS**

234  
235 Authors have declared that no competing interests exist.

## 236 237 **REFERENCES**

- 238  
239  
240 1. Oliveira LSB, Marangon LC, Feliciano ALP, Cardoso MO, Silva VF. Floristic, successional classification and dispersion  
241 syndromes in a remnant of Atlantic Forest, Moreno - PE. Brazilian Journal of Agricultural Sciences. 2011; 6 (3): 502-507.  
242 DOI: <http://dx.doi.org/10.5039/agraria.v6i3a1384>
- 243 2. Eisenlohr PVLF, Alves LC, Bernacci MCG, Padfurschi RB, Torres BEM, Silver FAM, Santos MA et al. Disturbances,  
244 elevation, topography and spatial proximity drive vegetation patterns along an altitudinal gradient of a top biodiversity  
245 hotspot. Biodiversity Conservation. 2013; 22 (12): 2767-2783.
- 246 3. Forzza RC, Baumgratz JFA, Bicudo CEM, Canals DAL, Carvalho-Junior AA, Rabbit MAN, Costa AF et al. New Brazilian  
247 floristic list highlights conservation challenges. Bioscience. 2012; 62 (1): 39-45.
- 248 4. Lucas EJ, Bünge. Myrtaceae in the Atlantic forest: their role as a "model" group. Biodiversity and Conservation. 2015;  
249 24 (9): 2165-2180. DOI: <http://dx.doi.org/10.1007/s10531-015-0992-7>
- 250 5. Aparicio WCS, Marangon LC, Ferreira RLC, Aparício OS, Costa Júnior RF. Structure of natural regeneration of tree  
251 species in a fragment of Atlantic Forest, Pernambuco. Brazilian Journal of Agricultural Sciences. 2011; 6 (3): 483-488.  
252 DOI: <http://dx.doi.org/10.5039/agraria.v6i3a791>
- 253 6. Souto MAG, Boeger MRT. Structure and composition of the regeneration stratum and associated vegetation of  
254 different successional stages in eastern Paraná. Forest Science. 2011; 21 (3): 393-406. DOI:  
255 <http://dx.doi.org/10.5902/198050983798>.
- 256 7. Range JRV, Botelho SA, Bentes-Range MM. Floristic composition and structure of the natural regeneration of Lower  
257 Várzea Secondary Forest in the Amazon Estuary. Tree Review. 2002; 26 (5): 559-556.
- 258 8. Santos KF, Ferreira TS, Higuchi P, Silva AC, Vandresen PB, Costa A et al. Natural regeneration of the arboreal  
259 component after mortality of a taquara massif in a fragment of mixed ombrophylous forest in Lages - SC. Forest Science.  
260 2015; 25 (1): 107-117.
- 261 9. Chazdon RL. Regeneration of tropical forests. Bulletin of the Museu Paraense Emílio Goeldi. Natural Sciences. 2012; 7  
262 (3): 195-238
- 263 10. Chadzon RL, Letcher SG, Van Breugel M, Martínez-Ramos M, Bongers F, Finegan B. Rates of change in tree  
264 communities of secondary Neotropical forests following major disturbances. Philosophical Transactions of the Royal  
265 Society. 2007; 362 (1): 273 – 289. DOI:  
266 <https://dx.doi.org/10.1098%2Frstb.2006.1990><https://dx.doi.org/10.1098%2Frstb.2006.1990>



- 267 11. Marangon LC, Soares JJ, Feliciano ALP, Brandão CFLS. Natural regeneration in a fragment of Seasonal Semideciduous  
268 Forest in Viçosa, Minas Gerais. *Tree Review*. 2008; 32 (1): 183-191.
- 269 12. Silva SO, Ferreira RLC, Silva JAA, Lira MA, Alves-Júnior FT, Cano MOO, Torres JEL. Natural regeneration in a remnant  
270 of caatinga with different use histories in Pernambuco. *Tree Review*. 2012; 36 (3): 441-450. DOI:  
271 <http://dx.doi.org/10.1590/S0100-67622012000300006>
- 272 13. Pinto JRR, Oliveira-Filho AT, Hay JDV. Influence of soil and topography on the composition of a tree community in  
273 Central Brazilian valley forest. *Edinburgh Journal of Botany*. 2005; 61 (1): 69-90.
- 274 14. Silva WC, Marangon LC, Ferreira RLC, Feliciano ALP, Costa Júnior RF. Study of the natural regeneration of tree species  
275 in fragments of Ombrophilous Dense Forest, Mata das Galinhas, in the municipality of Catende, Zona Sul Mata de  
276 Pernambuco. *Forest Science*. 2007; 17 (4): 321-331.
- 277 15. Pinto AVF, Silva MAM, Leite AVL, Nascimento LM, Lins e Silva ACB, Wheel MJN. Floristic and structure of the arboreal  
278 Community of a regeneration fragment of Atlantic Forest, Igarassu, Pernambuco, Brazil. *Revista Agro @ mbiental*. 2018;  
279 12 (2): 145-155.
- 280 16. Gurevitch J, Scheiner SM, Fox GA. *Plant Ecology*. 2nd ed. Porto Alegre: Artmed, 2009. 592 p.
- 281 17. Garcia CC, MGF Kings, Reis GG, Pezzopane JEM, Lopes HNS, Ramos DC. Natural regeneration of arboreal species in a  
282 semideciduous Montana seasonal forest fragment, in the Atlantic Forest domain, in Viçosa, MG. *Forest Science*. 2011; 21  
283 (4): 677-688.
- 284 18. Moura CHR, Bezerra OG, Silva JM. The natural values of the conservation units of Recife: Mata de Dois Irmãos and  
285 Mata do Engenho Uchôa. *NEMO*. 2018; 10 (1): 131-155
- 286 19. Köppen W. Translation Corrêa ACB. *Geographic System of Climates. Notes and Communication of Geography - Series*  
287 *B: Didactic texts nº 13*. Ed. University - UFPE, Department of Geographical Sciences, 1996. 31 p.
- 288 20. Moreira EBM, Galvêncio JD. Spatialization of surface temperatures in the city of Recife using UTM images -  
289 LANNSAT 7. *Revista de Geografia*. 2007; 24 (1): 101-115.
- 290 21. Jacomine PKT et al. Exploratory survey: soil recognition in the state of Pernambuco, Recife. Ministry of Agriculture /  
291 Northeast Development Superintendence. (Technical Bulletin nº 26; Pedological series nº 14). 1973. 359 p.
- 292 22. Machado IC, Lopes AV, Porto KC. Ecological Reserve of Dois Irmãos: studies on a remnant of Atlantic Forest in an  
293 urban area. Recife: University Publishing House of UFPE. 1998. 326 p.
- 294 23. Lima MS, Freire FJ, Marangon LC, Almeida BG, Ribeiro EP, Santos RL. Forest soils in a fragment of urban forest in the  
295 Dois Irmãos Forest, Recife, Pernambuco, Brazil. *Floresal Science*. 2018; 28 (2): 542-555.
- 296 24. APG IV. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants:  
297 APG IV. *Botanical Journal of the Linnean Society*. 2016.
- 298 25. Rodal MJN, Sampaio EVSB, Figueiredo MA. *Manual on methods of floristic and phytosociological study: caatinga*  
299 *ecosystem*. Brasília: Botanical Society of Brazil. 1992.

- 300 26. Alencar AL, Marangon LC, Feliciano ALP, Ferreira RLC, Teixeira LJ. Advanced natural regeneration of native tree  
301 species in the understory of Eucalyptus saligna Smith stands, in the Zona Sul Mata de Pernambuco. *Forest Science*. 2011;  
302 2 (2): 183-192.
- 303 27. Santiago DS, Fonseca CR, Carvalho FA. Phytosociology of the natural regeneration of an urban fragment of the  
304 Semideciduous Seasonal Forest (Juiz de Fora, MG). *Brazilian Journal of Agricultural Sciences*. 2014; 9 (1): 117-123.
- 305 28. Silva RKS, Feliciano ALP, Marangon LC, Lima RBA. Floristic and ecological succession of the arboreal vegetation in  
306 spring area of a fragment of Atlantic Forest, Pernambuco. *Brazilian Journal of Agricultural Sciences*. 2010; 5 (4): 550-559.
- 307 29. Rocha KD, Chaves LFC, Marangon LC, Silva ACBL. Characterization of adult arboreal vegetation in a fragment of  
308 Atlantic forest, Igarassu - PE. *Brazilian Journal of Agricultural Sciences*. 2007; 3 (1): 35-40.
- 309 30. Silva-Júnior JF, Marangon LC, Ferreira RLC, Feliciano ALP, Brandão CFLS, Alves-Júnior FT. Phytosociology of the tree  
310 component in a remnant of Atlantic Forest in the municipality of Cabo de Santo Agostinho, PE. *Brazilian Journal of*  
311 *Agricultural Sciences*. 2008; 3 (3): 276-282.
- 312 31. Brandão CFLS, Marangon LC, Ferreira RLC, Silva ACBL. Phytosociological structure and successional classification of  
313 the tree component in a fragment of Atlantic forest in Igarassu - Pernambuco. *Brazilian Journal of Agricultural Sciences*.  
314 2009; 4 (1): 55-61.
- 315 32. Imaña-Encinas J, Santana OA, Macedo LA, Paula JE. Diametric distribution of a section of the Semideciduous Seasonal  
316 Forest in the area of the Cerrado Ecomuseum. *Cerne*. 2008; 14 (1): 33-45.
- 317 33. Budowski G. The distinction between old secondary and climax species in tropical central American lowland  
318 rainforest. *Tropical Ecology*. 1970; 11 (1): 44-48.
- 319 34. Fonseca CR, Carvalho FA. Floristic and phytosociological aspects of the tree community of an urban fragment of  
320 Atlantic forest (Juiz de Fora, MG, Brazil). *Bioscience Journal*. 2012; 28 (5): 820-832.
- 321 35. Lima RBA, Marangon LC, Freire FJ, Feliciano ALP, Silva RKS. Regenerative potential of arboreal species in a fragment  
322 of Atlantic Forest, Pernambuco, Brazil. *Green Magazine on Agroecology and Sustainable Development*. 2017; 12 (4): 666-  
323 673.
- 324 36. Silva WC, Marangon LC, Ferreira RLC, Feliciano ALP, Aparício PS, Costa Júnior RF. Horizontal and vertical structure of  
325 arboreal component in natural regeneration phase of Santa Luzia forest, in the municipality of Carende - PE. *Tree*  
326 *Review*. 2010; 34 (5): 57-66.
- 327 37. Higuchi P, Reis GG, Pinheiro AL, Silva CT, Oliveira CHR. Floristic composition of the natural regeneration of tree  
328 species over a period of eight years in a semideciduous seasonal forest fragment, in Viçosa, MG. *Tree Review*. 2006; 30  
329 (6): 893-904.
- 330 38. Trentin BE, Estevan DA, Rossetto EFS, Gorenstein MR, Brizola GP, Bechara FC. Forest restoration in the Atlantic  
331 forest: passive, nucleation and high diversity planting. *Forest Science*. 2018; 28 (1): 160-174.
- 332 39. Rech CCC, Silva AC, Higuchi P, Schimalski MB, Pscheidt F, Schmit AB, Ansolin RD et al. Evaluation of the forest  
333 restoration of a degraded APP in Santa Catarina. *Forest and Environment*. 2015; 22 (2): 194-203.
- 334 40. Sizenando-Filho FA, Maracajá PB, Diniz-Filho ET, Freitas RAC. Floristic and phytosociological study of the herbaceous  
335 flora of the municipality of Messias Targino, RN / PB. *Journal of Biology and Earth Sciences*. 2007; 7 (2).

- 336 41. Lima RBA, Silva RKS, Paula MD, Guimarães ETR, Braga ECB. Phytosociological and diametric structure of a fragment  
337 of Atlantic forest, Pernambuco, Brazil. *Challenges Magazine*. 2017; 4 (4): 143-153.
- 338 42. Lopes CGR, Ferraz EMN, Araújo EL. Physiological characterization of a fragment of Atlantic Forest in the municipality  
339 of São Vicente Férrer, PE, Brazil. *Brazilian Journal of Bioscience*. 2007; 5 (2): 1174-1176
- 340

UNDER PEER REVIEW