

## Original Research Article

# Occurrence of different kinds of diseases in sesame cultivation in Myanmar and their impact to sesame yield

### ABSTRACT

We surveyed diseases of sesame in 10 farmers' fields at Nay Pyi Taw, Myanmar and did interviews with 25 farmers for the occurrence of diseases and its impact on yield in Magway, the major sesame growing area in Myanmar. We found phyllody, charcoal rot (root and stem rot), *Alternaria* leaf blight, powdery mildew, and leaf curl, based on on-site symptoms and their microscopic observation in Nay Pyi Taw. The disease incidence ranged from 5% to 30% in phyllody, from 10% to 30% in charcoal rot (root and stem rot) and 10% to 40% in *Alternaria* blight, while leaf curl and powdery mildew were not observed abundantly. According to interviews conducted in Magway, 60% of the farmers suffered from phyllody disease symptoms, 80% from charcoal rot, 48% from *Cercospora*, 28% bacterial leaf spot and 24% diseases with the symptoms of leaf roll. Most farmers (84%) noticed combinations of diseases symptoms either phyllody or charcoal rot/black and stem rot or *Cercospora* leaf spot and/or bacterial leaf spot. Yield losses ranged from 10 to 50% by phyllody, from 15 to 100% by charcoal rot (root and stem rot) and from 0 to 50% by leaf spots. Other abnormal symptoms such as discoloring of root, seedling death and leaf yellowing were also observed. There were no significance relations between the actual yield and yield losses estimated by each disease. A half of farmers (54%) burnt the crop residues after harvest, while 45% directly buried them in their fields including disease infected plant parts in the soil. Although there was no difference in sesame yield between these two practices, the average yield was higher by 15% in farmers with the burnt practice. Only a few farmers applied fungicides. Potential constraints to cause yield reduction and necessary actions to increase sesame yield are discussed.

**Keywords:** *Alternaria*, charcoal rot, control measures, disease incidence, phyllody, yield loss,

### 1. INTRODUCTION

Sesame (*Sesamum indicum* L.) is the major oilseed crop in Myanmar for both export and domestic consumption. Myanmar stands the second largest producers of sesame in the world [1]. Sesame is cultivated two times a year: May to August as the monsoon crop and September or October to December as the cool season cultivation. The total sesame sown area in 2017-2018 was 1.6 million ha, in which 829,000 ton of sesame was produced with the yield of  $0.54 \text{ t ha}^{-1}$  [2]. The sesame yield in Myanmar is low compared with that in other major sesame producing countries such as  $1.41 \text{ t ha}^{-1}$  in Chia,  $0.90 \text{ t ha}^{-1}$  in Bangladesh,  $0.78 \text{ t ha}^{-1}$  in Ethiopia [1]. In Myanmar, 80% of the sown area is within the cereals zone, which comprises parts of Mandalay, Sagaing and Magway regions and makes up the major growing area for oilseed crops.

Sesame is damaged by a number of fungal, bacterial and viral diseases, such as *Cercosporas sesamei*, *Cylindrosporium sesamei*, *Alternaria seasami*, *Pseudomonas sesamei*, *Macrophomina phaseoli*, *Phytophthora parasitica*, *Oidium* spp., *Erysiphe circhoracearum*, *Phytophthora nicotianae* var. *sesame* and *Helminthosporium sesamiare* [3]. Considerable yield losses due to diseases have been reported; yield losses of 45% by the powdery mildew, *Luveillutataurica*, 55%-90% by phyllody disease caused by phytoplasma [4,5], 5-100% by *M. phasiolina*, 22-53% by

Comment [U1]: I guess China

30 Cercospora leaf spot [6]. A disease complex of powdery mildew with dry root rot disease, which is caused by the fungi  
31 *Macrophomina* and *Oidium*, results in low productivity of sesame in India [7].

32 In Myanmar, different symptoms in sesame cultivation have been reported by the Department of Plant Pathology, Yezin  
33 Agricultural University: phyllody disease, virescence, yellowing floral sterility and stem proliferation in infected plants,  
34 *Alternaria* leaf blight, dark brown rounded to irregular lesions on leaves, powdery mildew, dirty white powder. However, in-  
35 depth investigations have not been undertaken in Myanmar until now and there is no sufficient information on the major  
36 threat of sesame cultivation, making it difficult to develop suitable disease control measures. Available information in  
37 Myanmar is only the studies by Win et al. [8] who worked on phyllody disease and by Wai et al. (2007)[9] who did varietal  
38 screening against charcoal rot disease caused by *M.phaseolina*.

Comment [U2]: Be consistent in the citation

39 In order to manage diseases in the field, it is necessary to know the incidence and distribution of each sesame disease  
40 and their impact on production. In Myanmar, sesame has been cultivated by small holders' farmers with minimum inputs.  
41 Therefore, expanding knowledge of sesame diseases to local farmers is very important for the development of effective  
42 disease control tactics in sustainable agriculture. The aims of this study were to observe the prevalence of sesame  
43 diseases and evaluate its control practices by small holders' farmers. In this study, we firstly identified different diseases  
44 through microscopic examination of specimens collected from sesame fields in Nay Pyi Taw and then evaluated their  
45 incidence. Then, we interviewed on disease occurrence, impact on sesame yield and managements practiced with small  
46 holder's farmers in Magway regions, one of the major sesame growing areas.

Comment [U3]: Delete 's'

## 48 2. MATERIALS AND METHODS

### 49 2.1 Disease Specimen Collection

51 Specimens were collected from four different sesame fields at the flowering time of sesame, in the middle of June, 2016 in  
52 Yezin, Pynmana located in Nay Pyi Taw areas. The appearance of aboveground disease symptoms which were likely to  
53 be charcoal rot, *Alternaria* blight, powdery mildew, phyllody and leaf curl was carefully observed and recorded in each  
54 field with photographs of the disease symptoms. Then, the specimens collected were put into a plastic bag for  
55 microscopic examination and brought to the laboratory.

#### 56 2.1.1 Microscopic Examination

57 Microscopic examination was done in the laboratory of Plant Pathology, Yezin Agricultural University. Specimens were  
58 collected from symptoms on the fresh leaf specimens with dark brown color round to irregular zonate lesions, dirty white  
59 powdery substance on leaves and black dots on stems and carefully checked for the presence of spores and spore  
60 fruiting body of pathogens under low and high power objective lens (10x and 40x) [10].

61 For charcoal rot disease, firstly the bark of diseased stems of sesame was peeled off and placed onto a drop of sterilized  
62 water on a slide and checked under microscope. For mycelium observation, a drop of sterilized water was placed on a  
63 clean glass slide and the bark of diseased stems of sesame was peeled off and cut into small pieces and surface  
64 sterilized with 95% ethyl alcohol for 30 seconds and transferred into 10% sodium hypochloride-hypochlorite for 1 minute.  
65 Then, the specimens were washed into sterilized water for 3 times and finally the pieces were dried and transferred onto  
66 water agar and incubate at room temperature. After 3 days of incubation, the mycelium formation of  
67 *Macrophominaphaseolina* name was confirmed.

68 For *Alternaria* leaf blight, a drop of sterilized water was placed on a clean glass slide. A small amount of fungal spores  
69 taken from the dark brown irregular lesions on the leaves was placed into a drop of water drop and covered with a cover  
70 slip. Morphological characteristics of conidia and conidiophores of the causal organism were checked and then identified.

71 For powdery mildew, a drop of sterilized water was placed on a clean glass slide. Three to five pieces of thin cross section  
72 leaves collected from plant parts showing dirty whitish fungal patches symptoms were cut with sterilized razor blade and  
73 put in a water drop on the slide and covered with a cover slip. Shape, color, spore size of conidia and conidiophore were  
74 observed under microscope.

#### 75 2.1.2 Occurrence of Sesame Diseases in Nay Pyi Taw


76 Total 10 sesame fields were surveyed for the occurrence of phyllody, charcoal rot, *Alternaria* leaf blight in northern part of  
77 Nay Pyi Taw area in 2016. Out of 10 fields, 4 fields from Pynmanar, 3 from Pobbathri and 3 from Zeyathiri township were  
78 selected. Aboveground symptoms, in particular appearance of each disease were carefully checked across the field

Comment [U4]: You previously mentioned four (4) fields were visited. Please, reconcile

79 (diagonal). For each disease, plants were randomly checked across the field from 7m apart (14 x14 cm row spacing) and  
80 recorded the percentage of infested plants based on the visual examination. Disease incidence was calculated on the  
81 basis of per cent plant infected in total plant populations. Symptoms of charcoal rot/root and stem rot disease; black dots  
82 on the stems, Alternaria leaf blight, powdery mildew, phyllody, leaf curl were found out in several sesame fields as shown  
83 in plates (1, 2, 3 and 4). Except for phyllody and leaf curl disease, microscopic examination of above ground symptoms of  
84 charcoal rot/root and stem rot, Alternaria blight and powdery mildews were checked using leaves and stems samples. The  
85 presence of pathogen spores: pycnidia and pycnidiospores of *M.phaseolina* on the bark of sesame stems (Plate 1),  
86 Alternaria spore when fungal spores taken from black color lesions on leaves (Plate 2) and pathogens of *Oidium* with  
87 hyaline barrel shaped spore after making cross section of white powdery mildews appearance on sesame leaves (Plate 3)  
88 were observed.

## 89 2.2 Evaluation of Awareness of Sesame Diseases and Control Measures by Smallholder Farmers in 90 Magway

91 The large proportion of farmers is engaged in sesame production throughout the year in Magway Township. Photographs  
92 were prepared for typical disease symptoms of Charcoal rot, phyllody, Alternaria leaf blight, leaf curl, Cercospora leaf spot  
93 and bacterial leaf spot with their causal agents. Using the photographs, a disease questionnaires' survey was conducted in  
94 three villages in Magway Township in the harvesting period, August, 2016. Survey was done in a total of 25 farmers on  
95 their knowledge of sesame diseases. If the disease symptoms were noticed in their fields, effect on sesame yield and  
96 control measures were interviewed. Additional information on farm experience, cultivation practices, inputs and  
97 constraints on sesame production and management practices was also collected.



98 Plate 1. (A) Elongated brownish dark  
99 lesions on stem, (B) Pycnidia formation on bark of sesame stem, (C) Pycnidia and pycnidiospores of  
100 *Macrophomina phaseolina* (40x), (D) Hypha of *Rhizoctonia bataticola* (40x).




102 Plate 2. (A) Dark brown irregular lesions with concentric ring on sesame leaves, (B) Conidia of *Alternaria sesami*  
103 taken from dark brown lesions leaves (40x)  
104

105  
106 **3. RESULTS**

107  
108 **3.1 Occurrence of Sesame Diseases and Its Incidence in Nay Pyi Taw**

109 Disease symptoms of phyllody, charcoal rot and Alternaria blight were most frequently observed in all survey fields. The  
110 disease incidence (DI) % of phyllody, charcoal rot (stem rot), and Alternaria blight ranged from 5%-30%, 10%-30% and  
111 10%-40%, respectively (Table 1). Symptoms of leaf curl, powdery mildew was not abundantly observed in Nay Pyi Taw  
112 area and their incidence were not assessed.



117  
118 **Plate 3. (A) Dirty white podwery on sesame, (B) Conidia and conidiophores of *Oidium* spp. (40x) after cross**  
119 **section of leaves with dirty white spot**



**Plate 4. Transformation of flora parts into green leafy structure**

**Table 1. Occurrence of sesame diseases and their incidence (%) in Nay Pyi Taw area**

Field No.	Phyllody disease	Black stem rot	Alternaria leaf blight
1	0	20	10
2	20	15	30
3	10	10	20
4	5	25	15
5	10	30	5
6	20	25	20
7	30	15	30
8	20	10	40
9	10	20	30
10	15	30	25
Avg.	14	20	22.5

130  
131 **3.2 Observation of sesame diseases and control measures by smallholder farmers in Magway region**

132 **3.2.1 Cultivation Practice and Cropping Pattern**

Formatted: Font: (Default) Helvetica, Not Bold

Formatted: Normal

Formatted: Right

P


P

133 In Magway region, sesame has been rotated with pulses for more than 20 years and is the major income source for small  
 134 scale farmers. It is grown as a single crop or intercropped with pigeon pea. Most of farmers used sources of sesame  
 135 seeds from their local market while others purchased seeds from the Department of Agriculture (DoA) in Magway  
 136 township. Cowdung-cow dung manure and basal fertilizers were applied during land preparation. Insecticides to control for  
 137 leaf roll were sprayed within 15-45 days after sowing if necessary. Although the recommended rate of balanced fertilizers  
 138 was 125 kg urea (58 kg N) ha<sup>-1</sup>, 125 kg triple super phosphate (23.5 kg P) ha<sup>-1</sup> and 62 kg potash (32.5 kg K) ha<sup>-1</sup> for  
 139 sesame production, the practical application rates varied among the farmers.


140 It was noticed that 84% of the farmers (21 out of 25) cultivated sesame alone in the monsoon cultivation and 16% of  
 141 farmers did sesame with pigeon pea or other kinds of pulses. Black sesame (Sinyadar 13) was the most popular variety  
 142 with yields of 245-735 kg ha<sup>-1</sup>.

### 143 3.3 Incidence of Sesame Diseases and Yield Loss

144 More than a half of farmers (60%, 15 out of 25) suffered phyllody disease symptoms in their fields, 80% suffered from  
 145 charcoal rot, 48% from Cercospora, 28% from bacterial leaf spot and 24% from diseases with the symptoms of leaf roll  
 146 (Fig. 1). Most farmers (84%) answered that their sesame fields suffered combinations of disease symptoms either  
 147 phyllody or charcoal rot/black and stem rot or Cercospora leaf spot and/or bacterial leaf spot (Fig. 2). All disease  
 148 symptoms were apparently noticed by farmers 30 days onwards after sowing. In case of phyllody, yield losses ranged  
 149 from 10-50%, while stem rot symptoms brought yield losses ranging from 15 to 100% and little yield reduction to up 50%  
 150 yield losses were observed by the symptoms of leaf  
 151 spots.



158  
 159 **Fig. 1. Percentage of farmers suffering from sesame diseases**  
 160 **Results of interviewing with a total of 25 farmers in Magway**



164  
**Fig. 2. Ratio (%) of farmers (total 25) who suffered from sesame diseases complex**


167 No one answered that they noticed the symptoms  
 168 appearance of powdery mildew and Alternaria leaf blight. Instead, some farmers additionally informed their experiences in  
 169 the occurrence of abnormal symptoms showing root red and seedling death, root rot, and leaf yellowing around 20 to 40  
 170 days after sowing and which seemed to reduce sesame yield by 10 to 40% if these symptoms were observed in their  
 171 fields. No significance relation was found between the actual yield and yield loss estimated for each disease; phyllody,  
 172 charcoal rot, Cercospora, bacterial leaf spot and other diseases answered by farmers in Magway (Fig. 3a,b,c,d,e).

### 173 3.4 Actions Taken for Sesame Diseases by Farmers in Magway Region and Control Measures

174 In survey areas, 54% of farmers burned sesame crop residues while 45% farmers directly buried them in the soil including  
 175 disease infected plant parts. Although there was no difference in sesame yield between these two practices, the average  
 176 yield was higher by 15% in farmers with the burnt practice. Only a few farmers applied fungicides such as carbendazim for  
 177 charcoal rot/black rot of stem and amidachloroit for phyllody diseases to the field.

#### 179 4. DISCUSSION

180  
 181 The present study revealed that sesame cultivation in Nay Pyi Taw and Magway, Myanmar was damaged from mixtures  
 182 of various fungal, bacterial and viral diseases that are caused by *Candidatus Phytoplasma*, *Macrophomina*, *Alternaria*,  
 183 *Oidium*, *Cercospora* and *Xanthomonas* etc. Among the diseases, phyllody and charcoal rot diseases were prominent in  
 184 both areas. Farmers noticed most of the disease symptoms around one month after sowing. According to Gupta et al. [6],  
 185 sesame phyllody disease caused by *Phytoplasma* can be prevalent throughout the year and is transmitted by the insect  
 186 vector (*Orosius albicinctus*). Although their study did not mention disease incidence and yield losses, the present study  
 187 showed disease incidence of phyllody ranging from 5-30% and yield losses by up to 50% in Magway.



188  
 189 **Fig. 3. Estimated yield loss due to (a) Phyllody, (b) Charcoal rot, (c) Cercospora leaf spot, (d) Bacterial leaf spot,**  
 190 **(e) Other diseases by farmers and their actual sesame yield in Magway**

191 In general, root rot caused by *M. phaseolina* is the most devastating disease in many growing countries such as India and  
 192 Pakistan [11,12,13]. Murugesan et al. [14] reported that 1% increase in the incidence of *M. phaseolina* reduced seed yield  
 193 by 1.8 kg ha<sup>-1</sup>. In this study, 30% of disease incidence was recorded by charcoal rot disease and almost all the farmers  
 194 experienced in a yield reduction of up to 50% to 100% when they noticed charcoal rot symptoms in their fields, indicating  
 195 that the disease incidence of *M. phaseolina* and its yield losses would be the largest in sesame cultivation.

196 We did not identify the actual disease incidence and severity of each disease in farmer fields in Magway, instead, we  
 197 asked farmers for their yield and knowledge on sesame diseases. Most of the farmers in Myanmar are living in rural areas  
 198 and outskirts and thus understanding of their perceptions and knowledge on constraints of sesame production are very  
 199 important to analyze the impact of diseases on sesame yield and efficiency of their control measures. However, it was  
 200 difficult to conclude that estimation of yield loss caused by each disease was directly related with their actual yield in this  
 201 study, because there were no significant relations between actual sesame yield and yield losses caused by each disease.

Comment [U5]: Re-write this statement

202 Low yields have been attributed to several factors, e.g. variety, agronomic practices, soil salinity, poor drainage, poor  
203 planting methods (broadcasting), weeds, diseases and insect pests [4,15]. One or several of these reasons could  
204 probably affect low yields in Myanmar. Most of the farmers seemed to lack in the knowledge on the proper use of fertilizers  
205 throughout the crop growing season. Recently, the presence of pigeon pea cyst nematode *Heterodera cajani* has been  
206 confirmed in sesame cultivated fields in Magway (unpublished data). Therefore, improper fertilization and/or the nematode  
207 might cause damage and thus yield reduction. Further investigation of densities of cyst nematodes in sesame cultivated  
208 fields relation to yield is now under planning.

209 Although farmers noticed abnormal symptoms, sometimes seriously ones, in their fields, most farmers did not take an  
210 action. This is because farmers did not either pay attention to diseases or know effective control measures. At present,  
211 chemical fungicides are the first choice for farmers to combat diseases because of their easy applicability and immediate  
212 therapy [7]. In the present study, farmers did not clearly recognize the name of chemicals even though they used.  
213 Fungicides still play a vital role in the control of the disease due to the lacks of a high quality of seeds and proper  
214 sanitation in their fields.

Comment [U6]: Misspelt

#### 215 4. CONCLUSION AND RECOMMENDATION

216  
217 Based on the microscopic examination of symptoms appearance, diseases of charcoal rot, Alternaria blight, powdery  
218 mildew and their causal agents were detected. Limited knowledge on sesame diseases was noticed among farmers  
219 except on charcoal rot disease (locally called 'Yoeme'). There is a need for more studies on crop loss assessment by  
220 individual diseases in Myanmar. The use of chemical fungicide is the only way control to practice so far. The interest of  
221 farmers in managing infested crop residues and enough knowledge on proper application of using fungicides are lacking.  
222 Therefore, proper trainings or extension service for sesame diseases and awareness of fungicide use should be provided  
223 to those rural areas by various organizations. The research attention to developing holistic disease control measures (use  
224 of the resistant variety, biocontrol agents, and proper cultural practices) for the improvement of sesame yield qualitatively  
225 and quantitatively should be investigated as of its being a vital role in earning foreign income.

#### 226 227 228 229 COMPETING INTERESTS

230  
231 Authors have declared that no competing interests exist.

#### 232 233 234 REFERENCES

- 235 1. FAOSTAT (2017) <http://www.fao.org/faostat/en/#data/QC>
- 236 2. MOALI. Myanmar Agriculture in Brief. Ministry of Agriculture, Livestock and Irrigation, Nay Pyi Taw,  
237 Myanmar.2018.
- 238 3. Weiss EA. 2000.Oilseed Crops. 2<sup>nd</sup> edition. Blackwell Science, UK.
- 239 4. Egonyu JP, Kyamanywa S, Anyanga W, Ssekabembe CK. Review of pests and diseases of sesame in Uganda.  
240 African Crop Conference Proceedings. 2005;7:1411-1416.
- 241 5. Gopal K, Jagadeswar R, Prasad G. Evaluation of sesame (*Sesamum indicum*) genotypes for their reaction to  
242 powdery mildew and phyllody diseases. Plant Disease Research. 2005;20(2):126-130.
- 243 6. Gupta KN, Naik, KRR, Bisen R. Status of sesame diseases and their integrated management using indigenous  
244 practices. International Journal of Chemical Studies. 2018;6(2):1945-1952.
- 245 7. Jealakshmi C, Rettinassababady C, Nema S. Integrated management of sesame diseases. Journal of  
246 Biopesticides.2003;691:68-70.
- 247 8. Win NKK, Back C-G, Jung H-Y. Phyllody phytoplasma infecting sesame (*Sesamum indicum*) in Myanmar. Tropical  
248 Plant Pathology. 2010;35(5):310-313.DOI: 10.1590/S1982-56762010000500006

- 249 9. Wai KPP. Pycnidium formation in vitro of *Macrophominaphaseolina* (*Rhizoctoniabataticola*), causal agent of stem  
250 and root rot of sesame (*Sesame indicum* L.), and reaction of some sesame varieties to the disease. Thesis, Plant  
251 Pathology Department, Yezin Agricultural University. 2007.
- 252 10. Yu KH. Methods in plant pathology and integrated disease management. Lecture book, Department of Plant  
253 Pathology, Yezin Agricultural University, Myanmar. 2013.
- 254 11. Choudhary CS, Arun A, Prasad SM. Management of stem and root rot of sesame. International Journal of  
255 Agricultural Sciences. 2014;10(2):755-760.
- 256 12. Meena B, Indiragandhi P, Ushakumari R. Screening of sesame (*Sesamum indicum* L.) germplasm against major  
257 diseases. Journal of Pharmacognosy and Phytochemistry. 2018;7(1S):1466-1468.
- 258 13. Bedawy IMA, Moharm MHA. Reaction and performance of some sesame genotypes for resistance to  
259 *Macrophominaphaseolina*, the incitant of charcoal rot disease. Alexandria Science Exchange Journal.  
260 2019;40(1):12-18.
- 261 14. Murugesan M, Shanmugam MM, Menon PP, Arokiaraj A, Dhamu, KP and Kochubabu M. Statistical assessment  
262 of yield loss of sesamum due to insect pests and diseases. Madras Agricultural Journal. 1978;65:290-295.
- 263 15. El-Bramawy MASA, Abd Al-Wahid OA. Evaluation of resistance of selected sesame (*Sesamum indicum*)  
264 genotypes to Fusarium wilt disease caused by *Fusarium oxysporum* f. sp. *sesami*. Tunisian Journal of Plant  
265 Protection. 2009;4(1):29-39.  
266