

# Resistance of *Malpighia emarginata* genotypes to *Meloidogyne enterolobii* parasitism

## ABSTRACT

**Objective:** Considering the inexistence of *M. emarginata* cultivars resistant to *M. enterolobii* available for cultivation, and the scarcity of information about the severity of its parasitism in *M. emarginata*, the present study investigated the response of genotypes from the active germplasm bank of Universidade Federal Rural de Pernambuco to *M. enterolobii* parasitism, aiming the selection of resistant genotypes for use as rootstocks for commercial varieties.

**Study design:** The experimental design was completely randomized, with 21 genotypes and one independent matrix (control), with six replicates each. The experimental unit was represented by one plant per plot.

**Place and Duration of Study:** Department of Agronomy, Universidade Federal Rural de Pernambuco – UFRPE - Brazil between June 2013 and July 2014.

**Methods:** In the experiment a completely randomized design was adopted, with 21 genotypes from the AGB and one as a control for susceptibility. The *M. emarginata* cuttings were inoculated with 10,000 nematode eggs, and after 150 days, they were evaluated for the following parameters: egg mass index, gall index, reproduction factor, number of eggs per gram of root, number of eggs per root system.

**Results:** Twenty out of the twenty-two genotypes analyzed were susceptible. The genotypes 021-CMF and 037-CMF were considered resistant. To our knowledge, this is the first identification of *M. emarginata* genotypes resistant to *M. enterolobii*.

**Conclusions:** These results are of great importance for the breeding and cultivation of the species, since these two genotypes can be indicated for use as rootstocks and for breeding programs aimed at transferring resistance to other cultivars with desirable production characteristics that are susceptible to the phytonematode.

**Keywords:** *Acerola*, *Brabados cherry*, *root-knot nematode*, *rootstocks*.

**Comment [A1]:** Minimum keywords should be five and meaningful.

## 1. INTRODUCTION

*Malpighia emarginata*, also known as Barbados cherry and Acerola, has been gaining space in the fruit-growing sector due to its high amount of vitamin C. Its cultivation has shown a great potential for expansion, whether for fresh consumption, juice industry or for the pharmaceutical industry [1]. Although it is a rustic plant, *M. emarginata* is sensitive to root-knot nematodes, which are its main limiting factor, negatively influencing the production and the quality of fruits. In the last years, orchards have shown a considerable decrease in production due to these phytonematodes [2]. One of the main species that has been shown to be very harmful to the crop is the *Meloidogyne enterolobii* Yang & Eisenback, described in China in 1983 [3,4].

The main symptoms caused by the attack of this phytonematodes are the small, deformed and yellowing leaves; delay and reduction in the seedlings development, and in cases of high infestations, poor plant development and declining production may occur [5-8].

A survey carried out in several irrigated perimeters in the São Francisco Valley Region, in northwestern Brazil, revealed a high percentage of infected *M. emarginata* trees, raising the concern that this phytonematode may turn the cultivation unfeasible, as it has happened to guava orchards in that same region [7-9]. Genotypes with resistance or tolerance to phytonematodes may be used as rootstocks, as a low cost and sustainable alternative to chemical control methods and, can easily be adopted by growers without environmental and sanitary risks [9-11]. The use of resistant rootstocks could provide an effective control and significantly reduce the damages caused by *M. enterolobii*, allowing the recovery of infested

**Comment [A2]:** The full plant authority should be mentioned when discussing about the plant material. In this regard, in each case that you aim to discuss about your plant, it should be reported for example as: *Borago officinalis* L. (not *Borago officinalis*).

53 areas[12]. The knowledge of how *M. emarginata* genotypes respond to *M. enterolobii* infection  
54 is also of great importance, since in perennial crops the management of these organisms is  
55 even more difficult. Therefore, for new orchards, it is essential to choose non-infested areas or  
56 the use of resistant genotypes [2]. Thus, in this work we evaluated 21 different *M. emarginata*  
57 genotypes from the active germplasm bank of UFRPE, aiming at the selection and indication of  
58 genotypes resistant to the phytonematode *M. enterolobii*.  
59

## 60 2. MATERIAL AND METHODS

61 The experiment was conducted in a greenhouse located at the Agronomy Department of the  
62 Universidade Federal Rural de Pernambuco – UFRPE - Brazil. Twenty-two *M. emarginata*  
63 genotypes were utilized, of which twenty-one belong to the Active Bank of Germplasm (AGB) of  
64 the Carpina Sugarcane Experimental Station (E.E.C.A.C./UFRPE), located in the municipality of  
65 Carpina – PE and the variety BRS Sertaneja, which was selected due to its susceptibility to *M.*  
66 *enterolobii* [2,11].

### 67 2.1 Collection and propagation of the plant material

68 Semi-woody cuttings with three nodes and two pairs of leaves were obtained from the evaluated  
69 genotypes which were between thirteen and fifteen years old. The cuttings were planted in a  
70 mini-tunnel containing commercial Brasplant® substrate; the depth of planting was 1/3 of the  
71 length of the stake. In order to maintain humidity, the mini tunnel was covered with transparent  
72 white plastic and a 50% shade was used for shading. Irrigation was performed daily early in the  
73 morning and late in the afternoon.

### 74 2.2 Inoculum source

75 *M. enterolobii* inoculums were obtained from Embrapa Semiárido - CPATSA - Petrolina, PE,  
76 and kept in tomato and multiplied in tomato plants (*Solanum lycopersicon* L.), lineage 684,  
77 known as resistant to *M. incognita* and *M. javanica* [7]. Two months after the inoculation, the  
78 tomato roots were carefully removed from the substrate, then washed and cut into 1-2cm  
79 segments. Eggs were then extracted according to the technique described by [13]. After  
80 obtaining the nematode egg suspension, eggs were counted from 1 mL samples on photon  
81 microscopes and the concentration of the suspension was adjusted to 1000 eggs/mL using  
82 distilled water.

83  
84  
85  
86  
87

### 88 2.3 Evaluation of genotypes for resistance to *M. enterolobii*

89  
90 Sixty-day seedlings were transplanted to 10 L plastic pots containing commercial Brasplant®  
91 substrate and placed in a greenhouse. During the conduction of the experiment the average  
92 temperature was  $26 \pm 2$  ° C and relative humidity of  $65 \pm 5\%$ . Water irrigation was performed  
93 daily in the early morning and late afternoon; irrigation and fertilization with was done weekly  
94 [14].

95 After 120 days of planting, inoculation was done at a concentration of 10,000 eggs/plant. The  
96 egg suspension was deposited in three small holes in the soil around the plant's neck, with an  
97 automatic graduated pipette. 150 days after inoculation, the root system of each genotype was  
98 carefully washed and evaluated according to the following parameters: gall index (GI) and egg  
99 mass index (EMI), both determined according to the scale proposed by [15].

100 Subsequently, the eggs were extracted following the methodology described by Hussey and  
101 Barker (1973) [13]. The number of eggs per root system (NER) was estimated with a photonic  
102 microscope. In addition, the number of eggs per gram of root (NEGR) and reproduction factor  
103 (RF), obtained by the ratio of the final number of eggs to the initial number of inoculated eggs,  
104 were also estimated [16].

105 With the RF, the highest value was taken as the susceptibility standard and, from this, the  
106 percentages of RF reduction was obtained by the formula:  $(RRF) = \frac{Frp-Frt}{Frp} \times 100$  were

107 calculated, where: Frp = reduction in the reproduction factor in the standard and Frt =  
 108 reproduction factor in the treatment [16]. According to Moura and Regis (1987)[17], it is possible  
 109 to classify genotypes for resistance or susceptibility by considering RRF values. Thus, RRF = 0-  
 110 25 (Highly Susceptible-HS); RFR = 26-50 (Susceptible-S); RFR = 51-75 (Little Resistant-LR);  
 111 RRF = 76-95 (Moderately Resistant-RM); RRF = 96-99 (Resistant-R); RRF = 100 (Highly  
 112 Resistant-HR or Immune-I). Relative weight of the shoots (RWS), the relative weight of the roots  
 113 (RWR), as well as the shoots dry biomass (SDB) were also calculated.

114

## 115 2.4 Experimental Design

116

117 The experimental design was completely randomized, with 21 genotypes and one independent  
 118 matrix (control), with six replicates each. The experimental unit was represented by one plant  
 119 per plot.

120 Analysis of variance was performed using the Sisvar software. The data was transformed into  
 121 log, for the variable number of eggs per root grass, in square root of x, for number of eggs per  
 122 root system and the reproduction factor, the other variables did not undergo any transformation.  
 123 Subsequently, the means were compared by the Scott-Knott test, at 5% probability.

124

## 125 3. RESULTS

126 Five months after inoculation with *M. enterolobii*, root galls and egg masses were detected in all  
 127 inoculated plants. Significant differences based on the analysis of variance were found by the  
 128 Scott-Knott test to the following variables: GI, EMI, RF, NER, NEGR, RWS, RWR and SDB  
 129 (Table 1).

130

131 **Table 1. Analysis of variance for the resistance indexes of *M. emarginata* to the parasitism of *M.***  
 132 ***enterolobii***

	DF	MS							
		GI	EMI	RF	NER	NEGR	RWS	RWR	SDB
Genotypes	21	1.14**	1.68**	4.05**	1928.40**	728.28**	425.04**	787.57**	146.56**
Residual	110	0.11	0.14	0.78	70.99	204.77	48.97	94.72	10.02
CV (%)		17.65	21.75	48.21	48.14	49.43	16.23	24.30	18.75

133 *DF*: Degrees of freedom; *MS*: mean square; *GI*: gall index; *EMI*: egg mass index; *RF*: reproduction factor; *NER*: number of  
 134 eggs per root system; *NEGR*: Number of eggs per root grain; *RWS*: Relative weight of the shoots; *RWR*: Relative weight of  
 135 the roots; *SDB*: Shoots dry biomass; *Cv*: coefficient of variation. \*\*  $p < 0,05$  by the scott-knott test.

136

137 The lowest values for the GI and EMI variables and according to the criterion of Sasser (1980)  
 138 [18] were observed in 018-CMF and 37-CMF genotypes (Table 2).

139

140

141

142 **Table 2. *M. emarginata* genotypes response in relation to the parasitism of *M. enterolobii***

Genotypes	GI	EMI	SR	RF	RRF	DR
SERTANEJA	4,7 <sup>1</sup>	4,2 <sup>1</sup>	S <sup>1</sup>	8,50 <sup>1</sup>	41,13	S
002-SPE	5	5	S	10,34	28,39	S
016-CMF	4,2	4	S	1,78	87,67	MR
017-CMF	4	3,8	S	3,42	76,32	MR
018-CMF	1,3	0,8	R	1,30	91,00	MR
021-CMF	2,7	2	S	0,56	96,12	R

022-CMF	3	2,7	S	4,56	68,42	LR
023-CMF	5	4,5	S	2,38	83,52	MR
024-CMF	2,8	1,8	S	1,34	90,72	MR
025-CMF	3,8	2,8	S	2,46	82,96	MR
028-CMF	4,3	4,3	S	2,89	79,99	MR
033-CMF	5	5	S	14,44	-	S
036-CMF	4,8	4,3	S	2,04	85,87	MR
037-CMF	1	0,4	R	0,50	96,54	R
038-CMF	4,8	4,3	S	5,97	58,66	LR
039-CMF	4,6	3,8	S	2,77	80,82	MR
040-CMF	3	2	S	2,43	83,17	MR
041-CMF	4,4	3,4	S	6,98	51,66	LR
042-CMF	4,7	4,5	S	7,27	49,65	S
043-UFRPE	4,2	2,8	S	2,30	84,07	MR
044-APE	5	4,7	S	10,50	27,29	S
045-APE	5	5	S	7,77	46,19	S

143 <sup>1</sup>mean value for the six replicates; <sup>2</sup> negative values compared to the control; GI= 0 to 5 according to  
144 Sasser (1980); EMI= Egg Mass Index (0-5); SR= Susceptibility reaction: S= susceptible (IG≥3); R=  
145 resistant (IG≤3); RF= Reproduction; RRF = reduction in the reproductive factor compared to the control  
146 DR= Differential Reaction: HS= highly susceptible; S = susceptible; LR= low resistance; MR= moderately  
147 resistant.  
148

149 The observed RF values ranged from 0.50 (037-CMF) to 14.44 (033-CMF) (Table 2). The  
150 variable RFR represents how much each genotype differed in its RF in relation to the most  
151 susceptible genotype. The 033-CMF was the most susceptible genotype observed, with RF  
152 even greater than the susceptibility control, the Sertaneja cultivar. The highest percentages  
153 were obtained by the genotypes 21 and 37 which obtained a RRF of 96.12 and 96.54,  
154 respectively, and they were classified as resistant (R) (Table 2).

155 Regarding the amount of eggs per root gram (ERG) and the amount of eggs per root system  
156 (ERS), the genotypes 21 and 37 were characterized by the lowest values for (15.55 and  
157 11.77) and (64.73 and 64.6), resulting in promising genotypes regarding resistance (Table  
158 3).  
159

160 **Table 3. Reaction of *M. emarginata* genotypes to *M. enterolobii*, for the indicator variables of**  
161 **susceptibility evaluated**

Genotypes	NEGR <sup>1</sup>	NER <sup>2</sup>	RWR (g)	RWS (g)	SDB (g)
Sertaneja	41.19 b	263.74 b	39.01 b	42.00 b	16.89 b
002-SPE	46.31 b	331.19 b	47.00 c	46.66 b	19.66 c
016-CMF	17.86 a	124.63 a	48.94 c	41.33 a	15.66 b
017-CMF	27.81 a	175.68 a	42.54 c	56.66 d	25.34 d
018-CMF	19.18 a	93.71 a	21.97 a	36.66 a	11.94 a

021-CMF	15.55 a	64.73 a	25.66 a	37.66 a	11.66 a
022-CMF	38.54 a	201.69 b	29.22 a	33.33 a	12.95 a
023-CMF	24.92 a	149.52 a	39.10 b	45.74 b	15.86 b
024-CMF	20.51 a	109.45 a	30.66 a	37.33 a	11.33 a
025-CMF	18.48 a	130.57 a	48.28 c	46.66 b	19.72 c
028-CMF	27.38 a	161.67 a	34.66 b	40.66 a	15.33 b
033-CMF	42.13 b	357.32 b	73.00 d	61.66 d	23.00 c
036-CMF	23.82 a	136.87 a	32.54 b	51.24 c	24.35 d
037-CMF	11.77 a	64.6 a	25.11 a	37.83 a	15.81 b
038-CMF	37.63 b	223.86 b	33.85 b	36.33 a	11.66 a
039-CMF	17.84 a	143.37 a	55.94 c	36.60 a	14.51 b
040-CMF	20.12 a	137.69 a	42.29 c	48.00 b	21.64 c
041-CMF	36.05 b	247.94 b	45.65 c	59.33 d	28.01 d
042-CMF	39.79 b	235.71 b	37.66 b	38.33 a	14.00 b
043-UFRPE	18.91 a	130.98 a	46.60 c	40.00 a	17.27 b
044-APE	44.13 b	297.23 b	45.33 c	39.00 a	12.00 a
045-APE	46.88 b	276.33 b	36.01 b	36.77 a	12.79 a

MEGR= Number of eggs per grain of root and NER= number of eggs per root system; RWR= Relative weight of the roots; RWS = Relative weight of the shoots; SDB= Shoots dry biomass. <sup>1</sup>log<sub>x</sub> turned variables, <sup>2</sup>√x turned variables. <sup>x</sup> average values followed by the same letter in the columns do not differ by the scott-knott test at the 5% probability.

165  
166  
167  
168

#### 4. DISCUSSION

169 The concept of resistance used in plant nematology describes the ability of a given plant to  
170 suppress the development and reproduction, or even the infection process of a nematode [19].  
171 To detect the root-knot nematodes, the symptoms can be evaluated with ease, but it is common  
172 for symptoms caused by these parasites to be confused with physiological problems such as  
173 nutritional deficiency and hydric stress, or even with other pests and diseases [20], and is also  
174 common for some plant species the absence of any apparent symptoms, despite the infection of  
175 its roots, therefore, the term resistance is also used to describe the capacity of a host to  
176 suppress the disease [19, 21]. The Gall index and the degree of galling may be used to  
177 measure the ability of a plant to lessen or overcome the attack by the root-knot nematode.  
178 However, these indexes do not indicate the occurrence of nematode reproduction directly, while  
179 the reproduction factor, is a variable that allows the direct measurement of the nematode's  
180 reproductive capacity in the host [22]. The GI is usually used in germplasm tests to address the  
181 type of host reaction and the percentage of reduction of the parasite's reproduction rate in  
182 relation to the most susceptible cultivar, allowing the epidemiological characterization of the  
183 nematode-host interaction [2, 17, 23].  
184 Considering only the GI criterion, 018-CMF and 37-CMF genotypes could be classified as  
185 resistant (GI < 2), but the evaluation of nematode parasitism resistance based solely the  
186 development of galls may lead to inaccurate results due to the potential subjectivity and

Comment [A3]: Correct word

187 empiricism of the counting methodology [24]. In this study, the genotypes exhibited an  
188 expressive variation in their susceptibility to the phytonematode considering the RF criterion.  
189 Genotypes 21-CMF and 37-CMF, in addition to exhibiting low values in relation to the GI, EMI,  
190 RWS and RWR parameters, presented RF <1, with values of 0.56 and 0.50 respectively (Table  
191 1), therefore being indicated as rootstocks resistant to *M. enterolobii* parasitism. The genotype  
192 018-CMF despite being considered resistant by the GI criteria, exhibited a RF of 1.3 and RFR of  
193 91, and therefore classified only as moderately resistant.

194 Regarding the RWS and RWR variables (Table 3), it was verified that there was a significant  
195 difference between the studied genotypes. The genotypes 017-CMF and 033-CMF exhibited the  
196 higher RWS values in relation to the others showing good development of the shoots even  
197 when parasite by *M. enterolobii*. The genotype 033-CMF showed the higher value of RWR, and  
198 also the higher GI and RF values, being the most susceptible of the observed genotypes. [11]  
199 evaluated the responses of eleven UFRPE-AGB *M. emarginata* genotypes to the parasitism of  
200 *M. enterolobii*. Regarding the variables RWS and RWR, the authors verified a significant  
201 difference only for the 028-CMF genotype, which exhibited the higher shoot and roots  
202 development of the evaluated genotypes and was classified as moderately resistant. For the  
203 SDB variable, the highest values were observed for genotypes 017-CMF, 036-CMF and 041-  
204 CMF, which were classified little or moderately resistant to *M. enterolobii*. Considering another  
205 species of the *Meloidogyne* genus, [10] did not find significant difference for the parameters  
206 RWR and RWS, in *M. emarginata* UFRPE AGB genotypes parasited by *M. incognita*.

207 The evaluation of RWS and RWR may contribute to the selection of tolerant genotypes, to root-  
208 knot nematodes, since the absorption and distribution of the nutrients are highly related to the  
209 growth rate of the plants, and may be impaired by parasites in the root system [25], but only the  
210 observation of developmental characteristics are not sufficient to the determination of resistance  
211 or long term tolerance to these parasites.

## 213 5. CONCLUSIONS

- 214 1. The genotypes 021-CMF and 037-CMF were resistant to *M. enterolobii* and could be  
215 indicated as rootstocks.  
216 2. The genotype 033-CMF is indicated as susceptibility control to *M. enterolobii*  
217 parasitism, exhibiting higher values of RF than the commercial variety Sertaneja.

## 218 6. COMPETING INTERESTS

219 Authors have declared that no competing interests exist.  
220  
221  
222

## 223 7. REFERENCES

- 224  
225  
226 1. Almeida JPN, Dantas LGR. Fungo micorrízico arbuscular e extrato de algas no  
227 crescimento inicial de porta-enxerto de aceroleira. Revista de Ciências Agrárias. 2014;  
228 57(1):22-28. <http://dx.doi.org/10.4322/rca.2013.061>.  
229 2. Cavichioli JC, Garcia MJM, Brida AL & Wilcken SRS. Reação de aceroleira (*Malpighia*  
230 *emarginata* D.C.) à *Meloidogyne enterolobii*. Rev. Bras. Frutic. 2014; 36(1):156-160.  
231 <http://dx.doi.org/10.1590/0100-2945-429/13>.  
232 3. Hunt DJ, Handoo ZA. Taxonomy, identification and principal species. In: Moens M,  
233 Perry RN e Starr JL, editors. Root-Knot Nematodes. 1 st ed. Wallingford. CAB  
234 International; 2009.  
235 4. Moens M, Perry RN, Starr JL. *Meloidogyne* species - a diverse group of novel and  
236 important species. In: Moens, M.; Perry RN, Starr JL (ed) Root-Knot Nematodes.  
237 Wallingford. CAB International; 2009.  
238 5. Ritzinger R, Noronha ACS, Farias ARN, Ritzinger CHSP, Nascimento AS. Pragas em  
239 viveiro de mudas de aceroleira. Embrapa Mandioca e Fruticultura Tropical; 2006.  
240 (Acerola em Foco, 12).  
241 6. Castellano G, Quijada O, Jiménez N, Crozzoli R, Hernández V, Marin RC. Reacción de  
242 cultivares de cerecita (*Malpighia glabra*) a *Meloidogyne enterolobii* (nematoda:

Comment [A4]: Conclusion is too short

- 243 *Meloidogynidae*. Fitopatol. venez. 2011;24(1):25-27.  
244 <http://www.ovefit.com.ve/boletines/24-1/DOC5.pdf>.
- 245 7. Martins LSS, Musser RS, Souza AG, Resende LV, Maluf WR. Parasitismo de  
246 *Meloidogyne enterolobii* em espécies de myrtaceae. Rev. Bras. Frutic. 2013;35(2):477-  
247 484. <http://dx.doi.org/10.1590/S0100-29452013000200017>.
- 248 8. Souza AG, Resende LV, Lima IP, Martins LSS, Techio VH. Chromosome number and  
249 nuclear DNA amount in *Psidium* spp. resistant and susceptible to *Meloidogyne*  
250 *enterolobii* and its relation with compatibility between rootstocks and commercial  
251 varieties of guava tree. Plant Syst Evol. 2015;301(1):231-237.  
252 <https://doi.org/10.1007/s00606-014-1068-y>.
- 253 9. Souza FF, Deon MDI, Castro JMC, Lima MAC, Ribka ACP, Freitas ST. Principais  
254 Variedades de Aceroleiras Cultivadas no Submédio do Vale do São Francisco.  
255 *Documentos online* 255, 2013; 1:21. Accessed 28 September 2014. Available:  
256 <http://www.infoteca.cnptia.embrapa.br/infoteca/handle/doc/982090>.
- 257 10. Rossiter JGA, Musser RS, Martins LSS, Pedrosa EMR, Medeiros JE. Seleção de  
258 genótipos de aceroleira assistida por marcadores isoenzimáticos visando à resistência  
259 a *Meloidogyne incognita* raça 2. Rev. Bras. Frutic. 2008;30(4):1057-1064.  
260 <https://dx.doi.org/10.1590/S0100-29452008000400036>
- 261 11. Moreira AA, Martins LSS, Musser RS, Moraes Filho RS, Maranhão W, et al. Response  
262 of *Malpighia emarginata* active germplasm bank accessions to *Meloidogyne enterolobii*  
263 parasitism. Gen. Mol. Res. 2016;15:4. <http://dx.doi.org/10.4238/gmr15048868>.
- 264 12. Freitas VM. Resistência genética de goiabeira e reação de espécies frutíferas visando  
265 o manejo de *Meloidogyne enterolobii*. 92f. Ph.D. Thesis (Faculdade de Agronomia e  
266 Medicina Veterinária). Universidade de Brasília; 2012.
- 267 13. Hussey RS, Barker KRA. Comparison of methods collecting inocula of *Meloidogyne*  
268 spp. including a new technique. Plant Disease Reporter, 1973;57:1025-1028.
- 269 14. Hoagland DR, Arnon DI. The water culture method of growing plants without soil.  
270 Berkeley: University of California. 1950;347:2-32.
- 271 15. Taylor AL, Sasse JN. Biology, identification and control of root-knot nematodes  
272 (*Meloidogyne* species). Raleigh: International *Meloidogyne* Project. NCSU & USAID  
273 Coop.Plub. 1978:111.
- 274 16. Salgado SML, Resende MLV, Campos VP. Reprodução de *Meloidogyne exigua* em  
275 cultivares de cafeeiros resistentes e suscetíveis. Fitopatol. Bras. 2005;30:413-415.
- 276 17. Moura RM, Régis EMO. Reações de cultivares de feijoeiro comum (*Phaseolus vulgaris*)  
277 em relação ao parasitismo do *Meloidogyne javanica* e *M. incognita*. Brasília: Nematol.  
278 Bras.1987;11:215-225.
- 279 18. Sasser JN Root-knot nematodes: a global menace to crop production. Plant Dis.  
280 1980;64(1):36-41.
- 281 19. Roberts PA. Concepts and consequences of resistance. In: Starr JL, Cook R, Bridge J,  
282 editors. Plant Resistance to Parasitic Nematodes. Wallingford: CAB International; 2002.
- 283 20. RITZINGER CHSP, RITZINGER R. Nematoides em acerola. Cruz das Almas: Embrapa  
284 Mandioca e Fruticultura, 2002. (Embrapa Mandioca e Fruticultura. Acerola em Foco, 2).
- 285 21. Sasser JN, Carter CC, Hartman KM. Standardization of Host Suitability Studies and  
286 Reporting of Resistance to Root-knot Nematodes. North Carolina State University,  
287 Raleigh and United States Agency for International Development; 1984.
- 288 22. Dong W, Holbrook CC, Timper P, Brennenman TB, Mullinix BG. Comparison of Methods  
289 for Assessing Resistance to *Meloidogyne arenaria*. In: Peanut. J. Nematol.  
290 2007;39(2):169-75.
- 291 23. Santos JLF. (2017). Levantamento de nematóides-das-galhas e identificação de fontes  
292 de resistência a *Meloidogyne enterolobii* em aceroleira no submédio do Vale do São  
293 Francisco. 2017. Master's Dissertation (Mestrado em Agronomia – Produção Vegetal) –  
294 Universidade Federal do Vale do São Francisco, Campus Ciências Agrárias, Petrolina-  
295 PE; 2017.
- 296 24. Costa Filho JH. Coleta e reação de acessos de melancia a *Meloidogyne enterolobii*.  
297 Master's thesis (Fitotecnia), Universidade Federal Rural do Semi-Árido, Mossoró; 2012.
- 298 25. Hillocks RJ, Thresh JM, Bellotti AC. Cassava: biology, production, and utilization.  
299 London: CAB International; 2002.