

Short communication

Effects of gamma rays on cocona (*Solanum sessiliflorum*)

Comment [A1]: The full plant authority should be mentioned when discussing about the plant material. In this regard, in each case that you aim to discuss about your plant, it should be reported for example as: *Borago officinalis* L. (not *Borago officinalis*).

ABSTRACT

Aims: The present work aims to assess the gamma rays effects on cocona (*Solanum sessiliflorum*).

Study design: Cocona seeds of CUB-08 genotype were exposed to gamma rays (0, 100, 150, 200, 300 and 400 Gy) and sowed in styrofoam seedling trays. Seedlings were then planted, in the open field, three months later, following randomized block design with four blocks and seven plants per plot.

Place and Duration of Study: The present study was developed in Instituto Nacional de Pesquisas da Amazônia at the Agricultural experimental station, which is located on Km 14 AM-10 roadway, from April 2016 to December 2016

Methodology: Fruits were harvested from November to December 2016. Fruit characteristics such as yield, pilosity (1=No pilosity and 3=Excessive pilosity), weight, diameter, number per plant, and ripening after seven days (1=firm; 3=rotten) were assessed. Furthermore, stem pilosity was assessed, as well.

Results: Results showed 200 Gy dose to decrease the germinating vigor, whereas 300 and 400 Gy were highly deleterious and prevented seeds from germinating. In contrast, 100 and 150 Gy doses did not affect germinating vigor, but just created fruit pilosity variability.

Conclusion: The gamma ray-150 dose may be used to enhance this species' genetic diversity.

Keywords: *cubiu*; irradiation; mutation; plant breeding

Comment [A2]: Minimum keywords should be five and meaningful

1. INTRODUCTION

Cocona (*Solanum sessiliflorum*) is a traditional Amazonian fruit, its flavor is like a unknown citrus mixture. This fruit displays pilosity, which is a typical characteristic of this species, making it harder to be harvested on account of the itching it inflicts onto the pickers' exposed skin parts. This phenotype would have to keep on exhibiting its natural genetic variability so as to overcome this drawback, which will enable it to be improved through manual crossings or backcrossings. However, cocona's fruit pilosity variability must be induced since it is hard to be found naturally. The use of gamma radiation, to come up with desirable mutants, has shown to be the successful way of increasing any crop's genetic diversity [1,2,3,4].

Radiation dose and number of genetic mutations have shown to be directly related, yet higher doses induce cromossomical mutations [5], preventing the seeds from germinating and, thus leading the embryo to die. Conversely, lower doses have shown to be unable to induce significant mutations [2]. Therefore, it is essential to be able to calculate the accurate

32 dose to be used. In the case of cocona, there seems to be no published information
33 concerning the optimal dose to be used, therefore, one must seek for it on kin species.

34
35 Eggplant, a species phylogenetically related to cocona, has enabled to determine that 300
36 Gy of gamma radiation increases the fruit size and mass, when it is applied to its seeds [6].
37 In another similar work in eggplant, 160 Gy displayed to be appropriate dose for induce
38 favorable mutations [2].

39
40 Gamma radiation that is used in plants, is generally produced by radioactive elements such
41 as cobalt-60 and cesium-137 [7]. Its short waves, at picometer level, have high penetration
42 power, which produce ions inside the cells causing irreparable damage to cellular nucleus. It
43 is considered the mutations main inductor for creating new varieties from 1971 to 2008 [7],
44 its genetic effects on cellular nucleus go from genome mutations to gene mutations [8]. The
45 gene mutations for plant breeding are more useful because they don't damage most of the
46 genetic material.

47
48 In plant breeding programs have been used different plant tissue for irradiation, such as
49 whole plants, seeds, tubers, stems, buds, bulbs, pollen, in vitro plants, embryos,
50 microspores and callus [8]. Yet, seeds have been preferred for their easy to transport,
51 handle and store, i.e. can be irradiated, stored and germinated at any time.

52
53 The present study has irradiated cocona seeds with different doses ranging from 0 to 400
54 Gy for determining the optimal dose to bring about their variability.

55 56 **2. MATERIAL AND METHODS**

57
58 The experiment was carry out at Instituto Nacional de Pesquisas da Amazônia (Manaus,
59 AM) in the Agronomical Experimental Station (Manaus, AM), which is located on Km 14 AM-
60 10 roadway (2.9964 S and 60.0236 W).

61 Cocona seeds of "CUB-08" genotype were gamma-irradiated at Centro de Energia Nuclear
62 Aplicada à Agricultura (CENA/USP, Piracicaba, SP) with different doses i.e. 0, 100, 150,
63 200, 250, 300 and 400 Gy with a cobalt-60 source, which itself emitted 5010 Gy/h.

64 These seeds were sent to Manaus-AM and sown in 128-cell styrofoam trays filled with
65 Plantmax® mineral-organic substrate. Following three weeks, the seedlings were
66 transplanted to plastic cups filled with this same substrate, then, three months later; they
67 were transplanted to the open field in randomized complete block design, with four blocks
68 and seven plants per plot. Plants were grown for five months and their fruits harvested every
69 week for two months

70 The assessed fruit characteristics were yield, pilosity (1=No pilosity and 3=Excessive
71 pilosity), weight, diameter, number per plant, and ripening after seven days (1=firm;
72 3=rotten). Furthermore, stem pilosity was assessed, as well.

73 Data were submitted to analyses of variance (ANOVA) and means were compared by the
74 Duncan Test ($P < 0.05$). These analyses were performed in SAS 9.1.3, procedure PROC
75 GLM (SAS Institute, Cary NC). Moreover, using JMP 10 software was done an biplot graphic
76 based on correlations and principal components analysis.

77
78

79 **3. RESULTS AND DISCUSSION**

80
81 Mutagenic agents which are grouped into chemical, physical and biological ones may be
82 recommended for inducing genetic variability in plants (Bado et al. 2015). Physical radiations
83 have been a very popular, plant breeding mutagenic agent, and gamma radiation has shown
84 to be the most utilized one [7]. However, there are no former studies addressing radiation
85 dosages in cocona. On account of both cocona and eggplant being phylogenetically related
86 [9], this research was oriented through a previous work on eggplant [6], which demonstrated
87 that 300 Gy of gamma radiation would induce non deleterious variability, thus the tested
88 dosages for cocona were determined from 0 to 400 Gy.

89 The experiment was initially planned to be conducted with 20 plants per plot, yet only seven
90 plants per plot were used due to their seeds' general germinating vigor loss, which was
91 evident in the control treatment (0 Gy). This indicates the seeds to have likely been
92 submitted to temperature stress during their transport (Manaus-São Paulo-Manaus). Studies
93 have shown that cocona seeds are thermosensible [10, 11], where variations of temperature
94 from 20 to 30 °C increase the germination percentage. This reduced plot size is close to that
95 being used for other Solanaceae family species: *Capsicum annuum* (10 plants) [12,13].
96 Therefore, the plot plant number used in this study is acceptable.

97 Dose of 150 Gy significantly decreased fruit pilosity (Table 1). Furthermore, it induced no
98 effect on the other traits, such as the number of plants per plot, the stem's pilosity, fruit yield,
99 mass, length (L), diameter (D), L/D ratio and ripening. This conclusion is also shown on
100 biplot graphic (Figure 1), which explains 89% of total variation, where 150 Gy is found in the
101 fruit pilosity vector opposite direction, i.e. this dose decreases the character expression.
102 Moreover, fruit yield and number vectors orthogonal projections point toward 150 Gy,
103 suggesting this dose to tend to increase these characters. Therefore, this dose shows to be
104 more adequate for inducing the fruit pilosity's genetic variability and increasing fruit yield.

105 **Table 1. Effect of different gamma radiations on the morphoagronomical trait means of cocona**
106 **(*Solanum sessiliflorum*). Manaus 2016.**

107 ¹ 1=No pilosity e
108 3=Excessive
109 pilosity. ² 1=firm;
110 3=rotten. ³
111 Coefficient of
112 variation
113 ⁴Equal letters
114 indicate there not

Radiation dosages	Plant number per plot	Stem pilosity	Fruit pilosity ¹	Fruit Yield (t ha ⁻¹)	Mean fruit mass	Fruit length (L)	Fruit diameter (D)	L/D ratio	Ripening (at 7 days) ²
0 Gy	3.00 b ¹	2.08 a	2.28 a	7.19 a	152.66 a	8.60 a	6.40 a	1.34 a	1.46 a
100 Gy	5.75 a	2.20 a	2.18 ab	6.65 a	167.44 a	8.48 a	6.62 a	1.31 a	1.62 a
150 Gy	4.00 ab	2.05 a	2.00 b	7.42 a	146.90 a	7.98 a	6.08 a	1.35 a	1.18 a
200 Gy	2.75 b	2.00 a	2.25 a	2.07 a	132.85 a	7.62 a	6.22 a	1.22 a	1.13 a
Mean	3.87	2.08	2.16	5.83	149.96	8.17	6.33	1.29	1.34
CV ³ (%)	30.10	9.31	5.80	59.95	13.93	9.19	5.40	6.93	25.01

115 to have been significant radiation differences, by the Duncan test (P=.05)

116
117 These findings shown to be similar to found for plant dry and fresh weights in a eggplant
118 genotype (160 Gy) [2]. But these researchers observed low effect of the same radiation dose
119 in another eggplant genotype, displaying the presence of genotype x radiation interaction.
120 Thus, our results would be valid mainly for CUB-08 cocona genotype. Other genotypes can
121 need different doses.


122 Given these four treatments, our findings have shown 100 and 150 Gy to contribute to
123 significantly increase the germinating vigor as compared with 0 Gy (Table 1). In contrast,
124 200 Gy decreases that vigor. The rest of characters such as, stem, fruit's pilosity, number,
125 yield, mass, length, diameter, C/D ratio and ripening time have shown no significant
126 radiation dose-induced differences. They suggest 100 to 200 Gy doses not to be deleterious
127 for these characters. However, 200 Gy tends to reduce fruits number per plant and yield
128 (Table 1).

129 Biplot graphic (Figure 1) explained about 89% of total variation. In consequence, this graphic
 130 represents most part of data and their interactions. It showed three equidistant radiation
 131 doses (100, 150 and 200 Gy), suggesting their different and contrasting effects. Yet, 0 Gy is
 132 near the vector origin, indicating its lower effect for modifying the morphoagronomic
 133 characters. Hence, gamma rays have shown to be able to modify agronomical characters in
 134 cocona. On the other hand, 150 Gy was related with high fruit yield and fruit number per
 135 plant. Whereas 100 Gy was related with high values of fruit mass and diameter. These
 136 findings indicate 100 to 150 Gy being the optimal range of doses that creates variability for
 137 fruit yield and its components.

138 This biplot has also shown 200 Gy being in opposite direction of characteristics vectors. This
 139 indicates that their effect is deleterious for cocona both plant developing and fruit yielding.

140 Dose recommendation of gamma ray to induce non-deleterious mutations depends on
 141 species and vegetative material. For example, in tomato seeds was recommended 50 to 150
 142 Gy [14], in grasspea seeds 100 Gy [15]. Yet, in banana plants and shoots was
 143 recommended 20 to 30 kGy [16], in fig plantlets 30 kGy [17] and in potato meristems 2.5 Gy
 144 [18]. Therefore, each species and material needs a specific recommendation.

145



146

147 **Fig. 1. Biplot graphic of gamma radiation doses over cocona (*Solanum sessiliflorum*)**
 148 **seeds and some morphoagronomic characters (vectors)**
 149

150 Doses from 250 to 400 Gy were highly deleterious as shown by their seeds'germinating
 151 vigor loss and, experimental treatments number reduction to four (0, 100, 150 and 200 Gy).
 152 These findings suggest genetic variability in cocona to be able to be induced by using doses
 153 of less than 250 Gy.

154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175

176
177
178

179
180
181

182
183
184

185
186
187
188
189
190
191
192

193
194
195

196
197

4. CONCLUSION

Gamma radiation doses applied on cocona seeds above 250 Gy showed to be highly deleterious on account of hampering the seed germination. The optimal dosages for inducing mutations on cocona have shown to range from 100 to 200 Gy, yet, if the objective is to induce variability, 150 Gy has shown to be more effective, in addition to not modifying any of the fruit's agronomical traits such as, yield, mass, length and diameter.

COMPETING INTERESTS

Authors have declared there not to be any competing interests.

REFERENCES

1. Taheri S, Abdullah TL, Ahmad Z, Sahebi M, Azizi P. Phenotypic and molecular effects of chronic gamma irradiation on *Curcuma alismatifolia*. *European Journal of Horticultural Science*. 2016; 81(3): 137-147.
2. Ulukapi K, Ozdemir B, Onus AN. Determination of proper gamma radiation dose in mutation breeding in eggplant (*Solanum melongena*). *Advances in environmental and Agricultural Science*. 2015;1(1): 149-153.
3. Dhillon RS, Saharan RP, Jattan M, Rani T, von Wuehlisch G. Molecular characterization of gamma-rays induced mutants in *Jatropha curcas* L. *Indian Journal of Biotechnology*. 2014;13(1): 67-74.
4. Shin JM, Kim BK, Seo SG, Jeon SB, Kim JS, Jun BK, et al. Mutation breeding of sweet potato by gamma-ray radiation. *African Journal of Agricultural Research*. 2011;6(6):1447-1454.
5. Zhang J, Jiang Y, Guo YL, Li GR, Yang ZJ, Xu DL, Xuan P. Identification of Novel Chromosomal Aberrations Induced by Co-60-gamma-Irradiation in Wheat-*Dasyphyrum villosum* Lines. *International Journal of Molecular Sciences*. 2015;16(12): 29787-29796.
6. Ramaswamy N, Sayed S. Studies on the effect of gamma rays on eggplant. *Progressive Horticulture*. 1977; 9:77-79.
7. Mba C, Shub QY. Gamma Irradiation. In: Shu QY, Forster BP, Nakagawa H, editors. *Plant mutation breeding and Biotechnology*. Rome: FAO; 2011.
8. Bado S, Foster BP, Nielen S, Ali AM, Lagoda PJJ, Till BJ, Laimer M. Plant mutation breeding: current progress and future assessment. p. In: Janick J, editor. *Plant breeding reviews*, Volume 39. Hoboken: John Wiley & Sons; 2015.
9. Bohs L. A chloroplast DNA phylogeny of *Solanum* section *Lasiocarpa*. *Systematic Botany*. 2004; 29(1):177-187.

- 198 10.Lopes JC, Pereira MD. Germinação de sementes de cubiu em diferentes substratos e
199 temperaturas. Revista Brasileira de Sementes. 2005. 27(2):146-150. Portuguese.
- 200 11.Santos LA, Bueno CR, Clement CR. Influencia da temperatura na germinação de
201 sementes de cubiu (*Solanum sessiliflorum* Dunal) no escuro. 2000; 30(4):671-675.
202 Portuguese.
- 203 12.Lorentz LH, Lúcio AD. Tamanho e forma de parcela para pimentão em estufa plástica.
204 Ciência Rural. 2009; 39(8): 2380-2387. Portuguese.
- 205 13.Padrón RAR, Lopes SJ, Renedo VSG. Estimation of the optimal plot size and number of
206 replications in a field pepper crop experiment with varying irrigation depths and
207 application. Scientia Horticulturae. 2018; 237:96-104.
- 208 14.Sikder S, Biswas P, Hazra P, Akhtar S, Chattopadhyay A, Badigannavar A M, D'souza
209 SF. Induction of mutation in tomato (*Solanum lycopersicum* L.) by gamma irradiation and
210 EMS. Indian Journal of Genetics and Plant Breeding. 2013; 73(4): 392-399.
- 211 15.Ramezani P, Siavoshi, M, More Ad, Ebrahimi M, Dastan S. Gamma rays and EMS
212 induced flower color mutation in grasspea (*Lathyrus sativus* Linn.). Tarim Bilimleri
213 Dergisi. 2017; 23(4): 423-427.
- 214 16.Pestana RKN; Amorim EP, De Oliveira e Silva S, Neto AT. Gamma radiation for in vitro
215 mutagenesis in banana, cultivar Terra Maranhão. Pesquisa Agropecuaria Brasileira.
216 2010; 45(11): 1328-1330.
- 217 17.Ferreira EA, Pasqual M, Tulmann Neto, A. 2009. In vitro sensivity of fig plantlets to
218 gamma rays. Scientia Agricola. 2009; 66: 540-542.
- 219 18.Issa, F. H.; Najialhasnawi, A.; Shehab Sabah, S. Influence of gamma radiation on in vitro
220 growth microtubersation and hormonal content of some potato (*Solanum tuberosum* L.)
221 cultivars. Plant Archives. 2018; 18(2): 2317-2323.