

Original Research Article

EXTRACTIVISM OF THE UMBUZEIRO IN THE PERCEPTION OF THE FARMERS OF THE COMMUNITY RIACHO DA SERRA, IN SÃO JOSÉ DO SABUGI-PB

Farmers' perception on the extractivism of Umbuzeiro (*Spondias tuberosa*, Arr. Câmara) in Riacho Da Serra, SAO JOSE DO SABUGI-PB

ABSTRACT

The present work aimed to diagnose the perception of the farmers of the community of Riacho da Serra, in São José do Sabugi-PB. Initially visits were made in the community with the intention of exposing the idea and performing. The research was conducted during the period from march/2016 to may/2016. We applied structured questionnaires, previously made with the objective of involving and extracting the maximum information from the whole community. With the data in hand, figures were produced that demonstrate the perception of the cooperative farmers about the extraction of umbu. Incomplete first grade (47%), followed by high school (40%). As regards the process of extraction of umbu by Community producers, a number of uses may be were indicated, with the main destinations being the pulp industry (34%), household consumption (33%), food for animals (19%) and direct sales (14%). Regarding marketing, when the sale was made, the bag price of 3.0 kg ranged between R \$ 2.00 and R \$ 2.50, and 62% of the producers consulted stated that they sell for a price of R \$ 2.00, while 31% sell at R\$ 2.50 and 7% at R\$ 2.50. Of the producers that who produce one hectare, 100% have another activity to supplement the income, among which: rice (85%), livestock (85%), handicrafts (7.5%) and others (7.5%).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Key-words: Umbu. Cooperativa. Sustainability.

1. INTRODUCTION

The umbuzeiro (*Spondias tuberosa*, Arr. Câmara), belonging to the family Anacardiaceae, is an endemic species of the semi-arid region of the Brazilian Northeast with great socioeconomic and environmental importance. Its fruits are marketed by small farmers and extractivists, especially those considered most needy [5].

Currently, there has been intense extractive exploitation, which can generate losses of genetic material, since almost all high quality fruits originated from superior genotypes are collected, preventing natural reproduction from these genotypes. In addition, the expansion of agricultural frontiers with the constant deforestation of native vegetation also contributes to this loss of genetic diversity [3, 7]

Extractivism, when practiced in a sustainable manner, can generate income for many families and contribute to the conservation of the Caatinga, protecting the diversity of plants and animals, water courses and the cultural wealth of its peoples. **With this** **The** income are **use to** purchased food, household goods, clothes for children and school supplies, once the harvest period coincides with beginning of school term in rural schools. The valorization of the umbu can strengthen the traditions of the people and the permanence in the field from the generation of complementary income [3].

Formatted: Highlight

The fruits of the umbuzeiro are drupe glabrous or slightly hairy and rounded, weighing around 10 to 20 g. They have a smooth surface or exhibit 4 to 5 small lumps in the distal portion. The characterization of the fruits has evidenced the existence of a high phenotypic correlation, in descending order, for pulp, bark and core weight, total soluble solids and total acidity [9]. When ripe, the fruit has a juicy, slightly acidic and pleasant-tasting pulp containing 14.2 mg of ascorbic acid per 100 mL, fiber, reducing sugars and tannin.[ref?](#)

The organized collective work can be a good strategy for the best use of the fruits and the improvement of the conditions of collection, storage, processing and commercialization of umbuzeiro. An example of a social and productive organization of umbu extractivism is the work of Cooperatives and Resident Associations.

In the Brazilian semi-arid region, this activity has promoted the emergence of entrepreneurs able to create economic bases for family agriculture in areas dependent on rainfall in the Brazilian Northeast. However, there is a decrease in fruit production, with a marked decrease in recent years [5].

This reduction may be related to the water deficit and to the death of centennial plants of umbuzeiro, worsening even more with the absence of descendants, because there are no young umbuzeiros in pasture areas of the Caatinga [5, 8].

The objective of this research was to diagnose the perception of farmers in the community of [??????](#). **In the state of. On the importance of extractivism of umbuzeiro.**

Formatted: Highlight

2. MATERIAL AND METHODS

The municipality of São José do Sabugi is located in the Center-North region of the State of Paraíba, Mesorregião Borborema and Microregion Serido Ocidental Paraibano. It is bordered to the north by Ouro Branco (RN) and Santana do Serido (RN), east by Santana do Serido (RN) and Ecuador (RN), south by Ecuador (RN) and Santa Luzia (PB) with Santa Luzia and Ouro Branco (RN). It has the municipality of São José do Sabugi area of 215.4 km² and the municipal seat is located at an altitude of 333 meters [6].

The access from João Pessoa is made through the BR-230 (Figure 1), east-west direction, in a course of 299 km to the city of Santa Luzia, passing through Campina Grande, Soledade, Juazeirinho and Junco do Serido. From Santa Luzia it follows, to the northeast, in an 18 km stretch to the city of São José de Sabugi through Highway PB 221.




Figure 1. Map of location of São José do Sabugi in the State of Paraíba (Adapted from Google maps, 2017).

Initially visits were made to the community with the intention of exposing the idea to the producers (Figure 2). The research was carried out during the period from March 2016 to May 2016. For the study was used as an anthropological research tool a structured questionnaire, in which the topics are defined previously by the researcher [10].



Figure 2. Visit to the producers of the Riacho da Serra community, in São José do Sabugi, State of Paraíba. All the three maps look the a same why not to use just one of them????????????

Instead out the figure of umbo and caatinga

The proposed strategy to diagnose farmers perceptions of the Riacho da Serra Community in São José do Sabugi was based on five main points of analysis: (a) evaluation of the most immediate reality of the community under study; (b) application of a structured questionnaire to local farmers' perceptions; (c) quantification of the umbuzeiro density in the study area; (d) incentives for farmers to grow umbu as a crop; (e) contribution to the growth of the population of umbuzeiros.

The research was qualitative, being conducted to the data collection during the visits in loco not clear???. After the researcher's familiarization with the environment, semi-structured questionnaires (appendix) were distributed to farmers. The data were tabulated in computerized worksheets and later analyzed through the relative frequencies of the answers.

Formatted: Highlight

The obtained results were submitted to analysis of variance by the test " F " and descriptive analyzes, for diagnoses of significant effects of each source of individual variation and of their respective variables. For the processing of the datas, was used the software statistical AGROESTAT [1, 2]. Not clear

Formatted: Highlight

3. RESULTS AND DISCUSSION

It can be seen in Figure 2, that part of the farmers consulted is literate; however, the incomplete elementary school predominates (47%), followed by the second grade (40%).

Formatted: Highlight




Figure 3. School situation of farmers interviewed in the Riacho da Serra Community, in São José do Sabugi (PB).

[7], developing a work in Campo Redondo (RN), in the community of Malhada Vermelha, found that the educational level of those who worked there was 31.11%, which is lower than that found in the Community where the present study.

It can be seen in Figure 3 that 25% of the community residents interviewed say that they would leave to seek better living conditions in another locality. The process of rural exodus has been mainly caused by the rural population who leave the countryside in search of better living conditions in the polo cities, or in the capitals of the States, attracted by the financial profit and the attractive factor that the cities exert in the low-income populations [4].

■ Would leave ■ I would not go out




Figure 4. Perspective of the residents of the community of Riacho da Serra in regard to leaving the activities of the field and move to another place.

It is known, however, that this process of rural exodus generates a number of social problems, with emphasis on unemployment and underemployment. In addition, in cities, activities such as street vendors, pickers of recyclable materials, flanelinhas, among others, are becoming more common every day [11].

For [12] another negative factor to mention is the swelling of cities, which in the absence of urban planning there is overpopulation of poor neighborhoods, housing in places without structure and the increase of favelas.

Concerning the source of income acquired by the interviewees in the Community (Figure 4), 61% of interview participants reported that they obtained from agricultural activities, followed by income from retirement and salaried service.

It can be seen that, in Figure 4, agricultural activities are still the main source of income for the respondents in the Riacho da Serra Community. This is evidenced by the data from [13], for the municipality of São José do Sabugi, in terms of the quantity of umbu fruit produced, which totaled 14 tons. This production provided an income of R \$ 14.000.00.




Figure 5. Sources of income obtained by the interviewees in the Riacho da Serra Community, in São José do Sabugi (PB).

The results of the research visualized in figure 5 indicated that in the homes of the interviewees in the Community Riacho da Serra all have electric power. However, in the middle of the 21st century, many homes still have no running water or sanitary facilities at home.






Figure 6. Resources available in the homes of the interviewees in the Riacho da Serra Community, in São José do Sabugi (PB).

Tanks are still the main way to store water for human consumption (Figure 6A). Already, when argued about the main source of water search in the soil, they answered that the tubular and amazon wells give support to domestic activities (Figure 6B).



Form of family supply

C



■ Can ■ Animales ■ Car kite ■ Piped

Figure 7. Situation of water supply in the residence of farmers interviewed in the Riacho da Serra Community, in São José do Sabugi, PB.

The construction of the cistern serves as an alternative for storing and supplying rainwater in rural areas, especially in the semi-arid region of Paraíba, where the available sources, such as wells and rivers, have a variable volume of water, under the effect of seasonality, and most often water with high levels of salts.

The quality of the water stored in the tanks depends fundamentally on a good system maintenance. This consists of the disposal of the first waters, inspection and cleaning of the roof, gutters, pipes and the tank itself [14].

As for the family supply (Figure 7C), it is noted that 65% of the water **is** channeled. However, 10% is still carried out by farmers using cans. [4] families spend up to 30 hours per month on water transportation. Cans, which contain up to 20 liters of water, are usually carried in the heads causing chronic negative effects, including spinal pain. This transportation, most of the times, **is** done by women, children and adolescents.

The Caatinga **is** still very much exploited for the purpose of removing the wood for domestic use (firewood) and making fences. However, it **is** noteworthy that there **is** also exploration for more noble purposes, such as the use of medicinal plants, the extraction of fruits for various purposes, such as umbu, and the diverse use of seeds (Figure 7).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight




Figure 8. Utilization of the caatinga of the according to farmers interviewed in the Riacho da Serra Community, in São José do Sabugi, PB.

In the process of extraction of umbu by Community producers, a number of uses can be indicated, with direct sales (14%), feed (19%), family consumption (33%) and pulp (34%) (Figure 8). Variability in the use of umbu by growers is observed, with several alternatives for final destination of the product.




Figure 9. Main uses of umbu by farmers interviewed in the Riacho da Serra Community, in São José do Sabugi, PB.

The number of people in the family who participate in the holding of umbu in the community is two to four people. Meanwhile, the distance from the residence in relation to the umbu tasting site is up to two leagues.

The vast majority (94%) of farmers have always had one. In order to verify the reduction of the population of umbuzeiro used by the community, it was asked if the farmer had already collected in some umbuzeiro that finished and the result was that 15% of the farmers already saw the umbuzeiros of the place finished (Figure 9).




Figure 10. Percentage of farmers who have always collected and collected in a place that ended up in the community of Riacho da Serra, in São José do Sabugi, PB.

This result may indicate the risk of extinction of this species, as well as the reduction of areas of exploitation of the activity. [15] in a study in the Soledade-PB region, observed that the farmers were already worried about the risk of extinction.

Evaluating the time farmers practice extractivism, it is observed that farmers have some experience in the activity, with 57% of them hiring more than 20 years, indicating that the farmers who harvest umbu have a family tradition. However, the other age groups present considerable values around 15% (Figure 10).




Figure 11. Time when the farmers interviewed hunt umbu in the community of Riacho da Serra, in São José do Sabugi, PB.

In addition to the extraction of umbu, the producers remove other products such as firewood (75%) and cutting (55%) from the caatinga (Figure 11). The extraction of umbu is a periodic activity and is not able to provide monthly income to the farmers and thus becomes a complementary activity, only at the time of umbuzeiro production. Therefore, it is necessary for the farmer to seek another way to acquire more resources, being the removal of firewood and cuttings the easiest way to extract products from the Caatinga. However, this activity should be used sustainably.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight




Figure 12. Percentage of farmers who make other uses of the Caatinga in the community of Riacho da Serra, in São José do Sabugi, PB.

The farmers who are currently hunting, 83% were influenced by their parents who already practiced this activity, and 17% of the producers consulted have no descendants of umbu pickers.




Figure 13. Descendancy and ancestry of the farmers in the activity of umbu culling in the Riacho da Serra Community, in São José do Sabugi, PB.

236 The producers of the Riacho da Serra Community have indicated that they work on
237 collecting umbu in areas of less than ten hectares, which contain a maximum of 15 feet of
238 umbuzeiro for extraction. As for the number of umbu feet needed to fill a 60 kg bag, farmers
239 have reported that up to two umbuzeiros are required.

240 In the umbuzeiro harvest, producers pick up to eight sacks of umbu per day, depending
241 on the production of each umbuzeiro and the productivity of each producer, that is, although
242 some farmers pick eight sacks in a day, others can taste much less.

243 The commercialization of umbu in the community of Riacho da Serra is very incipient,
244 being that the majority of the producers tastes only for the consumption of the family and
245 they sell in small quantities. In 2014, the consulted producers reported that they sold less
246 than 200 umbu bags, that is, when there was a sale, as most declared that they did not sell.

247 Still on the sale, when the sale was made, the price of the 3kg bag varied between R\$
248 2.00 and more than R\$ 2.50, and 62% of the producers consulted stated that they sell for the
249 price of R\$ 2, 00, while 31% sell at R \$ 2.50 and 7% at R\$ 2.50 (question IX.05, appendix I).
250 Of the producers that produce a crop, 100% have another activity to supplement the income,
251 among which are cited: rice (85%), livestock (85%), handicrafts (7.5%) and others (7.5%).

252 In addition to the activities described by the producers, 78% receive a grant, retirement or
253 government assistance. Therefore, the monthly income of the family of 68% of the producers
254 is less than a minimum wage, while 32% of the producers have income between one and
255 two wages (Figure 13).

Formatted: Highlight

Formatted: Highlight




Figure 14. Situation of the income distribution of farmers interviewed in the Riacho da Serra Community, in São José do Sabugi, PB.

The community farmers reported that they participated in two entities, being affiliated with the association of rural producers of the community and the union of rural producers of the municipality. When asked how they have access to the doctor and dentist, they have reported that they are served through the PSF (Family Health Program) accessible to community producers.

As for the children's education, farmers were asked if they are satisfied with their child's school and all respondents said they are satisfied. Among the main needs of the community were mentioned lack of employment and lack of water as major difficulties of the community. Farmers also reported that they did not participate in any social / environmental movement.

4. CONCLUSION

Environmental perception was characterized as a relevant instrument to stimulate popular participation;

So that the exploration of the umbuzeiro **is** configured as an activity of generating employment and income, it must be conceived and accepted by everyone in the Riacho da Serra Community. The community should become more involved in the activities, expose their expectations, wishes, visions and opinions.

The relevant contribution of the research was stimulated by encouraging the participation of the local community within a process with sustainable bases, considering that **it**, besides knowing the peculiarities of the environment in which it lives, must accept the exploration of umbu as a sustainable activity, so that negative impacts are minimized and potential positive aspects.

COMPETING INTERESTS

Authors have stated that there are no competitors.

INTERESTS EXIST

Authors have declared that no competing interests exist.

5. REFERENCES

1. Banzatto DA, Kronka SN. 2008. Experimentação Agrícola. 4 ed. Jaboticabal: Funep, 237p. biblioteca (s): Embrapa Hortaliças.
2. Barbosa JC, Maldonado Junior W. 2015. AgroEstat - sistema para análises estatísticas de ensaios agrônômicos. Jaboticabal: FCAV/UNESP.
3. Barreto LS, Castro MS. 2010. Boas práticas de manejo para o extrativismo sustentável do umbu. Brasília: *Embrapa Recursos Genéticos e Biotecnologia*, 64 p.
4. Casagrande AE, Souza EBCO. 2012. Espaço e a demografia: o planejamento regional em perspectiva nas margens paranaenses do Lago de Itaipu. *Sociedade e Território*, Natal, 24(1): 2-27.
5. Cruz FRS, Andrade LA, Feitosa RC. 2016. Produção de mudas de umbuzeiro (*Spondias tuberosa* Arruda Câmara) em diferentes substratos e tamanho de recipients. *Ciencia florestal*, 26(1): 69-80.

Formatted: Highlight

Formatted: Highlight

- 306 6. CPRM. 2005. Serviço Geológico do Brasil. Projeto cadastro de fontes de
307 abastecimento por água subterrânea. Diagnóstico do município de São José do
308 Sabugí, Estado da Paraíba. Mascarenhas, J.C.; Beltrão B.A.; Souza Junior LC,
309 Morais F, Mendes VA, Miranda JLF. Recife: CPRM/Prodeem. 22 p.
- 310
- 311 7. David AAA. 2015. biotecnologia na propagação e conservação do umbuzeiro
312 (*Spondias tuberosa* Arr. Cam.) e percepção sobre sua importância por agricultores
313 da Comunidade Malhada Vermelha, Campo Redondo (RN - Brasil). Dissertação.
314 86f. (Prodeem/Ufrn). Centro de Biociências, Universidade Federal do Rio Grande do
315 Norte.
- 316
- 317 8. Araújo FP, Matta VM, Monteiro RP, Melo NF. 2016. Extrativismo do umbu e
318 alternativas para a manutenção de áreas preservadas por agricultores familiares em
319 Uauá, BA. Petrolina: Embrapa Semiárido, 21p.
- 320
- 321 9. Silva CMS, Pires I, Silva HD. 1987. Caracterização dos frutos de umbuzeiro.
322 Petrolina: Embrapa-CPATSA, 17p. (Boletim de Pesquisa, 34).
- 323
- 324 10. Viertler RB. 2002. Métodos antropológicos como ferramenta para estudos em
325 etnobiologia e etnoecologia. In: Amorozo MCM, Ming LC, Silva SP. 2002. Métodos
326 de coleta e análise de dados em etnobiologia, etnoecologia e disciplinas correlatas.
327 Rio Claro: Unesp: 31-46.
- 328
- 329 11. Pereira, K. P.; Lopes, J. L. 2013. Pobreza x degradação ambiental: existe
330 correlação? Uma análise estatística para o Paraná. VIII Encontro de Produção
331 Científica e Tecnológica. 2013. In: O Método Científico.
- 332
- 333 12. Francisco, W. de C. "Êxodo Rural ": Brasil Escola. Disponível em
334 <<http://brasilecola.uol.com.br/geografia/exodo-rural.htm>>. Acesso em 06 de
335 fevereiro de 2017.
- 336
- 337 13. IBGE. 2016. Produção da Extração Vegetal e da Silvicultura 2015. Rio de Janeiro:
338 IBGE.
- 339
- 340 14. Andrade Neto, C. O. 2004. Proteção sanitária das cisternas rurais. In: SIMPÓSIO
341 Luso-brasileiro de Engenharia Sanitária e Ambiental, 11, 2004, Natal, RN. Anais...
342 Natal RN: ABES: APESB: APRH.
- 343
- 344 15. Farias LAO, Pereira FC, Rêgo VGS, Oliveira EM, Ferreira AC, Baracuh JGV. 2012.
345 Percepção dos agricultores do lagedode timbaúba, soledade (pb) quanto à
346 importância do umbuzeiro (*Spondias tuberosa* A. Câmara). Revista Educação
347 Agrícola Superior. 27(1): 39-44.
- 348
- 349
- 350
- 351
- 352
- 353