

Palynological characteristic of dark gray clays in the southern part of the city of Bingerville (Ivorian onshore basin)

ABSTRACT

Samples from two wells implanted at the level of the dark gray clay outcrop of Bingerville were the subject of this palynological study. The main objective of this work is to establish the age of these levels in order to understand the geographical distribution of tertiary formations north of the lagoon fault. The samples were processed according to the classical procedure of extraction and concentration of palynomorphs.

The high populations of dinocysts of the genus *Lejeunecysta lata*, *Opreculodinium centrocarpum* and *Selenopemphix quanta* as well as those of spores and pollens of the genre *Magnastriatites howardii*, *Perforicolpites digitatus*, *Pachydermites diederixi*, *Bombacidites bombax*, *Retitricolporites irregularis*, *Retitriporites sp.* *Verrucatosporites usmensis* are tributary to Oligocene.

At the palaeobotanical and palaeoecological level, the highlighted palynoflora made it possible to characterize three environments, including mangroves, coastal plain, and rainforests.

Keywords : Dinocyst, spore, pollen, Oligocene, onshore basin, Cote d'Ivoire

1. INTRODUCTION

Long remained unknown in the ivorian sedimentary basin, the Oligocene age formations were described for the first time by [1] southeast of the city of Bingerville. These results indicated lithologically that this stage is mainly composed of gray clays interspersed with thin joints of stratifications (hardground). The palynology data highlight the presence of characteristic dinocysts such as *Lejeunecysta communis*, *Lejeunecysta lata*, *Lejeunecysta pulchra*, *Lejeunecysta sp.*, *Lejeunecysta granosa*, *Lejeunecysta globosa*, *Lejeunecysta beninensis*, *Pheolodinium magnificum*, *Pheolodinium africanum*, *Selenopemphix nephroides* et *Cordosphaeridium inodes*. These dinocysts are associated with spores and pollen grains as *Magnastriatites howardii*, *Spirosyncolpites spiralis*, *Perforicolpites digitatus*, *Retitricolporites irregularis*, *Retimonocolpites irregularis*, *Pachydermites diederixii*, *Psilatricolporites operculatus* et *Punctodiporites harrisii*.

Beside these studies, no results exist on the mapping of Oligocene age formations in the ivorian sedimentary basin and in particular in the northern part of the lagoon fault.

Recently, as a result of the amenagement work, gray but darker clays located beneath the variegated clays of known Mio-Pliocene age have been exposed at the southwestern entrance to Bingerville. This study was undertaken to date these levels in order to contribute to the paleogeographic reconstruction of tertiary deposits.

2. MATERIAL AND METHODS

2.1. Presentation of the study area

Comment [fs1]: (Ivorian onshore basin)

Formatted: English (United Kingdom)

Comment [S2]: conclusions say other things

Formatted: English (United Kingdom)

Comment [fs3]: Materials and methods

Comment [S4]: NO ITALIC

Comment [S5]: OLIGOCENE

51 Bingerville area is located east of the city of Abidjan. This region is part of the onshore
52 sedimentary basin of Côte d'Ivoire. There are generally clay formation unconformity on
53 Meso-Cenozoic schist and granites.

Formatted: English (United Kingdom)

Comment [S6]: BIBLIOGRAPHY

Formatted: English (United Kingdom)

55 This very narrow onshore basin is crossed from west to east by a fault "Lagoon Fault" of a
56 rejection of several thousand meters separating two distinct zones [2] in [3] :

57
58 - South of this fault, a deep basin in which the base sinks at 4000 or 5000 m on the vertical of
59 the coast;

60 - north of this fault is the shallower basin where the sedimentary cover rarely reaches 300 m
61 thick. This onshore basin belongs to the lagoons region and covers an area of approximately
62 664 km².

63 This is the northern part of the Bingerville area. Sedimentation is dominated by clays and
64 sands or ferruginous sandstones. Two wells P1 and P2 of depth 5,5 m and 18,5 m respectively
65 were made in this study.

Comment [fs7]: the location is the same as that studied by Digbehi Z.B. et al. 2012. are they the same wells?


66 From a physical geography, the Bingerville area has relatively rugged terrain. It has numerous
67 lagoon water plans (the Ebrié South Lagoon and the Potou Lagoon in the North) around
68 which a mangrove forest has developed.

Comment [fs8]: indicate latitude and longitude

Formatted: English (United Kingdom)

Comment [S9]: BIBLIOGRAPHY

Formatted: English (United Kingdom)



74

75

76

77

78 **Figure 1: Location of the study area**

79

80

81 **2.2. Methods**

82 The material used consists of ten (10) cuttings samples (Table I) from two wells made in the
83 Bingerville area east of Abidjan (Côte d'Ivoire).

84
85
86
87

Table I: Number of Samples and Well Depth

Well	Number of sample	Well depth (m)
Well P1	4	5,5
Well P2	6	18,5

88
89
90
91
92
93
94
95
96
97
98
99

Each sample of cuttings collected underwent a palynological preparation. This preparation consists in destroying all the mineral phases of the sediment by the strong acids (HCl 30% and HF 70%) in order to preserve only the organic phase generally consisting of sporopollinic or palynomorphic materials. After this attack with strong acids, each sample is washed on a 10 micron canvas and the sporopollenic residue obtained is mounted between the blade and the lamella using a special resin to glue the coverslip. For each sample, a pair of slides is made and observed under a biological microscope. This observation aims to identify the palynomorphs present in the samples, to make a palynostratigraphy and to determine the depositional environment.

100
101
102

3. RESULTS

103
104
105


3.1. Lithological analysis of the wells

3.1.1. Lithology of well P1

106
107
108
109
110
111
112
113

The lithology indicates in the sense of sedimentation the presence of ferruginous sandstones surmounted by dark gray clays interspersed with past ferruginous cuirasses. These clay formations have an average thickness of 3 m. They are surmounted by variegated clays with pasts of ferruginous cuirasses on 1 m thick (Figure 2). At the top, sandstone (0,7 m) and lateritic clays (0,3 m) meet.

Comment [S10]: NO SEDIMENTOLOGICAL ANALYSIS HAS BEEN MADE



114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129


Figure 2: Lithological synthesis of the well P1



3.1.2. Lithology of the P2 well

This well P2 is distant from the well P1 of 5 Km. It reached a depth of 18,5 m. Sedimentation shows from the bottom to the top dark pyritic and micromicassed gray clays capped by silty dark gray clays (1m). Above this set are dark gray clays interspersed with ferruginous cuirasses on 9,5 m of power (Figure 3).

Above, ocher yellow clays (1 m) are in contact with variegated clays interspersed with past ferruginous cuirasses over 4 m. At the summit clay sands (1 m) and sands (0,3 m) meet.

Comment [fs11]: indicate latitude and longitude
Comment [fs12R11]:

Scale




130
131
132
133
134
135

Figure 3: Lithological synthesis of the well P2

136
137
138
139
140
141
142
143
144

3.2. Qualitative and quantitative palynological analysis of the studied wells

Quantitative analysis of these wells resulted in a total of 629 spores and pollen grains, 292 dinocysts and 43 foraminifera basals HAVE BEEN COUNTED (. The details of this quantitative study are given in Table II below.

Table II: Number of palynomorphs per well

Well	Dinocysts	Spores and pollen grains	Foraminifera basals
Well P1	54	153	4
Well P2	238	476	39
Totals	292	629	43

145
146
147
148
149
150
151
152
153
154
155

Qualitative analysis of the palynomorphs of these two wells revealed a low variety of dinocysts and many spores and pollen grains. The details of these palynomorphs are presented in tables III and IV below. ARE LISTED PALINOMORPHS

The analysis of these tables III and IV shows a poverty of samples in dinocysts and an exceptional richness in spores and pollen grains. These spores and pollen grains are numerous and varied.

3.3. Palynostratigraphy

The palynostratigraphic study of this well shows that the palynomorphs encountered extend over a single stage which is Oligocene. The different characteristic palynomorphs of this stage are illustrated on plates I, II and III.

Oligocene is characterized by the following dinocysts: *Operculodinium centrocarpum*, *Cordosphaeridium inodes*, *Spiniferites ramosus*, *Batiacasphaera sp.*, *Lejeunecysta globosa* and *Lejeunecysta lata*.

These dinocysts are associated with the following spores and pollen grains: *Verucatosporites usmensus*, *Laevigatosporites ovatus*, *Perforicolporites digitatus*, *Monocolpites sp.*, *Deltoidospora minor*, *pachydermites diderixii*, *Leiotriletes adriensis*, *polypodiaceosporites simplex*, *Monocolpopollenites sp.*, *Psilatricolporites laevigatus*, *Monocolpites irregularis*, *Triorites sp.*, *Momipites sp.*, *Striatopollis bellus*, *Crottricolites densus*, *Retitricolpites americana*, *Retitricolporites irregularis* and *Oculopollis magnoporus*.

Comment [S13]: NO QUANTITATIVE ANALYSIS, ONLY THE palynomorphs COUNT HAS BEEN CARRIED OUT

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Comment [S14]: ????

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Comment [S15]: WHICH TEXTS HAVE BEEN USED FOR TAXONOMIC DETERMINATION?

Comment [S16]: no italic

Comment [S17]: sp. no italic

Comment [S18]: sp.

Comment [S19]: sp.

Comment [S20]: and

178 4. DISCUSSION

179
180 4.1. Palynostratigraphy

181
182 Oligocene has long been considered absent throughout the ivorian sedimentary basin because
183 it is strongly eroded. This erosion is highlighted in the Port-Bouet 1 sounding where the
184 Miocene rests unconformably on the upper Cretaceous.

185 However, the recent palynological work of [4, 5] highlighted Oligocene in Bingerville east of
186 Abidjan. This study continued by [6], has actually revealed a palynological association in the
187 gray clays that overcome the variegated clays (Continental terminal). This palynological
188 association consists of pollens *Spirosyncolpites spiralis*, *Perforicolpites digitatus*.

189 The work of [7] in the same area, allowed to define two associations of palynomorphs:
190 *Crassoretitrites vanraadshooveni*, *Verrucatosporites usmensis* encountered in gray clays
191 and *Verrucatosporites laevigatus* encountered in peats are characteristic associations of the
192 lower Miocene

193 This work has highlighted most of the species cited by these authors. These are spore species
194 and pollen grains such as *Perforicolpites digitatus*, *Verrucatosporites usmensis*,
195 *Laevigatosporites ovatus*, *Pachydermites diederixii*, *Polypodiaceoisporites simplex*,
196 *Psilatricolporites laevigatus*, *Monocolpites irregularis*, *Tripurites sp.*, *Momipites sp.*,
197 *Striatopollis bellus*, *Striatopollis catatumbus*, *Spinizonocolpites echinatus* and *Occulopollis*
198 *magnoporus*.

199 These species are associated with dinocysts such as *Lejeunecysta pulchra*, *Lejeunecysta lata*,
200 *Lejeunecysta globosa* and *Selenopemphix nephroides* characteristic of the Oligocene in most
201 West African countries.

202 [8] have shown that the association of dinocysts consisting of *Lejeunecysta pulchra*,
203 *Lejeunecysta lata*, *Lejeunecysta globosa*, *lejeunecysta sp.*, *Selenopemphix nephroids*,
204 *Operculodinium centrocarpum*, *Selenopemphix quanta* and *Cordosphaeridium inodes*
205 characterizes the Oligocene in Nigeria.

206 [9] showed in Gabon that the association of spores and pollen grains such as *Perforicolpites*
207 *digitatus*, *Magnastriatites howardii*, *Praedapollis africanus*, *Psilatricolporites operculatus*,
208 *Pachydermites diederixii*, *Verrucatosporites usmensis*, *Striatopollis bellus*, *Retitricolporites*
209 *irregularis* and *Occulopollis magnoporus* characterizes the Oligocene.

210 [10] indicate that the species *Bombacidites bombax*, *Psilastephanocolporites perforatus*.
211 *Psilatricolporites operculatus*, *Magnastriatites howardii*, *Pachydermites diederixii*,
212 *Perforicolpites digitatus*, *Praedapollis africanus*, *Retitricolporites sp.* and *Verrucatosporites*
213 *usmensis* characterize the upper Oligocene-lower Miocene passage in most African
214 sedimentary basins.

215 Most of these palynomorphs cited by these authors have been highlighted in work. The
216 Oligocene age was retained and confirmed in view of the presence of *Lejeunecysta lata* and
217 *Lejeunecysta globosa* in the studied wells.

218
219
220
221
222
223
224
225
226
227

Comment [fs23]: see Digbehi Z.B., Doukoure, M., Tea Y. J., Yao R.K., Yao N.J-P., Kangah K.D. & Tahi I. Palynostratigraphy and palaeoenvironmental characterization and evidence of Oligocene in the terrestrial sedimentary basin, Bingerville area, Southern Côte d'Ivoire, Northern Gulf of Guinea. African journal of Environmental Science and Technology, 6, 1 : (2012) 28-42.

Comment [S24]: Why?

Comment [S25]: NO ITALIC

Formatted: English (United Kingdom)

Comment [S26]: BIBLIOGRAPHY

Formatted: English (United Kingdom)

Comment [S27]: NO ITALIC

Formatted: English (United Kingdom)

Comment [S28]: BIBLIOGRAPHY

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Comment [S29]: NO ITALIC

Formatted: English (United Kingdom)

228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277

4.2. Paleobotany and paleoecology

The paleobotanical study of the wells shows the presence of the pollen grains of the Arecaceae (*Retitricolporites irregularis*, *Monocolpopollenites sp.*), *Nypa* (*Monocolpites marginatus*, *Racemonocolpites hians*), Meliaceae (*Psilastephanocolporites punctatus*), Moraceae (*Momipites sp.*), Polygalaceae (*Psilastephanocolporites perforatus*), Caesalpiniaceae (*Striatopollis bellus*), Alchorneas (*Psitricolporites operculatus*), Apocynaceae (*Margocolporites rauwolfii*, *Psilatricolporites crassus*) and Fabaceae (*Crototricolporites densus*). These pollen grains are associated with spores of Polypodiaceae (*Laevigatosporites ovatus*, *verrucatosporites usmensis*) and Cyatheaceae (*Deltoidospora minor*).

These different botanical groups can be divided into three paleoecological groups that are all mangroves group, rainforest group and Coastal plain group (including Swamp Species).

4.2.1. Mangrove group

The main mangrove elements identified in this study are *Psilatricolporites crassus* (Apocynaceae), *Psilatricolporites laevigatus*, *Cyathidites minor* and *polypodiaceosporites regularis*.

Psilatricolporites crassus and *Psilatricolporites laevigatus* are important elements of mangrove widespread throughout tropical Africa and South America [11, 12, 13, 14].

The *Psilatricolporites crassus* pollen is believed to be derived from the mangrove plant *Pelliceria* [13, 15].

Ferns of the genus *Cyathidites minor* and *polypodiaceosporites regularis* present in this group are typical of the dense forests of the coastal plains and are also found in mangrove areas [14,16, 17].

4.2.2. Coastal Plain Group (including Swamp Species)

This group is represented by an assemblage consisting of *Pachydermites diederixi*, *Retitricolporites irregularis*, *Verrucatosporites usmensis*, *Laevigatosporites ovatus*, *Monocolpopollenites sp.*, *Polypodiaceosporites regularis*, *Momipites sp.* and *Cyathidites minor*.

The genus *Pachydermites diederixi* shows a constant occurrence in both wells studied and associated with the pteridophyte of the genus *Polypodiaceosporites regularis*. This fern spore inhabits coastal wetlands and wetlands [18].

The species *Pachydermites diederixi*, which belongs to the family *Symphonia globulifera*, is known to be a dominant species in the coastal marshes of Africa [11]. The species *Retitricolporites irregularis* has been identified as a taxon present in coastal swamp environments [11].

[19] attribute this same pollen to freshwater swamp forests. Tree ferns such as *cyathidites minor* (cyatheacea) and *Polypodiaceosporites regularis* (Pteridaceae) inhabit thick tropical forests [16, 20].

The association of *Pachydermites diederixi*, *Verrucatosporites usmensis* and *Laevigosporites ovatus* indicate a freshwater or brackish swamp environments [21]. This could probably happen in the freshwater marsh behind the mangrove.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom),
Highlight

Formatted: English (United Kingdom)

Comment [S30]: NO ITALIC

Formatted: English (United Kingdom)

278
279
280
281
282
283
284
285
286

287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321

4.2.3. Rainforest group

This group consists of *Psilastephanocolporites punctatus*, *Striatopollis bellus*, *Psilatricolporites operculatus*, *Margocolporites rauvolfii*, *Racemonocolpites hians*, and *Psilatricolporites laevigatus*. [14] have shown that *Psilastephanocolporites punctatus*, *Racemonocolpites hians*, and *Psilatricolporites laevigatus* are indicative of dense moist forest. [22] attribute the species *Striatopollis bellus* and *Racemonocolpites hians* to rainforest. [23] confirms that *Striatopollis bellus* is a characteristic species of rainforest. [13] also indicates that the species *Psilatricolporites laevigatus* is a characteristic species of rainforest.

CONCLUSION

a study was conducted on The biostratigraphic analysis of Tertiary deposits in the Bingerville region through two wells P1 and P2 allowed to carry out the lithostratigraphic description of the formations present and their contents in palynomorphs of...XXXXX samples taken from two wells located in...XXXXX.....

Sedimentologically, the sediments are mainly composed of lateritic clays, variegated, dark gray with pasts of ferruginous cuirasses. There are also ferruginous sandstones, sands and reddish clay sands.

From the stratigraphic point of view the study revealed a palinoflora attributable to the Oligocene Palynologically, the study of palynoflor showed Oligocen.e, it is characterized characterized by the following dinocysts: *Lejeunecysta pulchra*, *Lejeunecysta lata*, *Lejeunecysta globosa*, *Selenopemphix nephroids*, *Selenopemphix quanta*, *Operculodinium centrocarpum* and *Cordosphaeridium inodes*.

These dinocysts are associated with the following spores and pollen grains: *Magnastriatites howardii*, *Perforicolpites digitatus*, *Pachydermites diderixi*, *Bombacidites bombax*, *Retitricolporites irregularis*, *Retitriporites sp.* *Verrucatosporites usmensis etc.*

from the palaeo-environmental point of view the data obtained allowed to identify three At the palaeobotanical and palaeoecological level, the highlighted palynoflora made it possible to characterize three environments, including mangroves, coastal plain, and rainforests: the first...XXXXX.....
the second.....XXXXX.....

Comment [fs31]: correlations should be made with the levels studied by Digbehi Z.B., Doukoure, M., Tea Y. J., Yao R.K., Yao N.J-P., Kangah K.D. & Tahi I. Palynostratigraphy and palaeoenvironmental characterization and evidence of Oligocene in the terrestrial sedimentary basin, Bingerville area, Southern Côte d'Ivoire, Northern Gulf of Guinea. African journal of Environmental Science and Technology, 6, 1 : (2012) 28-42.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Highlight

Formatted: Highlight

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

322

323

324 **Plate 1: Oligocene Dinocysts of Bingerville**



1



2



3

325

326


327



4



5



6

328

329

330

331 *1- Operculodinium centrocarpum* ; *2- Spiniferites ramosus* ; *3- Lejeunecysta lata* ; *4-*
332 *Batiacasphaera* sp. ; *5- Lejeunecysta globosa* ; *6- foraminifera basal.*

333

334

335

336

337

338

339


340

341

342

343

344 **Plate 2: Oligocene spore and pollen grains of Bingerville**




345

346

1

2

3




347

348

4

5

6



349

350

7

8

9

351 | *1-Verrucatosporites usmensis* ; *2-Retitriporites* sp. ; *3- Retitricolporites irregularis* ; *4-*
352 *Cyathidites minor* ; *5- Pachydermites diderixii* ; *6- Laevigatosporites ovatus* ; *7-*
353 *Occulopollis magnoporus* ; *8- Psilastephanocolporites laevigatus* ; *9- Momipites* sp.

354


355

Formatted: Font: Not Italic

356

357

358 **Plate 3 : Oligocene spore and pollen grains of Bingerville**



359

1

2

3

360



361

4

5

6

362



363

7

8

9

364

365 | 1- *Leiotriletes adriensis* ; 2- *Tripurites* sp. ; 3- *Crototricolporites densus* ; 4- *Tricolpites* sp. ;
 366 | 5-*Tricolpites americana* ; 6 -*Striatopolis bellus*; 7- *Margocolporites rauwolfii* ; 8 -
 367 | *Psilatricolporites crassus* ; 9 - *Psilatricolporites operculatus*

368

369

Formatted: Font: Not Italic

Formatted: Font: Not Italic

370

371

372 **Plate 4 : Oligocene spore and pollen grains of Bingerville**



1



2



3

373

374

375



4



5

376

377

378 | 1- *Polypodiaceiosporites simplex*; 2 - *Psilastephanocolporites punctatus*; 3 -
379 | *Margocolporites rauwolfii*; 4 - *Monocolpites marginatus*; 5 - *Retitricolporites verrucatus*

380

381

382

383

384 **APPENDIX**

385

386 **Dinoflagellate cysts**

387 *Cordosphaeridium inodes* (Klumpp) Eisenack, 1963b

388 | *Lejeunecysta globosae* Biffi and Grignani, 1983

389 *Lejeunecysta lata* Biffi and Grignani, 1983

390 *Operculodinium centrocarpum* (O. Wetzel, 1933a) Deflandre and Cookson, 1955

391 *Spiniferites ramosus* (Ehrenberg, 1838) Mantell, 1854

392

393 **Pteridophyte and bryophyte spores**

394 *Laevigatosporites ovatus* Wilson & Webster, 1947

395 *Leiotriletes andriensis* Krutzsch, 1959

396 *Polypodiaceosporites simplex* Sah, 1967

397 | *Verrucatosporites usmensis* (Van der Hammen, 1956) Germeraad et al., 1968

398 *Deltoidospora minor* (Couper, 1953) Pocock, 1970

399

400 **Angiosperm pollen**

401 *Crotocolpites densus* Salard-Cheboldaeff, 1978

402 *Echitriporites trianguliformis* Van Hoeken-Klinkenberg, 1964

403 *Margocolporites rauwolfii* Salard, 1978

404 *Monocolpites marginatus* Van der Hammen, 1954

405 *Oculopollis magnoporus* Zaklinskaya, 1963

406 | *Pachydermites diderixi* Germeraad et al., 1968

407 *Perforicolpites digitatus* González Guzmán, 1967

408 *Praedapollis africanus* Boltenhagen & Salard, 1973

409 *Psilastephanocolporites perforatus* Salard-Cheboldaeff, 1978

410 *Psilastephanocolporites punctatus* Salard-Cheboldaeff, 1978

411 *Psilatrcolporites operculatus* Van Der Hammen and Wijmstra, 1964

412 *Psilatricolporites crassus* Van der Hammen and Wijmstra, 1964

413 *Psilatricolporites laevigatus* Van der Hammen and Wijmstra, 1964

414 *Retitricolpites americana* Wymstra, 1964

415 *Retitricolporites irregularis* Van Der Hammen and Wijmstra, 1964

416 *Spinizonocolpites cf. baculatus* Muller 1968

417 | *Spinizonocolpites echinatus* Müller Muller 1968

418 *Striatopollis bellus* Sah, 1967

419 | *Triorites festatus* Müller Muller, 1968

420

421

422

423

424

425

426

427

428

429

430

431

432

433

Comment [S32]: microandriensis

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Comment [S33]: Müller

434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482

REFERENCES

1. Digbehi Z.B., Doukoure, M., Tea Y. J., Yao R.K., Yao N.J-P., Kangah K.D. & Tahi I. Palynostratigraphy and palaeoenvironmental characterization and evidence of Oligocene in the terrestrial sedimentary basin, Bingerville area, Southern Côte d’Ivoire, Northern Gulf of Guinea. *African journal of Environmental Science and Technology*, 6, 1 : (2012) 28-42.

2. Charpy N. et Nahon D. Contribution à l’étude lithostratigraphique et chronostratigraphique du Tertiaire de Côte d’Ivoire. Ser. Doc. Dept. Sci-Terre, Univ. Abidjan, série 18 (1978).

3. Digbehi Z.B. Etude comparée de la sédimentation de premier stade d’ouverture atlantique : Golfe de Guinée-Golfe de Gascogne (Sédimentologie-Biostratigraphie). Thèse de Doctorat, Pau (1987) 366p.

4. Petroci. Sédimentologie et palynologie des argiles noires et grises de Bregbo (Rapport interne) ,(2000) 150 p.

5. Doukoure M. Biostratigraphie des dépôts tertiaires dans la région de Bingerville. DEA des Sciences de la Terre option Géologie Marine et Sédimentologie, UFR STRM, Univ. Cocody (Abidjan), (2006) 59p.

6. Digbehi Z. B., Doukoure M., Tea Y. J., Yao K. R., Yao N. J-P., Kangah K.D & Tahi I. Palynostratigraphy and paleoenvironmental characterization and evidence of Oligocene in terrestrial sedimentary basin, Bingerville area, Southern Côte d’Ivoire, Northern Gulf of Guinea. “*African Journal of Environmental Science and Technology*”. 6 (1) (2011) pp.28-42.

7. Ennin T. M. Etude sédimentaire et palynologique de quatre puits dans la région de Brégbo (Sud-Est d’Abidjan). DEA des Sciences de la Terre option Géologie marine et Sédimentologie, UFR STRM, Université d’Abidjan, (2003) 58p.

8. Uderico B. et Dario G. Peridinioid dinoflagellate cysts from the Oligocene of the Niger Delta; Nigeria. *Revue micropaleontology*, vol.29, n° 2, (1983) pp. 125-145.

9. Salard-Cheboldaeff. Paléopalynologie du bassin sédimentaire littoral du Cameroun dans ses rapports avec la stratigraphie et la paléocologie. Thèse de Doctorat d’Etat ès Sciences Naturelles, Univ. Pierre et Marie Curie, Paris VI, 8 (1977) 262 p.

10. Eisawi A. & Schrank E. Upper Cretaceous to Neogene palynology of the Melut Basin, Southeast Sudan. *Palynology* , vol.32, n°1, (2008) pp. 101-129.

11. Germeraad, J.H., Hopping, C.A. and Muller, J. Palynology of Tertiary sediments from tropical areas. *Review of Palaeobotany and Palynology*, 6, (1968) 189-348.

12. Graham A. New records of Pelliceria (Theaceae/Pelliceriaceae) in the Tertiary of the Caribbean. *Biotropica*, 9, (1977) 48-52.

13. Rull V. A quantitative palynological record from the Early Miocene of western Venezuela, with emphasis on mangroves. *Palynology*, 25, (2001) 109-126.

Comment [S34]: CHECK THE DISTANCE BETWEEN THE WORDS AND MAKE UNIFORM

Comment [S35]: make uniform

Formatted: Highlight

Formatted: Highlight

483 **14. Bankole S., Schrank E. and Adeonipekun P.** Paleocology of the neogene agbada
484 formation, niger delta, Nigeria. *Ife Journal of Science*, vol. 18, no. 4, (2016) pp 845-855.

Comment [S36]: make uniform

485
486 **15. Graham A.** Diversification of Gulf/Caribbean mangrove communities through Cenozoic
487 time. *Biotropica*, 27(1), (1995) 20-27.

Formatted: Highlight

Formatted: Highlight

488
489 **16. Graham A.** Studies in neotropical paleobotany. VI. The Lower Miocene communities of
490 Panama -The Curacha Formation. *Ann. Missouri Bot. Gard.*, 75, (1988b) 1467-1479.

Comment [S37]: make uniform

Formatted: Highlight

Formatted: Highlight

491
492 **17. Samant B. and Phadtare N.R.** Stratigraphic palynoflora of the early Eocene Rajparadi
493 lignite, Gujarat and the lower age limit of the Tareswar Formation of south Cambay
494 Basin, India. *Palaeontographica Abt. B*, 245, (1997) Lfg., 1-6, 1-108.

Comment [S38]: ?????

495
496 **18. Graham A.** Studies in neotropical paleobotany. X. The Pliocene communities of
497 Panama-composition, numerical representations and paleocommunity paleoenvironmental
498 reconstructions. *Ann. Missouri Bot. Gard.*, (1991) 78, 465-475.

499
500 **19. Armentrout J.M., Fearn L.B., Rodgers K., Root S., Lyle W.D, Herrick D.C., Bloch**
501 **R.B., Snedden J.W. and Nwankwo B.** High-resolution sequence biostratigraphy of a
502 lowstand prograding deltaic wedge: Oso field (Late Miocene), Nigeria. In: Jones, R. W.
503 & Simmons, M. D. (eds.) *Biostratigraphy in production and development geology.*
504 *Geological Society, London, Special Publications*, 152, (1999) 259-290.

Comment [S39]: make uniform

505
506 **20. Samant, B. and Phadtare, N. R.** Stratigraphic palynoflora of the early Eocene Rajparadi
507 lignite, Gujarat and the lower age limit of the Tareswar Formation of south Cambay Basin,
508 India. *Palaeontographica Abt. B*, 245, Lfg., (1997) 1-6, 1-108.

Comment [S40]: ??????

509
510
511 **21. Rull V.** Sequence analysis of Western Venezuela of cretaceous to oligocene sediments
512 using palynology: Chronopaleoenvironmental and paleovegetational approaches. *Palynology*,
513 n° 21, (1997) pp 79-90.

Comment [S41]: make uniform

514
515 **22. Morley R. J. and Richards K.** Gramineae cuticles: a key indicator of Late Cenozoic
516 climatic change in the Niger Delta. *Review of Palaeobotany and Palynology*, (1993) 77, 119-
517 127.

518
519 **23. Cécile O.M.** Limite Paléocène-Eocène dans le bassin de Douala : biostratigraphie et essai
520 de reconstitution des paléoenvironnements. Thèse de doctorat, faculté des Sciences, Unité de
521 Paléobiogéologie, Paléobotanique, Paléopalynologie, Université de Liège, (2013) 221p.

522
523
524
525
526
527
528
529
530