

1 | **Palynological characteristic of dark gray clays in the Ivorian onshore basin**
2 | **at southern part of the city of Bingerville (Ivorian onshore basin)**

7 | **ABSTRACT**

8 | Samples from two wells implanted at the level of the dark gray clay outcrop of Bingerville
9 | were the subject of this palynological study. The main objective of this work is to establish
10 | the age of these levels in order to understand the geographical distribution of tTertiary
11 | formations north of the lagoon fault. The samples were processed according to the classical
12 | procedure of extraction and concentration of palynomorphs.

13 | The high populations of dinocysts of the genus *Lejeunecysta lata*, *Opreculodinium*
14 | *centrocarpum* and *Selenopemphix quanta* as well as those of spores and pollens of the genre
15 | *Magnastriatites howardii*, *Perfotricolpites digitatus*, *Pachydermites diderixi*,
16 | *Bombacidites bombax*, *Retitricolporites irregularis*, *Retitriporites sp. Verrucatosporites*
17 | *usmensis* are tributary to Oligocene.

18 | At the palaeobotanical and palaeoecological level, the highlighted palynoflora made it
19 | possible to characterize three environments, including mangroves, coastal plain, and
20 | rainforests.

23 | **Keywords :** Dinocyst, spore, pollen, Oligocene

25 | **1. INTRODUCTION**

26 | Long remained unknown in the ivorian sedimentary basin, the Oligocene age formations were
27 | described for the first time by [1] southeast of the city of Bingerville. These results indicated
28 | lithologically that this stage is mainly composed of gray clays interspersed with thin joints of
29 | stratifications (hardground). The palynology data highlight the presence of characteristic
30 | dinocysts such as *Lejeunecysta communis*, *Lejeunecysta lata*, *Lejeunecysta pulchra*,
31 | *Lejeunecysta sp.*, *Lejeunecysta granosa.*, *Lejeunecysta globosa*, *Lejeunecysta beninensis*,
32 | *Pheolodinium magnificum*, *Pheolodinium africanum*, *Selenopemphix nephroides* et
33 | *Cordosphaeridium inodes*. These dinocysts are associated with spores and pollen grains as
34 | *Magnastriatites howardii*, *Spirosyncolpites spiralis*, *Perfotricolpites digitatus*,
35 | *Retitricolporites irregularis*, *Retimonocolpites irregularis*, *Pachydermites diderixii*,
36 | *Psilatricolporites operculatus* et *Punctodiporites harrisii*.

37 | Beside these studies, no results exist on the mapping of Oligocene age formations in the
38 | ivorian sedimentary basin and in particular in the northern part of the lagoon fault.

39 | Recently, as a result of the amenagement work, gray but darker clays located beneath the
40 | variegated clays of known Mio-Pliocene age have been exposed at the southwestern entrance
41 | to Bingerville. This study was undertaken to date these levels in order to contribute to the
42 | paleogeographic reconstruction of tTertiary deposits.

46 | **2. MATERIALS AND METHODS**

48 | **2.1. Presentation Introduction of the study area**

50 Bingerville area is located east of the city of Abidjan. This region is part of the onshore
51 | sedimentary basin of Côte d'Ivoire. There are generally clay formation unconformity on
52 Meso-Cenozoic schist and granites.

53
54 This very narrow onshore basin is crossed from west to east by a fault "Lagoon Fault" of a
55 rejection of several thousand meters separating two distinct zones [2] in [3] :


56
57 - South of this fault, a deep basin in which the base sinks at 4000 or 5000 m on the vertical of
58 the coast;

59 - north of this fault is the shallower basin where the sedimentary cover rarely reaches 300 m
60 thick. This onshore basin belongs to the lagoons region and covers an area of approximately
61 664 km².

62 This is the northern part of the Bingerville area. Sedimentation is dominated by clays and
63 sands or ferruginous sandstones. Two wells P1 and P2 of depth 5,5 m and 18,5 m respectively
64 were made in this study.

65 From a physical geography, the Bingerville area has relatively rugged terrain. It has numerous
66 lagoon water plans (the Ebrié South Lagoon and the Potou Lagoon in the North) around
67 which a mangrove forest has developed.

68
69
70
71
72



73
74
75
76
77
78
79
80

Figure 1: Location of the study area

2.2. Methods

81 | The material used consists of ten (10) cuttings samples (Table I1) from two wells made in the
82 | Bingerville area east of Abidjan (Côte d'Ivoire).

83
84

85 | **Table I1: Number of sSamples and wWell dDepth**

86

Well	Number of sample	Well depth (m)
Well P1	4	5,5
Well P2	6	18,5

87

88

89 | Each sample of cuttings collected underwent a palynological preparation. This preparation
90 | consists in destroying all the mineral phases of the sediment by the strong acids (HCl 30%
91 | and HF 70%) in order to preserve only the organic phase generally consisting of sporopollinic
92 | or palynomorphic materials. After this attack with strong acids, each sample is washed on a
93 | 10 micron canvas and the sporopollenic residue obtained is mounted between the blade and
94 | the lamella using a special resin to glue the coverslip. For each sample, a pair of slides is was
95 | made and observed under a biological microscope. This observation aimeds to identify the
96 | palynomorphs present in the samples, to make a palynostratigraphy and to determine the
97 | depositional environment.

98 |

99

100 | 3. RESULTS

101

102 | 3.1. Lithological analysis of the wells

103 | 3.1.1. Lithology of well P1

104

105 | The lithology indicates in the sense of sedimentation as the presence of ferruginous
106 | sandstones surmounted by dark gray clays interspersed with past ferruginous cuirasses. These
107 | clay formations have an average thickness of 3 m. They are surmounted by variegated clays
108 | with some pasts of ferruginous cuirasses on 1 m thick (Figure 2). At the top, sandstone (0,7
109 | m) and lateritic clays (0,3 m) meetintersect.


110

111

112

Formatted: Right

Formatted: English (United States)




113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128



Figure 2: Lithological synthesis of the well P1

3.1.2. Lithology of the P2 well

This well P2 is distant from the well P1 of 5 Km. It reached a depth of 18,5 m. Sedimentation shows from the bottom to the top dark pyritic and micromicassed gray clays capped by silty dark gray clays (1m). Above this set are dark gray clays interspersed with ferruginous cuirasses on 9,5 m of power (Figure 3).

Above, ocher yellow clays (1 m) are in contact with variegated clays interspersed with past ferruginous cuirasses over 4 m. At the summit clay sands (1 m) and sands (0,3 m) meet.

Scale




129
130
131
132
133
134

Figure 3: Lithological synthesis of the well P2

135
136
137
138
139
140
141
142
143

3.2. Qualitative and quantitative palynological analysis of the studied wells

Quantitative analysis of these wells resulted in a total of 629 spores and pollen grains, 292 dinocysts and 43 foraminifera basals. The details of this quantitative study are given in Table II 2 below.

Table II2: Number of palynomorphs per well

Well	Dinocysts	Spores and pollen grains	Foraminifera basals
Well P1	54	153	4
Well P2	238	476	39
Totals	292	629	43

Formatted: Font: Not Bold

Formatted: Font: Not Bold

144
145
146
147
148
149
150
151
152
153
154

Qualitative analysis of the palynomorphs of these two wells revealed a low variety of dinocysts and many spores and pollen grains. The details of these palynomorphs are presented in Ttables III 3 and IV 4 below.

The analysis of these tables III and IV shows a poverty of samples in dinocysts and an exceptional richness in spores and pollen grains. These spores and pollen grains are numerous and varied.

3.3. Palynostratigraphy

The palynostratigraphic study of this well shows that the palynomorphs encountered extend over thea single stage which is Oligocene. The different characteristic palynomorphs of this stage are illustrated on plates Figs. I, II and III.

Comment [GT1]: Continue with the numbering of figures

Oligocene is characterized by the following dinocysts: *Operculodinium centrocarpum*, *Cordosphaeridium inodes*, *Spiniferites ramosus*, *Baticasphaera sp.*, *Lejeunecysta globosa* and *Lejeunecysta lata*.

161 These dinocysts are associated with the following spores and pollen grains: *Verucatosporites usmensus*, *Laevigatosporites ovatus*, *Perfotricolporites digitatus*, *Monocolpites sp.*, *Deltoidospora minor*, *pachydermites diderixii*, *Leiotriletes adriensis*, *polypodiaceoisporites simplex*, *Monocolpopollenites sp.*, *Psilatricolporites laevigatus*, *Monocolpites irregularis*, *Triorites sp.*, *Momipites sp.*, *Striatopollis bellus*, *Crottricolites densus*, *Retitricolpites americana*, *Retitricolporites irregularis* and *Oculopollis magnoporus*.

177 **4. DISCUSSION**

178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226

4.1. Palynostratigraphy

Oligocene has long been considered absent throughout the ivorian sedimentary basin because it is strongly eroded. This erosion is highlighted in the Port-Bouet 1 sounding where the Miocene rests unconformably on the upper Cretaceous.

However, the recent palynological work of [4, 5] highlighted Oligocene in Bingerville east of Abidjan. This study continued by [6], has actually revealed a palynological association in the gray clays that overcome the variegated clays (Continental terminal). This palynological association consists of pollens *Spirosyncolpites spiralis*, *Perfotricolpites digitatus*.

The work of [7] in the same area, allowed to definition of two associations of palynomorphs: *Crassoretitriletes vanraadshooveni*, *Verrucatosporites usmensis* encountered in gray clays and *Verrutricolporites laevigatus* encountered in peats are characteristic associations of the lower Miocene

This work has highlighted most of the species cited by these authors. These are spore species and pollen grains such as *Perfotricolpites digitatus*, *Verucatosporites usmensis*, *Laevigatosporites ovatus*, *Pachydermites diederixii*, *Polypodiaceoisporites simplex*, *Psilatricolporites laevigatus*, *Monocolpites irregularis*, *Triporites sp.*, *Momipites sp.*, *Striatopollis bellus*, *Striatopollis catatumbus*, *Spinizonocolpites echinatus* and *Occulopollis magnoporus*.

These species are associated with dinocysts such as *Lejeunecysta pulchra*, *Lejeunecysta lata*, *Lejeunecysta globosa* and *Selenopemphix nephroides* characteristic of the Oligocene in most West African countries.

[8] have shown that the association of dinocysts consisting of *Lejeunecysta pulchra*, *Lejeunecysta lata*, *Lejeunecysta globosa*, *lejeunecysta sp.* *Selenopemphix nephroids*, *Operculodinium centrocarpum*, *Selenopemphix quanta* and *Cordosphaeridium inodes* characterizes the Oligocene in Nigeria.

[9] showed in Gabon that the association of spores and pollen grains such as *Perfotricolpites digitatus*, *Magnastriatites howardii*, *Praedapollis africanus*, *Psilatricolporites operculatus*, *Pachydermites diederixii*, *Verrucatosporites usmensis*, *Striatopollis bellus*, *Retitricolporites irregularis* and *Occulopollis magnoporus* characterizes the Oligocene.

[10] indicate that the species *Bombacidites bombax*, *Psilastephanocolporites perforatus*, *Psilatricolporites operculatus*, *Magnastriatites howardii*, *Pachydermites diederixii*, *Perfotricolpites digitatus*, *Praedapollis africanus*, *Retitricolporites sp.* and *Verrucatosporites usmensis* characterize the upper Oligocene-lower Miocene passage in most African sedimentary basins.

Most of these palynomorphs cited by these authors have been highlighted in this work. The Oligocene age was retained and confirmed in view of the presence of *Lejeunecysta lata* and *Lejeunecysta globosa* in the studied wells.

227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276

4.2. Paleobotany and paleoecology

The paleobotanical study of the wells shows the presence of the pollen grains of the *Arecaceae* (*Retitricolporites irregularis*, *Monocolpopollenites sp.*), *Nypa* (*Monocolpites marginatus*, *Racemonocolpites hians*), *Meliaceae* (*Psilastephanocolporites punctatus*), *Moraceae* (*Momipites sp.*), *Polygalaceae* (*Psilastephanocolporites perforatus*), *Caesalpinaceae* (*Striatopollis bellus*), *Alchorneas* (*Psitricolporites operculatus*), *Apocynaceae* (*Margocolporites rauvolfii*, *Psilatricolporites crassus*) and *Fabaceae* (*Crototricolporites densus*). These pollen grains are associated with spores of *Polypodiaceae* (*Laevigatosporites ovatus*, *verrucatosporites usmensis*) and *Cyatheaceae* (*Deltoidospora minor*).

These different botanical groups can be divided into three paleoecological groups that are all mangroves group, rainforest group and Coastal plain group (including Swamp Species).

4.2.1. Mangrove group

The main mangrove elements identified in this study are *Psilatricolporites crassus* (*Apocynaceae*), *Psilatricolporites laevigatus*, *Cyathidites minor* and *polypodiaceosporites regularis*.

Psilatricolporites crassus and *Psilatricolporites laevigatus* are important elements of mangrove widespread throughout tropical Africa and South America [11, 12, 13, 14].

The *Psilatricolporites crassus* pollen is believed to be derived from the mangrove plant *Pelliceria* [13, 15].

Ferns of the genus *Cyathidites minor* and *polypodiaceosporites regularis* present in this group are typical of the dense forests of the coastal plains and are also found in mangrove areas [14,16, 17].

4.2.2. Coastal Plain Group (including Swamp Species)

This group is represented by an assemblage consisting of *Pachydermites diderixi*, *Retitricolporites irregularis*, *Verrucatosporites usmensis*, *Laevigatosporites ovatus*, *Monocolpopollenites sp.*, *Polypodiaceosporites regularis*, *Momipites sp.* and *Cyathidites minor*.

The genus *Pachydermites diderixi* shows a constant occurrence in both wells studied and associated with the pteridophyte of the genus *Polypodiaceosporites regularis*. This fern spore inhabits coastal wetlands and wetlands [18].

The species *Pachydermites diderixi*, which belongs to the family *Symphonia globulifera*, is known to be a dominant species in the coastal marshes of Africa [11]. The species *Retitricolporites irregularis* has been identified as a taxon present in coastal swamp environments [11].

[19] attribute this same pollen to freshwater swamp forests. Tree ferns such as *cyathidites minor* (*cyatheacea*) and *Polypodiaceosporites regularis* (*Pteridaceae*) inhabit thick tropical forests [16, 20].

The association of *Pachydermites diderixi*, *Verrucatosporites usmensis* and *Laevigosporites ovatus* indicate a freshwater or brackish swamp environments [21]. This could probably happen in the freshwater marsh behind the mangrove.

277
278
279
280
281
282
283
284
285

286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305

306
307
308
309
310
311
312
313
314
315
316
317
318

4.2.3. Rainforest group

This group consists of *Psilastephanocolporites punctatus*, *Striatopollis bellus*, *Psilatricolporites operculatus*, *Margocolporites rauvolffii*, *Racemonocolpites hians*, and *Psilatricolporites laevigatus*. [14] have shown that *Psilastephanocolporites punctatus*, *Racemonocolpites hians*, and *Psilatricolporites laevigatus* are indicative of dense moist forest. [22] attribute the species *Striatopollis bellus* and *Racemonocolpites hians* to rainforest. [23] confirms that *Striatopollis bellus* is a characteristic species of rainforest. [13] also indicates that the species *Psilatricolporites laevigatus* is a characteristic species of rainforest.

CONCLUSION

The biostratigraphic analysis of Tertiary deposits in the Bingerville region through two wells P1 and P2 allowed the study to carry out the lithostratigraphic description of the formations present and their contents in palynomorphs.


Sedimentologically, the sediments are mainly composed of lateritic clays, variegated, dark gray with parts of ferruginous cuirasses. There are also ferruginous sandstones, sands and reddish clay sands.

Palynologically, the study of palynoflor showed Oligocene, characterized by the following dinocysts: *Lejeunecysta pulchra*, *Lejeunecysta lata*, *Lejeunecysta globosa*, *Selenopemphix nephroids*, *Selenopemphix quanta*, *Operculodinium centrocarpum* and *Cordosphaeridium inodes*.

These dinocysts are associated with the following spores and pollen grains: *Magnastriatites howardii*, *Perforicolpites digitatus*, *Pachydermites diderixi*, *Bombacacidites bombax*, *Retitricolporites irregularis*, *Retitriporites sp.* *Verrucatosporites usmensis* etc.

At the palaeobotanical and palaeoecological level, the highlighted palynoflora made it possible to characterize three environments, including mangroves, coastal plain, and rainforests.

319 | **Plate 1: Oligocene Dinocysts of Bingerville**




320
321 | **1A**

2B

3C

322



323
324 | **4D**

5E

6F

325 | **Fig. 4: Oligocene Dinocysts of Bingerville**

326 | **1A**- *Operculodinium centrocarpum* ; **2B**- *Spiniferites ramosus* ; **3C**- *Lejeunecysta lata* ; **4D**-
327 | *Batiacasphaera* sp. ; **5E**- *Lejeunecysta globosa* ; **6F**- foraminifera basal.

328

329

330

331

332

333

334

335


336

337

338

Formatted: Tab stops: 0.75", Left + 0.77", Left

339 Plate 2: Oligocene spore and pollen grains of Bingerville




340

341

1

2

3




342

343

4

5

6



344

345

7

8

9

346 *1-Verrucatosporites usmensis* ; 2-*Retitriporites sp.* ; 3- *Retitricolporites irregularis* ; 4-
347 *Cyathidites minor* ; 5- *Pachydermites diderixii* ; 6- *Laevigatosporites ovatus* ; 7-
348 *Occulopollis magnoporus* ; 8- *Psilastephanocolporites laevigatus* ; 9- *Momipites sp.*


349

350

351

352

353 **Plate 3 : Oligocene spore and pollen grains of Bingerville**



354

355

1

2

3



356

357

4

5

6



358

359

7

8

9

360 *1- Leiotriletes adriensis* ; 2- *Tripurites sp.* ; 3- *Crototricolporites densus* ; 4- *Tricolpites sp.* ;
361 *5-Tricolpites americana* ; 6 -*Striatopolis bellus*; 7- *Margocolporites rauwolfii* ; 8 -
362 *Psilatricolporites crassus* ; 9 - *Psilatricolporites operculatus*


363

364

365

366


367 **Plate 4 : Oligocene spore and pollen grains of Bingerville**



368
369 **1** **2** **3**

Comment [GT2]:

Comment [GT3R2]: Plate labels on the figures



370
371
372 **4** **5**

373 1- *Polypodiaceiosporites simplex* ; 2 - *Psilastephanocolporites punctatus* ; 3 -
374 *Margocolporites rauwolfii* ; 4 - *Monocolpites marginatus* ; 5 - *Retitricolporites verrucatus*

375
376
377
378

379 **APPENDIX**

380

381 **Dinoflagellate cysts**

382 *Cordosphaeridium inodes* (Klumpp) Eisenack, 1963b

383 *Lejeunecysta globose* Biffi and Grignani, 1983

384 *Lejeunecysta lata* Biffi and Grignani, 1983

385 *Operculodinium centrocarpum* (O. Wetzel, 1933a) Deflandre and Cookson, 1955

386 *Spiniferites ramosus* (Ehrenberg, 1838) Mantell, 1854

387

388 **Pteridophyte and bryophyte spores**

389 *Laevigatosporites ovatus* Wilson & Webster, 1947

390 *Leiotriletes andriensis* Krutzsch, 1959

391 *Polypodiaceosporites simplex* Sah, 1967

392 *Verrucatosporites usmensis* (Van der Hammen, 1956) Germeraad et al., 1968

393 *Deltoidospora minor* (Couper, 1953) Pocock, 1970

394

395 **Angiosperm pollen**

396 *Crotocolpites densus* Salard-Cheboldaeff, 1978

397 *Echitriporites trianguliformis* Van Hoeken-Klinkenberg, 1964

398 *Margocolporites rauwolfii* Salard, 1978

399 *Monocolpites marginatus* Van der Hammen, 1954

400 *Oculopollis magnoporus* Zaklinskaya, 1963

401 *Pachydermites diderixi* Germeraad et al., 1968

402 *Perfotricolpites digitatus* González Guzmán, 1967

403 *Praedapollis africanus* Boltenhagen & Salard, 1973

404 *Psilastephanocolporites perforatus* Salard-Cheboldaeff, 1978

405 *Psilastephanocolporites punctatus* Salard-Cheboldaeff, 1978

406 *Psilatricolporites operculatus* Van Der Hammen and Wijmstra, 1964

407 *Psilatricolporites crassus* Van der Hammen and Wijmstra, 1964

408 *Psilatricolporites laevigatus* Van der Hammen and Wijmstra, 1964

409 *Retitricolpites americana* Wymstra, 1964

410 *Retitricolporites irregularis* Van Der Hammen and Wijmstra, 1964

411 *Spinizonocolpites* cf. *baculatus* Muller 1968

412 *Spinizonocolpites echinatus* Muller 1968

413 *Striatopollis bellus* Sah, 1967

414 *Triorites festatus* Muller, 1968

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429 REFERENCES

- 430 |
- 431 **1. Digbehi Z.B., Doukoure, M., Tea Y. J., Yao R.K., Yao N.J-P., Kangah K.D. & Tahi I.**
- 432 Palynostratigraphy and palaeoenvironmental characterization and evidence of Oligocene
- 433 in the terrestrial sedimentary basin, Bingerville area, Southern Côte d'Ivoire, Northern Gulf of
- 434 Guinea. *African journal of Environmental Science and Technology*, 6, 1 : (2012) 28-42.
- 435
- 436 **2. Charpy N. et Nahon D.** Contribution à l'étude lithostratigraphique et chronostratigraphique
- 437 du Tertiaire de Côte d'Ivoire. Ser. Doc. Dept. Sci-Terre, Univ. Abidjan, série 18 (1978).
- 438
- 439 **3. Digbehi Z.B.** Etude comparée de la sédimentation de premier stade d'ouverture
- 440 atlantique : Golfe de Guinée-Golfe de Gascogne (Sédimentologie-Biostratigraphie).
- 441 Thèse de Doctorat, Pau (1987) 366p.
- 442
- 443 **4. Petroci.** Sédimentologie et palynologie des argiles noires et grises de Bregbo (Rapport
- 444 interne) ,(2000) 150 p.
- 445
- 446 **5. Doukoure M.** Biostratigraphie des dépôts tertiaires dans la région de Bingerville. DEA des
- 447 Sciences de la Terre option Géologie Marine et Sédimentologie, UFR STRM, Univ. Cocody
- 448 (Abidjan), (2006) 59p.
- 449
- 450 **6. Digbehi Z. B., Doukoure M., Tea Y. J., Yao K. R., Yao N. J-P., Kangah K.D & Tahi**
- 451 **I.** Palynostratigraphy and paleoenvironmental characterization and evidence of Oligocene in
- 452 terrestrial sedimentary basin, Bingerville area, Southern Côte d'Ivoire, Northern Gulf of
- 453 Guinea. "African Journal of Environmental Science and Technology". 6 (1) (2011) pp.28-42.
- 454
- 455 **7. Ennin T. M.** Etude sédimentaire et palynologique de quatre puits dans la région de Brégbo
- 456 (Sud-Est d'Abidjan). DEA des Sciences de la Terre option Géologie marine et
- 457 Sédimentologie, UFR STRM, Université d'Abidjan, (2003) 58p.
- 458
- 459 **8. Uderico B. et Dario G.** Peridinioid dinoflagellate cysts from the Oligocene of the Niger
- 460 Delta; Nigeria. *Revue micropaleontology*, vol.29, n° 2, (1983) pp. 125-145.
- 461
- 462 **9. Salard-Cheboldaeff.** Paléopalynologie du bassin sédimentaire littoral du Cameroun dans
- 463 ses rapports avec la stratigraphie et la paléocologie. Thèse de Doctorat d'Etat ès Sciences
- 464 Naturelles, Univ. Pierre et Marie Curie, Paris VI, 8 (1977) 262 p.
- 465
- 466 **10. Eisawi A. & Schrank E.** Upper Cretaceous to Neogene palynology of the Melut Basin,
- 467 Southeast Sudan. *Palynology* , vol.32, n°1, (2008) pp. 101-129.
- 468
- 469 **11. Germeraad, J.H., Hopping, C.A. and Muller, J.** Palynology of Tertiary sediments
- 470 from tropical areas. *Review of Palaeobotany and Palynology*, 6, (1968) 189-348.
- 471
- 472 **12. Graham A.** New records of Pelliceria (Theaceae/Pelliceriaceae) in the Tertiary of the
- 473 Caribbean. *Biotropica*, 9, (1977) 48-52.
- 474
- 475 **13. Rull V.** A quantitative palynological record from the Early Miocene of western
- 476 Venezuela, with emphasis on mangroves. *Palynology*, 25, (2001) 109-126.
- 477

Formatted: Centered

Comment [GT4]: List references according to the Journal format

- 478 **14. Bankole S., Schrank E. and Adeonipekun P.** Paleocology of the neogene agbada
479 formation, niger delta, Nigeria. *Ife Journal of Science*, vol. 18, no. 4, (2016) pp 845-855.
480
- 481 **15. Graham A.** Diversification of Gulf/Caribbean mangrove communities through Cenozoic
482 time. *Biotropica*, 27(1), (1995) 20-27.
483
- 484 **16. Graham A.** Studies in neotropical paleobotany. VI. The Lower Miocene communities of
485 Panama -The Curacha Formation. *Ann. Missouri Bot. Gard.*, 75, (1988b) 1467-1479.
486
- 487 **17. Samant B. and Phadtare N.R.** Stratigraphic palynoflora of the early Eocene Rajpardi
488 lignite, Gujarat and the lower age limit of the Tareswar Formation of south Cambay
489 Basin, India. *Palaeontographica Abt. B*, 245, (1997) Lfg., 1-6, 1-108.
490
- 491 **18. Graham A.** Studies in neotropical paleobotany. X. The Pliocene communities of
492 Panama-composition, numerical representations and paleocommunity paleoenvironmental
493 reconstructions. *Ann. Missouri Bot. Gard.*, (1991) 78, 465-475.
494
- 495 **19. Armentrout J.M., Fearn L.B., Rodgers K., Root S., Lyle W.D, Herrick D.C., Bloch**
496 **R.B., Snedden J.W. and Nwankwo B.** High-resolution sequence biostratigraphy of a
497 lowstand prograding deltaic wedge: Oso field (Late Miocene), Nigeria. In: Jones, R. W.
498 & Simmons, M. D. (eds.) *Biostratigraphy in production and development geology.*
499 *Geological Society, London, Special Publications*, 152, (1999) 259-290.
500
- 501 **20. Samant, B. and Phadtare, N. R.** Stratigraphic palynoflora of the early Eocene Rajpardi
502 lignite, Gujarat and the lower age limit of the Tareswar Formation of south Cambay Basin,
503 India. *Palaeontographica Abt. B*, 245, Lfg., (1997) 1-6, 1-108.
504
- 505
- 506 **21. Rull V.** Sequence analysis of Western Venezuela of cretaceous to oligocene sediments
507 using palynology: Chronopaleoenvironmental and paleovegetational approaches. *Palynology*,
508 n° 21, (1997) pp 79-90.
509
- 510 **22. Morley R. J. and Richards K.** Gramineae cuticles: a key indicator of Late Cenozoic
511 climatic change in the Niger Delta. *Review of Palaeobotany and Palynology*, (1993) 77, 119-
512 127.
513
- 514 **23. Cécile O.M.** Limite Paléocène-Eocène dans le bassin de Douala : biostratigraphie et essai
515 de reconstitution des paléoenvironnements. Thèse de doctorat, faculté des Sciences, Unité de
516 Paléobiogéologie, Paléobotanique, Paléopalynologie, Université de Liège, (2013) 221p.
517
518
519
520
521
522
523
524
525