

Review Paper

Role of Bamboo forest for mitigation and adaptation to Climate Change challenges in China

ABSTRACT

Bamboo is one of the fastest growing plants on the planet, with many attributes which make it a useful potential resource for humankind. Though having fast growth and good regeneration performance after harvesting is a unique characteristic of the specie. It enhances a high carbon storage potential particularly when the harvested culms are transformed into durable products. China has many bamboo species with distribution and area coverage's, and highly connected in using the production of bamboo resources. Its characteristics make it an ideal solution for the environmental and social consequences of tropical deforestation. The aim of this review paper is to assess the contribution of bamboo in mitigating and adapting impacts of climate change and its importance in terms of ecological and socio-economic benefits. The review summarized the role of bamboo forests towards mitigating and adapting its potential to overcome the impacts of climate change currently seen globally and particularly to China. Therefore, advancing bamboo farming systems in different levels it's advantages to reduce greenhouse gas in atmosphere and expanding bamboo forests in future under wider use and intensive management is recommended.

Key words: - Bamboo; carbon sink; climate change; greenhouse gas

1. INTRODUCTION

Bamboo is a grass type of Gramineae family, and it is an important component of many forest environments. It adapts easily to a range of climatic and soil conditions, and is widely distributed in the tropical and subtropical zones approximately between 46° N and 47° S latitude, covering a total area of about 31.5 million ha, and accounted for about 0.8% of the World's total forested area in

Comment [AB1]: ,

42 2010 [1]. It has unique features that distinguish it from most other woody plants. For example: culms
43 that are connected by an extensive system of rhizomes, leading to emerge new culms by rapid
44 asexual reproduction [2, 3, and 4].

45 The bamboo resources are distributed in many countries in the World, majorly found in Asia, Africa,
46 and Latin America; however, the origin of most of species lie in Southeast Asia. Worldwide bamboo
47 families categorized with more than 107 genera and 1300 species [6]. Along with this all China has the
48 highest bamboo species diversity, with 39 genera and 509 species, accounting for 36% and 39%,
49 respectively of the total bamboo genera and species with the rest of the World [6]. Besides that,
50 China's bamboo forests cover an area of 4.84 million ha in 2005, which accounts for 2.8% of China's
51 forested area and 15.4% of the World's area of bamboo [7]. Bamboo forests is the most important
52 non-wood forest product and substitute wood products. As a **resource**, bamboo forests are an
53 important part of **eco-systems, which** provide a number of basic environmental services. Bamboo
54 provides food and raw materials, reduces water erosion on slopes, regulates water flows and it act
55 like windbreak in shelterbelts which offer protection against storms [1]. In addition to this because of
56 its special root re-sprouting regeneration strategy bamboo forests it generates a good potential of
57 carbon storage mechanisms, and water and soil conservation and many more advantageous [5].

58 Many studies have shown that well managed and regularly harvested bamboo can sequester more
59 carbon than bamboo in natural state. Despite that, it can sequester more carbon than fast-growing
60 tropical and subtropical trees under comparable conditions. If bamboo forests **were** not managed
61 through annual harvesting practices, they would be significantly less effective in carbon sequestration.
62 The aim of this review paper is to assess the contribution of bamboo in mitigating and adapting
63 impacts of climate change and its importance in terms of ecological and socio-economic benefits

Comment [AB2]: ,

64 2. ROLES OF FORESTS IN MITIGATING CLIMATE CHANGE

65 Forests are one of the biggest reservoirs of carbon, so they help to keep the carbon cycle and other
66 natural processes working and help reduce climate change. However, forests can also be one of the
67 biggest sources of CO₂ emissions [8]. In addition, forests provide a wide range of ecological, social,
68 and economic benefits, in the form of goods and services to society, that are much less easier to
69 quantify. Besides **that**, the demand for timber and related products will require more efficient and
70 sustainable use of natural resources. Forests are the most vulnerable climate dependent systems, but
71 have also been recognized to have significant and crucial contribution to address the challenges of
72 mitigation and adaptation in tandem with the issues of livelihoods, economic growth and development.

73 However, the most recent report from the International Union of Forest Research Organizations
74 (IUFRO) indicates the gloomy picture about the future of the World forests in changed climate, as it
75 suggests that in a warmer World, the current carbon regulating services of forests as carbon sinks
76 may be entirely lost as land ecosystems could turn into a net source of carbon dioxide later in the
77 century [9]. It plays in combating climate change impact through reducing the emissions from
78 deforestation and forest degradation has become a fundamental issue to international dialogues on
79 preventing the current global temperature increases [10]. The challenges of climate change seen in

Comment [AB3]: Long phrase

80 affecting forest ecosystems in their structure and morphology, thus causing an implication on
81 functionality of forests in every corner of the World [11, 12]. Beside that, it is considered one of the
82 greatest threats facing humanity in the current global situations. According to the Intergovernmental
83 Panel on Climate Change (IPCC), global warming is unambiguous, with evidence towards the
84 increases of average air and ocean temperatures, which leads to aggravate melting of snow and ice
85 and sea levels [8, 13].

86 3. BAMBOOS FOR CLIMATE CHANGE ADAPTATION

87 Bamboos are one of the World's strongest and fastest growing woody plants capable of providing
88 ecological, economic and livelihood security to the people, distributed over ranges of climate from mild
89 temperate to tropical. Bamboo's fast growth ability to grow on varied soils and climate, renewability
90 and positive socio-economic impacts make them an excellent alternative for combating climate
91 change [14]. The high growth potential and ability to store large amounts of carbon make
92 sequestration and on the other hand, their environmental and socio-economic services can help
93 communities in developing countries to adapt to the climate change impacts. According to the
94 research result by International Network of Bamboo and Rattan (INBAR) had shown that well
95 managed bamboos could be an effective in carbon sink and perform better than other species like
96 Chinese fir and eucalyptus growing under similar conditions. Furthermore, the necessities of bamboo
97 management with period, sustainable way and selective harvesting mechanism of stem which are
98 turned into products that can hold carbon for many years. The increasing popularity of durable
99 bamboo products ensures that for the foreseeable future, productive bamboo systems can be
100 considered as an important carbon sink [15].

101 3.1 BAMBOO FOR TIMBER DEMAND AND CLIMATE CHANGE

102 The demand for timber and different agricultural products will continue to increase with the global
103 population. Instead of satisfying the increasing demand for different commodities, the global policies
104 must need to shift towards using efficient and sustainable production systems [13]. Bamboo is one of
105 an alternative resource that used to play an important contribution towards reducing the direct
106 pressure on forest resources [13, 24]. Furthermore, one of the best example in China, following the
107 logging bans of certain forests resources came into effect in 1998, while bamboo forests has been
108 used as a possible substitute to timber.

Comment [AB4]: Erase the next line

109
110 Magel et al [16] argues that the growth of new shoots in a bamboo plantations occur because of
111 transfer of the energy accumulated in culms through photosynthesis in the previous year. The result
112 of such growth of bamboo culms is not driven by its own carbon sequestration, but by sequestration in
113 previous seasons in other parts of the bamboo system, and such growth of new shoots is not an
114 indicator of sequestration rate. Another report by Zhou [17] show that as bamboo system requires
115 more inputs in the shooting period of young culms, which means when new shoots developed during
116 that time high growth of bamboo shoots, can be equated with a high rate of carbon sequestration. The

117 maturity period of most bamboo culms estimated between 7-10 years approximately, after that they
118 deteriorate rapidly, releasing carbon from the aboveground biomass back into the atmosphere [18].
119 Therefore, at a natural circumstance, bamboo will reach a stable level of above ground carbon
120 relatively quickly, even though carbon accumulation through sequestration is offset by carbon release
121 through deterioration of old culms.

Comment [AB5]: ,

122 3.2 SUITABLE ECOLOGICAL GROWING CONDITIONS FOR BAMBOO SPECIES

123 Naturally, bamboo species has a potential to grow at different altitudinal range from 0 to 4000 meter
124 above sea level. It prefers well-drained sandy loams to loamy clay types of soils originated from river
125 alluvium or underlying rock. In most of the bamboo thrive well at annual average temperatures range
126 of 8.8 - 36°C and annual rainfall of 1270 – 4050 mm. On the other hand, some bamboo species are
127 also growing under high rainfall areas, while some can tolerate limited winter frost [19]. ICFRE [20]
128 reported as one of the fastest growing species in the planet, under ideal environment; it can be
129 growing up to one meter a day. The biomass of freshly planted bamboo plantation increases rapidly
130 for the early six to eight years after which emergence and death of culms tend to become equal.

Comment [AB6]: 36 °C

131 4. MANAGEMENT AND HARVESTING OF BAMBOO

132 Bamboos have well-developed rhizomes with good root systems which help to obtain strengthen
133 during their existence. Its culms mature within three to four years and naturally die after eight to ten
134 years, if not used the products [21]. The periodic removal of mature culms from each bamboo clump
135 and this cycle of removal may vary from two to four year. Thus, provide a highly renewable resource
136 with a high degree of sustainability. It makes bamboos acquiescent to sequester accumulated CO₂
137 from the atmosphere throughout the lifetime [22]. The sequestration rate of bamboo is higher during
138 the initial eight to ten years period of fast growth. Numerous studies categorize the production
139 management of bamboo management practices in different five major points: timber, shoot, pulp,
140 ornamental and water/soil conservation benefits.

141 5. BAMBOOS FOR CLIMATE CHANGE MITIGATION

142 Bamboo grows more rapidly than any other trees and reach to give yield within three to four years
143 after planting. Hence, it is one of the fast growing and responding well against drought, which can
144 make the species more acceptable in making evergreen environment in addition to soil and water
145 conservation, carbon storage and rehabilitation of degraded lands [23]. It offers one of the quickest
146 ways to remove vast amounts of that CO₂ from the atmosphere. Lou et al [24] report that at the age of
147 9 -10 years old in moso bamboo (*Phyllostachys pubescens*) plantation the above ground carbon stock
148 ranges between 25 to 32t ha⁻¹ in China. Furthermore, another study show that in *Phyllostachys*
149 *pubescens* and *Phyllostachys bambusoides* from natural forests in Japan have an aboveground carbon
150 stock of 78.6 t ha⁻¹ and 52.3 t ha⁻¹, respectively [25]. A four year mixed bamboo plantation (*Bambusa*
151 *vulgaris*, *B. blcooa*, *B. cacharensis*) in India shows that the aboveground carbon stock is about 61.05t
152 ha⁻¹ [26]. Despite that, it provides a minimum CO₂ gas and generates up to 25% more oxygen than

Comment [AB7]: Phyllostachys

Comment [AB8]: P.

Comment [AB9]: P.

Comment [AB10]: balcooa

153 other trees within the same level. One hectare of bamboo can sequester up to 62 t of CO₂ yr⁻¹,
154 whereas equivalent of young forest sequesters 15 t of CO₂ yr⁻¹. The *Guadua* plantations in Costa Rica
155 estimated to absorb 17 t of CO₂ ha yr⁻¹ as the study showed by Janssen [27]. Another research study
156 by INBAR states that over the past 15 years, areas under bamboos in Asia grew by 10%. Studies
157 have estimated that the carbon stored in Chinese bamboo forests will increase from 727.08 Tg C in
158 2010 to 1,017.54 Tg C in 2050, which equates to an increase of nearly 40% in 40 years. This
159 represents a significant contribution to the Chinese forest carbon stock and a range that shows that
160 policies aiming at combating climate change with bamboos can indeed have significant promise [28].
161 For example: by INBAR's, as modeling shows that a managed moso bamboo forest accumulates
162 about 300 t of carbon ha⁻¹ after 60 years. As well, it does also produce more biomass under well
163 managed and regular harvesting of mature culms. Another report by Lou et al [24], confirmed that the
164 amount of carbon sequestration between a fast growing Chinese Fir plantation and monopodial
165 (*Phyllostachys pubescens*) plantation modeled for subtropical agro-ecologies in South East China and
166 the results showed that, bamboo sequestered more carbon than the Chinese Fir in the first 5 years.
167 This might be due to rapid early growth; bamboos sequester more carbon in the early years of a
168 plantation than comparable forest trees. In the other way, unmanaged bamboo stands do not store
169 high levels of carbon, as their productivity is low and the accumulated carbon returns quickly to the
170 atmosphere as the older culms decompose [28].

Comment [AB11]: P.


171 5.1 CARBON SEQUESTRATION POTENTIAL OF BAMBOO

172 Carbon sequestration is the process of capturing and storing atmospheric carbon dioxide. Naturally,
173 the performance of bamboo plantation is different comparing with other tree species since it has a fast
174 growth rate with high re-growth behavior after harvesting. Despite that, it has a high carbon storage
175 potential according to Zhou and Jiang studies [29], especially when the harvested culms are
176 transformed into durable products. The increased lifespan of durable bamboo products made possible
177 by modern technology can ensure that the sequestered carbon will not return quickly in to the
178 atmosphere, thereby prolonging the carbon storage of bamboo. According to Agarwal and his
179 colleague [30] research result in the Mid Himalayan region conducting in comparing the carbon
180 sequestration potential under different bamboo species, Monopodial species (*Phyllostachys nigra*)
181 has showed a high potential to sequester carbon than other Sympodial species this might be due to
182 high density of culms and high percent dry matter in the Himalayan region [31]. Currently in China,
183 about 53 M ha of forest plantation is there with a volume stock of 1.5 billion m³. Between 2005 and
184 2020, China has pledged to establish more than 40 million ha of plantations, referred to as carbon
185 sink forest. As plantations have been recognized as the national strategy for mitigating atmospheric
186 CO₂, it is essential to assess the potential of fast-growing and high yield plantations in carbon storage
187 and sequestration at stand, regional and national scales [32].

Comment [AB12]: P.

Comment [AB13]: . This

Comment [AB14]: Figure 1?...must be "called" before



188
189

190 Figure 1. Contribution of bamboo on adding C sink and reducing C emission (Source: Yuen [33].

Comment [AB15]:)


191

192 The area coverage of bamboo forests in China is about 6 million ha^2 , which stores about 780 Tg
 193 carbon, accounting for 14% of total forest carbon stock in China. Beside the carbon density of
 194 bamboo forest ecosystems in China, the estimated global bamboo carbon stock is about 4 Pg,
 195 accounting for 0.43%-0.61% of total global forest carbon stock [33].

196 To combat climate change, bamboo should be a core development resource – providing countries
 197 and development partners with a wealth of practical solutions to reduce the negative effects that
 198 changing climate patterns have on millions of rural communities.

Comment [AB16]: Figure 2???

199



200

201 Figure 2. The five key functions of Bamboo help to mitigate/adapt the impacts of Climate Change.

202 **6. SUMMARY**

203 Currently the concern of climate change is a very serious and burning issue of global agendas. In this
 204 **paper**, we tried to review the contribution of bamboo forests towards mitigating the impacts of climate
 205 change and the versatility of its ecological and socioeconomic development benefits. It offers one of
 206 the quickest ways to remove huge amounts of CO₂ from the atmosphere. It minimizes CO₂ gas and
 207 generates more amount of oxygen than an equivalent stand of other tree species. Many scholars
 208 suggested that bamboo forest ecosystems provide significant services for human adaptation and
 209 development at the same time mitigate climate change impacts through carbon sequestration. Under

210 well-managed bamboo plantations, it shows an effective carbon sink and better performance than
211 other tree species growing under similar conditions. Despite this, it is a source of income in rewarding
212 the diverse requirements at small and large-scales in rural areas. Generally, this review prepared to
213 demonstrate the role of bamboo forests towards mitigating and adapting potential to overcome the
214 impacts of climate change seen in the world and particularly to China. Therefore, advancing bamboo
215 farming systems in different levels it's advantageous in reducing greenhouse gas in atmosphere and
216 expanding bamboo forests in future under wider use and intensive management is recommended.

217
218

REFERENCES

- 219 1. FAO. 2010. Global Forest Resources Assessment 2010: Main Report. Food and Agriculture
220 Organization of the United Nations, Rome.
- 221 2. Janzen D.H. 1976. Why bamboos wait so long to flower. *Annu. Rev. Ecol. Syst.* 7(1): 347–391.
222 doi:10.1146/annurev.es.07.110176.002023.
- 223 3. Isagi Y. Kawahara T. Kamo K. and Ito H. 1997. Net production and carbon cycling in a bamboo
224 *Phyllostachys pubescens* stand. *Plant Ecol.* 130: 41–52. doi:10.1023/A:1009711814070.
- 225 4. Li R. Werger M.J.A. During H.J. and Zhong Z.C. 1998. Carbon and nutrient dynamics in relation to
226 growth rhythm in the giant bamboo *Phyllostachys pubescens*. *Plant Soil*, 201(1): 113–123.
227 doi:10.1023/A:1004322812651.
- 228 5. Lobovikov M. Lou Y.P. et al. 2009. The poor man's carbon sink. Bamboo in climate change and
229 poverty alleviation. Non-Wood Forest Products. Working Document (FAO), no 8, FAO, Rome
230 (Italy), Forestry Dept. 68p., FAO.
- 231 6. Zhu Z.H. 2001. The development of bamboo and rattan in tropical China. Beijing, China: China
232 Forestry Publishing House.
- 233 7. SFAPRC. 2006. Statistics of Forest Resources in China (1999–2003). State Forestry
234 Administration, P.R. China.
- 235 8. IPCC 2007. Solomon S. Qin D. Manning M. Chen Z. Marquis M. Averyt K.B. M. Tignor and H.L.
236 Miller (eds). The Physical Science Basis, Contribution of Working Group I to the Fourth
237 Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge and New
238 York, NY, USA, 996.
- 239 9. Seppala R. Buck A. Katila. 2009. Eds. Adaptation of forests and people to climate change: A global
240 assessment report. IUFRO World Series Vol. 22. Vienna, IUFRO.
- 241 10. Wei S.J. L. Sun S.W. Wei et al. 2013: Effects of climate changes on forest disasters and the
242 preventive measures. *Journal of Catastrophology*, 28(1), 36-38.
- 243 11. MoEF (India. Ministry of Environment and Forests). 2012. India Second National Communication
244 to the United Nations Framework Convention on Climate Change. New Delhi, Ministry of
245 Environment and Forests, Government of India.
- 246 12. Hal Stern. 2006. Reviewed work(s): Source: The American Statistician, Vol. 60, No. 4 (Nov.,
247 2006), pp. 328-331.
- 248 13. Eliasch, J., 2008. Climate change: financing global forests: The Eliasch Review.
- 249 14. Dransfield S. Widjaja E. Eds. 1995. Bamboo Plant Resources of South-East Asia.
250 No. 7. Leiden, Backhuys Publishers. 190p.
- 251 15. INBAR. 2009. Bamboo: A strategic resource for countries to reduce the effects of climate change.
252 INBAR Policy Synthesis Report
- 253 16. Magel E. Kruse S. Lutje G. Liese W. 2005. Soluble Carbohydrates and Acid lyertases involved in
254 the rapid growth of the developing culms in *Sasa palmate* (Bean) Camus. *Bamboo Science and*
255 *Culture*, Baton Rouge/USA 19(2005) 1, S. 23-29.
- 256 17. Zhou G.M. Jiang P.K. Mo L.F. 2009. Bamboo: a possible approach to the control of global
257 warming. *International Journal of Nonlinear Sciences and Numerical Simulation*, 10(5): 547-550.

Comment [AB17]: Annual?

Comment [AB18]: erase

Comment [AB19]: erase

- 258 18. Liese W. 2009. Bamboo as Carbon sink fact or fiction? VIII World Bamboo Congress Proceedings.
259 3: 71-77.
- 260 19. FSI (Forestry Survey of India). 2011. India state of forestry report 2011. Dehradun, Forest Survey
261 of India. 286p.
- 262 20. ICFRE (India Council of Forestry, Research and Education). 1992. Bamboo. Dehradun, ICFRE.
- 263 21. Banik RL. 2008. Issues of production of bamboo planting materials-Lessons and strategies. Indian
264 For (Bamb Iss) 134(3):291-304.
- 265 22. Fu Maoyi and R.L. Banik. 1995. Bamboo, people and the management. Proceedings of the Vth
266 International Bamboo Workshop and the IV International Bamboo Congress Ubud, Bali,
267 Indonesia 10-22 June 1995. Volume I Propagation and management INBAR Technical Report
268 No.8p 18-30.
- 269 23. Terefe R. Samuel D. Sanbeto M. Daba M. 2016. Adaptation and growth performance of different
270 lowland bamboo species in Bako, West Shoa, Ethiopia. Journal of Natural Sciences Research.
271 2016; 6(9):61-65.
- 272 24. Lou Y. Li Y. Kathler B. Giles H. Zhou G. 2010. Bamboo and climate change mitigation:
273 Comparative analysis of carbon sequestration. INBAR. Technical Report No. 32. 2010; 1-20.
- 274 25. Isagi Y. 1994. Carbon stock and cycling in a bamboo *Phyllostachys bambusoides* stand. Ecol.
275 Res. 9. 47-55.
- 276 26. Nath A.J. Lai R. Das A.K. 2009. Above ground standing biomass for carbon storage in village
277 bamboos in **northeast** India. Bio. Bioeng 33, 1188-1196.
- 278 27. Janssen J. J. A. 1996. Building codes for bamboo housing. In Ganapathy, M. Janssen, J.A.;
279 Sastry, C.B. ed., Bamboo, people and the environment, Vol. 3, Engineering and utilization.
280 Proceedings of the Vth International Bamboo Workshop.
- 281 28. Kuehl Y. Yiping L. 2012. Carbon off- setting with bamboo. INBAR Working paper No. 71. Beijing,
282 INBAR.
- 283 29. Zhou G.M. and Jiang P.K. 2004. Density, storage and spatial distribution of carbon in
284 *Phyllostachys pubescens* forest. Sci. Sil. Sin. 40: 20-25.
- 285 30. Agarwal A. Purwar J.P. 2009. Evaluation of above ground biomass produced by *Dendrocalamus*
286 *asper* in North Western Himalayan region of India. In: Proceedings of VIII World Bamboo
287 Congress, 4, 91-96.
- 288 31. Agarwal A. Purwar J.P. 2012. Biomass Production and Carbon Sequestration Potential of Various
289 Bamboo Species in the Mid Himalayan Region of India. Paper Presented at World Bamboo
290 Congress.
- 291 32. Chen X. Zhang X. Zhang Y. Booth T& He X. 2009. Changes of carbon stocks in bamboo stands in
292 China during 100 years. *Forest Ecology & Management* 258: 1489-1496.
- 293 33. Yuen JQ. Fung T. Ziegler AD. (2017) Carbon stocks in bamboo ecosystems worldwide: estimates
294 and uncertainties. For Ecol Manag 393:113-138.

Comment [AB20]: erase

Comment [AB21]: erase

Comment [AB22]: 2017.