

EFFECTS OF NATURAL AND INOCULATION FERMENTATION ON THE ANTINUTRIENT COMPOSITION OF AFRICAN WALNUT ENRICHED WITH CARROT

ABSTRACT

The effects of fermentation on the antinutrient composition of African walnut enriched with carrot and without carrot were carried out in this study. The submerged natural fermentation of non- enriched and carrot enriched (walnut 70% ; carrot 30%) African walnut was carried out and lasted for 5 days during which microorganisms were isolated and identified. Species of *Lactobacillus* and *Bacillus* that are most commonly used as probiotics in foods, drugs, and dietary supplements, which was isolated during the natural fermentation of African walnut, were then used singly and in consortium during inoculation fermentation. During the fermentation processes, the antinutritional analysis were carried out on the fermented samples on daily basis. Result revealed that the bacteria *B. subtilis* strain b17a and *L. lactis* SFL8 were the most predominant bacteria during the natural fermentation. Raw African walnut and carrot have tannin, saponins, Phytate and oxalate of 2.47 ± 0.10 , 1.46 ± 0.10 , 14.41 ± 0.10 , $3.29 \pm .10$ and $2.28 \pm .08$, $2.10 \pm .10$, $11.50 \pm .05$, $0.37 \pm .03$ respectively. Raw walnut generally had a higher anti-nutrient composition than carrot except in saponins. Fermentation led to the reduction in antinutrient composition in the fermented sample. Natural fermented African walnut (NFW) and natural fermented African walnut enriched with carrot (NFWC) resulted in the highest reduction in antinutrient composition. NFW and NFWC had a reduction in tannin, saponins, Phytate and oxalate from (2.47 ± 0.008^b , 1.46 ± 0.010^a , 14.41 ± 0.026^d , 3.29 ± 0.05^c) to (0.98 ± 0.05^a , 0.61 ± 0.05^a , 4.35 ± 0.03^a , 0.27 ± 0.05^a) and (0.91 ± 0.005^a , 0.45 ± 0.005^a , 3.61 ± 0.03^a , 0.17 ± 0.005^a) respectively. Inoculation fermentation in which a monoculture of *L. lactis* or *B. subtilis* was used as starter culture resulted in the least reduction in the antinutrient content. Carrot enrichment of fermented African walnut had a reducing effect on the antinutrient composition of African walnut as the fermentation enriched with carrot all had a reduced tannin, Phytate and oxalate composition compared to those that were not enriched. Hence, fermentation and enrichment with carrot can be employed in improving the nutritional quality of African walnut thereby helping in effectively utilising their potential as human food.

Keywords: Fermentation; antinutrient; natural; inoculation; composition

1. INTRODUCTION

African walnut (*T. conophorum*) is a tropical perennial woody plant of the family Euphorbiaceae that is widely distributed and consumed by the inhabitants of Africa (1). The African walnut is often found growing wild as a climber in the forest regions of Africa and India continents (2). African walnut has a long history as food plant and is grown by peasant farmers. Its plant is mostly cultivated for its nuts which can be cooked or consumed as snacks. In Nigeria, it is called “Asala or Awusa” in Yoruba, “Ukpa” in Igbo and “Okhue or Okwe” in Edo. The plant is decorative and possesses nutritive, medicinal, agricultural and industrial values (1).

African walnuts are rich in protein, carbohydrate, fats and oils, and minerals (1, 3). Its seed are rich in fat, nearly eighty per-cent of unpolysaturated fat with proven cholesterol lowering properties. They can be cooked, roasted or sun dried and the roasted seeds could be ground and used as a thickener in soup preparation. Dried walnuts can be grounded and turned into flour which can be used as composite flour during baking or in place of milk in tea preparation (4). However, despite its nutritive values, African Walnut is still included in the list of lesser known food stuff in Nigeria. African walnuts have been demonstrated to have high anti-nutrient content (5). Anti-nutrients are protective, synthetic or natural compounds produced by plants against harsh environments or damage by microbes as a defence mechanism (6). Antinutrient, when consumed can upset the digestive process and interfere with the

absorption of beneficial essential organic nutrients and inorganic minerals (7). They either bind to nutrients to prevent their absorption or react with nutrient so it can't be used by the body (6). Removals of undesirable components are very essential in improving the nutritional quality thereby helping in effectively utilise their potential as human food. Several food processing methods such as germination, soaking, cooking and fermentation are known to reduce antinutritional factors effectively and upgrade their nutritional quality (8). Among the Several food processing methods which can reduce anti-nutritional factors, fermentation has been demonstrated to be the most effective (9).

Fermentation is the chemical breakdown of substance by bacteria, yeast or other microorganisms into alcohol, carbon dioxide or organic acids (9). Fermentation in food processing serves five main purposes which are: to enrich the diet through development of a diversity of flavours, aromas, and textures in food substrates; to preserve substantial amounts of food through lactic acid, alcohol, acetic acid, and alkaline fermentations; to reduce cooking time and the associated use of fuel; to retain food substrates with protein, enzymes, essential amino acids, vitamins and other nutrients that are usually destroyed by food processing since fermentation doesn't involve the use of heat and; to eliminate antinutrient (10). Fruits have been used to improve human diet for centuries, enriching it medicinally, nutritionally and sensorially. Carrot has been demonstrated to have smaller amount of anti-nutrient, zero cholesterol, low to no fat and high levels of vitamins, minerals and fibre (11). Hence, this project was aim at determining the effect of fermentation on the antinutrient composition of African walnut enriched with carrot.

2. MATERIALS AND METHODS

2.1. Sample materials

The African walnuts used for this study were obtained from a farm in Itapa, Ekiti State, Nigeria while the carrots used were bought from Oja Oba Market, Akure, Ondo State, Nigeria. The apparently healthy African walnut and Carrots were transported to the laboratory in separate clean and sterile black opaque polythene bags sealed at the tip and kept at refrigerated temperature (5°C) in the laboratory

2.2. Preparation of samples

The fruits (raw carrot and raw African walnut with pod) were gently washed with running tap water to remove dust and debris, then successively surface sterilized by washing with sterile distilled water and dipping in sodium hypochlorite. After which, the nuts were removed from the pod with the aid of a knife and the unshelled walnuts were then divided into two parts. The first part was cracked open and blended into powder while the second part was roasted in an oven at 130°C for one hour and later blended into powder. The unroasted blended walnuts were further subdivided into two parts; one portion was enriched with blended carrot while the other subdivision was not enriched with carrot. While the roasted blended walnut was further subdivide into six portions (three portions were enriched with carrot while the other three portions weren't enriched with carrot).

2.3. Preparation of the starter cultures

The microorganisms used were the most predominant microorganisms isolated from the natural fermentation of African walnut and they include *B. subtilis* b17a and *L. lactis* SFL8. Fresh starter cultures were prepared by reactivating the stock cultures of the pre-isolated microorganisms on Nutrient agar and Man- Rogosa-Sharpe (MRS) agar for *B. subtilis* and *L. lactis* respectively and cultured at 37°C for 48 hours followed by two successive rounds of sub-culturing in their respective broth with incubation at 37°C for 24 hours. The organisms were harvested by centrifugation at 4000 for 20min and pellets were added prior to fermentation in cell concentrations of 10^8 cfu/g (12). Mixed fermentation by both organisms was initiated by using equal proportion of each pure strain (12).

2.4. Submerged Fermentation of Substrates

Natural fermentation of unroasted walnut (NFW) and carrot enriched unroasted walnut (NFWC) was carried out in an enclosed plastic container by placing 40.0g of blended African walnut inside 200.0ml of water for the non- enriched fermentation while for the enriched sample, 28.0g (70%) gram of African walnut and 12.0g (30%) of carrot was placed in 200.0ml of water. The fermentation lasted for 5 days during which microorganisms were isolated and identified using both cultural and molecular methods.

The most predominant microorganisms having probiotic potentials which was isolated during the natural fermentation (*B. subtilis* strain b17a and *L. lactis* strain SFL8) were then used singly and in consortium during inoculation fermentation of non-enriched and carrot enriched African walnut (roasted in an oven at 130°C for 1hr) which was also carried out in an enclosed plastic container by putting the same grams as used in natural fermentation for the enriched and non- enriched fermentation [roasted African walnut fermented with *L. lactis* (WFL); roasted African walnut fermented with *B. subtilis* (WFB); roasted African walnut fermented with the combination of both *Bacillus subtilis* and *L. lactis* (WFBL); carrot enriched roasted African walnut fermented with *L. lactis* (WCFL); carrot enriched roasted African walnut fermented with *B. subtilis* (WCFB); carrot enriched roasted African walnut fermented with the combination of both *B. subtilis* and *L. lactis* (WCFBL)]. In order to prevent contamination, a container per day was used for different samples. During the five days fermentation, the antinutrient analysis was carried out on daily basis.

2.5. Isolation and conventional identification of microorganisms during fermentation

Fermented broth cultures were cultured on nutrient agar, Man de, Rogosa and Sharpe (MRS) agar, yeast extract agar and potato dextrose agar. Pure isolates were subjected to some biochemical tests and identified according to Cheesbrough (13). The identification of fungi in fermented samples was also carried out using standard morphological characteristics, which include morphology, surface characteristic and microscopic examination (14).

2.6. Molecular identification of bacteria isolate

Extraction of DNA using CTAB method was done according to Akinrotaye (15), PCR analysis was run with a universal primer for fungi called ITS1 and ITS4 and bacteria which was run with a universal primer called 16S rRNA. The amplicon was further purified before the sequencing using 2M Sodium Acetate wash techniques.

2.7. Antinutrient determination

Tannin and Phytate was determining according to the method of Wheeler and Ferrel (16). Oxalate was determined using the method of Day and Underwood (17). The spectrophotometric method of Brunner (18) was used for Saponin determination.

2.8. Statistical analysis

Data are presented as mean \pm standard error (SE). Significance of difference between different treatment groups was tested using one-way analysis of variance (ANOVA) using SPSS (Statistical Package for Social Science) version 20 software. For all tests, the significance was determined at the level of $P < 0.05$

3. RESULTS

The organisms isolated from raw African walnut before fermentation include *Bacillus subtilis* and *Micrococcus roseus* for bacteria and *Rhizopus oryzae*, *Candida albicans* and *Penicillium chrysogenum* for fungi. While the organisms isolated from raw carrot include *Micococcus roseus*, *Lactobacillus plantarum* and *Lactobacillus fermentum* for bacteria and *Rhizopus stolonifer*, *Candida albicans* and *Alternaria alternate* for fungi.

The array of bacteria and fungi that was isolated from the fermented broth cultures during the five days fermentation period revealed that the bacteria *Bacillus subtilis*, *Lactobacillus Planterium*,

Micrococcus roseus, *Lactobacillus lactis*, and *Lactobacillus fermentum* while the fungi *Penicillium notatum*, *Rhizopus oryzae*, *Penicillium chrysogenum*, *Candida albicans* *Alternaria alternata* and *Rhizopus stolonifer* were associated with the fermentation of African walnut.

The biochemical characteristics of the bacterial isolates are presented in Table 3. All the isolates showed different morphological features and biochemical properties. The Morphological characteristics features of fungi isolated from the fermented broth cultures are given in Table 4

The frequency distribution of bacterial and fungal isolates in the fermented samples are represented in Tables 1 and 2 respectively. The result revealed that *B. subtilis* was the most predominant bacteria in the fermented samples, while *Rhizopus oryzae* was the most predominant fungi isolated from the fermented sample.

Molecular identification of the bacterial isolates is shown in Table 5. The lengths of amplified products were 1453, 1133, 1450, 1525 and 1509 base pair for *B. subtilis*, *L. lactis*, *L. plantarum*, *Micrococcus roseus* and *L. fermentum* respectively Plate 2 and base pair 510 for *Candida albicans* Plate 1. The sequence obtained was BLAST in National Centre for Biotechnology Information (NCBI) database. Based on the 16SrRNA sequences, the bacteria *B. subtilis*, *L. lactis*, *L. plantarum*, *M. roseus* and *L. fermentum* were confirmed to be *B. subtilis* b17a, *L. lactis* SFL8, *L. paracasei* KLDS 1.0651, *M. luteus* NCTC and *L. casei* JCM 8621 while the yeast *Candida albicans* was confirmed to be *C. albicans* h70b. The result also revealed a difference in cultural identification of *L. plantarum*, *M. roseus* and *L. fermentum*. The Phylogenetic tree of the organisms isolated is shown in Figure 1

The anti-nutrient composition of raw walnut, raw carrot and roasted walnut are shown in figure 2. Roasted African walnut had the highest anti-nutrient composition of tannin 3.45 ± 0.053^b , Phytate 16.5314 ± 0.025^c and oxalate 3.5317 ± 0.025^b while raw carrot had the highest saponins content 2.05 ± 0.050^b . Raw walnut generally had a higher anti-nutrient composition than carrot, except in saponins. Fermentation led to daily reduction in anti-nutrient composition in the fermented sample with the least reduction recorded on the last day of fermentation. Natural fermented African walnut (NFW) and carrot enriched natural fermented African walnut (NFWC) resulted in the highest reduction in anti-nutrient composition. NFW and NFWC had a reduction in tannin, saponins, Phytate and oxalate from 2.47 ± 0.008^b , 1.46 ± 0.010^a , 14.41 ± 0.026^d , 3.29 ± 0.05^c to 0.98 ± 0.05^a , 0.61 ± 0.05^a , 4.35 ± 0.03^a , 0.27 ± 0.05^a and 0.91 ± 0.005^a , 0.45 ± 0.005^a , 3.61 ± 0.03^a , 0.17 ± 0.005^a respectively Figure 3 to 6. Inoculation fermentation in which a monoculture of *L. lactis* or *B. subtilis* was used as starter culture resulted in the least reduction in the anti-nutrient content. Carrot enrichment of fermented African walnut had significant effect $p < 0.05$ in reducing the antinutrient composition of African walnut as all the fermentation enriched with carrot all had a reduced tannin, Phytate and oxalate composition compared to the corresponding natural and inoculation fermentation of African walnut that were not enriched Figure 3 to 6.

Table 1: Frequency distribution of bacteria in the fermented samples

Isolates	NFW	NFWC	Occurrence (%)
<i>Bacillus subtilis</i>	+	+	65
<i>Lactobacillus plantarum</i>	-	+	5
<i>Lactobacillus lactis</i>	+	+	20
<i>Micococcus roseus</i>	+	+	5
<i>Lactobacillus fermentum</i>	+	+	5

Keys: + = Present, - = Absent, **NFW:** Natural fermentation of African walnut, **NFWC:** Natural fermentation of African walnut enriched with carrot,

Table 2: Frequency distribution of fungi isolate in the fermented samples

Isolates	NFW	NFWC	Occurrence (%)
<i>Rhizopus oryzae</i>	+	+	50
<i>Penicillium notatum</i>	+	+	15
<i>Penicillium chrysogenum</i>	+	+	15
<i>Candida albicans</i>	+	+	10
<i>Alternaria alternata</i>	-	+	5
<i>Rhizopus stolonifer</i>	-	+	5

Keys: + = Present, - = Absent, **NFW:** Natural fermentation of African walnut , **NFWC:** Natural fermentation of African walnut enriched with carrot

Table 3: Colonial, morphological and biochemical characteristics of bacterial isolates from fermented broth cultures

Isolate No	Colony Morphology	Gram's Reaction	Catalase	Coagulase	Motility	Mannitol	Glucose	Fructose	Maltose	Lactose	Sucrose	Citrate	Indole	Spore Forming	Methyl Starch	Urease	Probable Identity	
1	Cream, circular, opaque, flat, rough	+	+	NA	+	+	A G	A G	A G	A G	AG	+	-	+	-	+	-	<i>Bacillus subtilis</i>
2	Circular, opaque, convex, cream, smooth colonies	+	-	-	-	-	A	A G	A G	A	AG	-	-	-	+	-	+	<i>Lactobacillus plantarum</i>
3	Cream, circular, smooth, entire	+	+	NA	-	-	A G	A	-	-	AG	-	-	-	-	-	-	<i>Lactobacillus lactis</i>
4	Cream, circular, raised and smooth	-	+	-	+	-	A	A	-	-	A	-	-	NA	+	-	-	<i>Micococcus roseus</i>
5	Circular, translucent, convex, creamy, smooth colonies	+	-	-	-	-	A G	A	A G	A G	AG	-	-	-	-	-	-	<i>Lactobacillus fermentum</i>

Keyword: (+) = positive, (AG) = Acid and Gas, (-) = negative, (A) = Acid, (NA) = not applicable

Table 4: Morphological characteristics of fungal isolates from fermented broth cultures

Isolate No	Cultural and Microscopy description	Probable Identity
1	Yellowish green to dark green hyphae. Conidiophores arising from the mycelium singly or less often in synnemata, branched near the apex, penicillate, ending in a group of phialides	<i>Penicillium notatum</i>
2	Colonies are very fast growing with some tendency to collapse, white cottony at first becoming brownish grey to blackish-grey, sporangiophores are non-septate, arising from stolons opposite rhizoids usually in groups of 3 or more. Sporangia are globose with a flattened base, greyish black, powdery in appearance, Columellae and apophysis together are globose. Sporangiospores are angular, subglobose to ellipsoidal, with ridges on the surface.	<i>Rhizopus oryzae</i>
3	Greenish mycelium , Conidiophore hyaline, slender with spairing upper part, branched conidia and septate hyphae	<i>Penicillium chrysogenum</i>
4	Greenish growth, Conidiophore hyaline, slender with spairing upper part, branched conidia and septate hyphae	<i>Candida albicans</i>
5	White greenish growth with Conidia have a globose shape with rough surface wall	<i>Alternaria alternata</i>
6	Hyphae broad, not or scarcely septate; rhizoids and stolons present; sporangiophores brown, solitary or in tufts on the stolons, diverging from the point at which the rhizoids form; sporangia rather round; apophysis absent or scarcely apparent; sporangiophores ovoid.	<i>Rhizopus stolonifer</i>




Plate 1: PCR amplification of genomic DNA targeted to amplify the 18S rRNA gene of Yeast isolate on 1.0% agarose gel electrophoresis.

Key: M = Molecular marker




Plate 2: PCR amplification of genomic DNA targeted to amplify the 16S rRNA gene of bacterial isolate on agarose gel electrophoresis.

Key: M = Molecular marker

1= *Micrococcus roseus*

2= *Lactobacillus fermentum*

3= *Lactobacillus plantarum*

4= *Bacillus subtilis*

5= *Lactobacillus lactis*

Table 5: Molecular identification of isolated bacteria and yeast

Cultural and biochemical identification	Gene sequence identification	Max Identity	Accession number
<i>Candida albican</i>	<i>Candida albican</i> h70b	100	KP674768.1
<i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> b17a	100	JX406823.1
<i>Lactobacillus lactis</i>	<i>Lactobacillus lactis</i> SFL8	99	KT757263.1
<i>Lactobacillus Plantarum</i>	<i>Lactobacillus paracasei</i> KLDS 1.0651	99	FJ861111.1
<i>Micrococcus roseus</i>	<i>Micrococcus luteus</i> NCTC 2665	100	NR075062.2
<i>Lactobacillus fermentum</i>	<i>Lactobacillus casei</i> JCM 8621	100	AB690208.1




Figure 1: Phylogenetic tree of the microorganism isolated




Figure 2: Antinutrient composition of raw walnut, roasted walnut and raw carrot




Figure 3: Effects of fermentation on the tannin composition fermented samples




Figure 4: Effects of fermentation on the saponins content of fermented samples




Figure 5: Effects of fermentation on the Phytate composition of fermented samples




Figure 6: Effects of fermentation on the oxalate composition of carrot enriched and non-enriched African walnut

Keys: NFW: Natural fermentation of African walnut , WCFBL: Carrot enriched African walnut fermented *B. subtilis* and *L. lactis*, NFWC: Natural fermentation of African walnut enriched with carrot , WFB: African walnut fermented with *B. subtilis* , WFB: African walnut fermented with *B. subtilis* and *L. lactis* , WFL: African walnut fermented with *L. lactis* , WCFB: Carrot enriched African walnut fermented with *B. subtilis* , WCFBL: Carrot enriched African walnut fermented with *L. lactis*, Control: Unfermented walnut.

4. Discussion

In this study, the effects of fermentation on the anti-nutritional composition of African walnut enriched with carrot were evaluated. In addition, the microorganisms associated with the fermentation of African walnut were also isolated and identified. Since the major constituents of African walnut are proteins, fats and carbohydrates, the organisms responsible for its fermentation must be capable of utilizing these three constituents. Most of the organisms isolated from the fermented seeds are known to possess such characteristics. The organism isolated from the fermented samples using biochemical test were identified to be; *B. cereus*, *M. roseus*, *L. lactis*, *L. plantarum* and *L. fermentum* and the fungi isolated include *R. oryzae*, *C. albicans*, *Penicillium chrysogenum*, *R. stolonifer*, *Alternaria alternate*, and *Penicillium notatum*. These microorganisms were found during fermentation due to their ability to utilise the high protein content in African walnut as a source of nutrient for growth. *B. subtilis* was the predominant microorganism present; *B. subtilis* have been known to have proteolytic ability and also possess the capacity to break down oils (19). *B. subtilis* has been associated with fermenting locust bean for iru production (20) and for fermenting soy bean for natto production (21). These observations are in agreement

with the observation of Isu and Njoku, (22) that *Bacillus* species constitute over 95% of the total microbial population density in ugba fermentation.

Molecular techniques have the advantage: they are rapid, less laborious, and more sensitive, specific and efficient compared to the conventional method (23, 24). This study revealed a difference in cultural identification of *L. plantarum*, *M. roseus* and *L. fermentum*. A similar observation was also reported by Frickmann *et al.* (25), who reported differences in conventional method and molecular method of bacteria identification. However, the results of this study demonstrate clearly the interest and feasibility to introduce the 16S rRNA gene sequencing method in identification of bacteria, combination of conventional techniques and molecular approach will improve bacteriological investigation and authentication, allowing specific and efficient identification of microorganisms as against cultural method.

Anti-nutrients are generally known to reduce nutrient utilization and or food intake (26). Roasting generally led to increase in the level of these anti-nutritional components; this pattern of results was similar to what was reported by Enujiugba (27). Ekwe and Ihemeje (28) have also reported that toasting caused an increase in the phytate and tannin content of African walnut. Fermentation significantly led to the reduction in anti-nutrient composition in the fermented sample. This could be as a result of the fact that some anti-nutrient like tannins are water soluble and also the ability of some microorganisms to produce enzyme which can break down some of these anti-nutrient e.g. Phytases which has the ability of breakdown phytate content (29). A similar observation was reported by Esekheigbe and Onimawo, (26) where fermentation resulted in the reduction of anti-nutrients. Natural fermentation resulted in the highest reduction in anti-nutrient which maybe as a result of the various array of microorganism present in the natural fermentation as compared to the inoculation fermentation. This was further corroborated as the inoculation fermentation with consortium of two organisms resulting in a better reduction in anti-nutritional composition than when a monoculture was used. Carrot enrichment of fermented African walnut significantly help in reducing the anti-nutrient composition of African walnut as all the fermentation enriched with carrot all had a reduced tannin, Phytate and oxalate composition compared to the corresponding natural and inoculation fermentation of African walnut that were not enriched which maybe as a result of the fact that raw carrot had a very low Phytate, oxalate and tannin content when compared to raw African walnut. By and large, carrot enrichment of African walnut and Natural fermentation brought about maximal anti-nutrient reduction followed by inoculation fermentation with consortium of two organisms. Roasting is definitely not a good choice as the anti-nutrients composition were generally increased.

3. CONCLUSION

In this study, it is concluded that fermentation and enrichment with carrot helps in significantly reducing the antinutrient composition of African walnut and hence, both methods can be employed in improving the nutritional quality of the food thereby helping in effectively utilising their potential as human food.

7. REFERENCES

1. Isu NR, Njoku HO. An evaluation of the micro flora associated with fermented oil bean (*Pentaclethra macrophylla* Benth) seeds during Ugba production. *Plant Foods for Human Nutr.* 1997; 51: 145-157.
2. Edem CA, Dosunmu MI, Bassey FI. “ Determination of proximate composition, ascorbic acid and heavy metal content of African walnut (*Tetracarpidium conophorum*)” *Acad R Int.* 2009; 4(6): 501 – 512
3. Ayoola PB, Adeyeye A, Onawumi OO, Faboya O.O. “Phytochemical and nutrient evaluation of *Tetracarpidium conophorum* (Nigerian walnut) root”, *IJRRAS*, 2011; 7(2): 197 –202.
4. Ekwe CC, Ihemeje A. “Evaluation of physiochemical properties and preservation of African walnut (*Tetracarpidium conophorum*)”, *Acad R Int.* 2013; 4(6): 501 –512.
5. Kanu AM, Kalu JE, Okorie AC. Nutritional and Health Values of African Walnut (*Tetracarpidium Conophorum*). *IJSTR.* 2015; 4 (9): 2277-8616.

6. Onawumi OOE, Faboya OOP, Ayoola PB. "Chemical evaluation and nutritive values of African walnut leaf (*Plukenetia conophora* Mull. Arg)". *Int J Herb Med.*, 2013; 1 (3): 122 –126.
7. Ayoola PB, Adeyeye A, Onawumi OO, Faboya O.O. "Phytochemical and nutrient evaluation of *Tetracarpidium conophorum* (Nigerian walnut) root", *IJRRAS*, 2011; 7(2): 197 –202.
8. Ferial M, AbuSalem RK, Mohamed AY, Gibriel NM, Rasmy H. Levels of Some Antinutritional Factors in Tempeh Produced From Some Legumes and Jojobas Seeds. *Inter J Nut Food Eng.* 2014; 8 (3): 1307-6892
9. Hasan MN, Sultan MZ, Mar MU. Significance of Fermented Food in Nutrition and Food Science. *J. Sci. Res.* 2014; 6 (2): 373-386
10. Soccol CR, Vandenberghe LPS, Rodrigue C, Pandey A. New perspectives for citric acid production and application. *Food Technol. Biotechnol.* 2006; 44:141–149.
11. Adepaju. Proximate composition and micronutrient potentials of three locally available Wild fruits in Nigeria. *Afr. J. Agric. Res.*, 2009; 4(9): 887 892.
12. Hassen Y, Muskisa IM, Kurabachew H, Desalegn BB. Evaluation of Yeast and Lactic acid Bacteria Starter Cultures for the Production of Rice Injera. *J Food Process Technol.* 2018; 9(3): 2157-7110.
13. Cheesbrough M. District Laboratory Practice in Tropical Countries, second ed, Cambridge University Press, Cambridge, UK, 2006.
14. Bertrand TF, Natalia BE, Gordon TN, Bongsiysi GN, Irene AA, Samuel W. Probiotic properties of lactic acid bacteria isolated from fermented sap of palm tree (*Elaeis guineensis*). *J Microbiol Antimicrob.*, 2015; 7(5): 42-52.
15. Akinrotoy KP. Effects of fermented palm wine on some diarrhoeagenic bacteria. *Elite Research J Biotech. Microbiol.* 2014; 2(1): 4 – 14.
16. Wheeler EL, Ferrel RA. A method for phytic acid determination in wheat and wheat flour. *Cereal Chem.* 1971; 48: 313-314.
17. Day RA, Underwood AL. Quantitative Analysis, fifth ed. Prentice Hall Publication, US, 1986.
18. Brunner JH. Direct spectrophotometer determination of saponin. *Animal Chemist.* 1984; 34: 1314- 1326.
19. Enujiugha N. Major Fermentative Organisms in Some Nigerian Soup Condiments Pakistan. 2009; *J Nut*, 8 (3): 279-283
20. Antai SP, Ibrahim S.H. Microorganisms associated with African locust bean (*Parkia filicoidea* Welw) fermentation for 'dadawa' production. *J. Applied Bacteriol.* 1986; 61: 145-148.
21. Hesseltine CW. A millennium of Fungi, Food and Fermentations. *Mycologia*, 1965; 57: 149-197
22. Isu NR, Njoku HO. An evaluation of the micro flora associated with fermented oil bean (*Pentaclethra macrophylla* Benth) seeds during Ugba production. *Plant Foods for Human Nutr.* 1997; 51: 145-157.
23. Keramas G, Bang DD, Lund M, Madsen M, Bunkenborg H, Telleman P, Christensen CB. Use of culture, PCR analysis, and DNA microarrays for detection of *Campylobacter jejuni* and *Campylobacter coli* from chicken faeces. *J Clin Microbiol.* 2004; 42(9): 3985-91.
24. Magistrado PA, Garcia MM, Raymundo AK. Isolation and polymerase chain reaction-based detection of *Campylobacter jejuni* and *Campylobacter coli* from poultry in the Philippines. *Int. J. Food Microbiol.* 2001; 70:197–206.
25. Frickmann H, Dekker D, Schwarz NG, Hahn A, Boahen K., Sarpong N. 16S rRNA Gene Sequence-Based Identification of Bacteria in Automatically Incubated Blood Culture Materials from Tropical Sub-Saharan African. *PLoS ONE*, 2015; 10(8): 01-923
26. Esekheigbe A. and Onimawo, I.A. 2015. Effect of Fermentation and heat Treatment on the Nutritient and Antinutrient Composition of African Walnut (*Tetracarpidium conophorum*). *Pak J Nut.* 2015; 14(12): 972-975.
27. Enujiugha VN. "Chemical and functional characteristics of conophor nut". *Pak J Nutri.* 2003; 2(6): 335 –338.
28. Ekwe CC, Ihemeje A. "Evaluation of physiochemical properties and preservation of African walnut (*Tetracarpidium conophorum*)", *Acad R Int.* 2013; 4(6): 501 –512.
29. Akande KE, Doma UD, Agu HO. and Adamu HM. Major Antinutrients Found in Plant Protein Sources: Their Effect on Nutrition. *Pak J of Nut.* 2010; 9 (8): 827-832.