

**PRIMING METHODS: ALTERNATIVE STRATEGY TO
IMPROVE SEED AND SEEDLING PERFORMANCE
OF SOURSOP (*Annona muricata*)**

D.J.A. Najorda¹ and R.J.G. Rosales²

¹*Graduate of Bachelor of Science in Agriculture major in Horticulture, Department of
Agricultural Sciences, College of Agriculture, Food and Sustainable
Development, Mariano Marcos State University,
City of Batac, Ilocos Norte, Philippines*

²*Faculty of Department of Agricultural Sciences, College of Agriculture, Food and
Sustainable Development, Mariano Marcos State University,
City of Batac, Ilocos Norte, Philippines*

Authors' contributions

*This work was carried out in collaboration between the authors. The authors read and
approved the final manuscript.*

ABSTRACT

The evaluation of seed priming methods on the seed and seedling performance of soursop was conducted January 9 to April 3, 2019 in San Nicolas, Ilocos Norte, Philippines. The study was conducted to investigate the effect of seed priming methods capable of breaking dormancy; improve germination rate; determine the seed vigor of soursop; and identify the best seed priming method that provide better seedling performance. The experimental treatments (unprimed and three priming methods, hydropriming, halopriming and hormonal priming) were laid out in Completely Randomized Design with three replications. A total of 20 polyethylene bags were used per treatment per replication with one seed sown in every bag.

Alternative way to improve seed and seedling performance is the use of these seed priming methods. The seed and seedling performance of soursop were significantly affected by priming methods. Primed seeds had higher percentage germination rate (PGR) than unprimed seeds. But numerically, the highest PGR was hydropriming. Hormonal priming produced significantly taller seedlings at 10 and 40 days after emergence (DAE) than unprimed seeds and more leaves per seedling at 30 to 50 DAE. Hormonal, hydropriming and halopriming produced significantly higher fresh weight of seedlings than unprimed seeds.

Keywords: *hydropriming, halopriming, hormonal priming, seed treatment*

INTRODUCTION

Soursop (*Annona muricata*) is one fruit crop being propagated in the Philippines particularly in Ilocos Norte. Its fruit is a good source of essential minerals and nutrients which are beneficial to human health [10]. The crop is commonly propagated by seed due to its easy procedure. However, the seeds have hard and thick seed coats which restrict the entry of moisture, thus delaying germination and resulting in uneven maturity[16]. The unprimed seeds emerge for almost two to three weeks. Because of this situation, mass propagation is done through vegetative method, which has shorter waiting time to flower. Still, not all growers have the capacity to access materials and tools to perform vegetative propagation which entails additional cost on their part; hence, seed propagation is still economical for them. An alternative way to better improve the seed and seedling performance of a crop is the utilization of seed priming methods. This is used to break seed dormancy and thus enhance the germination. Among the methods are hydropriming (soaking in water), halopriming (soaking in salt solution) and hormonal priming (soaking in gibberellic acid solution). These methods hydrate the seeds and reduce seedling emergence time [3] and the procedure is easy to perform.

The following are the effects of seed priming methods: allows some of the metabolic processes necessary for germination to occur without germination taking place; increase germination rate; uniform seedling emergence; enhance crop enzyme rate resulted to increase ration of crop development; and faster and better seedling development [12 and 9]. Moreover, those seeds that are different solutions with high osmotic potential prevent the seeds from absorbing in enough water for radicle protrusion which will suspend the seeds in the lag phase [12 and 9]. The faster and better seedling performance is a result of reduced time between sowing and seedling performance and to synchronized emergence [14].

In order to extend help to the growers, avoid the delay in the germination of the seeds and longer germination period, this experiment was conducted to improve the performance of soursop at seedling stage and evaluate the seedling performance of soursop using three seed priming methods.

Specifically, this study aimed to: determine if the seed priming methods are capable of breaking dormancy; determine if it will improve germination rate; determine the seed vigor of soursop; and identify the best seed priming method that provides the best seedling performance.

The results of the study will provide further information on how to improve the seedling performance of soursop which could help interested growers. Another advantage is that if seed priming methods shorten the time to germination then transplanting can be done much earlier.

2. MATERIALS AND METHODS

The study was conducted in a partially shaded area in Barangay 14 San Gregorio, San Nicolas, Ilocos Norte in the Philippines. The area is shaded due to trees of sugar apple or sweetsop (*Annona squamosa*), drumstick tree (*Moringa oleifera*) and mango (*Mangifera indica*).

Ripe soursop fruits were bought from the public market of Laoag City, Ilocos Norte. The seeds were freshly removed from the ripe fruits. Extracted seeds were soaked in water for a minute to see if they float or not. All floating seeds which are immature were discarded. The seeds were stored in a refrigerator to avoid rapid drying of the seeds.

Seeds were scarified by removing small portion of the seed coat using a nail cutter before they were sown. All the seeds were soaked in separate treatments such as hydropriming (water), halopriming (salt solution) and hormonal priming (GA₃) for 24 Hours. For GA₃, the concentration used was 500 ppm [17]. For hydropriming method, tap water was used to soak the seeds. Four teaspoons or 19.7 Grams of salt was dissolved in 500ml of water for the salt solution priming method.

Equal parts of garden soil, carbonized rice hull, and organic fertilizer were placed in polyethylene bag (2x6 inches size) used to grow the soursop seeds. One seed was sown for every polyethylene bag at 2-3 cm depth and irrigation followed after sowing. A total of 20 polyethylene bags were used per treatment per replication giving an overall of 240 bags.

Complete fertilizer (5 g per bag) was dissolved in a Liter of water and applied through fertigation after germination and at 21 days after germination (Pinto and Silva, 1994). Irrigation was done after fertilizer application. Follow-up irrigation was done depending on the soil and weather conditions.

Data Gathering Procedures

Percent Germination Rate

This was recorded at 10 days after emergence (DAE) up to 50 DAE with 10 days interval using the formula below.

$$\text{Percent Germination} = \frac{\text{Total Seeds Germinated}}{\text{Total Number of Seeds Sown}} \times 100$$

Days to Seedling Emergence

This was taken by counting the number of days from sowing up to the time when 50% of the population of the seeds have emerged.

Seed Vigor Index (SVI)

This was done by counting for five times starting at five days after emergence (DAE) up to 25 DAE within five days interval. The SVI was calculated using the formula below [3].

$$SVI = \frac{G1+G2+...+GL}{D1+D2+...+DL}$$

G1 = number of germinants (first count)

D1 = number of days to first count

G2 = number of germinants (second count)

D2 = number of days to second count

GL = number of germinants (last count)

DL = number of days to last count

Seedling Height

The seedling height of four sample seedlings per treatment was measured at 10, 20, 30, 40 and 50 DAE. The measurement was done from ground surface to the tip of the longest leaf using a ruler.

Number of Leaves per Seedling

The same sample seedlings in the measurement of plant height were used to count the number of leaves per seedling. Counting was done at 10 up to 50 DAE with 10 days interval.

Leaf Length

First shoot that appeared in the same sample seedling in measuring plant height was used in measuring leaf length. Measurement was done at 20 up to 50 DAE with 10 days interval.

Shoot and Root Length

Shoot and root length of four sample seedlings were measured from the collar region to the point of attachment of cotyledons and from collar region up to the tip of root, respectively using ruler. This was taken at 40 days after emergence (DAE). To facilitate easy data gathering, the media of the seedling was removed first and the roots are rinsed with water afterwards.

Seedling Length (shoot length plus root length)

This was taken by getting the sum of the measurement of shoot and root length using the same sample seedlings used in shoot and root length measurement.

Seedling Vigor Index

This was recorded by using the formula below used by Das *et al.* (2017).

Vigor Index = Seedling Length x Germination Percentage

Fresh and Oven Dry Weight of Seedling

The sample seedlings used in the measurement of shoot and root length were used to gather fresh and oven dry weight of seedling at 40 DAE. Fresh roots and shoots of the four sample seedlings per treatment were oven dried for two days with the temperature of 65 °C – 70°C [8]. Afterwards, these were weighed using an electronically digital weighing scale.

Data Analysis

All data gathered was analyzed using analysis of variance in a Completely Randomized Design. Where F-test was significant at 5% ($P = .05$) or 1% ($P = .01$), treatment mean difference was further tested using Least Significant Difference (LSD) test at ($P = .05$). Statistical analysis was done using STAR program.

3. RESULTS AND DISCUSSION

3.1 Percent Germination Rate

Seed priming methods significantly ($P = .01$) affected the percent germination rate (PGR) at 10 days after emergence (DAE). Table 1 shows that primed seeds had higher PGR irrespective of priming methods than unprimed seeds. According to Namaz *et al.* (2013) priming improves germination. There was no significant difference between priming methods but numerically, hydropriming (52%) had the highest PGR among the priming methods. Moreover, although germination rate increased from 20 to 50 DAE there were no significant differences between primed and unprimed seeds.

Percent germination was increased and shortened the seed germination period in Judas tree (*Cercis siliquastrum*) with the use of halopriming [7]. Furthermore, hydropriming improve improved seed germination and enhanced seedling emergence of soursop [13]. However, numerically, hydropriming had the highest PGR with 10%

difference from halopriming and hormonal priming. According to El-Barghathi and El-Bakosh (2005) the external application of GA₃ enhances the seed germination of Kermes oaks (*Quercus coccifera*). Similar results were reported by Gonzalez *et al.* (2005) and Singh and Maheswari (2017) that seeds soaked in 500 PPM GA₃ had best and highest seed germination of soursop. Moreover, seeds of tomato soaked in 900 MG L⁻¹ GA₃ have high percentage germination [2].

Also the experiment of Armin *et al.* (2010) on watermelon, KNO₃ increased germination and germination rate.

Table 1. Effects of seed priming methods on percent germination rate of soursop at different observation periods.

| TREATMENT | 10 DAE** | 20 DAE _{ns} | 30 DAE ^{ns} | 40 DAE ^{ns} | 50 DAE ^{ns} |
|-------------------------------------|-----------------|----------------------|----------------------|----------------------|----------------------|
| Unprimed | 22 ^b | 65 | 67 | 75 | 78 |
| Hydropriming (water) | 52 ^a | 72 | 75 | 77 | 80 |
| Halopriming (salt solution) | 42 ^a | 67 | 75 | 88 | 88 |
| Hormonal priming (GA ₃) | 42 ^a | 70 | 73 | 80 | 80 |
| CV (%) | 22.42 | 9.45 | 9.55 | 11.55 | 9.52 |

CV – coefficient of variance

** - significant at 1% level

3.2 Days to Seedling Emergence and Seed Vigor Index

There was no significant difference ($P = .01$) between primed and unprimed seeds on the days to emergence and seed vigor index of soursop (Table 2). Numerically, earliest seed emergence was observed from hormonal priming while unprimed seeds emerged last.

3.3 Seedling Height

At 10 DAE, seedling height was highly significantly affected ($P = .01$) by priming methods while at 40 DAE, seedling height was significantly ($P = .05$) affected by priming methods (Table 3). At 10 DAE, seeds primed with hormone (GA₃) produced highly significantly taller ($P = .01$) seedlings than the other methods and unprimed seeds. Heights of seedlings from hydroprimed and haloprimed seeds did not differ significantly from those of unprimed seeds.

At 40 DAE, hormonal and halopriming seeds had significantly taller ($P = .05$) seedlings than the unprimed seeds. Prasad *et al.* (2002) noted that gladioli are taller with the use of hormonal priming. This means that if the seedlings are taller, it could be transplanted early in the field.

3.4 Number of Leaves per Seedling

The number of leaves per seedling at 30 ($P = .01$) and 50 DAE ($P = .05$) was significantly higher by the use of priming methods than unprimed (Table 4). Generally, hormonal priming consistently produced higher number of leaves per seedling at 30 and 50 DAE than the other priming methods. According to Prasad *et al.* (2002), hormonal priming (250 PPM GA_3) increased the number of leaves in gladioli. Also there was more number of leaves in soursop using GA_3 at 500 ppm soaking in 24 hours than soaking in water for 24 hours [17]. If there is high number of leaves, there will be higher photosynthetic activity which will result in faster growth and development.

Table 2. Effects of seed priming methods on days to emergence and seed vigor index of soursop.

| TREATMENT | DAYS TO SEEDLING EMERGENCE ^{ns} | SEED VIGOR INDEX ^{ns} |
|-----------------------------|---|--------------------------------|
| Unprimed | 32 | 2 |
| Hydropriming (water) | 30 | 3 |
| Halopriming (salt solution) | 30 | 3 |
| Hormonal priming (GA_3) | 29 | 2 |
| CV (%) | 4.45 | 9.97 |

CV – coefficient of variance

ns – not significant

Table 3. Effect of seed priming methods on seedling height (cm) of soursop at different observation periods.

| TREATMENT | 10 DAE ^{**} | 20 DAE ^{ns} | 30 DAE ^{ns} | 40 DAE [*] | 50 DAE ^{ns} |
|-----------------------------|----------------------|-------------------------|----------------------|---------------------|-------------------------|
| Unprimed | 6.60 ^b | 10.13 | 11.46 | 12.14 ^b | 16.18 |
| Hydropriming (water) | 5.48 ^b | 10.15 | 11.99 | 12.87 ^b | 15.95 |
| Halopriming (salt solution) | 6.52 ^b | 11.43 | 12.43 | | 14.18 |
| Hormonal priming (GA_3) | 8.98 ^a | 12.89 | 13.85 | 15.48 ^a | 15.28 |
| CV (%) | 11.92 | 10.66 | 7.69 | 8.03 | 8.88 |

CV – coefficient of variance

** - significant at 1% level

ns – not significant

Table 4. Effect of seed priming methods on number of leaves per seedling of soursop at different observation periods.

| TREATMENT | 20 DAE | 30 DAE | 40 DAE | 50 DAE |
|-------------------------------------|--------|----------------|--------|-----------------|
| | ns | ** | ns | * |
| Unprimed | 2 | 3 ^b | 4 | 6 ^b |
| Hydropriming (water) | 3 | 3 ^b | 5 | 6 ^b |
| Halopriming (salt solution) | 3 | 4 ^b | 5 | 7 ^{ab} |
| Hormonal priming (GA ₃) | 4 | 6 ^a | 6 | 8 ^a |
| CV (%) | 20.38 | 20.41 | 17.04 | 9.33 |

CV – coefficient of variance

** - significant at 1% level

* - significant at 5% level

ns – not significant

3.5 Leaf Length

There was no significant difference in length of leaves produced by primed and unprimed seeds over the period (Table 5) though; hormonal priming consistently had the longest leaf length. Leaf length of gladioli is increased with the use of 500 PPM GA₃, hormonal priming [15].

Table 5. Effect of seed priming methods on leaf length (cm) of soursop at different observation periods.

| TREATMENT | 20 DAE ^{ns} | 30 DAE ^{ns} | 40 DAE ^{ns} | 50 DAE ^{ns} |
|-------------------------------------|----------------------|----------------------|----------------------|----------------------|
| Unprimed | 2.98 | 3.60 | 3.99 | 4.62 |
| Hydropriming (water) | 2.93 | 4.04 | 4.35 | 5.32 |
| Halopriming (salt solution) | 3.84 | 4.15 | 4.65 | 5.24 |
| Hormonal priming (GA ₃) | 4.62 | 6.14 | 4.95 | 5.69 |
| CV(%) | 22.29 | 28.37 | 12.32 | 8.87 |

CV – coefficient of variance

ns – not significant

3.6 Shoot Length and Root Length

Priming methods did not significantly affect the shoot and root length of soursop (Table 6). In the experiment of Singh and Maheswari (2017) soursop shoot and root length was high using 500 ppm GA₃ than using water. However, Matsushima and Sakagami (2013) found that the shoot length of rice increased using hydropriming due to rapid supply of nutrient required for cell growth. Balaguera-Lopez *et al* (2009) also reported that seeds of tomato soaked in 900 mg L⁻¹ GA₃ gave high root length.

Table 6. Effect of seed priming methods on shoot length and root length of soursop.

| TREATMENT | SHOOT LENGTH ^{ns} (cm) | ROOT LENGTH ^{ns} (cm) |
|-------------------------------------|---------------------------------|--------------------------------|
| Unprimed | 12.67 | 11.83 |
| Hydropriming (water) | 12.70 | 12.73 |
| Halopriming (salt solution) | 12.63 | 12.43 |
| Hormonal priming (GA ₃) | 11.60 | 12.73 |
| CV(%) | 5.39 | 10.81 |

CV – coefficient of variance

ns – not significant

3.7 Seedling Length

Seeds either primed or unprimed had comparable seedling length (Table 7). Numerically, hormonal priming had the lowest seedling length and seedling vigor index. KNO₃ has the most effective impact on the seedling growth compared with unprimed, PEG 6000 3%, HCL 0.1N, and NaCl 1.5N [1].

Table 7. Effect of seed priming methods on seedling length (cm) of soursop.

| TREATMENT | SEEDLING LENGTH ^{ns} (cm) |
|-------------------------------------|------------------------------------|
| Unprimed | 25 |
| Hydropriming (water) | 25 |
| Halopriming (salt solution) | 25 |
| Hormonal priming (GA ₃) | 24 |
| CV(%) | 5.85 |

CV – coefficient of variance

ns – not significant

3.8 Fresh Weight and Oven Dry Weight

Priming methods had significant ($P = .05$) influence on the seedling fresh weight but not on dry weight compared to unprimed seeds (Table 8). Primed seeds produced significantly higher seedling fresh weight than unprimed seeds. Among priming methods, hormonal priming had the highest seedling fresh weight. High fresh weight of hormonal priming means higher biomass than the other treatments.

Table 8. Effect of seed priming methods on fresh weight and oven dry weight of soursop.

| TREATMENT | FRESH WEIGHT* (g) | OVEN DRY WEIGHT ^{ns} (g) |
|-----------|-------------------|-----------------------------------|
|-----------|-------------------|-----------------------------------|

| | | |
|-------------------------------------|--------------------|-------|
| Unprimed | 3.26 ^b | 0.96 |
| Hydropriming (water) | 3.60 ^b | 1.16 |
| Halopriming (salt solution) | 4.10 ^{ab} | 1.36 |
| Hormonal priming (GA ₃) | 4.93 ^a | 1.10 |
| CV(%) | 15.64 | 19.92 |

CV – coefficient of variance

* - significant at 5% level

4. CONCLUSION

It can be concluded that priming methods significantly affected the seed and seedling performance of soursop. The seeds of soursop were able to break the dormancy either unprimed or primed taking 29 to 32 days to emerge. One advantage was that primed seeds produced significantly higher percent germination rate than unprimed seeds by 10 days after emergence especially with hydropriming. Both primed and unprimed seeds had the same seed vigor index. However, hormonal priming had the best effective impact due to significant influence because it consistently produced taller seedlings, more leaves per seedling, and comparable seedling fresh weight with other priming methods.

5. References

1. Armin, M., M. Asgharipour, and M. Razavi-Omrani. The effect of seed priming on germination and seedling of watermelon (*Citrullus lanatus*). *Advances in Environmental Biology*. 2010; 4(3): 501-505.
2. Balaguera-Lopez, H.E., J.G. Alvarez, and J.F. Cardenas-Hernandez. Effect of gibberellic acid (GA₃) on seed germination of tomato (*Solanum lycopersicum* L.). *Acta Horticulture*. 2009; 821(821):141-148.
3. Conway, K.E., R. Mereddy, B.A. Khan, Y. WU, S.W. Hallgren, and L. Wu. Beneficial effects of solid matrix chemo-priming in okra. *The American Phytopathological Society*. 2001; 85 (5): 535-537.
4. Das, M., M. Sharma, and P. Sivan. Seed germination and seedling vigor index in *Bixa orellana* and *Clitoria ternatea*. *International Journal of Pure and Applied Bioscience*. 2017; 5(5); 15-19.
5. EL-Barghathi, M.F. and El-Bakkosh, A. 2005. Effect of some mechanical and chemical pretreatments on seed germination and seedling growth of *Quercus coccifera* (Kermes oaks). *J. Jerash Private Univ*. 2005.
6. Gonzalez, J.M., S.G. Gonzalez, A.M. Rosales, F.R. Juarez, M.B. Tene, and F.T. Cervantes. Evaluation of Pregerminative Treatments in Soursop (*Annona muricata*) Seeds. *HORTSCIENCE*. 2005; 40 (4): 1023A-1023.

7. Haroni, N., M. Tabari, and D. Dey. Effect of halopriming treatment in seed germination and seedling emergence of judas tree (*Cercis siliquastrum* L. Caesalpiniaceae). AJAR. 2015; 10: 2355-2362.
8. Ibukunoluwa, M.J.E. 2016. What is the best temperature and duration for oven drying of leaves, stem or root portions of a plant for moisture content determination? Available https://www.researchgate.net/post/What_is_the_best_temperature_and_duartion_for_oven_drying_of_leaves_stem_or_root_portions_of_a_plant_for_moisture_content_determination.
9. Kaur, H., N. Chawla, G. Bassia, and M. Pathakb. Effect of different seed priming treatments on germination of okra. IJCS. 2015; 15: E51-58.
10. Link, R. Soursop (Graviola): health benefits and uses. Available: <https://www.healthline.com/nutrition/soursop-benefits>
11. Matsushima, K.I. and J.I. Sakagami. Effects of seed hydropriming on germination and seedling vigor during emergence of rice under different soil moisture conditions. American Journal of Plant Sciences. 2013; 4: 1584-1593.
12. Nawaz, J., M. Hussain, A. Jabbar, G. Nadeem, M.Sajid, M. Subtain, and I. Shabbir. Seed Priming A Technique. IJACS. 2013; 6-20: 1373-1381.
13. Okoli, N.A., J.C. Obiefuna, R.A. Alagba, and L.C. Emma-Okafor. Improvement of germination and seedling vigour of soursop (*Annona Muricata* L.) using hydro-priming and dehydration treatments. FUTOJNLS. 2013; 2: 57-61.
14. Parera, C.A. and D.J. Cantliffe. 2010. Presowing Seed Priming. Available: <file:///C:/Users/Win7/Downloads/PresowingSeedPriming.pdf>. DOI DOI: 10.1002/9780470650561.ch4
15. Prasad, A., R. Kumar, S. Arya, and K. Saxena. Varietal response of gladioli corms to GA₃ dipping. Journal of Ornamental Horticulture New Series. 2002; 5 (1): 69-70.
16. Rolston, M.P. Water impermeable seed dormancy. The Botanical Review. 1978; 44(3): 365-396.
17. Singh, S.S. and J.U. Maheswari. Influence of pre-sowing of seed treatments on the performance of soursop (*Annona muricata*) seedling. Plant Archives. 2014; 17: 1215-1218.