

1 **SOIL FERTILITY LEVELS IN BANGLADESH FOR RICE CULTIVATION**

2 Jatish C. Biswas<sup>1</sup>, N. Kalra<sup>2</sup>, M. Maniruzzaman<sup>3</sup>, M. M. Haque<sup>3\*</sup>,

3 U. A. Naher<sup>3</sup>, M. H. Ali<sup>4</sup>, W. Kabir<sup>4</sup>, S. Rahnamayan<sup>5</sup>

4  
5 <sup>1</sup>Soil Sci. Div., BRRI & KGF, Bangladesh

6 <sup>2</sup>Agricultural Physics, ICAR, India

7 <sup>3</sup>BRRI, Bangladesh

8 <sup>4</sup>Krishi Gobeshona Foundation, Bangladesh

9 <sup>5</sup>Nature Inspired Computational Intelligence, UOIT, Canada

10

11 \*Corresponding author

12 Md Mozammel Haque

13 E-mal: [mhaquesoil@yahoo.com](mailto:mhaquesoil@yahoo.com)

14 Soil Science Division, Bangladesh Rice Research Institute

15 Gazipur-1701, Bangladesh

16

17

18 **ABSTRACT**

19 **Determination of soil fertility with minimum data set for crop zoning and devising fertilizer**  
20 **recommendations as well as soil fertility evaluation method based on soil properties.** The data  
21 were collected from existing literatures and scoring was done on 0–100 scale. The lowest  
22 score was assigned for the minimum value of tested attributes and then gradually higher  
23 scoring values. Arithmetic, weighted, geometric and most minimum of mean scores were  
24 calculated and their performances were compared with grain yield of dry season irrigated  
25 (Boro) rice. Soil fertility in 10-12 and 39-52% **areas in Bangladesh** are very low and low,  
26 respectively. Medium fertile and fertile soils are distributed in 17-41% and in about 8% areas  
27 of the country. About 55% soils scored 70–95 (medium to high SOC) and the rest belongs to inferior  
28 quality. In some areas P build up has taken place (25% areas), but widespread K mining.  
29 Sulphur and Zn status in about 40% **areas** are low to very low (scored <35 and <40). Soils of  
30 the major areas of the country are with low pH (5.0-6.0) and CEC in the range of 15-25 cmol<sub>c</sub>  
31 kg<sup>-1</sup>. Weighted mean score and most minimum of eight attributes score showed good  
32 relationships with dry season irrigated rice yields than other tested methods indicating that  
33 this technique can be used for soil fertility rating in tropical countries.

34 **Key words:** Soil attributes, Score, Weighted mean, Most minimum mean, Maps

35

36

37

38

39

40

## 41 **1. INTRODUCTION**

42 Global population is increasing and so does the demand for food production, which has  
43 already created tremendous pressure on soil, a finite resource for mankind. It is our obligation  
44 to keep soil healthy and productive through appropriate amendments and crop management  
45 practices [1]. Indigenous nutrient supplying capacity and fertilizer management may make a  
46 soil fertile for one type of crop but could be deficient for the others. So, determination of soil  
47 fertility range would be important not only for producing healthy crops economically but also  
48 for maintaining its productivity for future generations. Soils in Bangladesh are exposed to  
49 high temperatures mostly; plenty of rainfall and greater pressure from growing two or more  
50 crops in a year with or without balanced fertilizations [2] and thus nutrients mining are  
51 widespread. New nutrient deficiencies are emerging [3], and there might be potential hidden  
52 hunger for many others that need to be identified for efficient crop production.

53 Soil fertility varies among regions indicating that variable amounts of fertilizers need to be  
54 applied for different types of crop production. Inadequate dose will impair crop yield, while  
55 overdose can cause not only economic losses but also could be responsible for environmental  
56 pollutions [4]. So, a broad knowledge on soil fertility can provide a better perception on  
57 current nutrient status, distribution patterns and trends [5] that can be obtained through geo-  
58 statistical and geospatial analyses [6,7]. Such analyses help in decision making processes for  
59 precision agriculture and thus for improvement of crop productivity [8,9].

60  
61 Soil fertility can be determined in different ways [10,11] by using soil pH, SOC, P, K,  
62 exchangeable calcium (Ca), magnesium (Mg) and aluminium (Al), S, etc [12,13]. Mbogoni et  
63 al [14] evaluated soil fertility by using average weighted data on SOC, soil pH, total N,  
64 electrical conductivity, C/N ration, available P, exchangeable Ca, Mg and texture for rice  
65 based system productivity improvement. Khaki et al [15] utilized square-root method as  
66 parametric approach and Joint Fuzzy Membership functions to compute soil fertility index

67 (SFI). They found both the system suitable for soil fertility mapping and showed good  
68 relations with rice yield. Desavathu et al [7] used soil pH, EC, N, P and for soil fertility  
69 evaluation through inverse distance weightage interpolation. Thus it is found that researchers  
70 had taken initiative for making soil fertility maps for specific locations or regions but a  
71 simple method for **the** country is still lacking. Therefore, the objective of this study was to  
72 use geo-referenced data on selected soil attributes for preparation of soil fertility maps using  
73 average, weighted mean, geometric mean and most minimum value techniques for  
74 Bangladesh and to establish their relationships with rice yields.

75

## 76 **2. MATERIALS AND METHODS**

77 Data on **SOC, P, S, Zn, and B, CEC**, soil pH and exchangeable K were collected from  
78 Bangladesh Agricultural Research Council website, Soil Resource Development Institute and  
79 existing available literatures. Average Boro clean rice (dry season irrigated crop, hereafter as  
80 Boro rice) yields from 2007 to 2013 were collected from different volumes of Bangladesh  
81 Bureau of Statistics and its relationships were established with soil fertility scores. Although  
82 crop yields vary depending on inherent soil fertility, some other factors like electrical  
83 conductivity, water quality (such as salinity) and its availability, agronomic management  
84 practices, other biotic and abiotic factors also greatly influences crop productivity.  
85 Nonetheless, inclusion of all those factors that influence soil fertility is beyond the scope of  
86 the present investigation.

### 87 **2.1. Scoring criteria and map preparation**

88 Soil nutrient status in Bangladesh has been classified as very low, low medium, optimum and  
89 high based on different ranges (Table 1). This classification system was considered for  
90 assigning scoring values (Table 2) against each selected soil attribute. The scoring scale, as  
91 considered in the present investigation, was 0–100. Attribute-wise soil fertility ratings over  
92 different locations of Bangladesh were made by using MS-Excel Macros and IDRISI3.2.

93  
 94 Soil fertility scores, as determined by arithmetic mean (AM), geometric mean (GM),  
 95 weighted mean and (WM) and most minimum attribute (MAtrib<sub>score</sub>) techniques, were used to  
 96 find out their relationships with Boro rice yields (64 districts of Bangladesh from 2007 to  
 97 2013) through regression analyses. Considering higher R<sup>2</sup> values, final soil fertility rating  
 98 maps were prepared based on weighted mean scores (Equa. I) and scores of the most  
 99 minimum of eight parameters for each district (Equa. II). Among soil attributes the most  
 100 limiting factors dictate crop yield, so we have provided weight to such factors in determining  
 101 WM as follows:

102  
 103 
$$WM = ([SOC_{score}] * [P_{score}] * [K_{score}] * [CEC_{score}] * [pH_{score}])^{(1/5)} * 0.5 + [S_{score}] * 0.25 +$$
  
 104 
$$([Zn_{score}] * [B_{score}])^{(1/2)} * 0.25 \dots\dots\dots (I)$$

105  
 106 where, SOC<sub>score</sub> is the soil organic carbon, P<sub>score</sub>, K<sub>score</sub>, CEC<sub>score</sub>, pH<sub>score</sub>, S<sub>score</sub>, Zn<sub>score</sub> and  
 107 B<sub>score</sub> stand for P, K, CEC, soil pH, S, Zn and B scores, respectively.

108 MAtrib<sub>score</sub> for selected eight soil parameters were determined as follows:

109 
$$MAtrib_{score} = \text{Geomean} (\text{Small}(\text{Atrib1:Atrib8,1}), \text{Small} (\text{Atrib1:Atrib8,2}), \dots\dots\dots, \text{Small}$$
  
 110 
$$(\text{Atrib1:Atrib8,8})) \dots\dots\dots (II)$$

111 where, Atrib1, 2, 3, ....., 8 are the soil parameters considered first, second, etc.

112  
 113 GM score was calculated as follows:

114 
$$GM = ([B_{score}] * [K_{score}] * [P_{score}] * [CEC_{score}] * [pH_{score}] * [SOC_{score}] * [S_{score}] * [Zn_{score}])^{(1/8)} \dots (III)$$

115  
 116 AM score was computed as follows:

117 
$$AM = ([B_{score}] + [K_{score}] + [P_{score}] + [CEC_{score}] + [pH_{score}] + [SOC_{score}] + [S_{score}] + [Zn_{score}]) / 8 \dots\dots (IV)$$

118

119 Scores for most minimum of 1, 2, 3, 4, 5, 6 and 7 soil attributes were also found out in similar  
120 fashion of Equa. II. The maps of tested attributes were prepared by using IDRISI3.2. Soil  
121 fertility rating maps on the basis of WM and most minimum of eight attributes were used for  
122 soil fertility delineation in Bangladesh. The other maps prepared based on different  
123 techniques were used as supplementary figures 1.

### 124 **3. RESULTS**

125 Soil organic carbon, a vital component of fertility index showed >95 score for about 25%  
126 areas in Bangladesh (Fig. 1a). About 55% soils had 70–95 score (medium to high SOC) and  
127 the rest belongs to inferior quality. The scores for soil P varied from <10 to >75 in which  
128 very low (<7 ppm), low (7-15 ppm), optimum (15-30 ppm) and high (>30 ppm) P levels  
129 covered about 22.64, 47.74, 12.98 and 16.64 percent areas in the country (Fig. 1b). About  
130 25% soils are with optimum/high (>80 score) K fertility. Majority areas (~43%) bear low K  
131 (0.091-0.18 meq 100 g<sup>-1</sup> soil) and the rest belong to very low (<10 score) and medium (40-80  
132 score) K categories y (Fig. 2a). The least score (<10) for S indicated that about 15.62% soils  
133 are very poor (<7.5 ppm); 26.04% low and 14.54% medium and 43.79% areas are with  
134 optimum/high S fertility status (Fig. 2b). In about 37.61% areas (score >75), soil Zn contents  
135 are optimum to high (>1.351 ppm), 20.77% areas (40-75 score) are with medium Zn  
136 containing soils and 41.61% soils scored <10 to 40 indicating (Fig. 3a) that Zn application is  
137 a must practice for Bangladesh. **The content of B** is very low to low in about 50% soils (score  
138 <10 to 40) and rest of the soils had medium B content (Fig. 3b).

139

140 Soil pH score varied from <25 to >85 depending on locations and soil types in the country.  
141 Maximum area coverage was 44.59% followed by 32.25% in the pH range of 5.0-6.0 and 6.5-  
142 7.5, respectively (Fig. 4a). Soil pH below 5.0 and above 7.5 covers about 7.29% areas of the  
143 country. The rest of the soils (15.22% areas) are with pH range of 6.0-6.5. The CEC scores  
144 ranged from <25 to >85 depending on location in the country (Fig. 4b). The CEC of major

145 soils (47.46%) are 15-25  $\text{cmol}_c \text{ kg}^{-1}$  followed by less than 15  $\text{cmol}_c \text{ kg}^{-1}$  in 37.72% areas of  
146 the country. Higher CEC ( $>25 \text{ cmol}_c \text{ kg}^{-1}$ ) was found in 14.81% areas only.

147  
148

### 149 **3.1. Soil fertility relationships with rice yield**

150 Soil fertility score based on different techniques and their relationships with clean rice yields  
151 are shown in Fig. 5. About 49% yield variabilities are explained by the WM and most  
152 minimum of eight tested soil attributes score (MAAttrib-8). The performances of AM and GM  
153 techniques in explaining yield variabilities were the least compared to others. Most minimum  
154 1-7 soil attributes score explained Boro rice yield variabilities by about 23-42%.

155

### 156 **3.2. Soil fertility status**

157 Soil fertility scores varied from  $<35$  to  $>60$  with WM score technique and it was  $<35$  to  $>55$  with the  
158 MAAttrib-8 (Fig. 6). In the lowest soil fertility score ( $<35$ ), area coverages are 10-12% of the country  
159 based on above stated two techniques. The largest areas (28-30%) fall within the score of 40-45 under  
160 both the techniques. Areas covered by higher scores ( $>55$ ) were only about 16% of the country. Soil  
161 fertility scores of 35-40 represented 9.41% and 24.08% areas under WM and MAAttrib-8 techniques,  
162 respectively. Similarly, 45-50 and 50-55 scores under WM and MAAttrib-8 represented about 20% and  
163 11-16% areas, respectively of the country. Based on GM, AM and MAAttrib-1 to MAAttrib-7 soil  
164 fertility score varied greatly and represented different areas of the country, but major areas showed  
165 low fertility score (data not shown). There were variations in the highest and the lowest scores  
166 because of method employed (Table 3). The standard deviations were  $\pm 8.52$ ,  $\pm 7.19$ , 7.73 and  $\pm 8.52$   
167 for GM, AM, WM and MAAttrib-8 means score, respectively having corresponding co-efficient of  
168 variations of 19.62%, 14.22%, 16.58% and 19.62%.

## 169 **4. DISCUSSION**

170 In about 29% areas of the country, the SOC was at medium category; although there are high  
171 and very high SOC in certain areas, especially with peat soils. In general, SOC was higher in  
172 low lying areas, the single cropped zones, which remain 5–6 months under water in a year.

173 This level of SOC specifically in about 18% areas of the country is still inadequate for  
174 satisfactory crop production [16]. As population pressure is increasing, farmers are using  
175 such lands to increase total production through cropping intensification resulting in depletion  
176 of SOC along with other plant nutrients. The decrease rate of SOC is comparatively faster  
177 with arable cropping over time [17] with or without addition of organic manures. So, we have  
178 found lower SOC rating in intensely cropping zones of Bangladesh. Partial productivity of  
179 applied fertilizers is also decreasing indicating that nutrients from organic matter (OM) need  
180 to be added that has been observed in our experiments at BRRI. Most soils showed good  
181 response when OM was incorporated either from poultry litter, cow dung, vermicompost  
182 [18,19] or green manuring because SOC influences soil pH, buffering capacity, nutrient  
183 supplies and soil biological activity [20].

184  
185 Although available P in the category of very low and low cover a larger area (about 70%), in  
186 some areas **its high** has taken place (Fig. 1b) because of cropping patterns followed, fertilizer  
187 management options and inherent characteristics of parent materials [16,21]. As a greater  
188 area suffers from available P, corrective measures have to be taken for profitable production  
189 [22]. This scenario is also true for global perspective in which P is depleting by  $5.1 \text{ kg ha}^{-1} \text{ yr}^{-1}$   
190 [23]. However, majority of the farmers in Bangladesh prefer to add N fertilizer because of  
191 its immediate visible effects [2] and thus nutrient imbalance impose negative impact on soil  
192 properties and crop production as a whole.

193  
194 Potassium levels in major areas were very low to low (Fig. 2a) indicating that K mining was  
195 taking place because of its substandard dose used by the farmers. Since farmers generally use  
196 more N fertilizer and minimum K rate, the later is depleting rapidly in many areas of  
197 Bangladesh [21, 22, 24 ]. In the global perspective, K is also depleting by  $38.8 \text{ kg ha}^{-1} \text{ yr}^{-1}$



198 [23]; although its build up is not either uncommon in some areas because of excessive use  
199 with certain crops [16].

200 Though S and Zn deficiencies are widespread in the country, its wet and dry depositions are  
201 also taking place because of industrial development [25]; but S fertilizer application still  
202 improves rice yields in many areas of the country [36].

203 The scenario of B fertility is not healthy because in some areas it has depleted severely over  
204 time [16]. Yields of wheat, mustard and papaya reduce greatly in many parts of the country  
205 without B application. The depletion of soil fertility in areas with high cropping intensities  
206 [26] indicated that replenishment of removed nutrients were not taking place or it is beyond  
207 the capacity of the soils to supply major nutrients for growing high yielding crop varieties.  
208 There are evidences that Zn and B contents have been depleted severely from 1991 to 2012 in  
209 some selected areas of Bangladesh and thus crop productivity is declining [16].

210  
211 Lower soil pH covers quite larger areas in the north and north-east part and higher pH in the  
212 southern part of the country where plant nutrients availability is a limiting factor for  
213 satisfactory crop production without proper amendment. In some cases soil pH is increasing,  
214 especially in northern part of the country and thus playing a negative role on nutrient  
215 availability. It was reported that nutrient availability from applied fertilizers may be  
216 unavailable by more than 33-75% if soil pH ranges from 4.5 to 5.5 [27]. Generally, major  
217 nutrients are available for plants when soil pH varies from 6.5 to 7.5 [28]. Among others, soil  
218 P and many micronutrients become unavailable when pH exceeds 7.5, but molybdenum (Mo)  
219 availability increases in alkaline pH. Moreover, CEC also depends on soil pH in a neutral soil  
220 will have higher CEC than acidic soils [29]. Low CEC indicates silty loam soils having  
221 tendency of K and Mg deficiencies and faster decrease in soil pH [29, 30]. In such situations  
222 frequent liming is needed for sandy type soils for profitable crop cultivation.

223

224 We have seen good relationships ( $R^2 = 0.49$ ) of WM and MAttrib-8 scores with rice yields,  
225 which is similar to the findings of Vasu et al. [31]. In Bangladesh, no grouping of soils has  
226 been made based on combine scores or combine effects of different soil attributes; but  
227 component-wise soil fertility delineations are available [22,26, 32]. So, our efforts are to  
228 group soil fertility status combining all tested attributes as score <35 (very low fertility), 35-  
229 45 (low fertility), 45-55 (medium fertile) and >55 (fertile). Accordingly, 10-12 and 39-52%  
230 areas of the country represented very low and low soil fertility, respectively (Fig. 6). Medium  
231 fertile and fertile soils are distributed in 17-41% and in about 8% areas of the country. These  
232 findings clearly indicate that special cares are needed for efficient and economic crop  
233 production in major areas of Bangladesh. To maintain soil fertility and to obtain optimum  
234 grain yields of rice emphasis should be provided for P management in north-east part (Fig.  
235 1b); K management in north and north-west regions (Fig. 2a); S management in north-west  
236 and south-central parts (Fig. 2b); Zn management in central and southern regions (Fig. 3a);  
237 and pH management in north-west part (Fig. 4a) of the country. However, crop yields not  
238 only depend on soil fertility, but also on other factors like water availability, temperature, and  
239 so on. Moreover, soil fertility scores alone cannot explain yield variability of a crop rather it  
240 can provide an indication for fertilizer rate determination and crop zoning for profitable  
241 farming.

242  
243 Population pressure is increasing in Bangladesh, while soil fertility is decreasing indicating  
244 that we are manipulating our soils beyond its bearing capacity. In general, nutrient mining is  
245 taking place in Bangladesh at about  $100 \text{ kg ha}^{-1} \text{ yr}^{-1}$  [22, 33] also reported low to very low  
246 soil fertility for most of the studied soils in Bangladesh. This scenario is also true in terms of  
247 global scale where soil fertility problems are associated with human-induced nutrient  
248 depletion [23]. Besides, soil nutrient availability is limiting in cultivated lands of tropical  
249 countries because of low inherent soil fertility [34]. Calcium deficiencies are emerging in

250 some agro-ecological zones (AEZ-3 and 21) of Bangladesh [3] and there might be hidden  
251 hunger for micronutrients and thus reducing soil fertility and ultimately crop yield, but not  
252 considered in the present investigation because of unavailability of data for the whole  
253 country. In time series analyses for nutrient depletion, it was found that the contents of  
254 exchangeable K, Ca and Mg have declined in all physiographic units except Old Himalayan  
255 Piedmont and Madhupur Tract after 27 years of crop cultivation [35]. In one of our study, it  
256 was also found that soil nutrient ratios have been changed in many places of Bangladesh  
257 because of over exploitation of inherent soil fertility and thus Ca:P and N:Zn were playing  
258 significant negative role with wet season rice yields under unfavourable ecosystems of  
259 Bangladesh [36]. Similarly P: K ratio was acting antagonistically in agricultural ecological  
260 zone 3, 18 and 26 of Bangladesh. All these factors indicate that we have to know our soils  
261 before its use for crop production. Determination of soil fertility status by combining  
262 important but minimum attributes can help in this regard for profitable farming and to  
263 recuperate soil fertility through crop and fertilizer management.

264

## 265 **5. CONCLUSION**

266 A simple method of soil fertility evaluation for a country with minimum data sets is very  
267 much desirable for proper crop zoning and delineating agronomic management options for  
268 satisfactory crop production. We have determined soil fertility scores using pH, CEC, SOC,  
269 available P, S, Zn, B and exchangeable K and following geometric, arithmetic, weighted and  
270 mean approaches along with most minimum of tested attributes score. Weighted mean and  
271 most minimum of soil attribute scoring methods showed better relationships with dry season  
272 irrigated rice yields in Bangladesh indicating that this technique can be employed for soil  
273 fertility assessment and its subsequent use for crop zoning and for determination of fertilizer  
274 rates in similar environments around the globe.

275

276 **ACKNOWLEDGEMENT**

277 We greatly acknowledge Krishi Gobeshona Foundation for their financial support in this  
278 research activity through Modelling Climate Change Impact Assessment in Bangladesh  
279 (CRP-II) project.

280

281 **REFERENCES**

282

- 283 1. MacCarthy DS, Agyare WA, Vlek PLG, Adiku SGK. Spatial variability of some soil  
284 chemical and physical properties on agricultural landscape. *West Afr. J. Appl. Ecol.*  
285 2013; 21: 47-61.
- 286 2. Biswas JC, Maniruzzaman M, Sattar MA, Neogi MG. Improvement of rice yield  
287 through fertilizer and cultural management at farmer's field. *Bangladesh Rice J.* 2008;  
288 13: 9-14.
- 289 3. Saha PK, Islam S, Islam MN, Biswas JC, Haque MM. Soil plant nutrient status under  
290 intensive rice-farming systems in unfavourable eco-system of Bangladesh. *Intl. J.*  
291 *Biol. Pharma. Sci.* 2016; 2: 1-11.
- 292 4. Yang Y, Zhang S. Approach of developing spatial distribution maps of soil nutrients.  
293 In L. Daoliang (ed.) *Proc IFIPTC on Computer and Computing Tech. Agric.* 2008;  
294 1:565-571.
- 295 5. Dafonte JD, Ulloa GM, Jorge P, Glécio S, Vázquez M, Vidal E. Mapping of soil  
296 micronutrients in an European Atlantic agricultural landscape using ordinary kriging  
297 and indicator approach. *Bragantia* 2010; 69: 175-186.
- 298 6. Behera SK, Shukla AK. Spatial distribution of surface soil acidity, electrical  
299 conductivity, soil organic carbon content and exchangeable potassium, calcium and  
300 magnesium in some cropped acid soils of India. *Land Degrad. Dev.* 2015; 26: 71-79.

- 301 7. Desavathu RN, Nadipena AR, Peddada JR. Assessment of soil fertility status in  
302 Paderu Mandal, Vaisakhapatnam district of Sndra Pradesh through Geopsatial  
303 techniques. *The Egyptian J. Remote Sensing Space Sci.* 2018; 21: 73-81.
- 304 8. Markoski M, Arsov S, Mitkova T, Stamenkovska J. The benefit of GIS technologies  
305 and precision agriculture principles in soil nutrient management for agricultural crop  
306 production. *Bulg. J. Agric. Sci.* 2015; 21: 554-559.
- 307 9. Rahman AMAE, Natarajan A, Hedge R. Assessment of land suitability and capability  
308 by integrating remote sensing and GIS for agriculture in Chamarajanagar district,  
309 Karnataka, India. *Egypt. J. Remote Sensing Space Sci.* 2016; 19: 125-141.
- 310 10. Khalid R, Mahmood T, Bibi R, Siddique MT, Alvi S, Naz SY. Distribution and  
311 indexation of plant available nutrients of rainfed calcareous soils of Pakistan. *Soil  
312 Environ.* 2012; 31: 146-151.
- 313 11. Salehi N, Sepanlow MG, Gorzin BJ. An evaluation of soil fertility using soil organic  
314 carbon, potassium, phosphorus and salinity factors for rice cultivation by fuzzy logic  
315 and AHP technique. *Int. J. Agri. Crop Sci.* 2013; 5: 2233-2241.
- 316 12. Moran EF, Brondizio ES, Tucker JM, da SilvaForsberg MC, McCracken S, Falesi I.  
317 Effects of soil fertility and land use on forest succession in Amazonia. *For. Ecol.  
318 Manag.* 2010; 139: 93-108.
- 319 13. Panwar P, Pal S, Reza SK, Shama B. Soil fertility index, soil evaluation factor, and  
320 microbial indices under different land uses in acidic soil of humid subtropical India.  
321 *Commun. Soil Sci. Plan Anal.* 2011; 42: 2724-2737.
- 322 14. Mbogoni JDJ, Kiwambo BJ, Urassa GJ, Assenga SV. Soil fertility appraisal for  
323 enhancing productivity in rice-based system of Ruvu basin, Agricultural Research  
324 Institute Mlingano, Tanga, Tanzania; 2011.

- 325 **15.** Khaki BD, Honarjoo N, Davatgar N, Jalalian A, Golsselfidi HT. Assessment of two  
326 soil fertility indexes to evaluate paddy fields for rice cultivation. *Sustainability* 2017;  
327 9: 1299b (doi:10.3390/su9081299).
- 328 **16.** Siddique MMNEA, Halim MA, Kararuzzaman M, Karim D, Sultana J. Comparative  
329 insights for investigation of soil fertility degradation in a piedmont area which cover  
330 the Anjamkhor union of Baliadangi upazila, Thakurgaon, Bangladesh. *IOSR J.*  
331 *Environ. Sci. Toxic Food Tech.* 2014; 8: 82-87.
- 332 **17.** Zahid AM, Hossain MB, Halim MA, Hossain MA, Shahreen F. Organic matter and  
333 plant nutrient depletion in major soil series in the high Ganges river floodplain. *Intl. J.*  
334 *Sustain. Agric. Tech.* 2011; 7: 30-37.
- 335 **18.** Saleque MA, Abedin MJ, Bhuiyan NI, Zaman SK, Panaullah GM. Long-term effects  
336 of inorganic and organic fertilizer sources on yield and nutrient accumulation of  
337 lowland rice. *Field Crop Res.* 2004; 86: 53-65.
- 338 **19.** Haque MM, Akter M, Biswas JC. Performance of vermicompost and poultry manure  
339 on rice yield and soil health. *Annual Research Review Workshop 2016-17. Soil*  
340 *Science Division, Bangladesh Rice Research Institute, Gazipur. 75pp; 2017.*
- 341 **20.** Karlen DL, Andrews SS, Wienhold BJ, Zobeck TM. Soil quality assessment: past,  
342 present and future. *J. Integrated Biosci.* 2008; 6: 3-14.
- 343 **21.** FRG (Fertilizer Recommendation Guide). Bangladesh Agricultural Research Council,  
344 Farmgate Dhaka, 1215. In press; 2018.
- 345 **22.** Shil NC, Saleque MA, Islam M R, Jahiruddin M. Soil fertility status of some of the  
346 intensive crop growing areas under major agro-ecological zones of Bangladesh.  
347 *Bangladesh J. Agril. Res.* 2016; 41: 735-757.
- 348 **23.** Tan ZX, Lal R, Wiebe KD. Global soil nutrient depletion and yield reduction. *J.*  
349 *Sustainable Agric.* 2005; 26: 123-146.

- 350 **24.** Rahman MH, Islam MR, Jahiruddin M, Rafii MY, Ismail MR, Malek MA.  
351 Fertilization for increased crop production and nutrient balance in the maize-legume-  
352 rice cropping pattern. *J. Food Agric. Environ.* 2013; 11: 653-656.
- 353 **25.** Biswas JC, Haque MM, Akter M, Hossain ATMS, Khan FH, Baki MZI, Sarker ABS,  
354 Islam MR. Element Composition of the Atmospheric Depositions in Bangladesh. *J.*  
355 *Environ. Protec.* 2018; 9:948–956.
- 356 **26.** Jahiruddin M, Satter MA. Land and soil resource management report, in: *Research*  
357 *Priority in Agriculture and Development Vision Document-2030 and Beyond*; 2010.
- 358 **27.** CU (Cornell University). How dose soil pH affect fertilizers and fertility. Cornell  
359 University Cooperative Extension of Suffolk County. Extension Education Center,  
360 423 Griffing Avenue, NY ([https://s3.amazonaws.com/assets.cce.cornell.edu/](https://s3.amazonaws.com/assets.cce.cornell.edu/attachments/3271/ow_does_soil_pH_affect_fertilizers_and_fertility.pdf?1413386058)  
361 [attachments/ 3271/ow\\_does\\_soil\\_pH\\_affect\\_fertilizers\\_and\\_fertility.pdf?1413386058](https://s3.amazonaws.com/assets.cce.cornell.edu/attachments/3271/ow_does_soil_pH_affect_fertilizers_and_fertility.pdf?1413386058)  
362 (Access on 6-2-2019).
- 363 **28.** IPNI (International Plant Nutrition Institute). Soil pH and the availability of plant  
364 nutrients. IPNI, Georgia 30092-2806 USA ([www.ipni.net](http://www.ipni.net)); 2010.
- 365 **29.** CU (Cornell University). Cation exchange capacity (CEC). Agronomy fact sheet  
366 series, Fact sheet number 22. Cornell University Cooperative Extension, College of  
367 Agriculture and Life Sciences (<http://nmisp.css.cornell.edu>); 2007.
- 368 **30.** Noble AD, Gillman GP, Ruaysoongnern S. A cation exchange index for assessing  
369 degradation of acid soil by further acidification under permanent agriculture in the  
370 tropics. *European J. Soil Sci.* 2007; 51: 233-243.
- 371 **31.** Vasu D, Singh SK, Ray SK, Duraisami VP, Tiwary P, Chandran P, Nimkar AM,  
372 Anantwar SG. Soil quality index (SQI) as a tool as a tool to evaluate crop productivity  
373 in semi-arid Deccan plateau, India. *Geoderma* 2016; 282: 70-79.

- 374 **32.** Rijmpa J, Jahiruddin M. Final; report on national strategy and p[lan for use of soil  
375 nutrient balance in Bangladesh. SFFP consultancy report, DAE, Khamarbari, Dhaka;  
376 2004.
- 377 **33.** Hossain M, Bayes A Islam SMF. A Diagnostic Study on Bangladesh Agriculture.  
378 RED Working Paper Series, BRAC; 2017.
- 379 **34.** FAO (Food and Agriculture Organization). Status and trends in land and water  
380 resources. Pages 19-60 in The State of the World's land and Water Resources for  
381 Food and Agriculture: Managing Systems At Risk. FAO, UN and Earthscan; 2011.
- 382 **35.** Ali MM, Shaheed SM, Kubota D, Masunaga T, Wakatsuki T. Soil degradation during  
383 the period 1967 – 1997 in Bangladesh. II. Selected chemical characters. Soil Sci.  
384 Plant Nut. 1997; 43: 879-890.
- 385 **36.** Biswas JC, Haque MM, Saha PK. Rice yield potential under unfavorable soil  
386 ecosystems in Bangladesh. Asian J. Soil Sci. Plant Nutr. 2017;1: 1-10.

387  
388

389 **Figure titles**

- 390 Fig. 1. Status of (a) soil organic carbon and (b) phosphorus in Bangladesh
- 391 Fig. 2. Status of (a) soil potassium and (b) sulphur in Bangladesh
- 392 Fig. 3. Status of (a) soil zinc and (b) boron in Bangladesh
- 393 Fig. 4. Distribution patterns of (a) soil pH (b) CEC in different parts of Bangladesh
- 394 Fig. 5. Relationships of clean Boro rice yields with scores of different soil attributes,  
395 Bangladesh
- 396 Fig. 6. Soil fertility variations in Bangladesh as per (a) weighted mean and (b) most minimum  
397 of eight soil-attributes scores
- 398 **Supplementary. Fig. 1.** Soil fertility for Bangladesh according to (a) arithmetic mean, (b)  
399 geometric mean, (c) one most minimum, (d) two most minimum, (e) three most minimum, (f)



400 four most minimum and (g) five most minimum, (h) six most minimum and (i) seven most

401 minimum attributes

402

403

404


405

406

407

408

409



410

411


Fig. 1 a & b

412

413

414

415



416


417

Fig. 2 a & b

418

419

420



421


422

Fig. 3 a & b

423

424

425



426

427

428

**Fig. 4 a & b**

429

430

431

432

433

434

435

436

437




438

439




440

441




442





443



444



445



446

447

448

**Fig. 5**

449


450

451

452

453

454



**Fig. 6 a & b**

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477 **Table 1. Soil nutrient status and its classifications in Bangladesh**

|  | Critical<br>limit | Very<br>low | Low         | Medium      | Optimum      | High        |
|--|-------------------|-------------|-------------|-------------|--------------|-------------|
| SOC (%)                                  | -                 | <0.336      | 0.337-0.574 | 0.575-1.148 | 1.489-2.308* | >2.308**    |
| Olsen P (mg dm <sup>-3</sup> )           | 10                | <7.50       | 7.51-15.00  | 15.1-22.5   | 22.51-30.00  | 30.1-37.50  |
| Bray P (mg dm <sup>-3</sup> )            | 7                 | <5.25       | 5.25-10.50  | 10.51-15.75 | 15.76-21.00  | 21.10-26.25 |
| S (mg dm <sup>-3</sup> )                 | 10                | <7.50       | 7.51-15.00  | 15.1-22.5   | 22.51-30.00  | 30.1-37.50  |
| K (cmol <sub>c</sub> dm <sup>-3</sup> )  | 0.12              | <0.09       | 0.091-0.18  | 0.181-0.27  | 0.271-0.36   | 0.361-0.45  |
| Ca (cmol <sub>c</sub> dm <sup>-3</sup> ) | 2                 | <1.50       | 1.51-3.00   | 3.1-4.50    | 4.51-6.00    | 6.1-7.50    |
| Mg (cmol <sub>c</sub> dm <sup>-3</sup> ) | 0.5               | <0.0375     | 0.376-0.75  | 0.751-1.25  | 1.16-1.50    | 1.51-1.875  |
| Cu (mg dm <sup>-3</sup> )                | 0.6               | <0.15       | 0.151-0.30  | 0.31-0.45   | 0.451-0.60   | 0.61-0.75   |
| Zn (mg dm <sup>-3</sup> )                | 0.2               | <0.45       | 0.451-0.90  | 0.91-1.35   | 1.351-1.81   | 1.81-2.25   |
| Fe (mg dm <sup>-3</sup> )                | 4                 | <3.00       | 3.10-6.00   | 6.1-9.00    | 9.1-12.00    | 12.1-15.00  |
| Mn (mg dm <sup>-3</sup> )                | 1                 | <0.75       | 0.756-1.50  | 1.51-2.25   | 2.56-3.00    | 3.1-3.75    |
| B (mg dm <sup>-3</sup> )                 | 0.2               | <0.15       | 0.151-0.30  | 0.31-0.45   | 0.451-0.60   | 0.61-0.75   |
| Mo (mg dm <sup>-3</sup> )                | 0.1               | <0.075      | 0.076-0.15  | 0.151-0.225 | 0.226-0.30   | 0.31-0.375  |

478 FRG, 2012; \*High and \*\*Very high

479  
480  
481  
482

483 **Table 2. Scoring criteria for different nutrient levels**

| Soil nutrients |       | Soil pH |       | SOC         |       | CEC   |       |
|----------------|-------|---------|-------|-------------|-------|-------|-------|
| Status         | Score | Range   | Score | Range       | Score | Range | Score |
| Very low       | 5     | <5.0    | 25    | <0.336      | 40    | <5    | 25    |
| Low            | 30    | 5.0-5.5 | 45    | 0.337-0.574 | 70    | 5-10  | 40    |
| Medium         | 70    | 5.5-6.0 | 65    | 0.575-1.148 | 85    | 10-20 | 65    |
| Optimum        | 100   | 6.0-6.5 | 75    | 1.489-2.308 | 95    | 20-30 | 75    |
| High           | 100   | 6.5-7.0 | 85    | >2.308      | 100   | 30-40 | 80    |
|                |       | 7.0-7.5 | 95    |             |       | 40-50 | 85    |
|                |       | >7.5    | 25    |             |       | >50   | 100   |

484

485

486

487

488 **Table 3. Soil fertility scoring variations due to methods**

|              | Geometric<br>mean | Arithmetic<br>mean | Weighted<br>mean | MAttrib-8 |
|--------------|-------------------|--------------------|------------------|-----------|
| Maximum      | 60.19             | 63.38              | 61.55            | 60.19     |
| Minimum      | 24.20             | 31.20              | 30.10            | 24.20     |
| Mean         | 43.42             | 50.57              | 46.60            | 43.42     |
| Sd ( $\pm$ ) | 8.52              | 7.19               | 7.73             | 8.52      |
| CV(%)        | 19.62             | 14.22              | 16.58            | 19.62     |

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506


507

508




509


510



511



512



513

514

515

516

517

518

Supplementary Fig. 1