

DYNAMIC BUCKLING OF A CLAMPED FINITE COLUMN RESTING ON A NON – LINEAR ELASTIC FOUNDATION

¹BASSEY, JULIUS E., ¹ETTE, ANTHONY M., * ¹CHUKWUCHEKWA, JOY U. AND ¹OSUJI, ATULEGGWU C.

¹ DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF TECHNOLOGY, OWERRI, IMO STATE NIGERIA.

E-MAIL: juli_real2k4@yahoo.com, tonimonsette@yahoo.com, joyuchekwa@gmail.com, aatac2000@gmail.com.

*CORRESPONDING AUTHOR - joyuchekwa@gmail.com

ABSTRACT: The analysis of the dynamic buckling of a clamped finite imperfect viscously damped column lying on a quadratic-cubic elastic foundation using the methods of asymptotic and perturbation technique is presented. The proposed governing equation contains two small independent parameters (δ and ϵ) which are used in asymptotic expansions of the relevant variables. The results of the analysis show that the dynamic buckling load of column decreases with its imperfections as well as with the increase in damping. The results obtained are strictly asymptotic and therefore valid as the parameters δ and ϵ become increasingly small relative to unity.

Keywords: Dynamic Buckling, Viscous damping, asymptotics and perturbation technique, Column-like elastic structures.

2010 AMS Classification: 00A69

1.0 INTRODUCTION

Buckling is a phenomenon associated with failure of column-like structures. Structures on non-linear elastic foundations are commonly used in engineering applications and occupy a prominent place in structural mechanics. These structures can also serve as simplified models for complex non-linear systems such as columns, shells and plates. Globally, collapse of buildings, bridges and other material structures are issues of concern. Structural failures are forms of material failures which are dangerous in nature and should be prevented by all cost. Series of investigations and studies have been done by Engineers and Applied Mathematicians to determine the maximum loads structures can carry before buckling occurs, yet buckling of elastic structures remain inevitable. Structural elastic materials normally display certain tendencies of failures and instability when loaded either statically or dynamically and one of the pre-occupations of the Structural Engineers and Applied Mathematicians is the determination of the load which a given elastic material can support prior to buckling.

A vast quantum of insights on dynamic stability of elastic structure has been achieved by subjecting these materials to diverse dynamic loading conditions. These loads include, step loading, impulsive loading, rectangular loading, triangular loading [1] and even periodic loading [1] and [2]. From these findings, it has become firmly established that initial imperfections, and to a lesser extent, the

loading duration, are some of the main factors that have been seriously implicated as causative agents of reduction of the elastic strength of these materials. [3] investigated the dynamic response of columns under impulsive axial compression. The investigation has been carried out on clamped specimens, made of metals and composite materials, loaded impulsively by a striking mass. In the theoretical study Rayleigh-type beam equations were assumed for a geometrically imperfect column of a linear-elastic anisotropic material, and the numerical solution, yielded buckling behaviour that correlated well with the experimental results. The results have shown that initial geometrical imperfections, duration of impulse and effective slenderness have a major influence on the buckling loads, whereas the effect of the material is secondary. Recent studies on dynamic buckling have been directed principally on columns, beams, plates, spherical shells and cylindrical shells, and so, extensive literatures (most often numerical approach), have since come to limelight. In this regard, mention must be made of [4], who studied some important parameters in dynamic buckling analysis of plated structures subjected to pulse loading, while [5] equally investigated the buckling of impulsively loaded prismatic cores. In the same token, [6] studied the dynamic buckling of thin-walled composite plates with varying width-wise material properties while [7] also investigated interactive dynamic buckling of thin-walled columns. We now mention [8], who studied the dynamic buckling of thin-walled viscoplastic columns, while [9] similarly investigated some aspects of dynamic buckling of plates under in-plane pulse compression. A study on longitudinal step-wise loading was undertaken by [10], while [11] investigated triply coupled vibrations of axially loaded thin-walled composite beams. An investigation on computational nonlinear stochastic dynamics was undertaken by [12], while [13] discussed nonlinear stochastic dynamical post buckling analysis of uncertain cylindrical shells. Similarly, [14] as well as [15], and [16] made excellent contributions to the dynamics of dynamic buckling. An investigation into the dynamic effect of lateral buckling of high temperature/high pressure offshore pipeline was carried out by [17]. In the same token, [18] investigated the dynamic buckling and fragmentation in brittle rods, while a study on the vibration of nonlocal Kelvin-Voight viscoelastic damped Timoshenko beams was undertaken by [19]. The study by [20] on non-linear analysis of viscoelastic rectangular plates subjected to in-plane compression was insightful. [21] also investigated the static buckling of infinitely column lying on quadratic-cubic elastic foundations using asymptotic approach, similarly [22] analyzed the dynamic stability of a simple quadratic elastic model structure that is pre-statically loaded but trapped by a step load using asymptotic approach. It is worthy of note the work of [25] in which the static buckling of the same structure discussed here was studied. The following important works on dynamic buckling analysis using asymptotic techniques are also worthy of note [26 – 29].

The dynamic buckling load of a viscously damped elastic structure trapped by a step load is a real life problem and the governing equation is the mathematical generalization of some of the physical structures encountered in engineering practice. This work aims at investigating, using asymptotic and perturbation procedures, the dynamic buckling of a viscously damped but clamped finite column lying on a quadratic-cubic nonlinear foundation. In addition, the effects of light viscous damping as well as imperfection on the dynamic stability of the structure are discussed.

The dynamic buckling load λ_D is defined as the maximum load parameter for which the displacement or solution of the governing equation remains bounded for all time and is obtained from the maximization [1],

$$\frac{d\lambda}{dU_a} = 0 \quad (1.1)$$

where λ is the load parameter and U_a is the maximum value of the displacement of the column.

2.0 FORMULATION OF THE PROBLEM

The usual dimensional differential equation satisfied by the deflection $W(X, T)$ of the column under consideration satisfies the following partial differential equation, as in [23] and [24],

$$m_0 W_{,TT} + c_0 W_{,T} + EI W_{,XXXX} + 2P(T) W_{,XX} + Wk_1 - k_2 W^2 - k_3 W^3 = -2P(T) \frac{d^2 W}{dx^2}, T > 0 \quad (2.2a)$$

$$0 < X < \pi \quad (2.2b)$$

$$W(X, 0) = 0 = W_{,T}(X, 0) = 0, 0 < X < \pi \quad (2.3)$$

$$W = W_{,X} = 0 \text{ at } X = 0, \pi \quad (2.4)$$

where, m_0 is the mass per unit length, c_0 is the damping coefficient, EI is the bending stiffness where, E and I are the Young's modulus and I is the moment of inertia respectively.

Here the nonlinear elastic foundation exerts a force per unit length given by

$Wk_1 - k_2W^2 - k_3W^3$ on the column where k_1, k_2 and k_3 are constants such that $k_1 > 0, k_2 > 0, k_3 > 0$. In this formulation, all nonlinearities higher than cubic are excluded, while all nonlinear derivatives of $W(X,T)$ are also excluded. Here, \bar{W} is the stress-free time independent twice-differentiable initial imperfection displacement and all aspects of axial inertia are neglected.

3.0 PERTURBATION PROCEDURE

To reduce equation (2.2) to (2.4) to non-dimensional form, we adopt the following quantities:

$$x = \left(\frac{k_1}{EI}\right)^{\frac{1}{4}} X, \quad \omega = \left(\frac{k_2}{k_1}\right)^{\frac{1}{2}} W, \quad \lambda f(t) = \frac{P(T)}{2(EIk_1)^{\frac{1}{2}}}, \quad t = \left(\frac{k_1}{m_0}\right)^{\frac{1}{2}} T, \quad \epsilon \bar{\omega} = \left(\frac{k_3}{k_1}\right)^{\frac{1}{2}} \bar{W}, \quad 2\delta = \frac{c_0}{(m_0k_1)^{\frac{1}{2}}}, \quad \alpha = \frac{k_2}{\sqrt{k_1k_2}},$$

$$\beta = \left(\frac{k_3}{k_1}\right)^{\frac{3}{2}} \quad (3.5a)$$

Here, we shall assume the following inequalities

$$0 < \delta \ll 1, \quad 0 < \epsilon \ll 1. \quad (3.5b)$$

On substituting (3.5a) in (3.2) and simplifying, the following is obtained

$$\omega_{,tt} + 2\delta\omega_{,t} + \omega_{,xxxx} + 2\lambda f(t)\omega_{,xx} + \omega - \alpha\omega^2 - \beta\omega^3 = -2\epsilon\lambda f(t) \frac{d^2\bar{\omega}}{dx^2} \quad (3.6)$$

$$t > 0, \quad 0 < x < \pi \quad (3.7a)$$

$$\omega(x, 0) = 0 = \omega_{,t}(x, 0), \quad 0 < x < \pi \quad (3.7b)$$

$$\omega = \omega_{,x} = 0 \text{ at } x = 0, \pi \quad (3.7c)$$

where, ω is the displacement, t is the time variable, δ is the damping coefficient, α and β are the imperfection – sensitivity parameters, ϵ is the amplitude of the imperfection, $\bar{\omega}$ is a stress-free time independent twice-differentiable imperfection and $f(t)$ is a time dependent loading function while λ is the nondimensional amplitude (or magnitude) of the loading.

Here, a subscript following a comma indicates partial differentiation while $\bar{\omega}$ is a twice-differentiable stress-free imperfection and $f(t)$ is a step load such that,

$$f(t) = \begin{cases} 1, & t > 0 \\ 0, & t < 0 \end{cases} \quad (3.8)$$

Here, it is assumed that δ and ϵ are two small but unrelated parameters that satisfy the inequalities as in (3.5b). Our ultimate aim is to determine the dynamic buckling load λ_D which is obtained by using the maximization (3.1).

Let,

$$\tau = \delta t \quad (3.9a)$$

$$\hat{t} = t + \frac{1}{\delta} [\omega_1(\tau)\epsilon + \omega_2(\tau)\epsilon^2 + \omega_3(\tau)\epsilon^3 + \omega_4(\tau)\epsilon^4 + \dots] \quad (3.9b)$$

where,

$$\omega_i(0) = 0, \quad i = 1, 2, 3, \dots \quad (3.10a)$$

Let,

$$\omega(x, t) = U(x, t, \tau, \epsilon, \delta) \quad (3.10b)$$

From equation (3.10b); we have;

$$\omega_{,t} = \left(\frac{\partial u}{\partial \hat{t}} \cdot \frac{\partial \hat{t}}{\partial t}\right) + \left(\frac{\partial u}{\partial \hat{t}} \cdot \frac{\partial \hat{t}}{\partial \tau} \cdot \frac{\partial \tau}{\partial t}\right) + \left(\frac{\partial u}{\partial \tau} \cdot \frac{\partial \tau}{\partial t}\right) \quad (3.11)$$

$$= U_{,\hat{t}} + (\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,t} + \delta U_{,\tau} \quad (3.12)$$

The following also follows:

$$\omega_{,tt} = U_{,\hat{t}\hat{t}} + (\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)^2 U_{,\hat{t}\hat{t}} + \delta^2 U_{,\tau\tau} + 2(\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,\hat{t}\tau} + 2\delta U_{,\hat{t}\tau} + 2\delta(\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,\hat{t}\tau} + \delta(\omega''_1\epsilon + \omega''_2\epsilon^2 + \omega''_3\epsilon^3 + \dots)U_{,\hat{t}} \quad (3.13)$$

Substituting (3.12) and (3.13) into equation (3.6) results to;

$$U_{,\hat{t}\hat{t}} + (\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)^2 U_{,\hat{t}\hat{t}} + \delta^2 U_{,\tau\tau} + 2(\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,\hat{t}\tau} + 2\delta U_{,\hat{t}\tau} + 2\delta(\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,\hat{t}\tau} + \delta(\omega''_1\epsilon + \omega''_2\epsilon^2 + \omega''_3\epsilon^3 + \dots)U_{,\hat{t}} + 2\delta[U_{,\hat{t}} + (\omega'_1\epsilon + \omega'_2\epsilon^2 + \omega'_3\epsilon^3 + \dots)U_{,\hat{t}} + \delta U_{,\tau}] + U_{,xxxx} + 2\lambda U_{,xx} + U + \alpha U^2 - \beta U^3 = -2\lambda\epsilon \frac{d^2\bar{\omega}}{dx^2} \quad (3.14)$$

Let,

$$\begin{aligned}
U(x, \epsilon, \tau) &= \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} U_n^{(i,j)}(x, t, \tau) \epsilon^i \delta^j \quad (3.15) \\
&= \epsilon (U^{(10)} + \delta U^{(11)} + \delta^2 U^{(12)} + \dots) + \epsilon^2 (U^{(20)} + \delta U^{(21)} + \delta^2 U^{(22)} + \dots) \\
&\quad + \epsilon^3 (U^{(30)} + \delta U^{(31)} + \delta^2 U^{(32)} + \dots) + \dots \quad (3.16)
\end{aligned}$$

Here, the ij in $U^{(ij)}$ are not powers but superscripts. Therefore, the following orders of equations are obtained

$$O(\epsilon) : U_{\hat{t}\hat{t}}^{(10)} + U_{,xxxx}^{(10)} + 2\lambda U_{,xx}^{(10)} + U^{(10)} = -2\lambda \frac{d^2 \bar{\omega}}{dx^2}, \quad (3.17)$$

$$O(\epsilon\delta) : U_{,\hat{t}\hat{t}}^{(11)} + U_{,xxxx}^{(11)} + 2\lambda U_{,xx}^{(11)} + U^{(11)} = -2U_{,\hat{t}\tau}^{(10)} - 2U_{,\hat{t}}^{(10)} \quad (3.18)$$

$$O(\epsilon\delta^2) : U_{,\hat{t}\hat{t}}^{(12)} + U_{,xxxx}^{(12)} + 2\lambda U_{,xx}^{(12)} + U^{(12)} = -2U_{,\hat{t}\tau}^{(11)} - 2U_{,\hat{t}}^{(11)} - U_{,\tau\tau}^{(10)}$$

$$O(\epsilon^2) : U_{,\hat{t}\hat{t}}^{(20)} + U_{,xxxx}^{(20)} + 2\lambda U_{,xx}^{(20)} + U^{(20)} = -(\alpha U^{(10)})^2 - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(10)} \quad (3.20)$$

$$O(\epsilon^2\delta) : U_{,\hat{t}\hat{t}}^{(21)} + U_{,xxxx}^{(21)} + 2\lambda U_{,xx}^{(21)} + U^{(21)} = -2\alpha U^{(10)} U^{(11)} - 2U_{,\hat{t}\tau}^{(20)} - 2U_{,\hat{t}}^{(20)} - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(11)} - \omega''_1 U_{,\hat{t}}^{(10)} - 2\omega'_1 U_{,\hat{t}}^{(10)} \quad (3.21)$$

$$O(\epsilon^2\delta^2) : U_{,\hat{t}\hat{t}}^{(22)} + U_{,xxxx}^{(22)} + 2\lambda U_{,xx}^{(22)} + U^{(22)} = -U_{,\tau\tau}^{(20)} - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(12)} - 2U_{,\hat{t}\tau}^{(21)} - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(12)} - 2\omega''_1 U_{,\hat{t}}^{(11)} - 2U_{,\hat{t}}^{(21)} - 2\omega'_1 U_{,\hat{t}}^{(11)} - \alpha \{ (U^{(11)})^2 + U^{(10)} U^{(12)} \} \quad (3.22)$$

$$O(\epsilon^3) : U_{,\hat{t}\hat{t}}^{(30)} + U_{,xxxx}^{(30)} + 2\lambda U_{,xx}^{(30)} + U^{(30)} = -(\omega'_1)^2 U_{,\hat{t}\hat{t}}^{(10)} - 2(\omega'_1 U_{,\hat{t}\hat{t}}^{(20)} + \omega'_2 U_{,\hat{t}\hat{t}}^{(10)}) - 2\alpha U^{(20)} U^{(12)} + \beta (U^{(10)})^3 \quad (3.23)$$

$$\begin{aligned}
O(\epsilon^3\delta) : U_{,\hat{t}\hat{t}}^{(31)} + U_{,xxxx}^{(31)} + 2\lambda U_{,xx}^{(31)} + U^{(31)} \\
= -(\omega'_1)^2 U_{,\hat{t}\hat{t}}^{(10)} - 2(\omega'_1 U_{,\hat{t}\tau}^{(21)} + \omega'_2 U_{,\hat{t}\tau}^{(11)}) - 2U_{,\hat{t}\tau}^{(30)} + 2(\omega'_1 U_{,\hat{t}\hat{t}}^{(20)} + \omega'_2 U_{,\hat{t}\hat{t}}^{(10)}) \\
- (\omega''_1 U_{,\hat{t}}^{(20)} + \omega''_2 U_{,\hat{t}}^{(10)}) - 2\{U_{,\hat{t}}^{(30)} + (\omega'_1 U_{,\hat{t}}^{(20)} + \omega'_2 U_{,\hat{t}}^{(10)})\} \\
- \alpha (U^{(10)} U^{(21)} + U^{(11)} U^{(20)}) + 3\beta (U^{(10)})^2 (U^{(11)}) \quad (3.24)
\end{aligned}$$

$$\begin{aligned}
O(\epsilon^3\delta^2) : U_{,\hat{t}\hat{t}}^{(32)} + U_{,xxxx}^{(32)} + 2\lambda U_{,xx}^{(32)} + U^{(32)} \\
= -(\omega'_1)^2 U_{,\hat{t}\hat{t}}^{(12)} - U_{,\tau\tau}^{(30)} - 2(\omega'_1 U_{,\hat{t}\hat{t}}^{(22)} + \omega'_2 U_{,\hat{t}\hat{t}}^{(12)}) - 2U_{,\hat{t}\tau}^{(31)} - 2(\omega'_1 U_{,\hat{t}\tau}^{(21)} + \omega'_2 U_{,\hat{t}\tau}^{(11)}) \\
- (\omega''_1 U_{,\hat{t}}^{(21)} + \omega''_2 U_{,\hat{t}}^{(11)}) - 2(U_{,\hat{t}}^{(31)} + \omega'_1 U_{,\hat{t}}^{(21)} + \omega'_2 U_{,\hat{t}}^{(11)}) - 2U_{,\tau}^{(30)} \\
- 2\alpha (U^{(10)} U^{(32)} + U^{(11)} U^{(21)} + U^{(12)} U^{(20)}) \\
+ \beta [(U^{(10)})^2 U^{(12)} + 3U^{(10)} (U^{(10)})^2] \quad (3.25)
\end{aligned}$$

The associated initial conditions are as follows:

$$O(\epsilon) : U^{(ij)}(x, 0, 0) = 0; i = 1, 2, 3 \dots, j = 1, 2, 3 \dots \quad (3.26)$$

$$O(\epsilon\delta) : U_{,\hat{t}}^{(11)}(x, 0, 0) + U_{,\tau}^{(10)}(x, 0, 0) = 0 \quad (3.27)$$

$$O(\epsilon\delta^2) : U_{,\hat{t}}^{(12)}(x, 0, 0) + U_{,\tau}^{(11)}(x, 0, 0) = 0 \quad (3.28)$$

$$O(\epsilon^2) : U_{,\hat{t}}^{(20)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(10)}(x, 0, 0) = 0 \quad (3.29)$$

$$O(\epsilon^2\delta) : U_{,\hat{t}}^{(21)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(11)}(x, 0, 0) + U_{,\tau}^{(20)}(x, 0, 0) = 0 \quad (3.30)$$

$$O(\epsilon^2\delta^2) : U_{,\hat{t}}^{(22)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(12)}(x, 0, 0) + U_{,\tau}^{(21)}(x, 0, 0) = 0 \quad (3.31)$$

$$O(\epsilon^3) : U_{,\hat{t}}^{(30)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(20)}(x, 0, 0) + \omega'_2(0) U_{,\tau}^{(10)}(x, 0, 0) = 0 \quad (3.32)$$

$$O(\epsilon^3\delta) : U_{,\hat{t}}^{(31)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(21)}(x, 0, 0) + \omega'_2(0) U_{,\hat{t}}^{(11)}(x, 0, 0) + U_{,\tau}^{(30)}(x, 0, 0) = 0 \quad (3.33)$$

$$O(\epsilon^3\delta^2) : U_{,\hat{t}}^{(32)}(x, 0, 0) + \omega'_1(0) U_{,\hat{t}}^{(22)}(x, 0, 0) + \omega'_2(0) U_{,\hat{t}}^{(12)}(x, 0, 0) + U_{,\tau}^{(31)}(x, 0, 0) = 0 \quad (3.34)$$

The associated Boundary Conditions are

$$U^{(ij)} = U_x^{(ij)} = 0; x = 0, \pi \quad (3.35)$$

4.0 DYNAMIC DEFORMATION OF THE COLUMN

Let

$$\bar{\omega} = \bar{a}_m(1 - \cos 2mx), \text{ where } \bar{a}_m \text{ is a constant,} \quad (4.1)$$

And let

$$U^{(ij)}(t, \tau, x) = \sum_{n=1}^{\infty} U_n^{(ij)}(\hat{t}, \tau)(1 - \cos 2nx) \quad (4.2)$$

Solution of equation of order $\epsilon \delta^j$, $j=0,1,2$

Substituting (4.1) and (4.2) into (3.17) gives

$$\begin{aligned} \sum_{n=1}^{\infty} (1 - \cos 2nx) U_{n,\hat{t}\hat{t}}^{(10)} + \{-16n^4 + 8\lambda n^2 + (1 - \cos 2nx)\} U_n^{(10)} \\ = -8\lambda m^2 \bar{a}_m \cos 2mx \end{aligned} \quad (4.3)$$

Multiplying (4.3) through by $\cos 2mx$ and integrating from 0 to π and for $n = m$, the result is,

$$\begin{aligned} \int_0^{\pi} \sum_{n=1}^{\infty} [\{(1 - \cos 2nx) \cos 2mx\} U_{n,\hat{t}\hat{t}}^{(10)} \\ + U_n^{(10)} \{-16n^4 + 8\lambda n^2\} \cos 2nx \cos 2mx + (1 - \cos 2nx) \cos 2mx] dx \\ = - \int_0^{\pi} 8\lambda m^2 \bar{a}_m \cos 2mxdx = -8\lambda m^2 \bar{a}_m \int_0^{\pi} \frac{(1 + \cos 4mx)}{2} dx = \frac{-8\lambda m^2 \bar{a}_m \pi}{2} \\ = -4\lambda m^2 \bar{a}_m \pi \end{aligned} \quad (4.4)$$

The left hand side vanishes for all n except where $n = m$. Thus, for $n = m$, it easily follows that

$$\begin{aligned} \int_0^{\pi} \sum_{n=1}^{\infty} [\{(1 - \cos 2nx) \cos 2mx\} U_{n,\hat{t}\hat{t}}^{(10)} \\ + \{U_n^{(10)}(-16n^4 + 8\lambda n^2) \cos 2nx \\ + (1 - \cos 2nx) U_n^{(10)}\} \cos 2mx] dx \end{aligned} \quad (4.5)$$

It is to be noted that, when $n = m$, then

$$\begin{aligned} \int_0^{\pi} U_m^{(10)}(-16m^4 + 8\lambda m^2) \cos 2mxdx \\ = U_m^{(10)}(-16m^4 + 8\lambda m^2) \int_0^{\pi} \cos^2 2mxdx \\ = \frac{\pi}{2} U_m^{(10)}(-16m^4 + 8\lambda m^2) \end{aligned} \quad (4.6)$$

Thus, substituting (4.6) into (4.4), gives,

$$-\frac{\pi}{2} U_{m,\hat{t}\hat{t}}^{(10)} + \frac{\pi}{2}(-16m^4 + 8\lambda m^2) U_m^{(10)} - \frac{\pi}{2} U_m^{(10)} = -8\lambda m^2 \bar{a}_m \left(\frac{\pi}{2}\right) \quad (4.7a)$$

And this yields,

$$U_{m,\hat{t}\hat{t}}^{(10)} + (16m^4 - 8\lambda m^2 + 1) U_m^{(10)} + U_m^{(10)} = 8\lambda m^2 \bar{a}_m \quad (4.7b)$$

Let,

$$16m^4 - 8\lambda m^2 + 1 = \theta^2 \quad (4.7c)$$

Then (4.7b) becomes

$$U_{m,\hat{t}\hat{t}}^{(10)} + \theta^2 U_m^{(10)} + U_m^{(10)} = 8\lambda m^2 \bar{a}_m \quad (4.7d)$$

Initial conditions are

$$U_m^{(10)}(0,0) = 0; U_{m,\hat{t}}^{(10)}(0,0) = 0$$

Therefore, the solutions of (4.7d) is

$$U_m^{(10)} = \alpha_1(\tau) \cos \theta \hat{t} + \beta_1(\tau) \sin \theta \hat{t} + B \quad (4.7e)$$

$$\text{where, } B = \frac{8\lambda m^2 \bar{a}_m}{\theta^2} \quad (4.7f)$$

The use of initial conditions gives

$$\alpha_1(0) = -\frac{8\lambda m^2 \bar{a}_m}{\theta^2}, \beta_1 = 0 \quad (4.7g)$$

Thus

$$U^{(10)} = U_m^{(10)}(1 - \cos 2mx) \quad (4.8)$$

From (3.18), we have,

$$O(\epsilon\delta) : U_{,\hat{t}\hat{t}}^{(11)} + U_{,xxxx}^{(11)} + 2\lambda U_{,xx}^{(11)} + U^{(11)} = -2U_{,\hat{t}\tau}^{(10)} - 2U_{,\hat{t}}^{(10)}$$

Let

$$\begin{aligned} U^{(11)} &= \sum_{n=1}^{\infty} U_n^{(11)}(\hat{t}, \tau)(1 - \cos 2nx) \\ \sum_{n=1}^{\infty} [U_{n,\hat{t}\hat{t}}^{(11)}(1 - \cos 2nx) + (-16n^4 + 8\lambda n^2)U_n^{(11)}\cos 2nx + (1 - \cos nx)U_n^{(11)}]U_n^{(10)} \\ &= -2[U_{m,\hat{t}\tau}^{(10)} + U_{m,\hat{t}}^{(10)}](1 - \cos 2mx) \end{aligned} \quad (4.9a)$$

Multiplying both sides of (4.9a) through by $\cos 2mx$ and integrating from 0 to π and for $n=m$, gives

$$\begin{aligned} &\int_0^{\pi} \sum_{n=1}^{\infty} [(1 - \cos 2nx)\cos 2mx]U_{n,\hat{t}\hat{t}}^{(11)} \\ &\quad + U_n^{(11)}\{(-16n^4 + 8\lambda n^2)\cos 2nx\cos 2mx + (1 - \cos 2nx)\}\cos 2mx]dx \\ &= -2[U_{m,\hat{t}\tau}^{(10)} + U_{m,\hat{t}}^{(10)}] \int_0^{\pi} (1 - \cos 2mx)\cos 2mxdx \\ &= -\frac{\pi}{2}U_{m,\hat{t}\hat{t}}^{(11)} + \frac{\pi}{2}(-16m^4 + 8\lambda m^2)U_m^{(11)} - \frac{\pi}{2}U_m^{(11)} = -2(U_{m,\hat{t}\tau}^{(10)} + U_{m,\hat{t}}^{(10)})\left(-\frac{\pi}{2}\right) \end{aligned} \quad (4.9b)$$

Further simplification gives

$$U_{m,\hat{t}\hat{t}}^{(11)} + (16m^4 - 8\lambda m^2 + 1)U_m^{(11)} = -2(U_{m,\hat{t}\tau}^{(10)} + U_{m,\hat{t}}^{(10)}) \quad (4.9c)$$

i.e.

$$U_{m,\hat{t}\hat{t}}^{(11)} + \theta^2 U_m^{(11)} = -2(U_{m,\hat{t}\tau}^{(10)} + U_{m,\hat{t}}^{(10)}) \quad (4.10)$$

The initial conditions are

$$U_m^{(11)}(0,0) = 0; U_{m,\hat{t}}^{(11)}(0,0) + U_{m,\tau}^{(10)}$$

Substituting for $U_m^{(10)}$ on the right hand side (RHS) of (4.10), from (4.7e) gives

$$\begin{aligned} U_{m,\hat{t}\hat{t}}^{(11)} + \theta^2 U_m^{(11)} &= -2[-\theta\alpha_1' \sin\theta\hat{t} + \theta\beta_1' \cos\theta\hat{t} + (-\theta\alpha_1 \sin\theta\hat{t} + \theta\beta_1 \cos\theta\hat{t})] \\ &= -2\theta[-(\alpha_1' + \alpha_1)\sin\theta\hat{t} + (\beta_1' + \beta)\cos\theta\hat{t}] \end{aligned} \quad (4.11a)$$

To ensure a uniformly valid solution in \hat{t} , implies equating to zero the coefficients of $\cos\theta\hat{t}$ and $\sin\theta\hat{t}$ on the RHS of (4.11a). Therefore, the coefficient of $\cos\theta\hat{t}$ gives

$$\beta_1' + \beta = 0 \quad (4.11b)$$

The integrating factor is e^{τ} , then,

$$\frac{d(e^{\tau}\beta_1)}{d\tau} = 0 \quad (4.11c)$$

This gives,

$$\beta_1(\tau) = Ae^{-\tau} \text{ and } \beta_1(\tau) = 0 \quad (4.11d)$$

Similarly, the coefficient of $\sin\theta\hat{t}$ gives,

$$\alpha_1' + \alpha_1 = 0 \quad (4.11e)$$

This gives,

$$\alpha_1'(0) = -\alpha_1(0) = B \text{ and } \alpha_1(\tau) = -Be^{-\tau} \quad (4.11f)$$

$$\therefore U_m^{(10)} = \alpha_1(\tau)\cos\theta\hat{t} + B \quad (4.11g)$$

The remaining equation in (4.11a) is;

$$U_{m,\hat{t}\hat{t}}^{(11)} + \theta^2 U_m^{(11)} = 0 \quad (4.11h)$$

$$U_m^{(11)} = \alpha_2(\tau)\cos\theta\hat{t} + \beta_2(\tau)\sin\theta\hat{t} \quad (4.12a)$$

From $U_m^{(11)}(0,0) = 0$,

$$\alpha_2(0) = 0 \quad (4.12b)$$

From $U_{m,\hat{t}}^{(11)}(0,0) + U_{m,\tau}^{(10)} = 0$,

$$\beta_2(0)\theta + \alpha_1'(0) = 0 \text{ and } \beta_2(0) = -\frac{\alpha_1'(0)}{\theta} = \frac{-B}{\theta} \quad (4.12c)$$

$$\therefore U_m^{(11)} = U_m^{(11)}(1 - \cos 2mx) \quad (4.12d)$$

From (3.19); the next equation is

$$O(\epsilon\delta^2): U_{,\hat{t}\hat{t}}^{(12)} + U_{,xxxx}^{(12)} + 2\lambda U_{,xx}^{(12)} + U^{(12)} = -2U_{,\hat{t}\tau}^{(11)} - 2U_{,\hat{t}}^{(11)} - U_{,\tau\tau}^{(10)}$$

Substituting for $U_m^{(11)}$ and $U_m^{(10)}$ from (4.11g) and (4.12a) respectively on the RHS of (3.19), gives

$$U_{m,\hat{t}\hat{t}}^{(12)} + \theta^2 U_m^{(12)} = -2[-\theta\alpha_2'(\tau)\sin\theta\hat{t} + \theta\beta_2'(\tau)\cos\theta\hat{t} + (-\theta\alpha_2(\tau)\sin\theta\hat{t} + \theta\beta_2(\tau)\cos\theta\hat{t})] - \alpha_1''(\tau)\cos\theta\hat{t} \quad (4.13a)$$

$$\begin{aligned} &= 2\theta\alpha_2'(\tau)\sin\theta\hat{t} - 2\theta\beta_2'(\tau)\cos\theta\hat{t} - \theta\alpha_2(\tau)\sin\theta\hat{t} + \theta\beta_2(\tau)\cos\theta\hat{t} - \alpha_1''(\tau)\cos\theta\hat{t} \\ &= (2\theta\alpha_2'(\tau) - 2\theta\alpha_2(\tau))\sin\theta\hat{t} + (2\theta\beta_2(\tau) - 2\theta\beta_2'(\tau) - \alpha_1''(\tau))\cos\theta\hat{t} \end{aligned} \quad (4.13b)$$

To remove secular terms in the solution of $U_m^{(12)}$, ie to ensure a uniformly valid solution in \hat{t} implies equating to zero the coefficients of $\cos\theta\hat{t}$ and $\sin\theta\hat{t}$ on the RHS. These respectively give

$$\cos\theta\hat{t}: -2(\theta\beta_2' + \theta\beta_2) - \alpha_1'' = 0$$

And

$$\sin\theta\hat{t}: -2(-\theta\alpha_2' - \theta\alpha_2) = 0$$

$$\therefore \beta_2' + \beta_2 = \frac{-\alpha_1''}{2\theta} \text{ and } [\beta_2'(0) = \frac{3B}{2\theta}] \quad (4.13c)$$

$$\therefore \alpha_2' + \alpha_2 = 0 \quad (4.13d)$$

Therefore, from (4.13),

$$\alpha_2(\tau) \equiv 0, \quad (4.13e)$$

And from (4.13d),

$$\beta_2(\tau) = e^{-\tau} \left[-\int_0^\tau \frac{e^s \alpha_1''}{2\theta} ds + \beta_2(0) \right] \quad (4.13f)$$

i.e

$$\beta_2(\tau) = e^{-\tau} \left[-\int_0^\tau \frac{e^s \alpha_1''}{2\theta} ds - \frac{B}{\theta} \right] \quad (4.13g)$$

$$\therefore U_m^{(11)} = \beta_2(\tau)\sin\theta\hat{t} \quad (4.13h)$$

Equating the left hand side (LHS) of (4.13a) to zero,

i.e

$$U_{m,\hat{t}\hat{t}}^{(12)} + \theta^2 U_m^{(12)} = 0$$

The Initial conditions are

$$U_m^{(12)}(0,0) = 0, \quad U_{m,\hat{t}}^{(12)} + U_{m,\tau}^{(11)}(0,0) = 0$$

$$\therefore U_m^{(12)}(\hat{t}, \tau) = \alpha_3(\tau)\cos\theta\hat{t} + \beta_3(\tau)\sin\theta\hat{t} \quad (4.13i)$$

Applying the initial conditions,

$$\alpha_3(0) = 0, \quad \beta_3(0) = 0 \quad (4.13j)$$

$$U_{m,\hat{t}}^{(12)}(0,0) = \theta\beta_3(0) = 0$$

$$\beta_3(0) = 0 \quad (4.13k)$$

$$\therefore U^{12} = U_m^{(12)}(1 - \cos 2m\hat{x}) \quad (4.13l)$$

From (3.20),

$$O(\epsilon^2): U_{,\hat{t}\hat{t}}^{(20)} + U_{,xxxx}^{(20)} + 2\lambda U_{,xx}^{(20)} + U^{(20)} = -(\alpha U^{(10)})^2 - 2\omega_1' U_{,\hat{t}\hat{t}}^{(10)}$$

Let,

$$U^{(20)} = \sum_{n=1}^{\infty} U_n^{(20)}(\hat{t}, \tau)(1 - \cos 2n\hat{x}) \quad (4.14)$$

Substituting (4.14) into (3.20) gives ;

$$\begin{aligned} &\sum_{n=1} \left[U_{n,\hat{t}\hat{t}}^{(20)}(1 - \cos 2n\hat{x}) + (-16n^4 + 8\lambda n^2)U_n^{(20)} \cos 2n\hat{x} \right] \\ &\quad + (1 - \cos 2n\hat{x})U_n^{(20)} \\ &= -\alpha(U_m^{(10)})^2 \left[\frac{3}{2} - 2\cos 2m\hat{x} + \frac{1}{2}\cos 4m\hat{x} \right] \\ &\quad - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(10)}(1 - \cos 2m\hat{x}) \end{aligned} \quad (4.15a)$$

Multiplying both sides of (4.15a) through by $\cos 2m\hat{x}$ and integrating from 0 to π and for $n=m$, the result gives;

$$\begin{aligned} &\left[-\frac{\pi}{2} U_{m,\hat{t}\hat{t}}^{(20)} + (-16m^4 + 8\lambda m^2)U_m^{(20)} \left(\frac{\pi}{2} \right) + \left(\frac{\pi}{2} U_m^{(20)} \right) \right] \\ &= -\alpha(U_m^{(10)})^2 \left[-2 \left(\frac{\pi}{2} \right) - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(10)} \left(\frac{-\pi}{2} \right) \right] \end{aligned} \quad (4.15b)$$

i.e,

$$\frac{\pi}{2} \left[-U_{m,\hat{t}\hat{t}}^{(20)} + (-16m^4 + 8\lambda m^2)U_m^{(20)} - U_m^{(20)} \right] = \frac{\pi}{2} \left[2\alpha(U_m^{(10)})^2 + 2\omega'_1 U_{m,\hat{t}\hat{t}}^{(10)} \right] \quad (4.15c)$$

Simplification of (4.15c) gives,

$$U_{m,\hat{t}\hat{t}}^{(20)} + (16m^4 - 8\lambda m^2 + 1)U_m^{(20)} = - \left[2\alpha(U_m^{(10)})^2 + 2\omega'_1 U_{m,\hat{t}\hat{t}}^{(10)} \right] \quad (4.15d)$$

And this further gives,

$$U_{m,\hat{t}\hat{t}}^{(20)} + \theta^2 U_m^{(20)} = - \left[2\alpha(U_m^{(10)})^2 + 2\omega'_1 U_{m,\hat{t}\hat{t}}^{(10)} \right] \quad (4.16a)$$

The initial conditions are,

$$U_m^{(20)}(0,0) = 0, U_{m,\hat{t}}^{(20)}(0,0) + \omega'(0)U_{m,\hat{t}}^{(10)}(0,0) = 0$$

Next multiplying equation (4.15a) by $\cos 4m\hat{t}$ and integrating from 0 to π and for $n=m$, the result gives;

$$-\frac{\pi}{2} U_{2m,\hat{t}\hat{t}}^{(20)} + (-256m^4 + 32\lambda m^2) \left(\frac{\pi}{2} \right) U_{2m}^{(20)} - \left(\frac{\pi}{2} \right) U_{2m}^{(20)} = -\alpha(U_m^{(10)})^2 \left(\frac{1}{2} \cdot \frac{\pi}{2} \right) \quad (4.16b)$$

Simplifying (4.16b) gives;

$$U_{2m,\hat{t}\hat{t}}^{(20)} + (256m^4 - 32\lambda m^2 + 1)U_{2m}^{(20)} = \frac{\alpha}{2} (U_m^{(10)})^2 \quad (4.16c)$$

Let,

$$\varphi^2 = (256m^4 + 32\lambda m^2 + 1) > 0 \quad (4.16d)$$

Therefore, (4.16c) becomes

$$U_{2m,\hat{t}\hat{t}}^{(20)} + \varphi^2 U_m^{(20)} = \frac{\alpha}{2} (U_m^{(10)})^2 \quad (4.17)$$

The initial conditions are,

$$U_{2m}^{(20)}(0,0) = 0; U_{2m,\hat{t}}^{(20)}(0,0) + \omega'_1 U_{2m,\hat{t}}^{(10)}(0,0) = 0$$

On substituting for $U_m^{(10)}$ on the RHS of (4.16a), the simplification is

$$U_{2m,\hat{t}\hat{t}}^{(20)} + \theta^2 U_m^{(20)} = - \left[\{ 2\alpha(\alpha_1 \cos \theta \hat{t} + B)^2 \} + \{ 2\omega'_1 (-\alpha_1 \cos \theta \hat{t}) \} \right] \\ - \left[2\alpha \left\{ \left(\frac{\alpha_1^2}{2} + B^2 \right) + 2B\alpha_1 \cos \theta \hat{t} + \frac{\alpha_1^2}{2} \alpha_1 \cos 2\theta \hat{t} \right\} + 2\omega'_1 (-\alpha_1 \theta^2 \cos \theta \hat{t}) \right] \quad (4.18a)$$

To ensure a uniformly valid solution in \hat{t} , we equate to zero, the coefficients of $\cos 2\theta \hat{t}$ on the RHS of (4.18a). That is,

$$- [2B\alpha_1 - 2\omega'_1 \theta^2 \alpha_1] = 0 \quad (4.18b)$$

$$\therefore \omega'_1 = \frac{B}{\theta^2}; \omega_1 = \int \frac{B}{\theta^2} d\tau \quad (4.18c)$$

The remaining part of equation (4.18a) for $U_m^{(20)}$ is

$$U_{m,\hat{t}\hat{t}}^{(20)} + \theta^2 U_m^{(20)} = r_0 + r_1 \cos 2\theta \hat{t} \quad (4.19a)$$

where, $r_0 = -2\alpha \left(\frac{\alpha_1^2}{2} + B^2 \right)$, $r_0(0) = -3\alpha B^2$

$$r_1 = -\alpha \alpha_1^2, r_1(0) = -\alpha B^2, r'_0(0) = 2\alpha B^2, r'_1(0) = 2\alpha B^2$$

$$\therefore U_m^{(20)}(\hat{t}, \tau) = \alpha_4(\tau) \cos \theta \hat{t} + \beta_4(\tau) \sin \theta \hat{t} + \frac{r_0}{\theta^2} - \frac{r_1 \cos 2\theta \hat{t}}{3\theta^2} \quad (4.19b)$$

From the initial condition,

$$U_m^{(20)}(0,0) = 0; \text{ i.e, } \alpha_4(0) + \frac{r_0(0)}{\theta^2} - \frac{r_1}{3\theta^2} = 0$$

$$\therefore \alpha_4(0) = \frac{r_1}{3\theta^2} - \frac{r_0(0)}{\theta^2} = \frac{8\alpha B^2}{3\theta^2} \quad (4.19c)$$

Applying the initial condition, $U_{m,\hat{t}}^{(20)}(0,0) + \omega'(0) + U_{m,\hat{t}}^{(10)}(0,0)$ yields,

$$\beta_4(0) = 0 \quad (4.19d)$$

Simplification of (4.17) yields,

$$U_{2m,\hat{t}\hat{t}}^{(20)} + \varphi^2 U_m^{(20)} = \frac{\alpha}{2} \left[\left(\frac{\alpha_1^2}{2} + B^2 \right) + 2B\alpha_1 \cos \theta \hat{t} + \frac{\alpha_1^2}{2} \cos 2\theta \hat{t} \right] \quad (4.20a)$$

$$\therefore U_{2m}^{(20)}(\hat{t}, \tau) = \alpha_5(\tau) \cos \varphi \hat{t} + \beta_5(\tau) \sin \varphi \hat{t}$$

$$+ \frac{\alpha}{2} \left[\frac{\left(\frac{\alpha_1^2}{2} + B^2 \right)}{\varphi^2} + \frac{2B\alpha_1 \cos \theta \hat{t}}{(\varphi^2 - \theta^2)} + \frac{\alpha_1^2 \cos 2\theta \hat{t}}{2(\varphi^2 - \theta^2)} \right] \quad (4.20b)$$

From the initial conditions,

$$U_{2m}^{(20)}(0,0) = 0; U_{2m,\hat{t}}^{(20)}(0,0) + \omega'_1 U_{2m,\hat{t}}^{(10)}(0,0) = 0$$

$$\begin{aligned} & \therefore \alpha_5(0) + \frac{\alpha}{2} \left[\frac{(\frac{\alpha_1^2}{2} + B^2)}{\varphi^2} + \frac{2B\alpha_1}{(\varphi^2 - \theta^2)} + \frac{\alpha_1^2}{2(\varphi^2 - 4\theta^2)} \right] = 0 \\ \therefore \alpha_5(0) &= -\frac{\alpha}{2} \left[\frac{(\frac{\alpha_1^2}{2} + B^2)}{\varphi^2} + \frac{2B\alpha_1}{(\varphi^2 - \theta^2)} + \frac{\alpha_1^2}{2(\varphi^2 - 4\theta^2)} \right] \text{ at } \tau=0 \end{aligned}$$

i.e

$$\alpha_5(0) = -\frac{\alpha}{2} \left[\frac{3B^2}{2\varphi^2} - \frac{2B^2}{(\varphi^2 - \theta^2)} + \frac{B^2}{2(\varphi^2 - 4\theta^2)} \right] = B^2 \alpha S_0 \text{ and } \beta_5(0) = 0 \quad (4.20c)$$

$$\text{where, } S_0 = \left(-\frac{3\alpha}{2\varphi^2} + \frac{\alpha}{(\varphi^2 - \theta^2)} - \frac{\alpha}{4(\varphi^2 - 4\theta^2)} \right)$$

$$\therefore U^{(20)} = U_m^{(20)}(1 - \cos 2mx) + U_{2m}^{(20)}(1 - \cos 4mx) \quad (4.20d)$$

From (3.21),

$$\begin{aligned} 0(\epsilon^2 \delta) : U_{,\hat{t}\hat{t}}^{(21)} + U_{,xxxx}^{(21)} + 2\lambda U_{,xx}^{(21)} + U^{(21)} \\ = -2\alpha U^{(10)} U^{(11)} - 2U_{,\hat{t}\tau}^{(20)} - 2U_{,\hat{t}}^{(20)} - 2\omega_1' U_{,\hat{t}\hat{t}}^{(11)} - \omega_1'' U_{,\hat{t}}^{(10)} - 2\omega_1' U_{,\hat{t}}^{(10)} \end{aligned}$$

i.e

$$\begin{aligned} U_{,\hat{t}\hat{t}}^{(21)} + U_{,xxxx}^{(21)} + 2\lambda U_{,xx}^{(21)} + U^{(21)} &= -2\alpha U_m^{(10)}(1 - \cos 2mx) U_m^{(11)}(1 - \cos 2mx) - 2U_{m,\hat{t}\tau}^{(20)}(1 - \cos 2mx) \\ &- 2U_{m,\hat{t}}^{(20)}(1 - \cos 2mx) - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)}(1 - \cos 2mx) - \omega_1'' U_{m,\hat{t}}^{(10)}(1 - \cos 2mx) - \\ &2\omega_1' U_{m,\hat{t}}^{(10)}(1 - \cos 2mx) \end{aligned} \quad (4.21)$$

Let

$$U^{(21)} = \sum_{n=1}^{\infty} U_n^{(21)}(\hat{t}\tau)(1 - \cos 2nx)$$

Substituting into (4.21) gives,

$$\begin{aligned} \sum_{n=1}^{\infty} \left[U_{n,\hat{t}\hat{t}}^{(21)}(1 - \cos 2nx) + (-16n^4 + 8\lambda n^2) U_n^{(21)} \cos 2nx + (1 - \cos 2nx) U_n^{(21)} \right] \\ = -2\alpha U_m^{(10)} U_m^{(11)} \left[\frac{3}{2} - 2\cos 2mx + \frac{1}{2} \cos 4mx \right] - 2U_{m,\hat{t}\tau}^{(20)}(1 - \cos 2mx) \\ - 2U_{m,\hat{t}}^{(20)}(1 - \cos 2mx) - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)}(1 - \cos 2mx) - \omega_1'' U_{m,\hat{t}}^{(10)}(1 - \cos 2mx) \\ - 2\omega_1' U_{m,\hat{t}}^{(10)}(1 - \cos 2mx) \end{aligned} \quad (4.22a)$$

Multiplying both sides of (4.22) through by $\cos 2mx$ and integrating from 0 to π and for $n=m$, gives;

$$\begin{aligned} \left[-\frac{\pi}{2} U_{m,\hat{t}\hat{t}}^{(21)} + (-16m^4 + 8\lambda m^2) U_m^{(21)} \left(\frac{\pi}{2} \right) + \left(-\frac{\pi}{2} U_m^{(21)} \right) \right] \\ = \left[\begin{aligned} & -2\alpha U_m^{(10)} U_m^{(11)} \left(-\frac{\pi}{2} \right) - 2U_{m,\hat{t}\tau}^{(20)} \left(-\frac{\pi}{2} \right) - 2U_{m,\hat{t}}^{(20)} \\ & \left(-\frac{\pi}{2} \right) - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)} \left(-\frac{\pi}{2} \right) - \omega_1'' U_{m,\hat{t}}^{(10)} \left(-\frac{\pi}{2} \right) - 2\omega_1' U_{m,\hat{t}}^{(10)} \left(-\frac{\pi}{2} \right) \end{aligned} \right] \end{aligned} \quad (4.22b)$$

Further simplification of (4.22b) yields,

$$\begin{aligned} U_{m,\hat{t}\hat{t}}^{(21)} + (16m^4 - 8\lambda m^2 + 1) U_m^{(21)} \\ = -2\alpha U_m^{(10)} U_m^{(11)} - 2U_{m,\hat{t}\tau}^{(20)} - 2U_{m,\hat{t}}^{(20)} - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)} - \omega_1'' U_{m,\hat{t}}^{(10)} \\ - 2\omega_1' U_{m,\hat{t}}^{(10)} \end{aligned} \quad (4.22c)$$

The above finally yields,

$$\begin{aligned} U_{m,\hat{t}\hat{t}}^{(21)} + \theta^2 U_m^{(21)} &= \\ -2\alpha U_m^{(10)} U_m^{(11)} - 2U_{m,\hat{t}\tau}^{(20)} - 2U_{m,\hat{t}}^{(20)} - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)} - \omega_1'' U_{m,\hat{t}}^{(10)} - \\ 2\omega_1' U_{m,\hat{t}}^{(10)} \end{aligned} \quad (4.23a)$$

The initial conditions for (4.33a) are,

$$U_m^{(21)}(0,0) = 0; U_{m,\hat{t}}^{(21)}(0,0) + \omega_1'(0) U_{m,\hat{t}}^{(11)}(x,0,0) + U_{m,\tau}^{(20)}(0,0) = 0$$

Next, multiplying (4.22a) by $\cos 4mx$ and integrating from 0 to π for $n=m$, gives

$$\begin{aligned} \left[-\frac{\pi}{2} U_{2m,\hat{t}\hat{t}}^{(21)} + (-256m^4 + 32\lambda m^2) U_{2m}^{(21)} \left(\frac{\pi}{2} \right) - \frac{\pi}{2} U_{2m}^{(21)} \right] &= -2\alpha U_m^{(10)} U_m^{(11)} \left(\frac{\pi}{2} \right) \left(\frac{1}{2} \right) - 2 \left(U_{2m,\hat{t}\tau}^{(20)} + \right. \\ & \left. U_{m,\hat{t}}^{(20)} \right) \end{aligned} \quad (4.23b)$$

$$U_{2m,\hat{t}\hat{t}}^{(21)} + \varphi^2 U_{2m}^{(21)} = \alpha U_m^{(10)} U_m^{(11)} + 2 \left(U_{2m,\hat{t}\tau}^{(20)} + U_{2m,\hat{t}}^{(20)} \right) \quad (4.24)$$

The initial conditions for (4.33b) are,

$$U_{2m}^{(21)}(0,0) = 0; U_{2m,\hat{t}}^{(21)}(0,0) = 0$$

Substituting for $U_m^{(10)}$, $U_m^{(11)}$ and $U_m^{(20)}$ in (4.24) yields

$$\begin{aligned} & U_{m,\hat{t}\hat{t}}^{(21)} + \theta^2 U_m^{(21)} = \\ & -2\alpha U_m^{(10)} U_m^{(11)} - 2U_{m,\hat{t}\tau}^{(20)} - 2U_{m,\hat{t}}^{(20)} - 2\omega_1' U_{m,\hat{t}\hat{t}}^{(11)} - \omega_1'' U_{m,\hat{t}}^{(10)} - \\ & 2\omega_1' U_{m,\hat{t}}^{(10)} \end{aligned} \quad (4.25a)$$

i.e,

$$\begin{aligned} & -2\alpha \left(\frac{\alpha_1 \beta_2}{2} \sin\theta\hat{t} + B\beta_2 \sin\theta\hat{t} \right) - 2 \left(-\theta\alpha_4' \sin\theta\hat{t} + \theta\beta_4' \cos\theta\hat{t} + \frac{2\theta r_1' \sin 2\theta\hat{t}}{3\theta^2} \right) - 2 \left(-\theta\alpha_4 \sin\theta\hat{t} + \right. \\ & \left. \theta\beta_4 \cos\theta\hat{t} + \frac{2\theta r_1 \sin 2\theta\hat{t}}{3\theta^2} \right) - 2\omega_1' (-\theta^2)\beta_2 \sin\theta\hat{t} - \omega_1'' (-\theta\alpha_1 \sin\theta\hat{t}) - \\ & 2\omega_1' (-\alpha_1 \theta \sin\theta\hat{t}) \end{aligned} \quad (4.25b)$$

To ensure a uniformly valid solution in \hat{t} , we equate to zero the coefficients of $\cos\theta\hat{t}$ and $\sin\theta\hat{t}$. This yields respectively,

$$-2\theta\beta_4' - 2\theta\beta_4 = 0 \quad (4.25c)$$

$$\alpha B\beta_2 + 2\theta\alpha_4' + 2\theta\alpha_4 + 2\theta^2\omega_1'\beta_2 + \omega_1''\theta\alpha_1 + 2\omega_1'\alpha_1\theta \quad (4.26d)$$

$$\therefore \beta_4' + \beta_4 = 0 \quad (4.26e)$$

Solving (4.26e) yields,

$$\beta_4(\tau) = \beta_4(0)e^{-\tau} = 0 \text{ since } \beta_4(0) = 0 \quad (4.26f)$$

Solving (4.25d) yields,

$$\alpha_4' + \alpha_4 = \rho_1(\tau) = \frac{1}{2\theta} (\alpha B\beta_2 - 2\theta^2\omega_1'\beta_2 - \omega_1''\theta\alpha_1 - 2\omega_1'\alpha_1\theta) \quad (4.26g)$$

$$\begin{aligned} \alpha_4(\tau) &= e^{-\tau} \left[\int_0^\tau e^s \rho_1(s) ds + \alpha_4(0) \right] \\ \therefore \alpha_4'(0) &= \rho_1(0) - \alpha_4(0) \end{aligned} \quad (4.26h)$$

where

$$\alpha_4(0) = \frac{8\alpha B^2}{3\theta^2} \quad (4.26i)$$

$$\alpha_4'(0) = \frac{-13\alpha B^2}{3\theta^2} + \frac{4B^2}{\theta} \quad (4.26j)$$

$$\therefore \rho_1(0) = \frac{-5\alpha B^2}{3\theta^2} + \frac{4B^2}{\theta} = B^2 \left(\frac{-5\alpha}{3\theta^2} + \frac{4}{\theta} \right) = B^2 V \quad (4.26k)$$

where, $V = \left(\frac{-5\alpha}{3\theta^2} + \frac{4}{\theta} \right)$

The remaining equation in (4.25a) becomes,

$$U_{m,\hat{t}\hat{t}}^{(21)} + \theta^2 U_m^{(21)} = r_2 + r_3 \cos 2\theta\hat{t} + r_4 \sin 2\theta\hat{t} \quad (4.26l)$$

with the initial conditions,

$$U_m^{(21)}(0,0) = 0; U_{m,\hat{t}}^{(21)}(0,0) + \omega_1'(0)U_{m,\hat{t}}^{(11)} + U_{m,\hat{t}}^{(20)}(0,0) = 0$$

where,

$$r_2 = \alpha_1\alpha_2; r_2(0) = \alpha_1(0)\alpha_2(0) = 0$$

$$r_3 = \alpha\alpha_1\alpha_2; r_3(0) = \alpha\alpha_1(0)\alpha_2(0) = 0; r_3'(0) = 0,$$

$$r_4 = \left[\alpha\alpha_1\beta_2 + \frac{8\alpha}{3\theta} (\alpha_1\alpha_1' + \alpha_1^2) \right]; r_4(0) = \left[\alpha(-B) \left(\frac{-B}{\theta} \right) + \frac{8\alpha}{3\theta} (-B \cdot B + B^2) \right] = \frac{\alpha B^2}{\theta}, r_4'(0) = \frac{-5\alpha B^2}{2\theta}$$

The solution of (4.26l) becomes

$$U_m^{(21)}(\hat{t}, \tau) = \alpha_6 \cos\theta\hat{t} + \beta_6 \sin\theta\hat{t} + \frac{r_2}{\theta^2} - \left(\frac{r_3 \cos 2\theta\hat{t} + r_4 \sin 2\theta\hat{t}}{3\theta^2} \right) \quad (4.27a)$$

with the initial conditions, $\alpha_6(0) + \frac{r_2}{\theta^2} - \frac{r_3}{3\theta^2} = 0$

$$\therefore \alpha_6(0) = \frac{r_3 - 3r_2}{3\theta^2} = 0, \quad \beta_6(0) = 0 \quad (4.27b)$$

From (4.24),

$$\begin{aligned}
U_{2m,\hat{t}\hat{t}}^{(21)} + \varphi^2 U_{2m}^{(21)} &= \alpha \left[\frac{\alpha_1 \beta_2}{2} \sin 2\theta \hat{t} + B \beta_2 \sin \theta \hat{t} \right] + 2 \left(U_{2m,\hat{t}\tau}^{(20)} + U_{2m,\hat{t}}^{(20)} \right) \\
&= \alpha \left[\frac{\alpha_1 \beta_2}{2} \sin 2\theta \hat{t} + B \beta_2 \sin \theta \hat{t} \right] \\
&\quad + 2 \left[-\varphi \alpha'_5 \sin \varphi \hat{t} + \beta'_5 \varphi \cos \varphi \hat{t} + \frac{\alpha}{2} \left\{ \frac{-2\theta \alpha'_1 B \sin \theta \hat{t}}{\varphi^2 - \theta^2} - \frac{2\theta (\alpha_1^2)' \sin 2\theta \hat{t}}{2(\varphi^2 - 4\theta^2)} \right\} \right. \\
&\quad + \left. \left\{ -\varphi \alpha_5 \sin \varphi \hat{t} + \beta_5 \cos \varphi \hat{t} \right. \right. \\
&\quad \left. \left. + \frac{\alpha}{2} \left\{ \frac{-2\theta \alpha_1 B \sin \theta \hat{t}}{\varphi^2 - \theta^2} - \frac{2\theta \alpha_1^2 \sin 2\theta \hat{t}}{2(\varphi^2 - 4\theta^2)} \right\} \right\} \right] \quad (4.28a)
\end{aligned}$$

To ensure a uniformly valid solution in \hat{t} , we equate to zero the coefficient of $\cos \varphi \hat{t}$ and $\sin \varphi \hat{t}$

$$2\varphi \beta'_5 + 2\varphi \beta_5 = 0 \Rightarrow \beta'_5 + \beta_5 = 0 \Rightarrow \beta'_5(0) = -\beta_5(0) \quad (4.28b)$$

$$-2\varphi \alpha'_5 - 2\varphi \alpha_5 = 0 \Rightarrow \alpha'_5 + \alpha_5 = 0 \Rightarrow \alpha'_5(0) = -\alpha_5(0) \quad (4.28c)$$

$$\beta_5 = \beta_5(0)e^{-\tau} = 0, \alpha_5 = \alpha_5(0)e^{-\tau} = 0 \quad (4.28d)$$

The remaining equation of (4.27a) is:

$$U_{2m,\hat{t}\hat{t}}^{(21)} + \varphi^2 U_{2m}^{(21)} = \left[\alpha B \beta_2 + \frac{\alpha}{2} \left(\frac{-2\theta B}{\varphi^2 - \theta^2} \right) (\alpha'_1 + \alpha_1) \right] \sin \theta \hat{t} + \left[\frac{\alpha \alpha_1 \beta_2}{2} + \frac{\alpha}{2} \left\{ \frac{-\theta (\alpha_1^2)' + \alpha_1^2}{2(\varphi^2 - 4\theta^2)} \right\} \right] \sin 2\theta \hat{t} = r_5 \sin \theta \hat{t} + r_6 \sin 2\theta \hat{t} \quad (4.29a)$$

where,

$$r_5 = \left[\alpha B \beta_2 - \frac{-2\theta B}{(\varphi^2 - \theta^2)} (\alpha'_1 + \alpha) \right] = \alpha B \beta_2, \text{ since } \alpha'_1 + \alpha = 0,$$

$$\alpha'_1 = B; r_5(0) = \frac{-\alpha B^2}{\theta}, r_6 = \left[\frac{\alpha \alpha_1 \beta_2}{2} + \frac{\alpha}{2} \left\{ \frac{-\theta (\alpha_1^2)' + \alpha_1^2}{2(\varphi^2 - 4\theta^2)} \right\} \right]$$

$$\therefore r_6(0) = \left[\frac{\alpha \alpha_1(0) \beta_2(0)}{2} + \frac{\alpha}{2} \left\{ \frac{-\theta (\alpha_1^2(0))' + \alpha_1^2(0)}{2(\varphi^2 - 4\theta^2)} \right\} \right] = \frac{B^2 \alpha}{2\theta} + \frac{B^2 \theta \alpha}{4(\varphi^2 - 4\theta^2)} = B^2 S_1 \quad (4.29b)$$

where, $S_1 = \left(\frac{\alpha}{2\theta} + \frac{\alpha \theta}{4(\varphi^2 - 4\theta^2)} \right)$

$$\therefore U_{2m}^{(21)} = \alpha_7(\tau) \cos \varphi \hat{t} + \beta_7(\tau) \sin \varphi \hat{t} + \frac{r_5 \cos \theta \hat{t}}{\varphi^2 - \theta^2} + \frac{r_6 \cos 2\theta \hat{t}}{\varphi^2 - 4\theta^2} \quad (4.30)$$

The initial conditions for (4.30) are

$$U_{2m}^{(21)}(0,0) = 0; U_{2m,\hat{t}}^{(21)}(0,0) + U_{2m,\hat{t}}^{(20)}(0,0) = 0;$$

$$\Rightarrow -\varphi \alpha_7(0) \sin \varphi \hat{t} + \varphi \beta_7(0) \cos \varphi \hat{t} + \frac{\theta r_5(0) \cos \theta \hat{t}}{\varphi^2 - \theta^2} + \frac{2\theta r_6(0) \cos 2\theta \hat{t}}{\varphi^2 - 4\theta^2} + \alpha'_5(0) \cos \varphi \hat{t} + \frac{\alpha}{2} \left[\frac{\alpha'_1 \alpha_1}{\varphi^2} + \frac{2B\theta \alpha'_1 \cos \theta \hat{t}}{\varphi^2 - \theta^2} + \frac{2\alpha'_1 \alpha_1 \cos 2\theta \hat{t}}{2(\varphi^2 - 4\theta^2)} \right] = 0 \quad (4.31a)$$

$$\therefore \alpha_7(0) = 0 \quad (4.31b)$$

Similarly, the following is obtained

$$\varphi \beta_7(0) + \frac{\theta r_5(0)}{\varphi^2 - \theta^2} + \frac{2\theta r_6(0)}{\varphi^2 - 4\theta^2} + \alpha'_5(0) + \frac{\alpha}{2} \left[\frac{\alpha'_1(0) \alpha_1(0)}{\varphi^2} + \frac{2B\theta \alpha'_1(0)}{\varphi^2 - \theta^2} + \frac{\alpha'_1(0) \alpha_1(0)}{(\varphi^2 - 4\theta^2)} \right] = 0 \quad (4.32a)$$

$$\beta_7(0) = -\frac{1}{\varphi} \left[\frac{\theta r_5(0)}{\varphi^2 - \theta^2} + \frac{2\theta r_6(0)}{\varphi^2 - 4\theta^2} + \alpha'_5(0) + \frac{\alpha}{2} \left(\frac{\alpha'_1(0) \alpha_1(0)}{\varphi^2} + \frac{2B\theta \alpha'_1(0)}{\varphi^2 - \theta^2} + \frac{\alpha'_1(0) \alpha_1(0)}{(\varphi^2 - 4\theta^2)} \right) \right] \quad (4.32b)$$

i.e

$$\beta_7(0) = B^2 \left(\frac{\alpha S_0}{\varphi} + \frac{\alpha}{2\varphi^3} + \frac{\alpha}{2\alpha(\varphi^2 - 4\theta^2)} - \frac{\alpha}{\alpha(\varphi^2 - \theta^2)} - \frac{2\theta \alpha S_1}{\varphi(\varphi^2 - 4\theta^2)} \right) \quad (4.32c)$$

So far, it follows that

$$U^{(21)} = U_m^{(21)}(1 - \cos 2mx) + U_{2m}^{(21)}(1 - \cos 4mx) \quad (4.33)$$

From (3.23),

$$\begin{aligned}
O(\epsilon^2 \delta^2) : U_{,\hat{t}\hat{t}}^{(22)} + U_{,xxxx}^{(22)} + 2\lambda U_{,xx}^{(22)} &= -U_{,\tau\tau}^{(20)} - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(12)} - 2U_{,\hat{t}\tau}^{(21)} - 2\omega'_1 U_{,\hat{t}\hat{t}}^{(12)} - 2\omega'_1 U_{,\hat{t}}^{(11)} - \\
&2U_{,\hat{t}}^{(21)} - 2\omega'_1 U_{,\hat{t}}^{(11)} - \alpha \left\{ (U^{(11)})^2 + U^{(10)} \right\} \\
\Rightarrow U_{,\hat{t}\hat{t}}^{(22)} + U_{,xxxx}^{(22)} + 2\lambda U_{,xx}^{(22)} &= - \left[U_{m,\tau\tau}^{(20)}(1 - \cos 2mx) + U_{2m,\tau\tau}^{(20)}(1 - \cos 4mx) + 2\omega'_1 U_{m,\hat{t}\hat{t}}^{(12)}(1 - \right. \\
&\cos 2mx) + 2 \left\{ U_{m,\hat{t}\tau}^{(21)}(1 - \cos 2mx) + U_{2m,\hat{t}\tau}^{(21)}(1 - \cos 2mx) \right\} + 2\omega'_1 U_{m,\hat{t}}^{(11)}(1 - \cos 2mx) + \\
&2 \left\{ U_{m,\hat{t}}^{(21)}(1 - \cos 2mx) + U_{2m,\hat{t}}^{(21)}(1 - \cos 4mx) \right\} + 2\omega'_1 U_{m,\hat{t}\hat{t}}^{(11)}(1 - \cos 2mx) + \alpha (U_m^{(11)})^2 \left. \right\} \frac{3}{2} -
\end{aligned}$$

$$2\cos 2mx + \frac{1}{2}\cos 4mx\} + 2\{U_m^{(10)}U_m^{(12)}\}\left\{\frac{3}{2} - 2\cos 2mx + \frac{1}{2}\cos 4mx\right\} \quad (4.34)$$

Let

$$U^{(22)} = \sum_{n=1}^{\infty} U_n^{(22)}(\hat{t}, \tau)(1 - \cos 2nx)$$

The LHS of (4.34) simplifies to,

$$\sum_{n=1}^{\infty} \left[U_{n,\hat{t}\hat{t}}^{(22)}(1 - \cos 2nx) + (-16n^4 + 8\lambda n^2)U_n^{(22)} + U_n^{(22)}(1 \cos 2nx) \right] = \text{RHS of (4.34)}$$

Multiplying (4.30) through by $\cos 2mx$ and integrating from 0 to π and for $n=m$, we have,

$$\begin{aligned} & -\frac{\pi}{2}U_{m,\hat{t}\hat{t}}^{(22)} + (-16m^4 + 8\lambda m^2)U_m^{(22)}\left(\frac{\pi}{2}\right) + \left(-\frac{\pi}{2}U_m^{(22)}\right) \\ & = -\left[\left(-\frac{\pi}{2}\right)U_{m,\tau\tau}^{(20)} + 2\omega_1'U_{m,\hat{t}\hat{t}}^{(12)}\left(-\frac{\pi}{2}\right) + 2U_{m,\hat{t}\tau}^{(12)}\left(-\frac{\pi}{2}\right) + 2\omega_1''U_{m,\hat{t}}^{(11)}\left(-\frac{\pi}{2}\right)\right. \\ & \quad \left.+ 2U_{m,\hat{t}}^{(21)}\left(-\frac{\pi}{2}\right) + 2\omega_1'U_{m,\hat{t}}^{(11)}\left(-\frac{\pi}{2}\right) + \alpha(U_m^{(11)})^2\left(-2\cdot\frac{-\pi}{2}\right)\right. \\ & \quad \left.+ \alpha U_m^{(10)}U_m^{(12)}\left(-2\cdot\frac{-\pi}{2}\right)\right] \quad (4.35a) \end{aligned}$$

Further simplification of (4.35a) gives

$$\begin{aligned} & -\frac{\pi}{2}U_{m,\hat{t}\hat{t}}^{(22)} + (16m^4 - 8\lambda m^2 + 1)U_m^{(22)} \\ & = -\frac{\pi}{2}\left[-U_{m,\tau\tau}^{(20)} - 2\omega_1'U_{m,\hat{t}\hat{t}}^{(12)} - 2U_{m,\hat{t}\tau}^{(21)} - 2\omega_1''U_{m,\hat{t}}^{(11)} - 2U_{m,\hat{t}}^{(21)} - 2\omega_1'U_{m,\hat{t}}^{(11)}\right. \\ & \quad \left.+ 2\alpha(U_m^{(11)})^2 + 2\alpha U_m^{(10)}U_m^{(12)}\right] \quad (4.35b) \end{aligned}$$

Further simplification of (4.35b) yields

$$U_{m,\hat{t}\hat{t}}^{(22)} + \theta^2 U_m^{(22)} = -\left[U_{m,\tau\tau}^{(20)} + 2\omega_1'U_{m,\hat{t}\hat{t}}^{(12)} + 2U_{m,\hat{t}\tau}^{(21)} + 2\omega_1''U_{m,\hat{t}}^{(11)} + 2U_{m,\hat{t}}^{(21)} + 2\omega_1'U_{m,\hat{t}}^{(11)} - 2\{(U_m^{(11)})^2 + U_m^{(10)}U_m^{(12)}\}\right] \quad (4.35c)$$

The initial conditions for (4.35c) are

$$U_m^{(22)}(0,0) = 0; \quad U_{m,\hat{t}}^{(22)}(0,0) + \omega_1'(0)U_{m,\hat{t}}^{(12)} + U_{m,\tau}^{(21)}(0,0) = 0$$

Next from equation (4.34) for $n=2m$, let

$$U^{(22)} = \sum_{n=1}^{\infty} U_n^{(22)}(1 - \cos 4mx)$$

Multiplying (4.34) through by $\cos 4mx$ and integrating from 0 to π and for $n = 2m$, gives

$$-\frac{\pi}{2}U_{2m,\hat{t}\hat{t}}^{(22)} + \frac{\pi}{2}(-256m^4 + 32\lambda m^2)U_{2m}^{(22)} - \frac{\pi}{2}U_{2m}^{(22)}\frac{\alpha}{2}\left\{(U_m^{(11)})^2 + (U_m^{(10)}U_m^{(12)})\left(\frac{\pi}{2}\right)\right\} \quad (4.36a)$$

This further gives

$$U_{2m,\hat{t}\hat{t}}^{(22)} + \varphi^2 U_{2m}^{(22)} = -\frac{\alpha}{2}\left\{(U_m^{(11)})^2 + (U_m^{(10)}U_m^{(12)})\right\} \quad (4.36b)$$

The initial conditions are

$$U_{2m}^{(22)}(0,0) = 0; \quad U_{2m,\hat{t}}^{(22)}(0,0) + \omega_1'(0)U_{2m,\hat{t}}^{(12)} + U_{2m,\tau}^{(21)}(0,0) = 0$$

On substituting for terms in (4.35c) and simplifying, the result is

$$\begin{aligned} & U_{2m,\hat{t}\hat{t}}^{(22)} + \theta^2 U_m^{(22)} = \\ & -\left[\left\{\alpha_4''\cos\theta\hat{t} + \frac{r_0''}{\theta^2} - \frac{r_1''\cos 2\theta\hat{t}}{3\theta^2}\right\} + 2\omega_1'\{-\theta^2\alpha_3\sin\theta\hat{t} + \beta_3\theta^2\cos\theta\hat{t}\} + 2\{-\theta\alpha_6'\sin\theta\hat{t} + \theta\beta_6'\cos\theta\hat{t} - \right. \\ & \quad \left. \frac{2\theta r_3'\sin 2\theta\hat{t} + 2\theta r_4'\cos 2\theta\hat{t}}{3\theta^2}\right\} + 2\omega_1'(\theta\beta_2\cos\theta\hat{t}) + 2\left\{-\alpha_6\theta\sin\theta\hat{t} + \beta_6\theta\cos\theta\hat{t} + \left(\frac{2\theta r_3\sin\theta\hat{t} - 2\theta r_4\cos\theta\hat{t}}{3\theta^2}\right)\right\} + \\ & \quad 2\omega_1'\{\theta\beta_2\cos\theta\hat{t}\} + \\ & \quad 2\alpha\left\{\frac{\beta_2}{2}(1 - \cos 2\theta\hat{t}) + \right. \\ & \quad \left. \left(\frac{\alpha_1\beta_2}{2}\sin 2\theta\hat{t} + B\beta_2\sin 2\theta\hat{t}\right)\right\} \quad (4.37a) \end{aligned}$$

To ensure a uniformly valid solution in \hat{t} ; equate to zero the coefficients of $\cos\theta\hat{t}$ and $\sin\theta\hat{t}$ of (4.37a) and this yields respectively

$$-\alpha_4'' + 2\omega_1'\theta\beta_3 - 2\theta\beta_6' - 2\omega_1''\theta\beta_2 - 2\beta_6\theta - 2\omega_1'\theta\beta_2 = 0 \quad (4.37b)$$

and

$$2\omega_1'\theta^2\alpha_3 + 2\theta\alpha_6' + 2\alpha_6\theta - 2\alpha B\beta_2 = 0 \quad (4.37c)$$

Simplification of (4.37b) gives

$$\beta_6' + \beta_6 = \frac{1}{2\theta}[\alpha_4'' - 2\omega_1'\theta^2\beta_3 + 2\omega_1''\beta_2 + 2\omega_1'\theta\beta_2] = \rho_2(\tau) \quad (4.37d)$$

$$\beta_6(\tau) = e^{-\tau}[\int e^{\tau} \rho_2(\tau) d\tau + \beta_6(0)] = e^{-\tau}[\int e^{\tau} \rho_2(\tau) d\tau] \quad (4.37e)$$

Similarly, simplification of (4.37c) yields

$$\alpha'_6 + \alpha_6 = \frac{1}{2\theta}[-2\omega'_1\theta^2\alpha_3 + 2\alpha B\beta_2] = \rho_3(\tau) \quad (4.37f)$$

Therefore

$$\alpha_6 = e^{-\tau}[\int e^{\tau} \rho_3(\tau) d\tau + \alpha_6(0)] \quad (4.37g)$$

The remaining part of equation (4.37a) is

$$U_{m,\hat{t}\hat{t}}^{(22)} + \theta^2 U_m^{(22)} = r_7 + r_8 \cos 2\theta \hat{t} + r_9 \sin 2\theta \hat{t} \quad (4.38)$$

$$r_7 = -\left[\frac{r'_0}{\theta^2} - \frac{2r'_4}{3\theta} + \alpha\beta_2\right], r_8 = \left[\frac{r'_1}{3\theta^2} - \frac{4r_4}{3\theta} - \alpha\beta_2\right], r_9 = -\left[\frac{2r'_3}{3\theta^2} - \frac{4r_3}{3\theta} - \alpha\alpha_1\beta_2\right]$$

It is to be recalled that, $r_0 = -2\alpha\left(\frac{\alpha_1^2}{2} + B^2\right)$

$$\begin{aligned} \therefore r'_0 &= -2\alpha\alpha_1\alpha'_1, r''_0 = -2\alpha(\alpha_1'^2 + \alpha_1\alpha''_1), r'_0(0) = 2\alpha B^2, r''_0(0) = -4\alpha B^2 \\ r_4 &= \alpha\alpha_1\beta_2 + \frac{8\alpha}{3\theta}(\alpha_1\alpha'_1 + \alpha_1'^2), r'_4 = \alpha(\alpha'_1\beta_2 + \alpha_1'\beta_2') + \frac{8\alpha}{3\theta}(\alpha'_1\alpha'_1 + \alpha_1\alpha''_1 + 2\alpha_1\alpha'_1) \\ r'_4(0) &= \frac{-5\alpha B^2}{2\theta}, r_7(0) = \frac{17\alpha B^2}{3\theta^2} - \frac{\alpha B}{\theta}, r_1 = -\alpha_1\alpha_1^2; r'_1 = 2\alpha\alpha_1\alpha'_1; r''_1 = -2\alpha(\alpha_1'^2 + \alpha_1\alpha''_1) \\ r'_1(0) &= -2\alpha\alpha_1(0)\alpha'_1(0) = 2\alpha B^2; r''_1(0) = -4\alpha B^2, r_8(0) = \left(\frac{-4\alpha B^2}{3\theta^2} + \frac{4\alpha B^2}{3\theta^2} + \frac{\alpha B}{3\theta}\right) = \frac{\alpha B}{\theta}, r_3 = \\ \alpha\alpha_1\alpha_2; r_3(0) &= 0, r'_3 = \alpha(\alpha'_1\alpha_1 + \alpha_1\alpha'_2); r'_3(0) = 0, \beta'_2 = \frac{3B}{2\theta}, r_9(0) = \frac{3\theta B^2}{2\theta} \\ \therefore U_m^{(22)} &= \alpha_8 \cos \theta + \beta_8 \sin \theta + \frac{r_7}{3\theta^2} + \frac{r_8 \cos 2\theta}{\theta^2} + \frac{r_9 \sin 2\theta \hat{t}}{\theta^2} \end{aligned} \quad (4.39a)$$

The initial conditions are

$$U_m^{(22)}(0,0) = 0; U_m^{(22)}(0,0) + \omega'_1(0)U_{m,\hat{t}}^{(12)}(0,0) + U_{m,\tau}^{(21)}(0,0) = 0$$

$$\alpha_8(0) = \left(\frac{r_8(0)}{3\theta^2} - \frac{r_7(0)}{\theta^2}\right) = \frac{4\alpha B}{3\theta^3} - \frac{17\alpha B^2}{3\theta^4} \quad (4.39b)$$

$$\text{Similarly, } \theta\beta_8(0) + \frac{2\theta r_9(0)}{\theta^2} = 0$$

$$\therefore \beta_8(0) = \frac{-2\theta r_9(0)}{\theta^2} = \frac{-3\alpha B^2}{\theta^3} \quad (4.39c)$$

From equation (3.23),

$$\begin{aligned} O(\epsilon^3) : U_{,\hat{t}\hat{t}}^{(30)} + U_{,xxxx}^{(30)} + 2\lambda U_{,xx}^{(30)} + U^{(30)} \\ = -(\omega'_1)^2 U_{,\hat{t}\hat{t}}^{(10)} - 2(\omega'_1 U_{,\hat{t}\hat{t}}^{(20)} + \omega'_2 U_{,\hat{t}\hat{t}}^{(20)}) - 2\alpha U^{(20)} U^{(10)} + \beta(U^{(10)})^3 \end{aligned}$$

Then, substituting on the RHS of (3.23) yields,

$$\begin{aligned} U_{,\hat{t}\hat{t}}^{(30)} + U_{,xxxx}^{(30)} + 2\lambda U_{,xx}^{(30)} + U^{(30)} \\ = -(\omega'_1)^2 U_{m,\hat{t}\hat{t}}^{(10)}(1 - \cos 2mx) \\ - 2\left[\omega'_1 U_{m,\hat{t}\hat{t}}^{(20)} + (1 - \cos 2mx) + \omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)}(1 - \cos 2mx)\right] - 2\omega'_2 U_{,\hat{t}\hat{t}}^{(10)}(1 - \cos 2mx) \\ - 2\alpha \left[U_m^{(10)}(1 - \cos 2mx)\right] \left\{U_m^{(20)}(1 - \cos 2mx) + U_{2m}^{(20)}(1 - \cos 4mx)\right\} + \beta(U_m^{(10)})^3 (1 - \cos 2mx)^3 \end{aligned} \quad (4.40)$$

Therefore, on further simplifications, (4.40) becomes

$$\begin{aligned} U_{,\hat{t}\hat{t}}^{(30)} + U_{,xxxx}^{(30)} + 2\lambda U_{,xx}^{(30)} + U^{(30)} = -(\omega'_1)^2 U_{m,\hat{t}\hat{t}}^{(10)}(1 - \cos 2mx) - 2\left[\omega'_1 U_{m,\hat{t}\hat{t}}^{(20)}(1 - \cos 2mx) + \omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)}(1 - \cos 4mx) + \omega'_2 U_m^{(10)}(1 - \cos 2mx)\right] - 2\alpha \left[U_m^{(10)} U_m^{(20)} \left\{\frac{3}{2} - 2\cos 2mx + \frac{1}{2}\cos 4mx\right\} + U_m^{(10)} U_m^{(20)} \left\{1 - \frac{1}{2}\cos 2mx - \cos 4mx + \frac{1}{2}\cos 6mx\right\}\right] + \beta(U_m^{(10)})^3 \left[\frac{5}{2} - \frac{15}{4}\cos 2mx + \frac{3}{2}\cos 4mx + \frac{1}{4}\cos 6mx\right] \end{aligned} \quad (4.41)$$

Let

$$U^{(30)} = \sum_{n=1}^{\infty} U_n^{(30)}(1 - \cos 2nx)$$

Therefore, (4.41) becomes

$$\sum_{n=1}^{\infty} \left[U_{n,\hat{t}\hat{t}}^{(30)}(1 - \cos 2nx) + (-16n^4 + 8\lambda n^2 + 1)U_n^{(30)} \cos 2nx\right] = RHS \quad (4.41)$$

Multiplying (4.41) through by $\cos 2mx$ and integrating from 0 to π and for $n=m$, the result is

$$\begin{aligned}
& -\frac{\pi}{2}U_{m,\hat{t}\hat{t}}^{(30)} + (-16m^4 + 8\lambda m^2 + 1)U_m^{(30)}\left(-\frac{\pi}{2}\right) = -\left[(\omega'_1)^2U_{m,\hat{t}\hat{t}}^{(10)}\left(-\frac{\pi}{2}\right) + 2\omega'_1U_{m,\hat{t}\hat{t}}^{(20)}\left(-\frac{\pi}{2}\right) + \right. \\
& 2\omega'_2U_{m,\hat{t}\hat{t}}^{(10)}\left(-\frac{\pi}{2}\right) + 2\alpha U_m^{(10)}U_m^{(20)}\left(-2, -\frac{\pi}{2}\right) - \alpha U_m^{(10)}U_{2m}^{(20)}\left(-\frac{\pi}{2}\right) - \\
& \left. \frac{15}{4}(U_m^{(10)})^3\left(-\frac{\pi}{2}\right)\right] \quad (4.42a)
\end{aligned}$$

i.e,

$$\begin{aligned}
& -\frac{\pi}{2}\left[U_{m,\hat{t}\hat{t}}^{(30)} + (16m^4 - 8\lambda m^2 + 1)U_m^{(30)}\right] = \\
& -\frac{\pi}{2}\left[-(\omega'_1)^2U_{m,\hat{t}\hat{t}}^{(10)} - 2\omega'_1U_{m,\hat{t}\hat{t}}^{(20)} - 2\omega'_2U_{m,\hat{t}\hat{t}}^{(10)} - 2\alpha[2U_m^{(10)}U_m^{(20)} + U_m^{(10)}U_{2m}^{(20)}] - \right. \\
& \left. \frac{15}{4}\beta(U_m^{(10)})^3\right] \quad (4.42b)
\end{aligned}$$

$$\begin{aligned}
\therefore U_{m,\hat{t}\hat{t}}^{(30)} + \theta^2U_m^{(30)} = & -(\omega'_1)^2U_{m,\hat{t}\hat{t}}^{(10)} - 2\omega'_1U_{m,\hat{t}\hat{t}}^{(20)} - 2\omega'_2U_{m,\hat{t}\hat{t}}^{(10)} - 2\alpha[2U_m^{(10)}U_m^{(20)} + U_m^{(10)}U_{2m}^{(20)}] - \\
& \frac{15}{4}\beta(U_m^{(10)})^3 \quad (4.43)
\end{aligned}$$

The initial conditions are

$$U_m^{(30)}(0,0) = 0; \quad U_{m,\hat{t}}^{(30)}(0,0) + \omega'(0)U_{m,\hat{t}}^{(20)}(0,0) + \omega'_2(0)U_{\tau}^{(10)}(0,0) = 0$$

Multiplying (4.41) through by $\cos 4mx$ and integrating from 0 to π and for $n=2m$, the result gives

$$\begin{aligned}
& -\frac{\pi}{2}\left[U_{2m,\hat{t}\hat{t}}^{(30)} + (256m^4 - 32\lambda m^2 + 1)U_{2m}^{(30)}\right] \\
& = 2\left[\omega'_1U_{2m,\hat{t}\hat{t}}^{(20)}\left(-\frac{\pi}{2}\right)\right] + 2\alpha\left[U_m^{(10)}U_m^{(20)}\cdot\frac{1}{2}\left(-\frac{\pi}{2}\right) + U_m^{(10)}U_{2m}^{(20)}\left(-\frac{\pi}{2}\right)\right] \\
& - \beta(U_m^{(10)})^3\cdot\frac{3}{2}\left(-\frac{\pi}{2}\right) \\
\therefore U_{2m,\hat{t}\hat{t}}^{(30)} + \varphi^2U_{2m}^{(30)} = & -2\left[-\omega'_1U_{2m,\hat{t}\hat{t}}^{(20)}\right] + 2\alpha\left[U_m^{(10)}U_m^{(20)} - U_m^{(10)}U_{2m}^{(20)}\right] - \frac{3}{2}\beta(U_m^{(10)})^3 \quad (4.44)
\end{aligned}$$

The initial conditions are

$$U_{2m}^{(30)}(0,0) = 0; \quad U_{2m,\hat{t}}^{(30)}(0,0) + \omega'_1(0)U_{2m,\hat{t}}^{(20)}(0,0) = 0$$

Multiplying (4.41) through by $\cos 6mx$ and integrating from 0 to π and for $n=3m$ and get,

$$U_{3m,\hat{t}\hat{t}}^{(30)} + (1296m^4 - 72\lambda m^2 + 1)U_{3m}^{(30)} = \alpha U_m^{(10)}U_{2m}^{(20)} - \frac{1}{4}\beta(U_m^{(10)})^3 \quad (4.45a)$$

Let

$$\Omega^2 = 1296m^4 - 72\lambda m^2 + 1 > 0 \text{ for all } m$$

$$\therefore U_{3m,\hat{t}\hat{t}}^{(30)} + \Omega^2U_{3m}^{(30)} = \alpha U_m^{(10)}U_{2m}^{(20)} - \frac{1}{4}\beta(U_m^{(10)})^3 \quad (4.45b)$$

The initial conditions for (4.45b) are

$$U_{3m}^{(30)}(0,0) = 0; \quad U_{3m,\hat{t}}^{(30)}(0,0) = 0$$

Further simplification of (4.43) gives

$$\begin{aligned}
& U_{m,\hat{t}\hat{t}}^{(30)} + \theta^2U_m^{(30)} = -(\omega'_1)^2(-\theta^2\alpha_1\cos\theta\hat{t}) - 2\omega'_1\left(-\alpha_4\theta^2\cos\theta\hat{t} + \frac{4r_1\cos 2\theta\hat{t}}{3}\right) - 2\omega'_2(-\theta^2\alpha_1\cos\theta\hat{t}) - \\
& 2\alpha\left[\left(\frac{\alpha_1\alpha_4}{4} + \frac{Br_0}{\theta^2}\right) + \left(\frac{\alpha_1r_0}{\theta^2} - \frac{\alpha_1r_1}{6\theta^2} + B\alpha_4\right)\cos\theta\hat{t} + \left(\frac{\alpha_1\alpha_4}{4} - \frac{Br_1}{3\theta^2}\right)\cos 2\theta\hat{t} - \frac{\alpha_1r_1}{6\theta^2}\cos 3\theta\hat{t}\right] - 2\alpha\left[\frac{\alpha\alpha_1^2B}{2(\varphi^2-\theta^2)} + \right. \\
& \left.\left\{\frac{\alpha\alpha_1^2\left(\frac{\alpha_1^2}{2} + B^2\right)}{2\varphi^2} + \frac{\alpha\alpha_1^3}{8(\varphi^2-4\theta^2)}\right\}\cos\theta\hat{t} + \frac{\alpha_1\alpha_5}{2}\cos(\varphi + \theta)\hat{t} + \frac{\alpha\alpha_1^2B\cos 2\theta\hat{t}}{2(\varphi^2-\theta^2)} + \frac{\alpha_1\beta_5}{2}\sin(\varphi + \theta)\hat{t} + \frac{\alpha_1\alpha_5}{2}\cos(\varphi - \right. \\
& \left. \theta)\hat{t} + \frac{\alpha_1\beta_5}{2}\sin(\varphi - \theta)\hat{t} + \frac{\alpha\alpha_1^3\cos 3\theta\hat{t}}{8(\varphi^2-4\theta^2)}\right] - \frac{15\beta}{4}\left[\left(B^3 + \frac{3\alpha_1^2B}{2}\right)\right] + 3\left(\frac{\alpha_1^3}{4} + \alpha_1B^2\right)\cos\theta\hat{t} + \frac{3\alpha_1^2}{2}B\cos 2\theta\hat{t} + \\
& \frac{\alpha_1^3}{4}\cos 3\theta\hat{t} \quad (4.45c)
\end{aligned}$$

To ensure a uniformly valid solution in \hat{t} , equate to zero the coefficients of $\cos\theta\hat{t}$ and this yields

$$\begin{aligned}
& (\omega_1')^2 \theta^2 + 2\omega_1' \alpha_4 \theta^2 + 2\omega_2' \theta^2 \alpha_1 - 2\alpha \left(\frac{\alpha_1 r_0}{\theta^2} - \frac{\alpha_1 r_1}{6\theta^2} + B\alpha_4 \right) - \left\{ \frac{\alpha^2 \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{\varphi^2} + \frac{\alpha_1^3 \alpha_1}{4(\varphi^2 - 4\theta^2)} \right\} - \\
& \frac{45\beta}{4} \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) = 0 \\
\therefore \omega_2' = & -\frac{1}{2\theta^2 \alpha_1} \left[(\omega_1')^2 \theta^2 + 2\omega_1' \alpha_4 \theta^2 - 2\alpha \left(\frac{\alpha_1 r_0}{\theta^2} - \frac{\alpha_1 r_1}{6\theta^2} + B\alpha_4 \right) - \left\{ \frac{\alpha^2 \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{\varphi^2} + \frac{\alpha_1^3 \alpha_1}{4(\varphi^2 - 4\theta^2)} \right\} - \right. \\
& \left. \frac{45\beta}{4} \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) \right] \tag{4.46}
\end{aligned}$$

The remaining equation in (4.45c) is

$$U_{m,\hat{t}\hat{t}}^{(30)} + \theta^2 U_m^{(30)} = r_{10} + r_{11} \cos 2\theta \hat{t} + r_{12} \cos 3\theta \hat{t} + r_{13} \cos(\varphi + \theta) \hat{t} + r_{14} \sin(\varphi + \theta) \hat{t} + r_{15} \cos(\varphi - \theta) \hat{t} + r_{16} \sin(\varphi - \theta) \hat{t} \tag{4.47}$$

Solving (4.47) gives

$$\begin{aligned}
U_m^{(30)}(\hat{t}, \tau) = & \alpha_9(\tau) \cos \theta \hat{t} + \beta_9(\tau) \sin \theta \hat{t} + \frac{r_{10}}{\theta^2} - \frac{r_{11} \cos \theta \hat{t}}{3\theta^2} - \frac{r_{12} \cos 3\theta \hat{t}}{8\theta^2} \\
& - \frac{1}{\varphi(2\theta + \varphi)} [r_{13} \cos(\varphi + \theta) \hat{t} + r_{14} \sin(\varphi + \theta) \hat{t}] \\
& + \frac{1}{\varphi(2\theta - \varphi)} [r_{15} \cos(\varphi - \theta) \hat{t} + r_{16} \sin(\varphi - \theta) \hat{t}] \tag{4.48}
\end{aligned}$$

The initial conditions are

$$U_m^{(30)}(0,0) = 0, \quad U_{m,\hat{t}}^{(30)}(0,0) + \omega_1'(0) U_{m,\hat{t}}^{(20)}(0,0) + \omega_2'(0) U_{m,\tau}^{(10)}(0,0) = 0$$

where

$$\alpha_9(0) = \left[-\frac{r_{10}}{\theta^2} + \frac{r_{11}}{3\theta^2} + \frac{r_{12}}{8\theta^2} + \frac{r_{13}}{\varphi(2\theta + \varphi)} - \frac{r_{15}}{\varphi(2\theta - \varphi)} \right] \text{ at } \tau = 0$$

and

$$\beta_9(0) = \frac{1}{\theta} \left[\frac{r_{14}(\varphi + \theta)}{\varphi(2\theta + \varphi)} - \frac{r_{16}(\varphi - \theta)}{\varphi(2\theta - \varphi)} \right] \text{ at } \tau = 0$$

and where,

$$\begin{aligned}
r_{10} = & - \left[2\alpha \left(\frac{\alpha_1 \alpha_4}{4} + \frac{B r_0}{\theta^2} \right) + \frac{\alpha^2 \alpha_1^2 B}{\varphi^2 - \theta^2} - \frac{15\beta}{4} \left(B^3 + \frac{3\alpha_1^2 B}{2} \right) \right] \\
r_{10}(0) = & B^3 \left(\frac{10\alpha^2}{3\theta^2} - \frac{\alpha^2}{\varphi^2 - \theta^2} + \frac{75\beta}{8} \right) \\
r_{10}'(0) = & B^3 \left[\frac{\alpha S_5}{2} - \frac{8\alpha^2}{3\theta^2} - \frac{4\alpha^2}{\theta} - \frac{2\alpha^2}{(\varphi^2 - \theta^2)} - \frac{45\beta}{4} \right] \\
r_{11} = & - \left(\frac{8r_1 \omega_1'}{3} + 2\alpha \left(\frac{\alpha_1 \alpha_4}{4} + \frac{B r_0}{3\theta^2} \right) + \frac{\alpha^2 \alpha_1^2 B}{\varphi^2 - \theta^2} + \frac{45\beta \alpha_1^2 B}{2} \right) \\
r_{11}(0) = & B^3 \left(\frac{8\alpha}{3\theta^2} + \frac{2\alpha^2}{3\theta^2} - \frac{45\beta}{8} + \frac{\alpha^2}{\varphi^2 - \theta^2} \right) \\
r_{11}'(0) = & B^3 \left[\frac{4\alpha^2}{3\theta^2} - \frac{16\alpha}{3\theta^2} - \frac{\alpha S_5}{2} - \frac{2\alpha}{(\varphi^2 - \theta^2)} - \frac{45\beta}{4} \right] \\
r_{12} = & \frac{\alpha \alpha_1 r_1}{3\theta^2} - \frac{\alpha_1^3 \alpha^2}{4(\varphi^2 - \theta^2)} - \frac{15\beta \alpha_1^3}{16}, \quad r_{12}(0) = B^3 \left[\frac{\alpha^2}{3\theta^2} + \frac{\alpha^2}{4(\varphi^2 - \theta^2)} + \frac{15\beta}{16} \right] \quad r_{12}'(0) = B^3 \left[\frac{3\alpha^2}{4(\varphi^2 - \theta^2)} - \frac{\alpha^2}{3\theta^2} + \frac{45\beta}{4} \right], \\
r_{13} = & -\alpha \alpha_1 \alpha_5 = r_{15}, \quad r_{13}(0) = r_{15}(0) = B^3 \alpha^2 S_0, \quad r_{13}'(0) = r_{15}'(0) = -2\alpha S_0 B^3 \\
r_{14} = & -\alpha \alpha_1 \beta_5 = r_{16}, \quad r_{14}(0) = r_{16}(0) = 0 \text{ since } \beta_5(0) = 0, \quad r_{16}'(0) = r_{14}'(0) = 0
\end{aligned}$$

Substituting in (4.44) gives

$$\begin{aligned}
& U_{2m,\hat{t}\hat{t}}^{(30)} + \varphi^2 U_{2m}^{(30)} = \\
& 2\omega_1' \left[-\varphi^2 \alpha_5 \cos \varphi \hat{t} - \varphi^2 \beta_5 \sin \varphi \hat{t} + \frac{\alpha}{2} \left\{ \frac{-2\theta^2 B \alpha_1 \cos \theta \hat{t}}{\varphi^2 - \theta^2} - \frac{2\alpha_1^2 \theta^2 \cos 2\theta \hat{t}}{\varphi^2 - 4\theta^2} \right\} \right] + \\
& 2\alpha \left[\left(\frac{\alpha_1 \alpha_4}{4} + \frac{B r_0}{\theta^2} \right) + \left(\frac{\alpha_1 r_0}{\theta^2} - \frac{\alpha_1 r_1}{6\theta^2} + B\alpha_4 \right) \cos \theta \hat{t} + \left(\frac{\alpha_1 \alpha_4}{4} + \frac{B r_1}{3\theta^2} \right) \cos 2\theta \hat{t} \right. \\
& \quad \left. - \frac{\alpha_1 r_1}{6\theta^2} \cos 3\theta \hat{t} \right] - 2\alpha \left[\frac{\alpha \alpha_1^2 B}{2(\varphi^2 - \theta^2)} + \left\{ \frac{\alpha \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{2\varphi^2} + \right. \right.
\end{aligned}$$

$$\left. \frac{\alpha_1^3 \alpha}{8(\varphi^2 - 4\theta^2)} \right\} \cos\theta \hat{t} + \frac{\alpha_1 \alpha_5}{2} \cos(\varphi + \theta) \hat{t} + \frac{\alpha \alpha_1^2 B \cos 2\theta \hat{t}}{2(\varphi^2 - \theta^2)} + \frac{\alpha_1 \beta_5}{2} \sin(\varphi + \theta) \hat{t} + \frac{\alpha_1 \alpha_5}{2} \cos(\varphi - \theta) \hat{t} + \frac{\alpha_1 \beta_5}{2} \sin(\varphi - \theta) \hat{t} + \left. \frac{\alpha_1^3 \alpha \cos 3\theta \hat{t}}{8(\varphi^2 - 4\theta^2)} \right] - \frac{3\beta}{2} \left[\left(B^3 + \frac{3\alpha_1^2 B}{2} \right) + 3 \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) \cos\theta \hat{t} + \frac{3\alpha_1^2 B}{2} \cos 2\theta \hat{t} + \frac{\alpha_1^3}{4} \cos 3\theta \hat{t} \right] \quad (4.49)$$

To ensure a uniformly valid solution in \hat{t} , needs equating to zero the coefficients of $\cos\varphi \hat{t}$ and $\sin\varphi \hat{t}$. A further simplification of (4.49) gives

$$U_{2m, \hat{t}\hat{t}}^{(30)} + \varphi^2 U_{2m}^{(30)} = r_{17} + r_{18} \cos\theta \hat{t} + r_{19} \cos 2\theta \hat{t} + r_{20} \cos 3\theta \hat{t} \quad (4.50)$$

where,

$$\begin{aligned} r_{17} &= \left[2\alpha \left(\frac{\alpha_1 \alpha_4}{4} + \frac{B r_0}{\theta^2} \right) - \frac{\alpha^2 \alpha_1^2 B}{\varphi^2 - \theta^2} - \frac{3\beta}{2} \left(B^3 + \frac{3\alpha_1^2 B}{2} \right) \right] \\ r_{17}(0) &= B^3 \left(-\frac{22\alpha^2}{3\theta^2} + \frac{\alpha^2}{\varphi^2 - \theta^2} + \frac{15\beta}{4} \right) \\ r_{17}'(0) &= B^3 \left[\frac{-S_5 \alpha}{2} + \frac{20\alpha^2}{3\theta^2} + \frac{2\alpha^2}{2(\varphi^2 - \theta^2)} + \frac{9\beta}{2} \right] \\ r_{18} &= \left[\frac{-2\theta^2 \omega_1' B \alpha_1 \alpha}{\varphi^2 - \theta^2} + 2\alpha \left(\frac{\alpha_1 r_0}{\theta^2} - \frac{\alpha_1 r_1}{6\theta^2} + B \alpha_4 \right) - \frac{\alpha^2 \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{\varphi^2} - \frac{\alpha_2 \alpha_1^3}{4(\varphi^2 - 4\theta^2)} - \frac{9}{2} \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) \right] \\ r_{18}(0) &= B^3 \left(\frac{2\alpha}{\varphi^2 - \theta^2} + \frac{18\alpha^2}{6\theta^2} + \frac{3\alpha^2}{\varphi^2} + \frac{\alpha^2}{4(\varphi^2 - 4\theta^2)} + \frac{45}{2} \right) \\ r_{18}'(0) &= B^3 \left[2\alpha S_5 - 2 - \frac{43\alpha^2}{\theta^2} - \frac{5\alpha^2}{2\varphi^2} - \frac{3\alpha^2}{4(\varphi^2 - \theta^2)} - \frac{63}{8} \right] \\ r_{19} &= \left[\frac{-2\omega_1' \alpha_1^2 \alpha \theta^2}{\varphi^2 - \theta^2} + 2\alpha \left(\frac{\alpha_1 \alpha_4}{4} - \frac{B r_1}{3\theta^2} \right) + \frac{\alpha^2 \alpha_1^2 B}{2(\varphi^2 - \theta^2)} - \frac{9\alpha_1^2 B \beta}{4} \right] \\ r_{19}(0) &= B^3 \left(\frac{-2\alpha}{\varphi^2 - \theta^2} - \frac{4\alpha^2}{3\theta^2} + \frac{2\alpha^2}{3\theta^2} + \frac{\alpha^2}{2(\varphi^2 - \theta^2)} - \frac{9\beta}{4} \right) \\ r_{19}'(0) &= B^3 \left[\frac{4\alpha}{B(\varphi^2 - 4\theta^2)} - \frac{S_5}{2} + \frac{4\alpha^2}{3\theta^2} - \frac{\alpha^2}{(\varphi^2 - 4\theta^2)} + \frac{9\beta}{2} \right] \\ r_{20} &= \left[-\frac{\alpha \alpha_1 r_1}{3\theta^2} - \frac{\alpha_1^3 \alpha^2}{4(\varphi^2 - 4\theta^2)} - \frac{3\beta \alpha_1^3}{8} \right], r_{20}(0) = B^3 \left(-\frac{\alpha^2}{3\theta^2} + \frac{\alpha^2}{4(\varphi^2 - \theta^2)} + \frac{3\beta}{8} \right) \\ r_{20}'(0) &= B^3 \left[\frac{\alpha^2}{\theta^2} + \frac{3\alpha^2}{4(\varphi^2 - 4\theta^2)} - \frac{9\beta}{8} \right] \end{aligned}$$

The solution of (4.50) is

$$U_{2m}^{(30)} = \alpha_{10} \cos\varphi \hat{t} + \beta_{10} \sin\varphi \hat{t} + \frac{r_{17}}{\varphi^2} + \frac{r_{18} \cos\theta \hat{t}}{(\varphi^2 - \theta^2)} + \frac{r_{19} \cos 2\theta \hat{t}}{(\varphi^2 - 4\theta^2)} + \frac{r_{20} \cos 3\theta \hat{t}}{(\varphi^2 - 9\theta^2)} \quad (4.51a)$$

The initial conditions for (4.51) are

$$\begin{aligned} U_{2m}^{(30)}(0,0) &= 0; U_{2m, \hat{t}}^{(30)}(0,0) + \omega_1'(0) U_{2m, \hat{t}}^{(20)}(0,0) = 0 \\ \therefore \alpha_{10}(0) &= - \left[\frac{r_{18}}{(\varphi^2 - \theta^2)} + \frac{r_{19}}{(\varphi^2 - 4\theta^2)} + \frac{r_{20}}{(\varphi^2 - 9\theta^2)} \right] \text{ at } \tau = 0, \quad \beta_{10}(0) = 0 \end{aligned} \quad (4.51b)$$

Substituting in (4.45b) the following is obtained

$$\begin{aligned} U_{3m, \hat{t}\hat{t}}^{(30)} + \Omega^2 U_{3m}^{(30)} &= \alpha \left[\frac{\alpha \alpha_1^2 B}{2(\varphi^2 - \theta^2)} + \left\{ \frac{\alpha \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{2\varphi^2} + \frac{\alpha_1^3 \alpha}{8(\varphi^2 - 4\theta^2)} \right\} \cos\theta \hat{t} + \frac{\alpha_1 \alpha_5}{2} \cos(\varphi + \theta) \hat{t} + \frac{\alpha \alpha_1^2 B \cos 2\theta \hat{t}}{2(\varphi^2 - \theta^2)} + \frac{\alpha_1 \beta_5}{2} \sin(\varphi + \theta) \hat{t} + \frac{\alpha_1 \alpha_5}{2} \cos(\varphi - \theta) \hat{t} + \frac{\alpha_1 \beta_5}{2} \sin(\varphi - \theta) \hat{t} + \frac{\alpha_1^3 \alpha \cos 3\theta \hat{t}}{8(\varphi^2 - 4\theta^2)} \right] - \frac{\beta}{4} \left[\left(B^3 + \frac{3\alpha_1^2 B}{2} \right) + 3 \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) \cos\theta \hat{t} + \frac{3\alpha_1^2 B}{2} \cos 2\theta \hat{t} + \frac{\alpha_1^3}{4} \cos 3\theta \hat{t} \right] \end{aligned} \quad (4.52a)$$

Rewriting (4.52a) gives

$$\begin{aligned} U_{3m, \hat{t}\hat{t}}^{(30)} + \Omega^2 U_{3m}^{(30)} &= r_{21} + r_{22} \cos\theta \hat{t} + r_{23} \cos 2\theta \hat{t} + r_{24} \cos 3\theta \hat{t} + r_{25} \cos(\varphi + \theta) \hat{t} + r_{26} \sin(\varphi + \theta) \hat{t} \\ &\quad + r_{27} \cos(\varphi - \theta) \hat{t} + r_{28} \sin(\varphi - \theta) \hat{t} \end{aligned} \quad (4.52b)$$

The initial conditions are

$$U_{3m}^{(30)}(0,0) = 0; U_{3m,\hat{t}}^{(30)}(0,0) = 0$$

where,

$$r_{21} = \left\{ \frac{\alpha^2 \alpha_1^2 B}{2(\varphi^2 - \theta^2)} - \frac{\beta}{4} \left(B^3 + \frac{3\alpha_1^2 B}{2} \right) \right\}, r_{21}(0) = B^3 \left(\frac{\alpha^2 \alpha_1^2}{2(\varphi^2 - \theta^2)} - \frac{5\beta}{8} \right)$$

$$r_{22} = \left\{ \frac{\alpha^2 \alpha_1 \left(\frac{\alpha_1^2}{2} + B^2 \right)}{2\varphi^2} + \frac{\alpha_1^3 \alpha^2}{8(\varphi^2 - 4\theta^2)} - \frac{3\beta}{4} \left(\frac{\alpha_1^3}{4} + \alpha_1 B^2 \right) \right\};$$

$$r_{22}(0) = B^3 \left(\frac{15\beta}{16} - \frac{3\alpha^2}{4\varphi^2} - \frac{\alpha^2}{8(\varphi^2 - 4\theta^2)} \right), r'_{22}(0) = B^3 \left(\frac{5\alpha^2}{4\varphi} + \frac{3\alpha^2}{8(\varphi^2 - 4\theta^2)} - \frac{21\beta}{16} \right)$$

$$r_{23} = \left\{ \frac{\alpha^2 \alpha_1^2 B}{2(\varphi^2 - \theta^2)} - \frac{3\alpha_1^2 B}{8} \right\}, r_{23}(0) = B^3 \left(\frac{\alpha^2}{2(\varphi^2 - \theta^2)} - \frac{3\beta}{8} \right)$$

$$r'_{23}(0) = B^3 \left(\frac{3\beta}{4} - \frac{\alpha^2}{(\varphi^2 - \theta^2)} \right), r_{24} = \left(\frac{\alpha^2 \alpha_1^3 B}{8(\varphi^2 - \theta^2)} - \frac{3\alpha_1^3 \beta}{16} \right)$$

$$r_{24}(0) = B^3 \left(\frac{\beta}{16} - \frac{\alpha^2}{8(\varphi^2 - 4\theta^2)} \right), r'_{24}(0) = B^3 \left(\frac{3\alpha^2}{8(\varphi^2 - \theta^2)} - \frac{3\beta}{16} \right), r_{25} = \frac{\alpha \alpha_1 \alpha_3}{2} = r_{27}, r_{25}(0) = r_{27}(0) =$$

$$B^2 \alpha^2 S_0, r'_{25}(0) = r'_{27}(0) = \alpha S_0 B^3$$

$$r_{26} = \frac{\alpha \alpha_1 \beta_5}{2} = r_{28}; r_{26}(0) = r_{28}(0) = 0, r'_{26}(0) = r'_{28}(0) = 0$$

$$\begin{aligned} \therefore U_{3m}^{(30)}(\hat{t}, \tau) = & \alpha_{11}(\tau) \cos \Omega \hat{t} + \beta_{11}(\tau) \sin \Omega \hat{t} + \frac{r_{22} \cos \theta \hat{t}}{\Omega^2 - \theta^2} + \frac{r_{23} \cos 2\theta \hat{t}}{\Omega^2 - 4\theta^2} + \frac{r_{24} \cos 3\theta \hat{t}}{\Omega^2 - 9\theta^2} \\ & + \left\{ \frac{r_{25} \cos(\varphi + \theta) \hat{t} + r_{26} \sin(\varphi + \theta) \hat{t}}{\Omega^2 - (\varphi + \theta)^2} \right\} \\ & + \left\{ \frac{r_{27} \cos(\varphi - \theta) \hat{t} + r_{28} \sin(\varphi - \theta) \hat{t}}{\Omega^2 - (\varphi - \theta)^2} \right\} \end{aligned} \quad (4.52b)$$

$$\alpha_{11}(0) = - \left[\frac{r_{22}}{\Omega^2 - \theta^2} + \frac{r_{23}}{\Omega^2 - 4\theta^2} + \frac{r_{24}}{\Omega^2 - 9\theta^2} + \frac{r_{25}}{\Omega^2 - (\varphi + \theta)^2} + \frac{r_{27}}{\Omega^2 - (\varphi - \theta)^2} \right] \tau = 0 \quad (4.52c)$$

$$\beta_{11}(0) = \frac{-1}{\Omega} \left[\frac{r_{26}(\varphi + \theta)}{\Omega^2 - (\varphi + \theta)^2} + \frac{r_{28}(\varphi - \theta)}{\Omega^2 - (\varphi - \theta)^2} \right] \tau = 0 \quad (4.53)$$

So far, it follows that

$$U^{(30)} = U_m^{(30)}(1 - \cos 2mx) + U_{2m}^{(30)}(1 - \cos 4mx) + U_{3m}^{(30)}(1 - \cos 6mx)$$

From (3.24),

$$\begin{aligned} O(\epsilon^3 \delta) : U_{,\hat{t}\hat{t}}^{(31)} + U_{,xxxx}^{(31)} + 2\lambda U_{,xx}^{(31)} + U^{(31)} \\ = -(\omega_1')^2 U_{,\hat{t}\hat{t}}^{(11)} - 2(\omega_1' U_{,\hat{t}\tau}^{(21)} + \omega_2' U_{,\hat{t}\tau}^{(11)}) - 2U_{,\hat{t}\tau}^{(30)} + 2(\omega_1' U_{,\hat{t}\hat{t}}^{(20)} + \omega_2' U_{,\hat{t}\hat{t}}^{(10)}) \\ - (\omega_1'' U_{,\hat{t}}^{(20)} + \omega_2'' U_{,\hat{t}}^{(10)}) - 2\{U_{,\hat{t}}^{(30)} + (\omega_1' U_{,\hat{t}}^{(20)} + \omega_2' U_{,\hat{t}}^{(10)})\} \\ - \alpha(U^{(10)} U^{(21)} + U^{(11)} U^{(20)}) + 3\beta(U^{(10)})^2 (U^{(11)}) \end{aligned}$$

Substituting on the RHS of (3.24) gives

$$\begin{aligned} U_{,\hat{t}\hat{t}}^{(31)} + U_{,xxxx}^{(31)} + 2\lambda U_{,xx}^{(31)} + U^{(31)} = - \left[(\omega_1')^2 U_{m,\hat{t}\hat{t}}^{(11)} (1 - \cos 2mx) + 2\{\omega_1' (U_{m,\hat{t}\tau}^{(21)} (1 - \cos 2mx) + \right. \\ \left. U_{2m,\hat{t}\tau}^{(21)} (1 - \cos 4mx) + \omega_2' U_{2m,\hat{t}\tau}^{(11)} (1 - \cos 2mx)\} + 2\{U_{m,\hat{t}\tau}^{(30)} (1 - \cos 2mx) + U_{2m,\hat{t}\tau}^{(30)} (1 - \cos 4mx) + \right. \\ \left. U_{3m,\hat{t}\tau}^{(30)} (1 - \cos 6mx)\} - 2\{\omega_1' (U_{m,\hat{t}\hat{t}}^{(20)} (1 - \cos 2mx) + U_{2m,\hat{t}\hat{t}}^{(20)} (1 - \cos 4mx)) + \omega_2' U_{m,\hat{t}\hat{t}}^{(10)} (1 - \right. \\ \left. \cos 2mx)\} + \{\omega_1'' U_{m,\hat{t}}^{(20)} (1 - \cos 2mx) + U_{2m,\hat{t}}^{(20)} (1 - \cos 4mx) + \omega_2'' U_{m,\hat{t}}^{(10)} (1 - \cos 2mx)\} + 2\{U_{m,\hat{t}}^{(30)} (1 - \right. \\ \left. \cos 2mx) + U_{2m,\hat{t}}^{(30)} (1 - \cos 4mx) + U_{3m,\hat{t}}^{(30)} (1 - \cos 6mx) + \omega_1' (U_{m,\hat{t}}^{(20)} (1 - \cos 2mx) + U_{2m,\hat{t}}^{(20)} (1 - \right. \\ \left. \cos 4mx) + \omega_2' U_{m,\hat{t}}^{(10)} (1 - \cos 2mx))\} + \alpha\{U_m^{(10)} (1 - \cos 2mx) (U_m^{(21)} (1 - \cos 2mx) + U_{2m}^{(21)} (1 - \right. \\ \left. \cos 4mx)) + U_m^{(11)} (1 - \cos 2mx) (U_m^{(20)} (1 - \cos 2mx) + U_{2m}^{(20)} (1 - \cos 4mx))\} - 3\beta\{(U_m^{(10)})^2 U_m^{(11)} (1 - \right. \\ \left. \cos 2mx)^3\} \right] \end{aligned} \quad (4.54)$$

Further simplification of (4.54) yields

$$\begin{aligned} U_{,\hat{t}\hat{t}}^{(31)} + U_{,xxxx}^{(31)} + 2\lambda U^{(31)} + U^{(31)} = - \left[(\omega_1')^2 U_{m,\hat{t}\hat{t}}^{(21)} (1 - \cos 2mx) + 2\{\omega_1' (U_{m,\hat{t}\tau}^{(21)} (1 - \cos 2mx) + \right. \\ \left. U_{2m,\hat{t}\tau}^{(21)} (1 - \cos 4mx)) + \omega_2' U_{m,\hat{t}\tau}^{(11)} (1 - \cos 2mx)\} + 2\{U_{m,\hat{t}\tau}^{(30)} (1 - \cos 2mx) + U_{2m,\hat{t}\tau}^{(30)} (1 - \cos 4mx) + \right. \\ \left. U_{3m,\hat{t}\tau}^{(30)} (1 - \cos 6mx)\} - 2\{\omega_1' U_{m,\hat{t}\hat{t}}^{(20)} (1 - \cos 2mx) + U_{2m,\hat{t}\hat{t}}^{(20)} (1 - \cos 4mx) + \omega_2' U_{m,\hat{t}\hat{t}}^{(10)} (1 - \cos 2mx)\} + \right. \\ \left. \{\omega_1'' U_{m,\hat{t}}^{(20)} (1 - \cos 2mx) + U_{2m,\hat{t}}^{(20)} (1 - \cos 4mx) + \omega_2'' U_{m,\hat{t}}^{(10)} (1 - \cos 2mx)\} \right] \end{aligned}$$

$$\begin{aligned}
& +2 \left\{ U_{m,\hat{t}}^{(30)}(1 - \cos 2mx) + U_{2m,\hat{t}}^{(30)}(1 - \cos 4mx) + U_{3m,\hat{t}}^{(30)}(1 - \cos 6mx) + \omega'_1 \left(U_{m,\hat{t}}^{(20)}(1 - \cos 2mx) + \right. \right. \\
& U_{2m,\hat{t}}^{(20)}(1 - \cos 4mx) \left. \right) + \omega'_2 \left(U_{m,\hat{t}}^{(10)}(1 - \cos 2mx) \right) \left. \right\} + \alpha \left\{ U_m^{(10)} U_m^{(21)} \left(\frac{3}{2} - 2\cos 2mx + \frac{1}{2}\cos 4mx \right) + \right. \\
& U_m^{(10)} U_{2m}^{(21)} \left(1 - \frac{1}{2}\cos 2mx - \cos 4mx + \frac{1}{2}\cos 6mx \right) \left. \right\} + \alpha \left\{ U_m^{(11)} U_m^{(20)} \left(\frac{3}{2} - 2\cos 2mx + \frac{1}{2}\cos 4mx \right) + \right. \\
& U_m^{(11)} U_{2m}^{(20)} \left(1 - \frac{1}{2}\cos 2mx - \cos 4mx + \frac{1}{2}\cos 6mx \right) \left. \right\} - 3\beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \left(\frac{5}{2} - \frac{15}{4}\cos 2mx - \frac{3}{2}\cos 4mx - \right. \\
& \left. \frac{1}{4}\cos 6mx \right) \left. \right] \quad (4.55)
\end{aligned}$$

Let

$$U^{(31)} \sum_{n=1}^{\infty} U^{(31)}(1 - \cos 2nx)$$

The LHS of (4.55) becomes

$$\sum_{n=1}^{\infty} \left[U_{n,\hat{t}\hat{t}}^{(31)}(1 - \cos 2nx) + (-16n^4 + 8\lambda n^2 + 1)U_n^{(31)} \cos 2nx \right]$$

Multiplying (4.55) through $\cos 2mx$ and integrating from 0 to π and from $n=m$, gives

$$\begin{aligned}
& -\frac{\pi}{2} \left[U_{m,\hat{t}\hat{t}}^{(31)} + (16m^4 - 8\lambda m^2 + 1)U_m^{(31)} \right] = \\
& - \left[(\omega'_1)^2 U_{m,\hat{t}\hat{t}}^{(11)} \left(-\frac{\pi}{2} \right) + 2 \left\{ \omega'_1 U_{m,\hat{t}\tau}^{(21)} \left(-\frac{\pi}{2} \right) + \omega'_2 U_{m,\hat{t}\tau}^{(11)} \left(-\frac{\pi}{2} \right) \right\} + 2 \left\{ U_{m,\hat{t}\tau}^{(30)} \left(-\frac{\pi}{2} \right) \right\} - \right. \\
& 2 \left\{ \begin{array}{l} \omega'_1 U_{m,\hat{t}\hat{t}}^{(20)} \\ + \omega'_2 U_{m,\hat{t}\hat{t}}^{(10)} \left(-\frac{\pi}{2} \right) \end{array} \right\} + \left\{ \omega'_1 U_{m,\hat{t}\tau}^{(20)} \left(-\frac{\pi}{2} \right) + \omega'_2 U_{m,\hat{t}\tau}^{(10)} \left(-\frac{\pi}{2} \right) \right\} + 2 \left\{ U_{m,\hat{t}\tau}^{(30)} \left(-\frac{\pi}{2} \right) + \omega'_1 U_{m,\hat{t}\tau}^{(20)} \left(-\frac{\pi}{2} \right) + \right. \\
& \left. \omega'_2 U_{m,\hat{t}\tau}^{(10)} \left(-\frac{\pi}{2} \right) \right\} + \\
& \alpha \left\{ -2U_m^{(10)} U_m^{(21)} \left(-\frac{\pi}{2} \right) - U_m^{(10)} U_m^{(21)} \left(-\frac{\pi}{2} \right) - 2U_m^{(11)} U_m^{(21)} \left(-\frac{\pi}{2} \right) - 2U_m^{(11)} U_m^{(20)} \left(-\frac{\pi}{2} \right) - \right. \\
& \left. U_m^{(11)} U_{2m}^{(20)} \left(-\frac{\pi}{2} \right) + 3\beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \left(-\frac{15}{4} \right) \right] \quad (4.56)
\end{aligned}$$

A further simplification of (4.56) yields

$$\begin{aligned}
& U_{m,\hat{t}\hat{t}}^{(31)} + \theta^2 U_m^{(31)} = (\omega'_1)^2 U_{m,\hat{t}\hat{t}}^{(11)} - 2 \left\{ \omega'_1 U_{m,\hat{t}\tau}^{(21)} + \omega'_2 U_{m,\hat{t}\tau}^{(11)} \right\} - 2 \left\{ U_{m,\hat{t}\tau}^{(30)} \right\} + 2 \left\{ \omega'_1 U_{m,\hat{t}\hat{t}}^{(20)} + \omega'_2 U_{m,\hat{t}\hat{t}}^{(10)} \right\} - \\
& \left\{ \omega'_1 U_{m,\hat{t}\tau}^{(20)} + \omega'_2 U_{m,\hat{t}\tau}^{(10)} \right\} - 2 \left\{ U_{m,\hat{t}\tau}^{(30)} + \omega'_1 U_{m,\hat{t}\tau}^{(20)} + \omega'_2 U_{m,\hat{t}\tau}^{(10)} \right\} + \alpha \left\{ 2U_m^{(10)} U_m^{(21)} + U_m^{(10)} U_m^{(21)} + 2U_m^{(11)} U_m^{(21)} + \right. \\
& \left. U_m^{(11)} U_{2m}^{(20)} \right\} - \frac{45}{4} \beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \quad (4.57)
\end{aligned}$$

The initial conditions for (4.57) are

$$\begin{aligned}
& U_m^{(31)}(0,0) = 0; \\
& U_{m,\hat{t}}^{(31)}(0,0) + \omega'_1(0)U_{m,\hat{t}}^{(21)}(0,0) + \omega'_2(0)U_{m,\hat{t}}^{(11)}(0,0) + U_{m,\tau}^{(30)}(0,0) = 0
\end{aligned}$$

Multiplying (4.55) through by $\cos 4mx$ and integrating from 0 to π and for $n=2m$ gives

$$\begin{aligned}
& -\frac{\pi}{2} \left[U_{m,\hat{t}\hat{t}}^{(31)} + (256m^4 - 32\lambda m^2 + 1)U_{2m}^{(30)} \right] \\
& = - \left[2\omega'_1 U_{2m,\hat{t}\tau}^{(21)} \left(-\frac{\pi}{2} \right) + 2U_{2m,\hat{t}\tau}^{(30)} \left(-\frac{\pi}{2} \right) - 2\omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} \left(-\frac{\pi}{2} \right) + \omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} \left(-\frac{\pi}{2} \right) \right. \\
& + 2U_{2m,\hat{t}\hat{t}}^{(30)} \left(-\frac{\pi}{2} \right) + 2\omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} \left(-\frac{\pi}{2} \right) \\
& + \alpha \left\{ \frac{1}{2} U_m^{(10)} U_m^{(21)} \left(-\frac{\pi}{2} \right) - U_m^{(10)} U_{2m}^{(21)} \left(-\frac{\pi}{2} \right) + \frac{1}{2} U_m^{(11)} U_m^{(20)} \left(-\frac{\pi}{2} \right) \right. \\
& \left. - U_m^{(11)} U_{2m}^{(20)} \left(-\frac{\pi}{2} \right) \right\} - 3\beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \left(\frac{3}{2} \right) \left. \right] \quad (4.58)
\end{aligned}$$

$$\begin{aligned}
& \Rightarrow U_{2m,\hat{t}\hat{t}}^{(31)} + \varphi^2 U_{2m}^{(30)} = - \left[2\omega'_1 U_{2m,\hat{t}\tau}^{(21)} + 2U_{2m,\hat{t}\tau}^{(30)} - 2\omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} + \omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} + 2U_{2m,\hat{t}\hat{t}}^{(30)} + 2\omega'_1 U_{2m,\hat{t}\hat{t}}^{(20)} + \right. \\
& \left. \alpha \left\{ \frac{1}{2} U_m^{(10)} U_m^{(21)} - U_m^{(10)} U_{2m}^{(21)} + \frac{1}{2} U_m^{(11)} U_m^{(20)} - U_m^{(11)} U_{2m}^{(20)} \right\} - \frac{9}{4} \beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \right] \quad (4.59)
\end{aligned}$$

The initial conditions are

$$U_{2m}^{(31)}(0,0) = 0; U_{2m}^{(31)}(0,0) + \omega_1'(0)U_{2m,\hat{t}}^{(20)}(0,0) = 0$$

Multiplying (4.56) through by $\cos 6m\hat{t}$ and integrating from 0 to π and for $n=3m$, the result is

$$\begin{aligned} & -\frac{\pi}{2} \left[U_{m,\hat{t}\hat{t}}^{(31)} + (1296m^4 - 72\lambda m^2 + 1)U_{2m}^{(31)} \right] = \\ & - \left[2U_{3m,\hat{t}\tau}^{(30)} \left(-\frac{\pi}{2} \right) + 2U_{3m,\hat{t}}^{(30)} \left(-\frac{\pi}{2} \right) + \alpha \left\{ \frac{1}{2}U_m^{(10)}U_{2m}^{(21)} \left(-\frac{\pi}{2} \right) + \frac{1}{2}U_m^{(11)}U_{2m}^{(20)} \left(-\frac{\pi}{2} \right) \right\} - \right. \\ & \left. 3\beta \left(-\frac{1}{4} \right) \left(U_m^{(10)} \right)^2 U_m^{(11)} \right] \quad (4.60) \end{aligned}$$

A further simplification of (4.60) yields

$$\begin{aligned} & U_{3m,\hat{t}\hat{t}}^{(31)} + \Omega^2 U_{3m}^{(30)} = \\ & - \left[2U_{3m,\hat{t}\tau}^{(30)} + 2U_{3m,\hat{t}}^{(30)} + \alpha \left\{ \frac{1}{2}U_m^{(10)}U_{2m}^{(21)} + \frac{1}{2}U_m^{(11)}U_{2m}^{(20)} \right\} + \frac{3}{4}\beta \left(U_m^{(10)} \right)^2 U_m^{(11)} \right] \quad (4.61) \end{aligned}$$

The initial conditions for (4.61) are

$$U_{3m}^{(31)}(0,0) = 0; U_{3m,\hat{t}}^{(31)}(0,0) = 0$$

Further simplification of terms in (4.57) yields

$$\begin{aligned} & U_{m,\hat{t}\hat{t}}^{(31)} + \theta^2 U_m^{(31)} = (\omega_1')^2 \theta^2 \beta_2 \sin \theta \hat{t} - 2 \left[\omega_1' (-\theta \alpha_6' \sin \theta \hat{t} + \theta \beta_6' \cos \theta \hat{t}) - \frac{(-2\theta r_3' \sin 2\theta \hat{t} + 2\theta r_4' \cos 2\theta \hat{t})}{3\theta^2} \right] - \\ & 2\omega_2' \beta_2' \theta \cos \theta \hat{t} - 2 \left[-\alpha_9' \theta \sin \theta \hat{t} + \beta_9' \theta \cos \theta \hat{t} + \frac{2r_{11}' \sin 2\theta \hat{t}}{3\theta} + \frac{3r_{12}' \sin 3\theta \hat{t}}{8\theta} - \frac{1}{\varphi(2\theta + \varphi)} \{ -r_{13}'(\varphi + \theta) \sin(\varphi + \right. \\ & \left. \theta) \hat{t} + r_{14}'(\varphi + \theta) \cos(\varphi + \theta) \hat{t} \} + \frac{1}{\varphi(2\theta - \varphi)} \{ -r_{15}'(\varphi - \theta) \sin(\varphi - \theta) \hat{t} + r_{16}'(\varphi - \theta) \cos(\varphi - \theta) \hat{t} \} \right] + \\ & 2\omega_1' \left\{ -\theta^2 \alpha_4 \cos \theta \hat{t} + \frac{4r_1 \cos 2\theta \hat{t}}{3} \right\} + 2\omega_2' (-\alpha_1 \theta^2 \cos \theta \hat{t}) - \omega_1' \left\{ -\theta \alpha_4 \sin \theta \hat{t} + \frac{2r_1 \sin 2\theta \hat{t}}{3\theta} \right\} - \\ & \omega_2'' (-\alpha_1 \theta \sin \theta \hat{t}) - 2 \left\{ -\alpha_9 \sin \theta \hat{t} + \beta_9 \cos \theta \hat{t} + \frac{2r_{11} \sin 2\theta \hat{t}}{3\theta} + \frac{3r_{12} \sin 3\theta \hat{t}}{8\theta} - \frac{1}{\varphi(2\theta + \varphi)} \{ -(\varphi + \theta) r_{13} \sin(\varphi + \right. \\ & \left. \theta) \hat{t} + r_{14}(\varphi + \theta) \cos(\varphi + \theta) \hat{t} \} + \frac{1}{\varphi(2\theta - \varphi)} \{ -(\varphi + \theta) r_{15} \sin(\varphi - \theta) \hat{t} + (\varphi - \theta) r_{16} \cos(\varphi - \theta) \hat{t} \} \right\} - \\ & 2\omega_1' \left\{ -\theta \alpha_4 \sin \theta \hat{t} + \frac{2r_1 \sin 2\theta \hat{t}}{3\theta} \right\} - 2\omega_2' (-\theta \alpha_1 \sin \theta \hat{t}) + 2\alpha \left\{ \left(\frac{\alpha_1 \alpha_6}{2} - \frac{Br_2}{\theta^2} \right) + \left(\frac{\alpha_1 r_2}{2} - \frac{Br_3}{6\theta^2} + B\alpha_6 \right) \cos \theta \hat{t} + \right. \\ & \left. \left(\frac{B\beta_6 - \alpha_1 \alpha_4}{6\theta^2} \right) \sin \theta \hat{t} + \left(\frac{\alpha_1 \alpha_6}{2} - \frac{Br_3}{3\theta^2} \right) \cos 2\theta \hat{t} + \left(\frac{\alpha_1 \beta_6}{2} - \frac{Br_4}{3\theta^2} \right) \sin \theta \hat{t} - \frac{\alpha_1 r_3}{6\theta^2} \cos 3\theta \hat{t} - \frac{\alpha_1 r_4}{6\theta^2} \sin 3\theta \hat{t} \right\} + \\ & \alpha \left\{ \left(\frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} + \frac{Br_5}{\varphi^2 - \theta^2} \right) \sin \theta \hat{t} + \left(\frac{\alpha_1 r_5}{2(\varphi^2 - 4\theta^2)} - \frac{Br_6}{\varphi^2 - 4\theta^2} \right) \sin 2\theta \hat{t} + \frac{\alpha_1 r_6 \sin 3\theta \hat{t}}{2(\varphi^2 - 4\theta^2)} + B\alpha_7 \cos \varphi \hat{t} + B\beta_7 \sin \varphi \hat{t} + \right. \\ & \left. \frac{\alpha_1 \alpha_7}{2} \cos(\varphi + \theta) \hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi + \theta) \hat{t} + \frac{\alpha_1 \alpha_7}{2} \cos(\varphi - \theta) \hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi - \theta) \hat{t} \right\} + 2\alpha \left\{ \frac{\beta_2 \alpha_4}{2} \sin 2\theta \hat{t} + \right. \\ & \left. \frac{\beta_2 r_0}{\theta^2} \sin \theta \hat{t} - \frac{\beta_2 r_1}{6\theta^2} (\sin 3\theta \hat{t} - \sin \theta \hat{t}) \right\} \quad (4.62) \end{aligned}$$

To ensure a uniformly valid solution in \hat{t} , demands equating to zero the coefficients of $\sin \theta \hat{t}$ and $\cos \theta \hat{t}$ in (4.62) as further expanded. The coefficient of $\sin \theta \hat{t}$ leads to

$$\alpha_9' + \alpha_9 = h_1(\tau) \quad (4.63a)$$

$$h_1(\tau) = -\frac{1}{2\theta} \left[2\omega_1' \theta \alpha_6' + \omega_2'' \alpha_1 \theta + \omega_1' \theta \alpha_4 + 2\omega_1' \theta \alpha_4 + 2\omega_2' \theta \alpha_1 + 2\alpha \left(\frac{B\beta_6 - \alpha_1 r_4}{6\theta^2} \right) \right] \quad (4.63)$$

$$\therefore \alpha_9 = e^{-\tau} \left[\int e^s h_1(s) ds + \alpha_9(0) \right] \quad (4.64)$$

The coefficient of $\cos \theta \hat{t}$ yields

$$\beta_9' + \beta_9 = h_2(\tau) \quad (4.65)$$

$$h_2(\tau) = -\frac{1}{2\theta} \left[2\omega_1' \theta \beta_6' + 2\omega_2' \beta_2' \theta + 2\omega_1' \theta^2 \alpha_4 + 2\omega_2' \alpha_1 \theta^2 - 2\alpha \left(\frac{\alpha_1 r_2}{\theta^2} - \frac{\alpha_1 r_3}{6\theta^2} + B\alpha_6 \right) \right] \quad (4.66)$$

$$\therefore \beta_9 = e^{-\tau} \left[\int e^s h_2(s) ds + \beta_9(0) \right] \quad (6.67)$$

The remaining equation in (4.63)

$$\begin{aligned} & U_{m,\hat{t}\hat{t}}^{(31)} + \theta^2 U_m^{(31)} = r_{29} + r_{30} \sin 2\theta \hat{t} + r_{31} \cos 2\theta \hat{t} + r_{32} \cos 3\theta \hat{t} + r_{33} \cos 3\theta \hat{t} + r_{34} \cos \varphi \hat{t} + r_{35} \sin \varphi \hat{t} \\ & + r_{36} \cos(\varphi + \theta) \hat{t} + r_{37} \sin(\varphi + \theta) \hat{t} + r_{38} \cos(\varphi - \theta) \hat{t} \\ & + r_{39} \sin(\varphi - \theta) \hat{t} \quad (4.68) \end{aligned}$$

The initial conditions are

$$U_m^{(31)}(0,0) = 0; U_{m,\hat{t}}^{(31)}(0,0) + \omega_1'(0)U_{m,\hat{t}}^{(21)}(0,0) + \omega_2'(0)U_{m,\hat{t}}^{(11)}(0,0) + U_{m,\tau}^{(30)}(0,0) = 0$$

where,

$$r_{29} = 2\alpha \left(\frac{\alpha_1 \alpha_6}{2} + \frac{r_2 B}{\theta^2} \right); r_{29}(0) = 0$$

$$\begin{aligned}
r_{30} &= \left[\frac{-4r'_3}{3\theta} - \frac{4r'_1}{3\theta} - \frac{4r'_{11}}{3\theta} - \frac{2\omega'_1 r_1}{3\theta} - \frac{4r_{11}}{3\theta} - \frac{4\omega'_1 r_1}{3\theta} + 2\alpha \left(\frac{\alpha_1 \beta_6}{2} - \frac{r_4 B}{3\theta^2} \right) + \alpha \left(\frac{\alpha_1 r_5}{2(\varphi^2 - \theta^2)} + \frac{Br_6}{\varphi^2 - 4\theta^2} \right) \right. \\
&\quad \left. + \alpha \beta_2 \alpha_4 + \frac{\alpha^2 \alpha_1 B \beta_2}{2(\varphi^2 - \theta^2)} - \frac{45}{4} \beta B \alpha_1 \beta_2 \right] \\
r_{30}(0) &= B^3 \left(\frac{-8\alpha}{3\theta B} - \frac{4S_{21}}{3\theta} + \frac{2\alpha}{3\theta^3} - \frac{4S_4}{3\theta} + \frac{4\alpha}{3\theta^3} - \frac{2\alpha^2}{3\theta^3} + \frac{\alpha^2}{2\theta(\varphi^2 - \theta^2)} + \frac{\alpha^2}{2\theta B(\varphi^2 - \theta^2)} - \frac{45}{4\theta} \right) = B^3 S_{25}, \\
S_{25} &= \left(-\frac{8\alpha}{3B\theta} - \frac{4S_{21}}{2\theta} + \frac{2\alpha}{\theta^3} - \frac{4S_4}{3\theta} - \frac{2\alpha^2}{\theta^3} + \frac{\alpha^2}{2\theta(\varphi^2 - \theta^2)} + \frac{S_1}{(\varphi^2 - \theta^2)} + \frac{\alpha^2}{2\theta B(\varphi^2 - \theta^2)} - \frac{45\beta}{4\theta} \right) \\
r_{31} &= \left[\frac{8\omega'_1 r_1}{3} + 2\alpha \left(\frac{\alpha_1 \alpha_6}{2} - \frac{Br_3}{3\theta^2} \right) \right], r_{31}(0) = \frac{-8\alpha B^3}{3\theta^2} \\
r_{32} &= \left[-\frac{\alpha \alpha_1 r_3}{3\theta^2} \right], r_{32}(0) = 0 \\
r_{33} &= \left[\frac{-3r'_{12}}{4\theta} - \frac{3r_{12}}{4\theta} - \frac{\alpha \alpha_1 r_4}{3\theta^2} + \frac{\alpha \alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} - \frac{\alpha \beta_2 r_1}{3\theta^2} + \frac{\alpha^2 \alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} \right. \\
&\quad \left. - \frac{45}{16} \beta B \alpha_1 \beta_2 \right] r_{33}(0) \\
&= B^3 \left(-\frac{3S_{61}}{4\theta} - \frac{3S_5}{3\theta} + \frac{\alpha^2}{3\theta^3} - \frac{\alpha S_1}{2\theta(\varphi^2 - \theta^2)} \right) = B^3 S_{26} \\
S_{26} &= \left(-\frac{3S_{16}}{3\theta} - \frac{3S_5}{3\theta} + \frac{\alpha^2}{3\theta^3} - \frac{\alpha S_1}{2(\varphi^2 - 4\theta^2)} \right)
\end{aligned}$$

$$\begin{aligned}
r_{34} &= [\alpha B \alpha_7], r_{34}(0) = 0, \\
r_{35} &= [\alpha B \beta_7], r_{35}(0) = \frac{\alpha^2}{3\varphi^3} + \frac{\alpha^2 S_0}{\varphi} + \frac{\alpha^2}{2\varphi(\varphi^2 - 4\theta^2)} - \frac{\alpha^2}{\varphi(\varphi^2 - 4\theta^2)} \\
r_{36} &= \left[\frac{2r'_{14}(\varphi + \theta)}{\varphi(2\theta + \varphi)} - \frac{2r_{14}(\varphi + \theta)}{\varphi(2\theta + \varphi)} + \frac{\alpha \alpha_1 \alpha_7}{2} - \frac{\alpha \beta_2 \beta_5}{2} \right], r_{36}(0) = 0 \\
r_{37} &= \left[-\frac{2r'_{13}(\varphi + \theta)}{\varphi(2\theta + \varphi)} + \frac{2r_{13}(\varphi + \theta)}{\varphi(2\theta + \varphi)} + \frac{\alpha \alpha_1 \beta_7}{2} + \alpha \left(\frac{\beta_2 \alpha_5}{2} \right) \right] \\
r_{37}(0) &= B^3 \left(\frac{6\alpha(\varphi + \theta)S_0}{\varphi(2\theta + \varphi)} + \frac{\alpha S_{23}}{2} - \frac{S_0 \alpha}{2\theta} \right) = B^3 S_{27} \\
S_{27} &= \left(\frac{6\alpha(\varphi + \theta)S_0}{\varphi(2\theta + \varphi)} - \frac{\alpha S_{43}}{2} - \frac{\alpha S_0}{2\theta} \right) \\
r_{38} &= \left[\frac{-2r'_{16}(\varphi - \theta)}{\varphi(2\theta - \varphi)} + \frac{2r_{16}(\varphi - \theta)}{\varphi(2\theta - \varphi)} + \frac{\alpha \alpha_1 \alpha_7}{2} + \frac{\alpha \beta_2 \beta_5}{2} \right], r_{38}(0) = 0 \\
r_{39} &= \left[\frac{2r'_{15}(\varphi - \theta)}{\varphi(2\theta - \varphi)} + \frac{2r_{15}(\varphi - \theta)}{\varphi(2\theta - \varphi)} + \frac{\alpha \alpha_1 \beta_7}{2} - \alpha \left(\frac{\beta_2 \alpha_5}{2} \right) \right] \\
r_{39}(0) &= B^3 \left(\frac{4\alpha(\varphi - \theta)S_0}{\varphi(2\theta + \varphi)} - \frac{2\alpha(\varphi - \theta)S_0}{\varphi(2\theta - \varphi)} - \frac{\alpha^2 S_3}{2\varphi} + \frac{\alpha S_0}{2\theta} \right) = B^3 S_{29} \\
S_{29} &= \left(-\frac{3\alpha}{\theta^2(\varphi^2 - \theta^2)} - \frac{2\alpha}{(\varphi^2 - \theta^2)} \right)
\end{aligned}$$

Solving (4.69), the following is obtained

$$\begin{aligned}
U_m^{(31)} &= \alpha_{12} \cos \theta \hat{t} + \beta_{12} \sin \theta \hat{t} + \frac{r_{29}}{\theta^2} + \frac{r_{30} \sin 2\theta \hat{t} + r_{31} \cos 2\theta \hat{t}}{\theta^2 - 4\theta^2} + \frac{r_{32} \cos 3\theta \hat{t} + r_{33} \sin 3\theta \hat{t}}{\theta^2 - 9\theta^2} + \frac{r_{34} \cos \varphi \hat{t} + r_{35} \sin \varphi \hat{t}}{\theta^2 - \varphi^2} + \\
&\quad \frac{r_{36} \cos(\varphi + \theta) \hat{t} + r_{37} \sin(\varphi + \theta) \hat{t}}{\varphi(2\theta - \varphi)} + \frac{r_{38} \cos(\varphi - \theta) \hat{t} + r_{39} \sin(\varphi - \theta) \hat{t}}{\varphi(2\theta - \varphi)} \quad (4.69)
\end{aligned}$$

$$\begin{aligned}
\alpha_{12}(0) &= - \left[\frac{r_{29}}{\theta^2} + \frac{r_{31}}{\theta^2 - 4\theta^2} + \frac{r_{32}}{\theta^2 - 9\theta^2} + \frac{r_{34}}{\theta^2 - \varphi^2} - \frac{r_{36}}{\varphi(2\theta - \varphi)} + \frac{r_{38}}{\varphi(2\theta - \varphi)} - \frac{2\alpha B^3}{3\theta^4} + \frac{1}{2\theta^2} \left(\frac{B^2}{\theta^4} + \frac{16\alpha B^3}{3\theta^2} - \frac{10\alpha^2 B^3}{\theta^2} + \right. \right. \\
&\quad \left. \left. \frac{3\alpha^2 B^3}{2\varphi^2} + \frac{\alpha^2 B^3}{4(\varphi^2 - 4\theta^2)} + \frac{225\beta B^3}{16} \right) + \left(\alpha'_9 + \frac{r'_{10}}{\theta^2} - \frac{r'_{11}}{3\theta^2} - \frac{r'_{12}}{8\theta^2} - \frac{r'_{13}}{\varphi(2\theta + \varphi)} + \frac{r'_{15}}{\varphi(2\theta - \varphi)} \right) \right] \tau = \\
&= 0 \quad (4.70a)
\end{aligned}$$

$$\begin{aligned}
\beta_{12}(0) &= \frac{-1}{\theta} \left[\frac{2\theta r_{30}}{\theta^2 - 4\theta^2} + \frac{3\theta r_{33}}{\theta^2 - 9\theta^2} + \frac{\varphi r_{35}}{\theta^2 - \varphi^2} + \frac{(\varphi + \theta)r_{37}}{\varphi(2\theta - \varphi)} + \frac{(\varphi - \theta)r_{39}}{\varphi(2\theta - \varphi)} \right] \text{ at } \tau \\
&= 0 \quad (4.70b)
\end{aligned}$$

Substituting in (4.59) gives

$$\begin{aligned}
U_{2m,\hat{t}\hat{t}}^{(31)} + \varphi^2 U_{2m}^{(31)} = & - \left[2\omega'_1 \left\{ -\varphi\alpha'_7 \sin\varphi\hat{t} + \varphi\beta'_7 \cos\varphi\hat{t} + \frac{\theta r'_5 \cos\theta\hat{t}}{\varphi^2 - \theta^2} + \frac{2\theta r'_6 \cos 2\theta\hat{t}}{\varphi^2 - 4\theta^2} \right\} + 2 \left\{ -\varphi\alpha'_{10} \sin\varphi\hat{t} + \right. \\
& \varphi\beta'_{10} \cos\varphi\hat{t} - \frac{\theta r'_{18} \sin\theta\hat{t}}{\varphi^2 - \theta^2} - \frac{2\theta r'_{19} \sin 2\theta\hat{t}}{\varphi^2 - 4\theta^2} - \frac{3\theta r'_{20} \sin 3\theta\hat{t}}{\varphi^2 - 9\theta^2} \left. \right\} - \alpha\omega'_1 \left\{ \frac{2B\alpha_1 \theta^2 \cos\theta\hat{t}}{\varphi^2 - \theta^2} - \frac{2\theta\alpha_1^2 \cos 2\theta\hat{t}}{\varphi^2 - 4\theta^2} \right\} + \alpha(\omega'_1 + \\
& 2\omega'_1) \left\{ \frac{B\alpha_1 \theta \sin\theta\hat{t}}{\varphi^2 - \theta^2} - \frac{\theta\alpha_1^2 \sin 2\theta\hat{t}}{\varphi^2 - 4\theta^2} \right\} + 2 \left\{ -\varphi\alpha_{10} \sin\varphi\hat{t} + \varphi\beta_{10} \cos\varphi\hat{t} - \frac{\theta r_{18} \sin\theta\hat{t}}{\varphi^2 - \theta^2} - \frac{2\theta r_{19} \sin 2\theta\hat{t}}{\varphi^2 - 4\theta^2} - \frac{3\theta r_{20} \sin 3\theta\hat{t}}{\varphi^2 - 9\theta^2} \right\} + \\
& \frac{\alpha}{2} \left\{ \left(\frac{\alpha_1 \alpha_6}{2} + \frac{r_2 B}{\theta^2} \right) + \left(\frac{\alpha_1 r_2}{\theta^2} - \frac{\alpha_1 r_3}{6\theta^2} + B\alpha_6 \right) \cos\theta\hat{t} + \left(B\beta_6 - \frac{\alpha_1 r_4}{6\theta^2} \right) \sin\theta\hat{t} + \left(\frac{\alpha_1 \beta_6}{2} - \frac{r_4 B}{3\theta^2} \right) \sin 2\theta\hat{t} - \right. \\
& \frac{\alpha_1 r_3}{6\theta^2} \cos 3\theta\hat{t} - \frac{\alpha_1 r_4}{6\theta^2} \sin 3\theta\hat{t} \left. \right\} - \alpha \left\{ \left(\frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} + \frac{B r_5}{\varphi^2 - \theta^2} \right) \sin\theta\hat{t} + \frac{\alpha_1 \alpha_7}{2} \cos(\varphi + \theta)\hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi + \theta)\hat{t} + \right. \\
& \frac{\alpha_1 \alpha_7}{2} \cos(\varphi - \theta)\hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi - \theta)\hat{t} + \left. \left(\frac{\alpha_1 r_5}{2(\varphi^2 - \theta^2)} + \frac{B r_6}{\varphi^2 - 4\theta^2} \right) \sin 2\theta\hat{t} + \frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} \sin 3\theta\hat{t} + B\alpha_7 \cos\varphi\hat{t} + \right. \\
& B\beta_7 \sin\varphi\hat{t} \left. \right\} + \frac{\alpha}{2} \left\{ \frac{\beta_2 \alpha_4 \sin 2\theta\hat{t}}{2} + \frac{\beta_2 r_0 \sin\theta\hat{t}}{\theta^2} - \frac{\beta_2 r_1}{6\theta^2} (\sin 3\theta\hat{t} - \sin\theta\hat{t}) \right\} - \alpha \left\{ \frac{\alpha\beta_2}{2} \left(\frac{\alpha_1^2 + B^2}{\varphi^2} \right) - \frac{\alpha\alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} \sin\theta\hat{t} + \right. \\
& \left. \frac{\alpha\alpha_1 B\beta_2}{2(\varphi^2 - \theta^2)} \sin 2\theta\hat{t} + \frac{\alpha\alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} \sin 3\theta\hat{t} \right\} + \\
& \left. \frac{9\beta}{4} \left[\left\{ \beta_2 \left(B^2 + \frac{\alpha_1^2}{2} \right) - \frac{\beta_2 \alpha_1^2}{4} \right\} \sin\varphi\hat{t} + \beta_2 B\alpha_1 \sin 2\theta\hat{t} + \frac{\beta_2 \alpha_1^2}{4} \sin 3\theta\hat{t} \right] \right] \quad (4.71)
\end{aligned}$$

To ensure uniformly valid solution in \hat{t} needs equating the coefficients of $\cos\varphi\hat{t}$ and $\sin\varphi\hat{t}$ to zero. Equating the coefficient of $\cos\varphi\hat{t}$ yields

$$\begin{aligned}
-2\omega'_1 \varphi \beta'_7 - 2\varphi \beta'_{10} - 2\beta_{10} \varphi + B\alpha_1 \alpha_7 &= 0 \\
\therefore \beta'_{10} + \beta_{10} &= \frac{1}{2\varphi} [-2\omega'_1 \varphi \beta'_7 + B\alpha_1 \alpha_7] \quad (4.72a)
\end{aligned}$$

$$\therefore \beta'_{10} + \beta_{10} = h_3(\tau) \quad (4.72b)$$

where,

$$h_3(\tau) = \frac{1}{2\varphi} [-2\omega'_1 \varphi \beta'_7 + B\alpha_1 \alpha_7] \quad (4.72c)$$

It therefore follows that,

$$\beta_{10} = e^{-\tau} \left[\int h_3(s) e^{s\tau} ds + \beta_{10}(0) \right] \quad (4.72d)$$

The coefficient of $\sin\varphi\hat{t}$ leads to

$$2\omega'_1 \varphi \alpha'_7 + 2\varphi \alpha'_{10} + 2\alpha_{10} \varphi + B\beta_7 \alpha = 0 \quad (4.72e)$$

$$\begin{aligned}
\alpha'_{10} + \alpha_{10} &= h_4(\tau) \\
h_4(\tau) &= -\frac{1}{2\varphi} [2\omega'_1 \varphi \alpha'_7 + B\beta_7 \alpha] \quad (4.72f)
\end{aligned}$$

$$\therefore \alpha_{10} = e^{-\tau} \left[\int h_4(s) e^{s\tau} ds + \alpha_{10}(0) \right] \quad (4.72g)$$

The remaining equation in (4.71) is

$$\begin{aligned}
U_{2m,\hat{t}\hat{t}}^{(31)} + \varphi^2 U_{2m}^{(31)} &= r_{40} \cos\theta\hat{t} + r_{41} \sin\theta\hat{t} + r_{42} \cos 2\theta\hat{t} + r_{43} \sin 2\theta\hat{t} + r_{44} \cos 3\theta\hat{t} + r_{45} \sin 3\theta\hat{t} + \\
& r_{46} \cos(\varphi + \theta)\hat{t} + r_{47} \sin(\varphi + \theta)\hat{t} + r_{48} \cos(\varphi - \theta)\hat{t} + r_{49} \sin(\varphi - \theta)\hat{t} \quad (4.73)
\end{aligned}$$

The initial conditions are

$$U_{2m}^{(31)}(0,0) = 0; U_{2m,\hat{t}}^{(31)}(0,0) + \omega'_1(0)U_{2m,\hat{t}}^{(21)}(0,0) + U_{2m,\tau}^{(30)}(0,0) = 0$$

where,

$$\begin{aligned}
r_{40} &= \frac{-2\theta r_5^1 \omega'_1}{\varphi^2 - \theta^2} + \frac{2\alpha\omega'_1 B\alpha_1 \theta^2}{\varphi^2 - \theta^2} - \frac{\alpha}{2} \left(\left(\frac{\alpha_1 \alpha_6}{2} + \frac{r_2 B}{\theta^2} \right) + \left(\frac{\alpha_1 r_2}{\theta^2} - \frac{\alpha_1 r_3}{6\theta^2} \right) + B\alpha_6 \right) \\
r_{40}(0) &= B^3 \left(\frac{-3\alpha}{\theta^2(\varphi^2 - \theta^2)} - \frac{2\alpha}{(\varphi^2 - \theta^2)} \right) \\
r_{41} &= \left[\frac{2\theta r_{18}^1}{\varphi^2 - \theta^2} + \frac{(\omega'_1 + 2\omega'_1)\alpha B\alpha_1 \theta}{\varphi^2 - \theta^2} + \frac{2\theta r_{18}}{\varphi^2 - \theta^2} - \frac{\alpha}{2} \left(B\beta_6 - \frac{\alpha_1 r_4}{6\theta^2} \right) + \alpha \left(\frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} + \frac{B r_5}{\varphi^2 - \theta^2} \right) \right. \\
& \left. - \frac{\alpha\beta_2 r_0}{2\theta^2} - \frac{\alpha\beta_2 r_1}{12\theta^2} + \left\{ \frac{\alpha^2 \beta_2}{2\varphi^2} \left(\frac{\alpha_1^2}{2} + B^2 \right) - \frac{\alpha^2 \alpha_1 B\beta_2}{8(\varphi^2 - 4\theta^2)} + \frac{9\beta}{4} \left(\beta_2 \left(B^2 + \frac{\alpha_1^2}{2} \right) \right) \right\} \right] \\
r_{41}(0) &= B^3 \left(\frac{2\theta S_{24}}{(\varphi^2 - \theta^2)} - \frac{2\alpha}{\theta(\varphi^2 - \theta^2)} + \frac{2\theta S_7}{\varphi^2 - \theta^2} - \frac{2\alpha^2}{9\theta^4} - \frac{\alpha S_1}{2(\varphi^2 - 4\theta^2)} + \frac{\alpha^2}{\theta(\varphi^2 - \theta^2)} - \frac{17\alpha^2}{12\theta^3} \right. \\
& \left. - \frac{3\alpha^2}{(4\theta - \varphi^2)} - \frac{\alpha^2}{8\theta(\varphi^2 - 4\theta^2)} - \frac{45\beta}{16\theta} \right)
\end{aligned}$$

$$\begin{aligned}
r_{42} &= \left[\frac{4\theta\omega'_1 r_6}{(\varphi^2 - 4\theta^2)} - \frac{2\theta\alpha\omega'_1 \alpha_1^2}{(\varphi^2 - 4\theta^2)} \right], r_{42}(0) = B^3 \left(\frac{2\alpha}{\theta(\varphi^2 - 4\theta^2)} - \frac{4S_{14}}{\theta(\varphi^2 - 4\theta^2)} \right) \\
r_{43} &= \left[\frac{4\theta\omega'_1 r_{19}}{(\varphi^2 - 4\theta^2)} + \frac{(\omega'_1 + 2\omega'_1)\alpha\theta\alpha_1^2}{\varphi^2 - 4\theta^2} + \frac{4\theta r_{19}}{(\varphi^2 - 4\theta^2)} - \frac{\alpha}{2} \left(\frac{\alpha_1\beta_6}{2} - \frac{Br_4}{3\theta^2} \right) + \alpha \left(\frac{\alpha_1 r_5}{2(\varphi^2 - \theta^2)} + \frac{Br_6}{\varphi^2 - 4\theta^2} \right) \right. \\
&\quad \left. + \frac{\alpha^2 \alpha_1 B \beta_2}{2(\varphi^2 - \theta^2)} + \frac{9}{4} \beta_2 B \alpha_1 \right] \\
r_{43}(0) &= B^3 \left(\frac{4\theta S_{10}}{(\varphi^2 - 4\theta^2)} + \frac{2\alpha}{\theta(\varphi^2 - \theta^2)} + \frac{4\theta S_8}{\varphi^2 - 4\theta^2} + \frac{\alpha^2}{6\theta^3} + \frac{2\alpha^2}{2\theta(\varphi^2 - \theta^2)} + \frac{\alpha S_1}{(\varphi^2 - \theta^2)} + \frac{9\beta}{4\theta} \right) \\
r_{44} &= \left[\frac{\alpha}{2} \left(\frac{\alpha_2 r_3}{6\theta^2} \right) \right], r_{44}(0) = 0 \\
r_{45} &= \left[\frac{6\theta r'_{20}}{(\varphi^2 - 9\theta^2)} + \frac{6\theta r_{20}}{(\varphi^2 - 9\theta^2)} + \frac{\alpha}{2} \left(\frac{\alpha_1 r_4}{6\theta^2} \right) + \frac{\alpha\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} + \frac{\alpha\beta_2 r_1}{12} + \frac{\alpha^2 \alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} + \frac{9}{16} \beta_2 \alpha_1^2 B \right] \\
r_{45}(0) &= B^3 \left(\frac{6\theta S_{34}}{(\varphi^2 - 4\theta^2)} + \frac{6\theta S_9}{(\varphi^2 - \theta^2)} - \frac{\alpha^2}{12\theta^3} - \frac{\alpha S_1}{2(\varphi^2 - \theta^2)} - \frac{\alpha^2}{12\theta} - \frac{9\beta}{16\theta} \right) \\
r_{46} &= \left[\frac{\alpha_1 \alpha_7}{2} \right], r_{46}(0) = 0, r_{47} = \left[\frac{\alpha\alpha_1 \beta_7}{2} \right], r_{47}(0) = -\alpha B^3 S_{43} \\
r_{48} &= \left[\frac{\alpha\alpha_1 \alpha_7}{2} \right], r_{48}(0) = 0, r_{49} = \left[\frac{\alpha\alpha_1 \beta_7}{2} \right], r_{49}(0) = -\alpha B^3 S_{43} \\
S_{43} &= \frac{\alpha S_0}{\varphi} + \frac{\alpha}{2\varphi^3} + \frac{\alpha}{2\alpha(\varphi^2 - 4\theta^2)} - \frac{\alpha}{\alpha(\varphi^2 - \theta^2)} - \frac{2\theta\alpha S_1}{\varphi(\varphi^2 - 4\theta^2)} \\
\therefore U_{2m}^{(31)} &= \alpha_{13} \cos\varphi\hat{t} + \beta_{13} \sin\varphi\hat{t} + \frac{r_{42} \cos\theta\hat{t} + r_{43} \sin\theta\hat{t}}{\varphi^2 - \theta^2} + \frac{r_{44} \cos 2\theta\hat{t} + r_{45} \sin 2\theta\hat{t}}{\varphi^2 - 4\theta^2} \\
&\quad + \frac{r_{46} \cos 3\theta\hat{t} + r_{47} \sin 3\theta\hat{t}}{\varphi^2 - 9\theta^2} - \frac{r_{48} \cos(\varphi + \theta)\hat{t} + r_{49} \sin(\varphi + \theta)\hat{t}}{\theta(2\varphi + \theta)} \\
&\quad + \frac{r_{50} \cos(\varphi - \theta)\hat{t} + r_{39} \sin(\varphi - \theta)\hat{t}}{\theta(2\varphi - \theta)} \quad (4.74)
\end{aligned}$$

where, from the first initial condition

$$\alpha_{13}(0) = - \left[\frac{r_{42}}{\varphi^2 - \theta^2} + \frac{r_{44}}{\varphi^2 - 4\theta^2} + \frac{r_{46}}{\varphi^2 - 9\theta^2} - \frac{r_{48}}{\theta(2\varphi + \theta)} + \frac{r_{50}}{\theta(2\varphi - \theta)} \right] \text{ at } \tau = 0 \quad (4.75)$$

and from the second initial condition, it follows that

$$\begin{aligned}
&\left[\beta_{13}(0)\varphi + \frac{\theta r_{43}}{\varphi^2 - \theta^2} + \frac{2\theta r_{45}}{\varphi^2 - 4\theta^2} + \frac{3\theta r_{47}}{\varphi^2 - 9\theta^2} - \frac{(\theta + \varphi)r_{49}}{\theta(2\varphi + \theta)} + \frac{(\theta - \varphi)r_{51}}{\theta(2\varphi - \theta)} + \alpha'_{10}(0) + \frac{r'_{17}}{\varphi^2} + \frac{r'_{18}}{\varphi^2 - \theta^2} + \frac{r'_{19}}{\varphi^2 - 4\theta^2} + \frac{r'_{20}}{\varphi^2 - 9\theta^2} \right] = \\
&0 \\
\therefore \beta_{13}(0) &= - \frac{1}{\varphi} \left[\frac{\theta r_{43}}{\varphi^2 - \theta^2} + \frac{2\theta r_{45}}{\varphi^2 - 4\theta^2} + \frac{3\theta r_{47}}{\varphi^2 - 9\theta^2} - \frac{(\theta + \varphi)r_{49}}{\theta(2\varphi + \theta)} + \frac{(\theta - \varphi)r_{51}}{\theta(2\varphi - \theta)} + \alpha'_{10}(0) + \frac{r'_{17}}{\varphi^2} + \frac{r'_{18}}{\varphi^2 - \theta^2} + \frac{r'_{19}}{\varphi^2 - 4\theta^2} + \frac{r'_{20}}{\varphi^2 - 9\theta^2} \right] \quad (4.75b)
\end{aligned}$$

Substituting in (4.61)

$$U_{3m,\hat{t}\hat{t}}^{(31)} + \Omega^2 U_{3m}^{(31)} = - \left[2U_{3m,\hat{t}\tau}^{(30)} + 2U_{3m,\hat{t}}^{(30)} + \alpha \left\{ \frac{1}{2} U_m^{(10)} U_{2m}^{(21)} + \frac{1}{2} U_m^{(11)} U_{2m}^{(20)} \right\} + \frac{3}{4} \beta (U_m^{(10)})^2 U_m^{(11)} \right] \quad (4.75c)$$

Further simplification of (4.75c) yields

$$\begin{aligned}
U_{3m,\hat{t}\hat{t}}^{(31)} + \Omega^2 U_{3m}^{(31)} &= - \left[2 \left\{ -\Omega\alpha'_{11} \sin\Omega\hat{t} + \Omega\beta'_{11} \cos\Omega\hat{t} - \frac{\theta r'_{22} \sin\theta\hat{t}}{\Omega^2 - \theta^2} - \frac{2\theta r'_{23} \sin 2\theta\hat{t}}{\Omega^2 - 4\theta^2} - \frac{3\theta r'_{24} \sin 3\theta\hat{t}}{\Omega^2 - 9\theta^2} - \right. \right. \\
&\quad \left. \frac{(\varphi + \theta)r'_{25} \sin(\varphi + \theta)\hat{t}}{\Omega^2 - (\varphi + \theta)^2} + \frac{(\varphi + \theta)r'_{26} \cos(\varphi + \theta)\hat{t}}{\Omega^2 - (\varphi + \theta)^2} - \frac{(\varphi - \theta)r'_{27} \sin(\varphi - \theta)\hat{t}}{\Omega^2 - (\varphi - \theta)^2} + \frac{(\varphi - \theta)r'_{28} \cos(\varphi - \theta)\hat{t}}{\Omega^2 - (\varphi - \theta)^2} \right\} + 2 \left\{ -\Omega\alpha_{11} \sin\Omega\hat{t} + \right. \\
&\quad \left. \Omega\beta_{11} \cos\Omega\hat{t} - \frac{\theta r_{22} \sin\theta\hat{t}}{\Omega^2 - \theta^2} - \frac{2\theta r_{23} \sin 2\theta\hat{t}}{\Omega^2 - 4\theta^2} - \frac{3\theta r_{24} \sin 3\theta\hat{t}}{\Omega^2 - 9\theta^2} - \frac{(\varphi + \theta)r_{25} \sin(\varphi + \theta)\hat{t}}{\Omega^2 - (\varphi + \theta)^2} + \frac{(\varphi + \theta)r_{26} \cos(\varphi + \theta)\hat{t}}{\Omega^2 - (\varphi + \theta)^2} - \right. \\
&\quad \left. \frac{(\varphi - \theta)r_{27} \sin(\varphi - \theta)\hat{t}}{\Omega^2 - (\varphi - \theta)^2} + \frac{(\varphi - \theta)r_{28} \cos(\varphi - \theta)\hat{t}}{\Omega^2 - (\varphi - \theta)^2} \right\} + \frac{\alpha}{2} \left\{ \left(\frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} + \frac{Br_5}{\varphi^2 - \theta^2} \right) \sin\theta\hat{t} + \frac{\alpha_1 \alpha_7}{2} \cos(\varphi + \theta)\hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi + \right. \\
&\quad \left. \theta)\hat{t} + \frac{\alpha_1 \alpha_7}{2} \cos(\varphi - \theta)\hat{t} + \frac{\alpha_1 \beta_7}{2} \sin(\varphi - \theta)\hat{t} + \left(\frac{\alpha_1 r_5}{2(\varphi^2 - \theta^2)} + \frac{Br_6}{\varphi^2 - 4\theta^2} \right) \sin 2\theta\hat{t} + \frac{\alpha_1 r_6}{2(\varphi^2 - 4\theta^2)} \sin 3\theta\hat{t} + \right. \\
&\quad \left. B\alpha_7 \sin\varphi\hat{t} + B\beta_7 \sin\varphi\hat{t} \right\} + \frac{\alpha}{2} \left\{ \frac{\beta_2 \alpha}{2} \left(\frac{B^2 + \frac{\alpha_1^2}{2}}{\varphi^2} \right) - \frac{\alpha\alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} \sin\theta\hat{t} + \frac{\alpha\alpha_1 B \beta_2}{2(\varphi^2 - \theta^2)} \sin 2\theta\hat{t} + \frac{\alpha\alpha_1^2 \beta_2}{8(\varphi^2 - 4\theta^2)} \sin 3\theta\hat{t} - \right. \\
&\quad \left. \frac{\beta_2 \beta_5}{2} \cos(\varphi + \theta)\hat{t} \right\} + \frac{3\beta}{4} \left\{ \left(\beta_2 \left(B^2 + \frac{\alpha_1^2}{2} \right) - \frac{\beta_2 \alpha_1^2}{4} \right) \sin\theta\hat{t} + \beta_2 B \alpha_1 \sin 2\theta\hat{t} + \frac{\beta_2 \alpha_1^2}{4} \sin 3\theta\hat{t} \right\} \quad (4.76)
\end{aligned}$$

To ensure uniformly valid solution in \hat{t} , needs equating the coefficients of $\cos\Omega\hat{t}$ and $\sin\Omega\hat{t}$ to zero. The coefficients of $\cos\Omega\hat{t}$ yields

$$-2\Omega\beta'_{11} - 2\Omega\beta_{11} - \frac{\alpha B\beta_7}{2} = 0 \quad (4.77a)$$

$$\therefore \beta'_{11} + \beta_{11} = -\frac{\alpha B\alpha_7}{2\Omega} = h_5(\tau) \quad (4.77b)$$

where

$$h_5(\tau) = -\frac{\alpha B\alpha_7}{2\Omega} \quad (4.77c)$$

$$\therefore \beta_{11} = e^{-\tau} [\int h_5(\tau) e^{\tau} ds + \beta_{11}(0)] \quad (4.77d)$$

The coefficients of $\sin\Omega\hat{t}$ yields

$$-2\Omega\alpha'_{11} - 2\Omega\alpha_{11} - \frac{\alpha B\beta_7}{2} = 0 \quad (4.77e)$$

where

$$h_6(\tau) = \frac{\alpha B\beta_7}{4\Omega} \quad (4.77f)$$

$$\therefore \lim \alpha_{11} = e^{-\tau} [\int h_6(\tau) e^{\tau} ds + \alpha_{11}(0)] \quad (4.77g)$$

The remaining equation (4.76) is:

$$U_{3m,\hat{t}\hat{t}}^{(31)} + \Omega^2 U_{2m}^{(31)} = r_{50} \sin\theta\hat{t} + r_{51} \sin 2\theta\hat{t} + r_{52} \sin 3\theta\hat{t} + r_{53} \cos(\varphi + \theta)\hat{t} + r_{54} \sin(\varphi + \theta)\hat{t} \\ + r_{55} \cos(\varphi - \theta)\hat{t} + r_{56} \sin(\varphi - \theta)\hat{t} \quad (4.78)$$

The initial conditions are

$$U_{3m}^{(31)}(0,0) = 0; U_{3m,\hat{t}}^{(31)}(0,0) + U_{3m,\tau}^{(30)}(0,0) = 0$$

$$r_{50} = \frac{2\theta r_{22}^1}{\Omega^2 - \theta^2} + \frac{2\theta r_{22}}{\Omega^2 - \theta^2} - \frac{\alpha\alpha_1 r_6}{4(\varphi^2 - 4\theta^2)} - \frac{B\alpha r_5}{2(\varphi^2 - 4\theta^2)} - \frac{\alpha^2 \beta_2}{4} \left(\frac{\alpha_1^2}{2} + B^2 \right) + \frac{\alpha^2 \alpha_1 \beta_2}{16(\varphi^2 - 4\theta^2)} \\ - \frac{3\alpha\beta\beta_2}{8} \left(B^2 + \frac{\alpha_1^2}{2} \right) + \frac{3\alpha\beta\alpha_1^2}{32}$$

$$r_{50}(0) = B^3 \left(\frac{2\theta S_{17}}{(\Omega^2 - \theta^2)} + \frac{2\theta S_{11}}{(\Omega^2 - \theta^2)} + \frac{\alpha S_1}{4(\varphi^2 - 4\theta^2)} + \frac{\alpha^2}{2\theta(\varphi^2 - 4\theta^2)} + \frac{3\alpha^2}{8\theta\varphi^2} + \frac{\alpha^2}{16B(\varphi^2 - 4\theta^2)} \right) \\ + \frac{9\alpha\beta}{16\theta} + \frac{3\alpha\beta}{32B}$$

$$r_{51} = \frac{4\theta r_{23}^1}{\Omega^2 - 4\theta^2} + \frac{4\theta r_{23}}{\Omega^2 - 4\theta^2} - \frac{\alpha\alpha_1 r_5}{4(\varphi^2 - \theta^2)} - \frac{B\alpha r_6}{2(\varphi^2 - 4\theta^2)} - \frac{\alpha^2 \alpha_1 B\beta_2}{4(\varphi^2 - 4\theta^2)} - \frac{\alpha\alpha_1 \beta_2 B}{2}$$

$$r_{51}(0) = B^3 \left(\frac{4\theta S_{18}}{(\Omega^2 - 4\theta^2)} + \frac{4\theta S_{12}}{(\Omega^2 - 4\theta^2)} + \frac{\alpha^2}{2\theta(\varphi^2 - \theta^2)} - \frac{\alpha S_1}{2(\varphi^2 - 4\theta^2)} - \frac{\alpha}{2\theta} \right)$$

$$r_{52} = \frac{6\theta r_{24}^1}{\Omega^2 - 9\theta^2} + \frac{6\theta r_{24}}{\varphi^2 - 9\theta^2} - \frac{\alpha\alpha_1 r_6}{4(\varphi^2 - 4\theta^2)} - \frac{\alpha^2 \beta_2 \alpha_1^2}{16(\varphi^2 - 4\theta^2)} + \frac{\alpha\alpha_1^2 \beta_2}{8}$$

$$r_{52}(0) = B^3 \left(\frac{6\theta S_{19}}{(\Omega^2 - 9\theta^2)} + \frac{6\theta S_{13}}{(\Omega^2 - 9\theta^2)} + \frac{\alpha S_1}{4(\varphi^2 - 4\theta^2)} - \frac{\alpha^2}{16\theta(\varphi^2 - 4\theta^2)} + \frac{\alpha}{8\theta} \right)$$

$$r_{53} = -\frac{2r'_{26}(\varphi+\theta)}{\Omega^2 - (\varphi+\theta)^2} - \frac{2r_{26}(\varphi+\theta)}{\Omega^2 - (\varphi+\theta)^2} - \frac{\alpha\alpha_1\alpha_7}{4}, r_{53}(0) = 0$$

$$r_{54} = \frac{2r'_{25}(\varphi+\theta)}{\Omega^2 - (\varphi+\theta)^2} + \frac{2r_{25}(\varphi+\theta)}{\Omega^2 - (\varphi+\theta)^2} - \frac{\alpha\alpha_1\beta_7}{4}, r_{54}(0) = B^3 \left(\frac{6\alpha S_0(\varphi+\theta)}{\Omega^2 - (\varphi+\theta)^2} + \frac{\alpha S_{43}}{4} \right)$$

$$r_{55} = \frac{-2r'_{28}(\varphi-\theta)}{\Omega^2 - (\varphi-\theta)^2} - \frac{2r_{28}(\varphi-\theta)}{\Omega^2 - (\varphi-\theta)^2} - \frac{\alpha\alpha_1\alpha_7}{4}, r_{55}(0) = \frac{-4\alpha S_0 B^3}{\Omega^2 - (\varphi-\theta)^2}$$

$$r_{56} = \frac{2r'_{27}(\varphi-\theta)}{\Omega^2 - (\varphi-\theta)^2} + \frac{2r_{27}(\varphi-\theta)}{\Omega^2 - (\varphi-\theta)^2} + \frac{\alpha\alpha_1\beta_7}{4}, r_{56}(0) = B^3 \left(\frac{6\alpha S_0(\varphi-\theta)}{\Omega^2 - (\varphi-\theta)^2} + \frac{\alpha S_{43}}{4} \right)$$

Therefore;

$$U_{3m}^{(31)} = \alpha_{14} \cos\Omega\hat{t} + \beta_{14} \sin\Omega\hat{t} + \frac{r_{50} \sin\theta\hat{t}}{\Omega^2 - \theta^2} + \frac{r_{51} \sin 2\theta\hat{t}}{\Omega^2 - 4\theta^2} + \frac{r_{52} \sin 3\theta\hat{t}}{\Omega^2 - 9\theta^2} + \left(\frac{r_{53} \cos(\varphi+\theta)\hat{t} + r_{54} \sin(\varphi+\theta)\hat{t}}{\Omega^2 - (\varphi+\theta)^2} \right) + \\ \left(\frac{r_{55} \cos(\varphi-\theta)\hat{t} + r_{56} \sin(\varphi-\theta)\hat{t}}{\Omega^2 - (\varphi-\theta)^2} \right) \quad (4.79)$$

Therefore,

$$\alpha_{14}(0) = - \left[\frac{r_{53}}{\Omega^2 - (\varphi + \theta)^2} + \frac{r_{55}}{\Omega^2 - (\varphi - \theta)^2} \right] \Big|_{\tau = 0} \quad (4.80a)$$

$$\Omega \beta_{14}(0) = - \frac{\theta r_{50}}{\Omega^2 - \theta^2} - \frac{2\theta r_{51}}{\Omega^2 - 4\theta^2} - \frac{3\theta r_{52}}{\Omega^2 - 9\theta^2} - \frac{(\varphi + \theta)r_{54}}{\Omega^2 - (\varphi + \theta)^2} - \frac{(\varphi - \theta)r_{56}}{\Omega^2 - (\varphi - \theta)^2} - \alpha'_{11} - \frac{r'_{22}}{\Omega^2 - \theta^2} - \frac{r'_{23}}{\Omega^2 - 4\theta^2} - \frac{r'_{24}}{\Omega^2 - 9\theta^2} - \frac{r'_{25}}{\Omega^2 - (\varphi + \theta)^2} - \frac{r'_{27}}{\Omega^2 - (\varphi - \theta)^2}$$

Therefore;

$$\beta_{14}(0) = \frac{-1}{\Omega} \left[\frac{(\theta r_{50} + r'_{22})}{\varphi^2 - \theta^2} + \frac{(2\theta r_{51} + r'_{23})}{\varphi^2 - 4\theta^2} + \frac{(3\theta r_{52} + r'_{24})}{\varphi^2 - 9\theta^2} - \frac{((\theta + \varphi)r_{54} + r'_{24})}{\Omega^2 - (\varphi + \theta)^2} + \frac{(\theta - \varphi)r_{56} + r'_{27}}{\Omega^2 - (\varphi - \theta)^2} + \alpha'_{11}(0) \right] \quad (4.80b)$$

So far, it follows that

$$U^{(31)} = U_m^{(31)}(1 - \cos 2mx) + U_{2m}^{(31)}(1 - \cos 4mx) + U_{3m}^{(31)}(1 - \cos 6mx) \quad (4.81)$$

The summary of the solution so far is,

$$U(x, t, \tau) = (U^{(10)} + \delta U^{(11)} + \delta^2 U^{(12)} + \dots) + \epsilon^2 (U^{(20)} + \delta U^{(21)} + \delta^2 U^{(22)} + \dots) + \epsilon^3 (U^{(30)} + \delta U^{(31)} + \delta^2 U^{(32)} + \dots) + \dots \quad (4.82)$$

4.2 Maximum Displacement of the Column

The dynamic buckling load is obtained from the maximization $\frac{d\lambda}{dU_a} = 0$, where U_a is the maximum displacement and λ is the load parameter. The conditions for maximum displacement are,

$$\frac{\partial U}{\partial x} = 0, \quad \frac{\partial w}{\partial t} = 0 \quad (4.83a)$$

But from (3.12), it follows that

$$\frac{\partial w}{\partial t} = U_{,\hat{t}} + (\omega'_1 \epsilon + \omega'_2 \epsilon^2 + \dots) U_{,\hat{t}} + \delta U_{,\tau} = 0 \quad (4.83b)$$

The aim is to determine the maximum displacement;

$$U_a = U(x_a, \hat{t}_a, t_a)$$

where x_a, t_a, τ_a and \hat{t}_a are the values of x, t, τ , and \hat{t} respectively at maximum displacement and are to be next determined before finally determining the maximum displacement.

From the first condition of maximization, $\frac{\partial U}{\partial x} = 0$, this means

$$\epsilon \left[\frac{\partial U^{(10)}}{\partial x} + \delta \frac{\partial U^{(11)}}{\partial x} + \dots \right] + \epsilon^2 \left[\frac{\partial U^{(20)}}{\partial x} + \delta \frac{\partial U^{(21)}}{\partial x} + \dots \right] + \epsilon^3 \left[\frac{\partial U^{(30)}}{\partial x} + \delta \frac{\partial U^{(31)}}{\partial x} + \dots \right] = 0 \quad (4.84)$$

i.e,

$$2m\epsilon [U_m^{(10)} \sin 2mx + \delta U_m^{(11)} \sin 2mx + \dots] + \epsilon^2 [2m U_m^{(20)} \sin 2mx + 4m U_{2m}^{(20)} \sin 4mx + \dots \delta \{2m U_m^{(21)} \sin 2mx + 4m U_{2m}^{(21)} \sin 4mx + \dots\}] + \epsilon^3 [2m U_m^{(30)} \sin 2mx + 4m U_{2m}^{(30)} \sin 4mx + 6m U_{3m}^{(30)} \sin 6mx + \dots \delta \{2m U_m^{(31)} \sin 2mx + 4m U_{2m}^{(31)} \sin 4mx + 6m U_{3m}^{(31)} \sin 6mx + \dots\}] + \dots = 0 \quad (4.85)$$

The equation (4.85) is satisfied if $\sin 2mx_a = 0$, where x_a is the value of x at maximum displacement.

This means, $2mx_a = \pi n$, $n = 0, 1, 2, 3, \dots$, set $n = 1$, $x_a = \frac{\pi}{2m}$, substituting, $x_a = \frac{\pi}{2m}$ in $U(x, \hat{t}, \tau)$, gives

$$U(x_a, \hat{t}, \tau) = 2\epsilon [U_m^{(10)} + \delta U_m^{(11)} + \dots] + 2\epsilon^2 [U_m^{(20)} + \delta U_{2m}^{(21)} + \dots] + 2\epsilon^3 [(U_m^{(30)} + U_{3m}^{(30)}) + \delta (U_m^{(31)} + U_{3m}^{(31)}) + \dots] \quad (4.86)$$

Let \hat{t}_a, t_a and τ_a be the values of \hat{t}, t and τ respectively at maximum displacement and let them be expanded asymptotically as

$$\hat{t}_a = \hat{t}_0 + \delta \hat{t}_{01} + \delta^2 \hat{t}_{02} + \epsilon (\hat{t}_{10} + \delta \hat{t}_{11} + \delta^2 \hat{t}_{12} + \dots) + \epsilon^2 (\hat{t}_{20} + \delta \hat{t}_{21} + \delta^2 \hat{t}_{22} + \dots) \quad (4.87a)$$

$$t_a = t_0 + \delta t_{01} + \delta^2 t_{02} + \dots + \epsilon (t_{10} + \delta t_{11} + \delta^2 t_{12} + \dots) + \epsilon^2 (t_{20} + \delta t_{21} + \delta^2 t_{22} + \dots) \quad (4.87b)$$

$$\tau_\alpha = \delta[t_0 + \delta t_{01} + \delta^2 t_{02} + \dots \dots + \epsilon(t_{10} + \delta t_{11} + \delta^2 t_{12} + \dots) + \epsilon^2(t_{20} + \delta t_{21} + \delta^2 t_{22} + \dots)] \quad (4.87c)$$

Evaluating (4.87c) at the maximum values and simplifying, the following are obtained:

$$\hat{t}_0 = \frac{\pi}{\theta}, \quad t_0 = \frac{\pi}{\theta}, \quad t_{10} = -\frac{t_0 B}{\theta^2}, \quad t_{20} = \hat{t}_{20} - \hat{t}_{10} \omega'_1(0) - t_0 \omega'_2(0) \text{ and}$$

$$\hat{t}_{20} = \frac{B^2 \alpha S_0 \sin \varphi \hat{t}_0}{\theta^2} \left[\frac{(\varphi - \theta)}{\Omega^2 - (\varphi - \theta)^2} - \frac{(\varphi + \theta)}{\Omega^2 - (\varphi + \theta)^2} - \frac{(\varphi + \theta)}{\varphi(2\theta + \varphi)} + \frac{(\varphi - \theta)}{\varphi(2\theta - \varphi)} \right]$$

Let U_a be the maximum displacement. We now substitute for x_a ;

$$U\left(\frac{\pi}{2m}, \hat{t}, \tau\right) = \epsilon[2U_m^{10} + 2\delta U_m^{(11)} \dots] + \epsilon^2[2U_m^{20} + 2\delta U_{2m}^{(21)} \dots] + \epsilon^3[(2U_m^{30} + 2U_{3m}^{(30)} \dots) + \delta(2U_m^{31} + 2\delta U_{3m}^{(31)} \dots)] \quad (4.88)$$

Expanding each of the terms in (4.88) and evaluating (4.88) at maximum values and noting that all $U_m^{(ij)}$ are evaluated at $(\hat{t}_0, 0)$, the following are obtained

Therefore,

$$\begin{aligned} U_a = 2\epsilon & \left[U_m^{(10)} + \delta \left\{ \hat{t}_0 U_{m,\hat{t}}^{(10)} + t_0 U_{m,\tau}^{(10)} + U_m^{(11)} \right\} + \dots \right] \\ & + 2\epsilon^2 \left[\hat{t}_{10} U_{m,\hat{t}}^{(10)} + U_m^{(20)} \right. \\ & + \delta \left\{ \hat{t}_{11} U_{m,\hat{t}}^{(10)} + t_{10} U_{m,\tau}^{(10)} + \hat{t}_{01} \hat{t}_{10} U_{m,\hat{t}\hat{t}}^{(10)} + \hat{t}_{10} t_0 U_{m,\hat{t}\tau}^{(10)} + \hat{t}_{10} U_{m,\hat{t}}^{(11)} + \dots \right\} + \dots \left. \right] \\ & + 2\epsilon^3 \left[\hat{t}_{20} U_{m,\hat{t}}^{(10)} + \frac{(\hat{t}_{10})^2}{2} U_{m,\hat{t}\hat{t}}^{(10)} + \hat{t}_{10} U_{m,\hat{t}}^{(20)} + (U_m^{(30)} + U_{3m}^{(30)}) \right. \\ & + \delta \left\{ t_{21} U_{m,\hat{t}}^{(10)} + \hat{t}_{20} U_{m,\tau}^{(10)} + \hat{t}_{10} \hat{t}_{11} U_{m,\hat{t}\hat{t}}^{(10)} + \hat{t}_{20} t_0 U_{m,\hat{t}\tau}^{(10)} + \hat{t}_{10} t_{10} U_{m,\hat{t}\tau}^{(10)} + \hat{t}_{20} U_{m,\hat{t}}^{(11)} \dots \right\} \\ & + \frac{1}{2} (t_{10})^2 U_{m,\hat{t}\hat{t}}^{(11)} + \hat{t}_{11} U_{m,\hat{t}}^{(20)} + t_{10} U_{m,\tau}^{(20)} + \hat{t}_{10} t_0 U_{m,\hat{t}\tau}^{(20)} + \hat{t}_{10} U_{m,\hat{t}}^{(21)} \\ & \left. + \hat{t}_{01} (U_m^{(30)} + U_{3m}^{(30)})_{\hat{t}} + t_0 (U_m^{(30)} + U_{3m}^{(30)})_{,\tau} + \dots \right] \text{ at } \tau = 0 \quad (4.89) \end{aligned}$$

Therefore,

$$U_a = 2\epsilon [U_m^{(10)} + \delta t_0 U_{m,\tau}^{(10)} + \dots] + 2\epsilon^2 [U_m^{(20)} + \delta t_{10} U_{m,\tau}^{(10)} + \dots] + 2\epsilon^3 [(U_m^{(30)} + U_{3m}^{(30)}) + \delta t_{20} U_{m,\tau}^{(10)} + \delta \hat{t}_{20} U_{m,\hat{t}}^{(11)} + \delta t_{10} U_{m,\tau}^{(20)} + \delta t_0 (U_m^{(30)} + U_{3m}^{(30)})_{,\tau} + \dots] \text{ at } \tau = 0 \quad (4.90)$$

In what follows, simplifications of the terms in (4.89)-(4.90) are carried out to obtain the following

$$U_m^{(10)}(\hat{t}_0, 0) = 2BU_{m,\tau}^{(10)}(\hat{t}_0, 0) = -B \quad (4.91)$$

$$U_m^{(20)}(\hat{t}_0, 0) = -\alpha_4(0) + \frac{r_0(0)}{\theta^2} + \frac{r_1(0)}{3\theta^2} = \frac{-r_1(0)}{3\theta^2} + \frac{r_0(0)}{\theta^2} + \frac{r_0(0)}{\theta^2} - \frac{r_1(0)}{\theta^2} = 2 \left[\frac{-r_1(0)}{3\theta^2} + \frac{r_0(0)}{\theta^2} \right] = 2 \left[\frac{\alpha B^2}{3\theta^2} - \frac{3\alpha B^2}{\theta^2} \right] = \frac{-16\alpha B^2}{3\theta^2} \quad (4.92)$$

$$\begin{aligned} U_m^{(30)}\left(\frac{\pi}{\theta}, 0\right) &= \frac{135B^3\beta}{8\theta^2} \left[1 + \frac{8\theta^2}{135} \left\{ \left(\frac{\alpha}{\beta}\right) S_0 \left(\frac{1}{\varphi(2\theta + \varphi)} - \frac{1}{\varphi(2\theta - \varphi)} \right) \cos\left(\frac{\varphi\pi}{\theta}\right) \right\} \right. \\ & \left. + \frac{2}{3\theta^2} \left(\frac{\alpha^2}{\beta}\right) \cdot \frac{8\theta^2}{135\beta} (3k_3 - k_4) \right] = \frac{135B^3\beta}{8\theta^2} (1 + A_{31}) \quad (4.93) \end{aligned}$$

where,

$$\begin{aligned} A_{31} &= \left[1 + \frac{8\theta^2}{135} \left\{ \left(\frac{\alpha}{\beta}\right) S_0 \left(\frac{1}{\varphi(2\theta + \varphi)} - \frac{1}{\varphi(2\theta - \varphi)} \right) \cos\left(\frac{\varphi\pi}{\theta}\right) \right\} + \frac{2}{3\theta^2} \left(\frac{\alpha^2}{\beta}\right) \cdot \frac{8\theta^2}{135\beta} (3k_3 - k_4) \right] \\ S_0 &= \left(\frac{\alpha}{\varphi^2 - \theta^2} - \frac{\alpha}{4(\varphi^2 - 4\theta^2)} - \frac{3\alpha}{4\varphi^2} \right), k_3 = \left(\frac{10}{3\theta^2} - \frac{1}{(\varphi^2 - \theta^2)} \right) \\ k_4 &= \left(\frac{2}{3\theta^2} - \frac{1}{(\varphi^2 - \theta^2)} + \frac{8}{3\theta\alpha} \right), k_5 = \left(\frac{1}{3\theta^2} + \frac{1}{4(\varphi^2 - \theta^2)} \right) \end{aligned}$$

Similarly,

$$U_{3m}^{(30)} = -B^3\beta \left(A_{32} + \left(\frac{\alpha}{\beta}\right) S_0 A_{33} \right) \quad (4.94)$$

where,

$$A_{32} = \left[\frac{\frac{15}{16} \left(1 - \frac{16\alpha^2 k_{11}}{15} \right) (1 + \cos \Omega \hat{t}_0)}{\Omega^2 - \theta^2} + \frac{\frac{3}{8} (1 - k_{12}) (1 - \cos \Omega \hat{t}_0)}{\Omega^2 - 4\theta^2} + \frac{(1 - k_{13}) (1 + \cos \Omega \hat{t}_0)}{16(\Omega^2 - 9\theta^2)} \right]$$

and

$$A_{33} = \left[\frac{1 + \cos \Omega \hat{t}_0}{\Omega^2 - (\varphi + \theta)^2} - \frac{1 + \cos \Omega \hat{t}_0}{\Omega^2 - (\varphi - \theta)^2} \right], k_{12} = \left[-\frac{4}{3} \left(\frac{\alpha^2}{\beta} \right) \left(\frac{1}{\varphi^2 - \theta^2} \right) \right], k_{13} = \left[2 \left(\frac{\alpha^2}{\beta} \right) \left(\frac{1}{\varphi^2 - 4\theta^2} \right) \right]$$

$$U_{m,\tau}^{(20)} \left(\frac{\pi}{\theta}, 0 \right) = -\alpha'_4(0) + \frac{r'_0(0)}{\theta^2} - \frac{r'_1(0)}{3\theta^2} \quad (4.95)$$

From (4.24h),

$$\alpha'_4(0) = -\alpha_1(0) + \frac{1}{2\theta} [\alpha B \beta_2(0) - 2\theta^2 \omega'_1(0) \beta_2(0) - \omega''_1(0) \theta \alpha_1(0) - 2\omega'_1(0) \alpha_1 \theta] \alpha_1(0) = \frac{-13\alpha B^2}{3\theta^2} + \frac{4B^2}{\theta} \quad (4.96)$$

$$U_{m,\tau}^{(20)} \left(\frac{\pi}{\theta}, 0 \right) = \left(\frac{13\alpha B^2}{3\theta^2} - \frac{4B^2}{\theta} \right) + \frac{r'_0(0)}{\theta^2} - \frac{r'_1(0)}{3\theta^2} = \left(\frac{13\alpha B^2}{3\theta^2} - \frac{4B^2}{\theta} \right) + \frac{2\alpha B^2}{\theta^2} - \frac{2\alpha B^2}{3\theta^2} = \frac{17\alpha B^2}{3\theta^2} - \frac{4B^2}{\theta}$$

$$= B^2 \left(\frac{17\alpha}{3\theta^2} - \frac{4}{\theta} \right) \quad (4.97)$$

Also,

$$\omega_2'' = -\frac{1}{2\theta^2} \left[\frac{(\omega'_1)^2 \theta^2}{\alpha_1} + \frac{2\omega'_1 \theta^2}{\alpha_1} - 2\alpha \left(\frac{r_0}{\theta^2} - \frac{r_1}{6\theta^2} + 3 \left(\frac{\alpha_4}{\alpha_1} \right) \right) - \left\{ \frac{\left(\frac{\alpha_1^2}{2} + B^2 \right)}{\varphi^2} + \frac{\alpha_1^2 \alpha^2}{4(\varphi^2 - 4\theta^2)} \right\} - \frac{45\beta}{4} \left(\frac{\alpha_1^2}{4} + B^2 \right) \right] \quad (4.98)$$

$$\therefore \omega_2''(0) = -\frac{1}{2\theta^2} \left[\frac{\theta^2 \{ \alpha'_1(0) (\omega'_1(0))^2 - 2\alpha_1 \omega''_1(0) \omega'_1(0) \}}{\alpha_1^2(0)} + 2\theta^2 \{ \alpha'_1(0) (\omega'_1(0) \alpha_4(0)) - \alpha_1(0) (\omega''_1(0) \alpha_4(0) + \omega'_1(0) \alpha'_4(0)) \} - 2\alpha \left(\frac{r_0(0)}{\theta^2} - \frac{r_1(0)}{6\theta^2} + B \left(\frac{\alpha_1(0) \alpha'_4(0) - \alpha_4(0) \alpha'_1(0)}{\alpha_1^2(0)} \right) \right) - \left\{ \frac{\alpha_1(0) \alpha_1(0) \alpha^2}{\varphi^2} + \frac{\alpha^2 \alpha_1(0) \alpha'_1(0)}{4(\varphi^2 - 4\theta^2)} - \frac{45\beta}{4} \left(\frac{\alpha_1(0) \alpha'_1(0)}{2} \right) \right\} \right]$$

$$= -\frac{1}{2\theta^2} \left[\theta^2 \left\{ \frac{B^3}{\theta^4 B^2} \right\} + 2\theta^2 \left\{ \frac{B^2}{\theta^2} \cdot \frac{8\alpha B^2}{3\theta^2} + B \left(\frac{B}{\theta^2} \cdot B^2 S_{51} \right) \right\} - 2\alpha \left\{ \frac{2\alpha B^2}{\theta^2} - \frac{2\alpha B^2}{6\theta^2} + B \left(\frac{-B \cdot B^2 S_{51} - \frac{8\alpha B^3}{3\theta^2}}{B^2} \right) \right\} - \left\{ \alpha^2 \left(\frac{-B^2}{\varphi^2} \right) + \frac{\alpha^2 B(-B)}{2(\varphi^2 - 4\theta^2)} \right\} - \frac{45\beta(-B^2)}{8} \right]$$

$$= -\frac{1}{2\theta^2} \left[\frac{B}{\theta^2} + 2 \left\{ \frac{B^4 \alpha}{3\theta^2} + B^4 S_{51} \right\} - 2\alpha^2 \left\{ \frac{5B^2}{3\theta^2} - B^2 \left(\frac{S_{51}}{\alpha} + \frac{8}{3\theta^2} \right) \right\} + \alpha^2 B^2 \left(\frac{1}{\varphi^2} - \frac{1}{2(\varphi^2 - 4\theta^2)} \right) + \frac{45\beta B^2}{8} \right] \quad (4.99)$$

$$\Rightarrow \omega_2''(0) = -\frac{1}{2\theta^2} \left[\frac{B}{\theta^2} - 2\alpha^2 B^2 \left\{ \frac{5}{3\theta^2} - \left(\frac{S_{51}}{\alpha} + \frac{8}{3\theta^2} \right) \right\} + \alpha^2 B^2 \left(\frac{1}{\varphi^2} - \frac{1}{2(\varphi^2 - 4\theta^2)} \right) + \frac{45\beta B^2}{8} + 2B^4 \alpha \left(\frac{1}{3\theta^2} + \frac{S_{51}}{\alpha} \right) \right] \quad (4.100)$$

Similarly,

$$U_{m,\tau}^{(30)} \left(\frac{\pi}{\theta}, 0 \right) = -B^3 S_{65} + \frac{B^3 S_{20}}{\theta^2} + \frac{B^3 S_{20}}{\theta^2} + \frac{B^3 S_{21}}{3\theta^2} + \frac{2\alpha B^3 S_0}{\theta^2} \cos \left(\frac{\varphi \pi}{\theta} \right) \left[\frac{1}{(2\theta - \varphi)} - \frac{1}{(2\theta + \varphi)} \right]$$

$$\Rightarrow U_{m,\tau}^{(30)}\left(\frac{\pi}{\theta}, 0\right) = B^3 S_{65} + \frac{B^3 S_{20}}{\theta^2} + \frac{B^3 S_{20}}{\theta^2} + \frac{B^3 S_{21}}{3\theta^2} + \frac{2\alpha B^3 S_0}{\theta^2} \cos\left(\frac{\varphi\pi}{\theta}\right) \left[\frac{1}{(2\theta - \varphi)} - \frac{1}{(2\theta + \varphi)} \right] \quad (4.101)$$

where,

$$S_{65} = -S_{64} + \frac{S_{20}}{\theta^2} + \frac{S_{21}}{3\theta^2} + \frac{2\alpha S_0}{\theta^2} \cos\left(\frac{\varphi\pi}{\theta}\right) \left[\frac{1}{(2\theta - \varphi)} - \frac{1}{(2\theta + \varphi)} \right]$$

where,

$$S_{64} = S_{62} - S_{63}, \quad S_{62} = -\frac{1}{2\theta^2} \left[\frac{6\alpha^2}{\theta^3} - 2\theta S_{49} + \frac{\alpha^2}{3\theta^3} - \frac{\theta \omega_2''(0)}{B^2} \right], \quad h_1(0) = B^3 S_{62},$$

$$S_{63} = \left(\frac{-S_3}{\theta^2} + \frac{S_4}{3\theta^2} + \frac{S_5}{8\theta^2} + \frac{\alpha S_0}{\varphi(2\theta - \varphi)} \right)$$

Also,

$$U_{3m,\tau}^{(30)} = \alpha'_{11}(0) \cos \Omega \hat{t}_0 + \beta'_{11}(0) \sin \Omega \hat{t}_0 + \frac{r'_{22}(0) \cos \theta \hat{t}_0}{\Omega^2 - \theta^2} + \frac{r'_{23}(0) \cos 2\theta \hat{t}_0}{\Omega^2 - 4\theta^2} + \frac{r'_{24}(0) \cos 3\theta \hat{t}_0}{\Omega^2 - 9\theta^2} + \frac{r'_{25}(0) \cos(\varphi + \theta) \hat{t}_0 + r'_{26}(0) \sin(\varphi + \theta) \hat{t}_0}{\Omega^2 - (\varphi + \theta)^2} + \frac{r'_{27}(0) \cos(\varphi - \theta) \hat{t}_0 + r'_{28}(0) \sin(\varphi - \theta) \hat{t}_0}{\Omega^2 - (\varphi - \theta)^2} \quad (4.102)$$

$$\alpha'_{11}(0) = h_6(0) - \alpha_{11}(0) = \frac{\alpha B^3 S_{43}}{4\Omega} - B^3 S_{48} = B^3 S_{66}, \quad S_{66} = \frac{\alpha S_{43}}{4\Omega} - S_{48}$$

Similarly, $\beta'_{11}(0) = h_5(0) - \beta_{11}(0) = -\beta_{11}(0) = 0$ since $h_5(0) = 0$

$$\therefore U_{3m,\tau}^{(30)}\left(\frac{\pi}{\theta}, 0\right) = B^3 S_{66} \cos \Omega \left(\frac{\pi}{\theta}\right) - \frac{B^3 S_{17}}{\Omega^2 - \theta^2} + \frac{B^3 S_{18}}{\Omega^2 - 4\theta^2} - \frac{B^3 S_{19}}{\Omega^2 - 9\theta^2} - \frac{2\alpha B^3 S_0 \cos\left(\frac{\varphi\pi}{\theta}\right)}{\Omega^2 - (\varphi + \theta)^2} - \frac{2\alpha B^3 S_0 \cos\left(\frac{\varphi\pi}{\theta}\right)}{\Omega^2 - (\varphi - \theta)^2}$$

i.e.,

$$U_{3m,\tau}^{(30)}\left(\frac{\pi}{\theta}, 0\right) = B^3 S_{67} \quad (4.103)$$

where,

$$S_{67} = S_{66} \cos \Omega \left(\frac{\pi}{\theta}\right) - \frac{S_{17}}{\Omega^2 - \theta^2} + \frac{S_{18}}{\Omega^2 - 4\theta^2} - \frac{S_{19}}{\Omega^2 - 9\theta^2} - 2\alpha S_0 \cos\left(\frac{\varphi\pi}{\theta}\right) \left[\frac{1}{\Omega^2 - (\varphi + \theta)^2} - \frac{1}{\Omega^2 - (\varphi - \theta)^2} \right]$$

Therefore, the maximum displacement is

$$U_a\left(\frac{\pi}{\theta}, 0\right) = 2\epsilon [2B - t_0 B \delta + \dots] + 2\epsilon^2 \left[\frac{-16\alpha B^2}{3\theta^2} - \frac{t_0 B(-B)\delta}{\theta^2} + \dots \right] + 2\epsilon^3 \left[\frac{135B^3 \beta(1+A_{31})}{8\theta^2} - B^3 \beta \left(A_{32} + \frac{\alpha}{\beta} S_0 A_{33} \right) \right] + \delta \left[-t_{20} B - \hat{t}_{20} B + t_{10} B^2 \left(\frac{17\alpha}{3\theta^2} - \frac{4}{\theta^2} \right) + t_0 B^2 (S_{65} + S_{67}) + \dots \right] \quad (4.104)$$

i.e.,

$$U_a\left(\frac{\pi}{\theta}, 0\right) = \left[4B\epsilon \left(1 - \frac{t_0 \delta}{2} \right) - \frac{32\alpha B^2 \epsilon^2}{3\theta^2} \left(1 - \frac{3\delta t_0}{16\alpha} + \dots \right) + \frac{135\beta(1+A_{31})B^3 \epsilon^3}{4\theta^2} \left\{ 1 - \frac{8\theta^2 \left(A_{32} + \frac{\alpha}{\beta} S_0 A_{33} \right)}{135(1+A_{31})} \right\} + \frac{8\delta \theta^2}{135\beta(1+A_{31})} \left\{ -\frac{t_{20}}{B^2} - \frac{\hat{t}_{20}}{B^2} + \frac{t_{10}}{B} \left(\frac{17\alpha}{3\theta^2} - \frac{4}{\theta} \right) + t_0 (S_{65} + S_{67}) \right\} \right] \quad (4.105)$$

A further simplification of (4.105) yields

$$U_a\left(\frac{\pi}{\theta}, 0\right) \equiv U_a = 4B\epsilon D_1 - \frac{32\alpha B^2 D_2 \epsilon^2}{3\theta^2} + \frac{135\beta(1+A_{31})B^3 \epsilon^3}{4\theta^2} [D_3 + D_4] + \dots \quad (4.106)$$

where,

$$D_1 = 1 - \frac{t_0\delta}{2}, \quad D_2 = 1 - \frac{3t_0\delta}{16\alpha}, \quad D_3 = 1 - \frac{8\theta^2(A_{32} + \frac{\alpha}{\beta}S_0A_{33})}{135(1 + A_{31})}$$

$$D_4 = \frac{8\delta\theta^2}{135\beta(1 + A_{31})} \left\{ \frac{-t_{20}}{B^2} - \frac{\hat{t}_{20}}{B^2} + \frac{t_{10}}{B} \left(\frac{17\alpha}{3\theta^2} - \frac{4}{\theta} \right) + t_0(S_{65} + S_{67}) \right\}$$

Equation (4.106) can be rewritten as

$$U_a = 4B\epsilon D_1 - \frac{32\alpha B^2 D_2 \epsilon^2}{3\theta^2} + \frac{135\beta(1 + A_{31})B^3 D_3 \epsilon^3}{4\theta^2} \left[1 + \frac{D_4}{D_3} \right] + \dots \quad (4.107)$$

Equation (4.107) can further be rewritten as,

$$U_a = \epsilon c_1 + \epsilon^2 c_2 + \epsilon^3 c_3 + \dots \quad (4.108a)$$

where,

$$c_1 = 4BD_1, \quad c_2 = -\frac{32\alpha B^2 D_2}{3\theta^2}, \quad c_3 = \frac{135\beta(1+A_{31})B^3 D_3}{4\theta^2} \left(1 + \frac{D_4}{D_3} \right) = \frac{135\beta(1+A_{31})B^3 D_3(1+D_5)}{4\theta^2}$$

where, $D_5 = \left(\frac{D_4}{D_3} \right)$

To reverse the series (4.108a) as in Ette (2007), we have

$$\epsilon = d_1 U_a + d_2 U_a^2 + d_3 U_a^3 + \dots \quad (4.108b)$$

By substituting for U_a in (4.108b) and equating the coefficients of powers of ϵ , (4.108b) becomes

$$\epsilon = d_1(\epsilon c_1 + \epsilon^2 c_2 + \epsilon^3 c_3 + \dots) + d_2(\epsilon c_1 + \epsilon^2 c_2 + \epsilon^3 c_3 + \dots)^2 + d_3(\epsilon c_1 + \epsilon^2 c_2 + \epsilon^3 c_3 + \dots)^3 \quad (4.109a)$$

$$O(\epsilon): 1 = d_1 c_1$$

$$\therefore d_1 = \frac{1}{c_1}$$

$$O(\epsilon^2): 0 = d_1 c_1 + d_2 c_1^2$$

$$\therefore d_2 = \frac{d_1 c_2}{c_1^2} = -\frac{c_2}{c_1^3}$$

$$O(\epsilon^3): 0 = d_1 c_3 + 2d_2 c_1 c_2 + d_3 c_1^3$$

$$\therefore d_3 = \frac{-(d_1 c_3 + 2d_2 c_1 c_2)}{c_1^3} = \frac{2c_2^2 - c_1 c_3}{c_1^5}$$

4.3 The Dynamic Buckling Load, λ_D of the Column

As in (3.1), the dynamic buckling load λ_D is now obtained from the maximization, $\frac{d\lambda}{dU_a} = 0$. This is easily done from (4.108a) to yield,

$$\frac{d\epsilon}{dU_a} = \left(\frac{d\epsilon}{d\lambda} \cdot \frac{d\lambda}{dU_a} \right) = 0$$

$$\therefore d_1 + 2U_{aD}d_2 + 3d_3U_{aD}^2 = 0 \quad (4.110)$$

Where, U_{aD} is the value of U_a at buckling and solving (4.110) yields,

$$U_{aD} = \frac{1}{3d_3} \left\{ -d_2 \pm (d_2^2 - 3d_1d_3)^{\frac{1}{2}} \right\} \quad (4.111)$$

The negative root sign in (4.111) is considered because the positive root sign is of no physical significance. Therefore,

$$U_{aD} = \frac{1}{3d_3} \left\{ -d_2 - (d_2^2 - 3d_1d_3)^{\frac{1}{2}} \right\} \quad (4.112)$$

Further simplification of (4.112) yields

$$U_{aD} = \frac{1}{\frac{-c_3}{c_1^4} \left(1 - \frac{2c_2^2}{c_1 c_3} \right)} \left[-\sqrt{\frac{3c_3}{c_1^5} \left(1 - \frac{5c_2^2}{3c_1 c_3} \right)} \left\{ \left(1 - \frac{c_2}{\sqrt{3c_1 c_3} \left(1 - \frac{5c_2^2}{3c_1 c_3} \right)^{\frac{1}{2}}} \right) \right\} \right] \quad (4.113)$$

i.e,

$$U_{aD} = \sqrt{\frac{c_1^3}{3c_3}} \left[\sqrt{\left(1 - \frac{5c_2^2}{3c_1 c_3} \right)} \left\{ \frac{1 - \frac{c_2}{\sqrt{3c_1 c_3} \left(1 - \frac{5c_2^2}{3c_1 c_3} \right)^{\frac{1}{2}}}}{\frac{2c_2^2}{c_1 c_3}} \right\} \right] \quad (4.114)$$

But,

$$\sqrt{\frac{c_1^3}{3c_3}} = \frac{1}{\sqrt{3}} \left\{ \frac{\{4BD_1\}^3}{3\{135\beta(1+A_{31})B^3D_3(1+D_5)\}} \right\}^{\frac{1}{2}} = \frac{1}{\sqrt{3}} \left\{ \frac{64B^3D_1^3.4\theta^2}{405\beta B^3D_3(1+D_5)(1+A_{31})} \right\}^{\frac{1}{2}} = \frac{16\theta D_1^{\frac{3}{2}}}{9\sqrt{15\beta D_3(1+D_5)(1+A_{31})}} =$$

$$\frac{16\theta}{9\sqrt{15\beta(1+D_5)(1+A_{31})}} \left(\frac{D_1^{\frac{3}{2}}}{D_3^{\frac{1}{2}}} \right) = \frac{16\theta D_6}{9\sqrt{15\beta}} = \frac{16\theta D_6}{9\sqrt{15\beta^2}} \quad (1.115)$$

where, $D_6 = \frac{\left(\frac{D_1^3}{D_3}\right)^{\frac{1}{2}}}{\sqrt{(1+D_5)(1+A_{31})}}$

Further simplification of terms in (4.114) yields

$$U_{ad} = \frac{16\theta D_6}{9\sqrt{15\beta^2}} \left[D_7^{\frac{1}{2}} \left\{ \frac{1-D_8}{D_9} \right\} \right] = \frac{16\theta D_6 D_{10}}{9\sqrt{15\beta^2}} \quad (4.116)$$

where,

$$D_7 = 1 - \frac{5c_2^2}{3c_1c_3} = \left[1 + \frac{1024\left(\frac{\alpha^2}{\beta}\right)D_2^2}{729\theta^2 D_1 D_3 (1+D_5)(1+A_{31})} \right]$$

$$D_8 = 1 - \frac{c_2}{\sqrt{3c_1c_3} \left(1 - \frac{5c_2^2}{3c_1c_3}\right)^{\frac{1}{2}}} = 1 + \frac{32\left(\frac{\alpha}{\beta^2}\right)D_2}{27\sqrt{5}\theta\sqrt{D_1 D_3 D_7 (1+D_5)(1+A_{31})}}$$

$$D_9 = \left(1 - \frac{2c_2^2}{c_1c_3}\right) = 1 - \frac{2048D_2^2\left(\frac{\alpha^2}{\beta}\right)}{1215\theta^2 D_1 D_3 (1+A_{31})(1+D_5)}$$

Writing, $D_{10} = \left[D_7^{\frac{1}{2}} \left\{ \frac{1-D_8}{D_9} \right\} \right]$, (4.116) becomes, $U_{ad} = \frac{16\theta D_6 D_{10}}{9\sqrt{15\beta^2}}$

To determine the dynamic buckling load, λ_D , (4.108a) is evaluated at buckling to get,

$$\epsilon = d_1 U_{ad} + d_2 U_{ad}^2 + d_3 U_{ad}^3 + \dots \quad (4.117)$$

Multiplying equation (4.117) by 3, the following is obtained

$$3\epsilon = 3d_1 U_{ad} + 3d_2 U_{ad}^2 + 3d_3 U_{ad}^3 + \dots = 3(d_1 U_{ad} + d_2 U_{ad}^2) + U_{ad}(3d_3 U_{ad}^2) + \dots \quad (4.118)$$

But from (4.110),

$$3d_3 U_{ad}^2 = -d_1 - 2d_2 U_{ad} \quad (4.118)$$

Substituting (4.118) for $3d_3 U_{ad}^2$ in (4.117) yields,

$$3\epsilon = 3(d_1 U_{ad} + d_2 U_{ad}^2 + U_{ad} + \dots) = 2d_1 U_{ad} + d_2 U_{ad}^2 = 2d_1 U_{ad} \left(1 + \frac{d_2 U_{ad}}{2d_1}\right) \quad (4.119)$$

On substituting for d_1, d_2 in equation (4.119), the following is obtained

$$3\epsilon = \frac{2}{c_1} U_{ad} \left(1 - \frac{c_2 U_{ad}}{2c_1^2}\right) \quad (4.120)$$

On substituting for c_1, c_2 and U_{ad} in equation (4.120), the following is obtained

$$3\epsilon = \frac{2\left(\frac{16\theta D_6 D_{10}}{9\sqrt{15\beta^2}}\right)}{4BD_1} \left[1 - \frac{\left(\frac{-32\alpha B^2 D_2}{3\theta^2}\right) \left\{ \frac{16\theta D_6 D_{10}}{9\sqrt{15\beta^2}} \right\}}{2(4BD_1)^2} \right] = \frac{8\theta D_6 D_{10}}{9\sqrt{15} D_1 \beta^{\frac{1}{2}} B} \left[1 + \left(\frac{\alpha D_2}{(D_1 \theta)^2}\right) \left(\frac{16\theta D_6 D_{10}}{9\sqrt{15\beta^2}}\right) \right] = \frac{8\theta D_6 D_{10}}{9\sqrt{15} D_1 \beta^{\frac{1}{2}} B} \left[1 + \frac{16\left(\frac{\alpha}{\beta^2}\right) D_2 D_6 D_{10}}{27\sqrt{15} D_1^2 \theta} \right] \quad (4.121)$$

$$\Rightarrow 3\epsilon = \frac{8(16m^4 - 8\lambda_D m^2 + 1)^{\frac{1}{2}} D_6 D_{10} (16m^4 - 8\lambda_D m^2 + 1)}{9\sqrt{15} D_1 \beta^{\frac{1}{2}} \cdot 8\lambda_D m^2 \bar{a}_m} \left[1 + \frac{16\left(\frac{\alpha}{\beta^2}\right) D_2 D_6 D_{10}}{27\sqrt{15} D_1^2 \theta} \right]$$

i.e,

$$3\epsilon = \frac{(16m^4 - 8\lambda_D m^2 + 1)^{\frac{3}{2}} D_6 D_{10}}{9\sqrt{15} D_1 \beta^{\frac{1}{2}} (\lambda_D m^2 \bar{a}_m)} \left[1 + \frac{16\left(\frac{\alpha}{\beta^2}\right) D_2 D_6 D_{10}}{27\sqrt{15} D_1^2 \theta (\lambda_D)} \right]$$

$$\therefore (16m^4 - 8\lambda_D m^2 + 1)^{\frac{3}{2}} = 27\sqrt{15}D_1(\lambda_D \epsilon) m^2 \bar{a}_m \left[1 + \frac{16 \left(\frac{\alpha}{\beta \left(\frac{1}{2} \right)} \right) D_2 D_6 D_{10}}{27\sqrt{15}D_1^2 \theta(\lambda_D)} \right]^{-1} \quad (4.122)$$

A simple computer programme, written on Q-basic, gives the values of the dynamic buckling loads, λ_D , at different values of ϵ and δ using equation (4.122).

Table 1: Relationship between the Dynamic Buckling Load and the Imperfection Parameters for different values of damping factors, using equation (4.122).

$\bar{a}_1 \epsilon$	λ_D for $\delta = 0$	λ_D for $\delta = 0.01$	λ_D for $\delta = 0.03$
0.01	1.87913	1.87789	1.87548
0.02	1.81858	1.81736	1.81496
0.03	1.77694	1.77571	1.77332
0.04	1.74427	1.74306	1.74068
0.05	1.71702	1.71582	1.71345
0.06	1.69344	1.69225	1.68989
0.07	1.67257	1.67138	1.66903
0.08	1.65376	1.65257	1.65023
0.09	1.63659	1.63541	1.63307
0.1	1.62076	1.61959	1.61726




Figure 1: Relationship between the Dynamic Buckling Load and the Imperfection Parameters for different values of damping factors, using equation (4.122).

Table 2: Relationship between the Dynamic Buckling Load and the damping factors for different values of Imperfection Parameters, using equation (4.122).

δ	λ_D for $a_1 \epsilon = 0.01$	λ_D for $a_1 \epsilon = 0.03$	λ_D for $a_1 \epsilon = 0.05$
0.01	1.87789	1.87789	1.87789

0.02	1.87667	1.87667	1.87667
0.03	1.87548	1.87548	1.87548
0.04	1.87431	1.87431	1.87431
0.05	1.87316	1.87316	1.87316
0.06	1.87204	1.87204	1.87204
0.07	1.87093	1.87093	1.87093
0.08	1.86985	1.86985	1.86985
0.09	1.86878	1.86878	1.86878
0.1	1.86773	1.86773	1.86773




Figure 2: Relationship between the Dynamic Buckling Load and the damping factors for different values of Imperfection Parameters, using equation (4.122).

4.4 Analysis of the Result

The analysis of the result of the simple elastic model column structure trapped by a step load and lying on a quadratic-cubic foundation is hereby presented. The dynamic buckling load decreases with increased imperfection amplitudes as can be clearly seen in Table 1 and Figure 1. This is equivalent to saying that, the nearer the structure is to a perfect nature, the more stable it is for a step load. In the same token, it can be clearly observed that there is decrease in the dynamic buckling load with increase in the damping parameters, as can be clearly seen in Table 2 and Figure 2.

5.0 Conclusion

The research discussed the analysis of the dynamic buckling of a clamped finite imperfect viscously damped column that is subjected to a step load lying on a quadratic-cubic elastic foundation, using the methods of asymptotics and perturbation technique. The formulation of the governing equation contains δ and ϵ independent parameters which are used in asymptotic expansions of the relevant variables. Through the research, two main conclusions are obtained: the dynamic buckling load decreases with increased imperfections, the dynamic buckling load decreases with increase in damping.

The perturbation and asymptotic techniques applied in this work made it possible to change ordinary differential equations to partial differential equations. This method turns ordinary differential equations into partial differential equations to solve them. This method is not limited to the elastic model structure, but can also be used in other structural forms such as cylindrical shells, plates, etc.

REFERENCES

- [1] Budiansky, B. (1966); Dynamic buckling of elastic structures; criteria and estimates in, *Dynamic Stability of Structures*, Pergamon, New York. 350-357.
- [2] Hutchinson, J.W. and Budiansky, B (1966), Dynamic buckling estimates, *A.I.A.A.J.* 4, 525-530.
- [3] Eglitis, E.(2011); Dynamic buckling of composite shells. Ph.D. dissertation at Riga Technical University.
- [4] Kowal-Michalska, K. (2010); About some important parameters in dynamic buckling analysis of plated structures subjected to pulse loading, *Mechanics and Mechanical Engineering*, 14, (2), 269-279.
- [5] Ferri, E., Antinucci, E., He, M.Y., Hutchinson, J.W., Zok, F.W. and Evans, A.G. (2006); Dynamic buckling of impulsively loaded prismatic cores, *J. of Mechanics of materials and Structures*, 1, (8), 1345-1365 .
- [6] Kubiak, T. (2005); Dyanmic buckling of thin-walled composite plates with varying widthwise material properties, *Int. J. of Solids and Structures*, 42, 555-567.
- [7] Kubiak, T. (2007);Interactive buckling of thin-walled columns “in Polish”, *Scientific Bulletin of Lodz Technical University, Lodz*.
- [8] Mania, R.J. (2010),Dynamic buckling of thin-walled viscoplastic columns, (in Polish), *Scientific Bulletin of Lodz Technical University, Lodz*.
- [9] Kowal-Michalska, K. and Mania, R. (2008) some aspects of dynamic buckling of plates under in-plane compression, *Mechanics and Mechanical Engineering*, 12,(8), 135-146.
- [10] Beylaev, A.K, Slim, D.N and Morozov, N.F. (2013); stability of transverse vibration of rod under longitudinal step-wise loading, *J. of Physics, conference series*, 451, 1-6.
- [11] Vo, T.P., Lee, J. and Lee, K. (2010); On triply coupled vibrations of axially loaded thin-walled composite beams, *Composite and Structures*, 88, 144-154.
- [12] Capiet-Learnout, E., Soize, C. and Mignolet, M.-P. (2003); Computational nonlinear stochastic dynamics with model uncertainties and nonstationary stochastic excitation, 11th International conference on structural safety and reliability, New York.
- [13] Capiez-Lernout, E. Soize, C. and Mignolef M.P. (2013); Nonlinear stochastic dynamical post buckling analysis of uncertain cylindrical shells, 11th International conference on Recent Advances in Structural Dynamics, RASD, 2013, Pisa, Italy.
- [14] Chitra, V. and Priyadasini, R.S. (2013); Dynamic buckling of composite cylindrical shells subjected to axial impulse, *International Journal of Scientific and Engineering Research*, 4(5),162-165.
- [15] Priyadarasini, R.S., Kalyangraman, V. and Srinvasam, S.M. (2012); Numerical and experimental stability of buckling of advanced fibre composite cylinders under axial compression *Int. J. of Structural Stability and Dynamics* 12, (4), 1-25.
- [16] Mcshane, G.J., Pingle, S.M., Deshpande, V.S. and Fleck, N.A (2012); Dynamic buckling of an inclined strut, *Int. J. of Solids and Structures*. 49, 2830-2838.
- [17] Reda, A.M. and Forbes, G.L. (2012); Investigation into the dynamic effect of lateral buckling of high temperature/ high pressure offshore pipelines, *proceedings of Acoustics*, paper 83, Australia.
- [18] Gladden, J.R., Hamdy, N.z., Belmonte, A. and Villermaux, E. (2005); Dynamic buckling and fragmentation in brittle rods, *Physical Review*, 94, 035503, 1-4.
- [19] Lei, Y. Adhikari, S. and Friswell, M.I. (2013); Vibration of nonlocal Kelvin-Viogt viscoelastic damped Timoshenko beams, *Int. J. of Engineering Science*, 66 – 67, 1-13.
- [20] Salahi, M. and Safi-Djahanshahi, A. (2010); Nonlinear Analysis of viscoelastic rectangular plates subjected to in-plane compression, *J. of Mechanical Research and Applications*, 2, (1), 11-21.
- [21] Ette, A.M., Chukwuchekwa, J.U., Osuji, W.I., Udo-Akpan I.U. and Ozoigbo, G.E.(2018); On the asymptotic analysis of the static buckling of infinitely long and harmonically imperfect column lying on quadratic-cubic elastic foundations, *Asian Journal of Mathematics and Computer Research* , 23 (4), 220-230.
- [22] Ette, A.M., Chukwuchekwa, J. U. and Udo-Akpan I.U. (2016); On the buckling of a clamped viscosly damped column trapped by a step load, *IJASM Vol.3, Issue3*, 117-123.
- [23] Amazigo. J.C and Frank, D.(1973);Dynamic buckling of an imperfect column on a nonlinear foundation, *Quart. Appl. Math.*, 31(6),1-9.
- [24] Amazigo, J.C and Ette, A.M. (1987); On a two-small nonlinear parameter differential equation with application to dynamic buckling, *J. of Nigeria Mathematics Society* 6, 91-102 .

- [25] Chukwuchekwa, J. U. and Ette, A. M. (2015). Asymptotic Analysis of an Improved Quadratic Model Structure subjected to Static loading. *Journal of the Nigerian Association of Mathematical Physics*, Vol. 32, 237 – 244.
- [26] Ette, A. M., Chukwuchekwa, J. U., W. I. Osuji, Udo-Akpan , I. U. and Ozoigbo, G. E. (2018). Asymptotic Investigation Of The Buckling Of A Cubic – Quintic Nonlinear Elastic Model Structure Stressed By Static Load And A Dynamic Step Load, *IOSR Journal of Mathematics (IOSR-JM)*, Volume 14, Issue 1 Ver. II (Jan. - Feb. 2018), PP 16-30.
- [27] Ette, A. M., Chukwuchekwa, J. U., W. I. Osuji, Udo-Akpan , I. U. and Ozoigbo, G. E. (2018). On The Asymptotic Analysis Of The Static Buckling Of Infinitely Long And Harmonically Imperfect Column Lying On Quadratic – Cubic Elastic Foundations. *Asian Journal of Mathematics and Computer Research*, 23(4): 220-230.
- [28] Ette, A. M., Chukwuchekwa, J. U., Ozoigbo G. E. and Udo-Akpan, I. U. (2017). A Two – Small – Parameter Dynamic Buckling Of A Pre – Statically Loaded Quadratic Model Structure With Nonlinear And Light Linear Damping But Struck By A Step Load. *Bulletin Of Mathematics And Statistics Research*, Vol. 5, Issue 4, pp. 88 – 107.
- [29] **A. M. Ette, J. U. Chukwuchekwa, I. U. Udo-Akpan and W. I. Osuji** (2019). Asymptotic Analysis of the Static and Dynamic Buckling of a Column with Cubic - Quintic Nonlinearity Stressed by a Step Load. *Journal of Advances in Mathematics and Computer Science*, 30(6): 1-35, Article no.JAMCS.46862