

# **Investigation of extent of Interaction of Thiobarbituric acid and two selected Amino acids with Three Divalent Metal ions in Aqueous-organic medium at different Temperatures**

## **Abstract**

The formation constants of binary and ternary complexes of Thiobarbituric acid as primary ligand and L-tyrosine and L-histidine as secondary ligands have been examined in 40% (v/v) ethanol-water mixture at 27°C and 35°C and at ionic strength of 0.02NaNO<sub>3</sub> by potentiometric method. The ligands formed 1:1binary complexes with the metal ions. The primary and secondary ligands simultaneously coordinated to the metal ions to form 1:1:1ternary complexes. The difference in stability constants of binary and ternary complexes were determined by  $\Delta \log K$  and RS%. Ternary complexes exhibited enhanced stability than the binary complexes. The stability of the complexes decrease with increase in temperature. The thermodynamic parameters such as Gibb's free energy change ( $\Delta G$ ), entropy change ( $\Delta S$ ) and enthalpy change ( $\Delta H$ ) accompany the interactions were evaluated. The interactions were found to be spontaneous, exothermic, and entropically favoured.

**Keyword:** Stability constants, Ternary complexes, binary complexes, Thermodynamic parameters

## **Introduction**

The study of coordination compounds in solution has provided prospects for using ligands and metal complexes in medicine, agriculture and industries(1-4). The study of complexes containing mixed ligands of amino acids and heteraromatic nitrogen base ligands with metal ions has been considered as a model for interaction between bioligands and metal ions occurring in the body system[5,6]. The knowledge of binding properties of ligands with metal ions in solution has been used as tool by chemists for quantitative and qualitative analysis of metals. Some first row transition metals are such as Co, Cu, Fe, Mn, V, and Zn are essential in sufficient quantity for

normal functioning of body system due to their existence in proteins and excess of them could result to metal poisoning while some metals like Pb, As, Hg and Cd are toxic when presence in body system[7]. Chelating agents have been applied as therapeutic chelators in transportation of metal to target sites, detoxification and demetallization but most of them exhibit slow kinetic of release and toxic which may also contribute to problem of contamination[8]. As a result of these problems, researchers have been studying chelation properties of bioligands with different metal ions under different experimental conditions with a view to finding chelators that are harmless and exhibiting ease of release. Literature survey reveals paucity of work on interaction mixed-ligand of amino acids(L-tyrosine and L-histidine) and thiobarbituric acid with metal ions in aqueous-organic medium at different temperatures. This work therefore reports interactions of mixed-ligand of L-tyrosine, L-histidine and thiobarbituric acid with metal ions in ethanol-water medium with a view to providing data on their binding properties in forming binary and ternary complexes for various biological applications.

## **Experimental**

### Materials

All the chemicals used in present study were of analytical grade and were used without further purification. They include Cobalt(II)nitrate hexahydrate, copper(II)nitrate hexahydrate, lead (II) dihydrate, sodium nitrate, nitric acid, sodium hydroxide, double distilled water, ethanol, oxalic acid, methyl orange indicator, EDTA, L-tyrosine, L-histidine and Thiobarbituric acid. Metal ion solutions ( $0.002 \text{ mol dm}^{-3}$ ) and ligand solutions ( $0.002 \text{ mol dm}^{-3}$ ) were prepared from metal nitrate and the ligands in double distilled water. The concentrations of metal solutions were standardized by titrating against EDTA solution. The stock solutions of  $0.01 \text{ mol dm}^{-3} \text{ HNO}_3$  was prepared by measuring and dissolving amounts of the acid in appropriate volume  $\text{CO}_2$ -free double distilled water. The concentration of the acid was standardized by titrating against NaOH solution. Solution of  $0.134 \text{ mol dm}^{-3}$  of  $\text{CO}_2$ -free NaOH solution was prepared and standardized with oxalic acid solution. The ionic strength of each solution was adjusted to  $0.02 \text{ mol dm}^{-3}$  by addition of  $\text{NaNO}_3$

### Apparatus

Potentiometric titrations were carried out with Mettler Toledo(FP20 Standard) meter with accuracy  $\pm 0.03$  units using combined electrode assembly. The pH meter was calibrated against standard buffers solutions (pH 4.00, 6.86 and 9.18). All the titrations were carried out within the slope of 99-105%. The temperature was maintained constant at  $27^{\circ}\text{C}(\pm 0.1)$  and  $35^{\circ}\text{C}(\pm 0.1)$  throughout the titrations. All the titrations were carried out in multiples. The pH measurements are corrected according to Van Uitert and Hass procedure[9].

### Method

For determination of proton-ligand stability constants and stability constants of binary and ternary complexes in 40 % ( v/v) ethanol-water, the following sets of solution were prepared and the volume was made up to 40ml and titrated against  $\text{CO}_2$ -free NaOH solution.

- (1) Free  $\text{HNO}_3$ (A)
- (2) Free  $\text{HNO}_3$  + Ligand Thiobarbituric acid
- (3) Free  $\text{HNO}_3$ + Ligand Thiobarbituric acid + Metal ion
- (4) Free  $\text{HNO}_3$  + Ligands (Ltyrosine or L-histidine)
- (5) Free  $\text{HNO}_3$  + Ligands (Ltyrosine or L-histidine) + Metal Ion
- (6) Free  $\text{HNO}_3$  + Ligand Thiobarbituric acid + Ligands (Ltyrosine or L-histidine) + Metal Ion

The obtained values of pH were plotted against volume of alkali added for above sets of titration and titration curves were obtained using origin 8 program. From the curves, the equilibria existing in solution were examined by comparing the curves. The protonation constants and the formation constants of binary complexes were determined using Point wise method by calculating the average number of metal ions associated with the ligand  $\overline{n_A}$  at different pH values, average number of ligand molecules attached per metal ion ( $\bar{n}$ ) and free ligand exponent (pL) and formation constants of ternary complexes were determined by using Ramamoorthy and Santapa method[10]. Distributions of species were obtained by Hyperquad Simulation and Speciation program. Thermodynamic parameters such as Gibb's free energy change ( $\Delta G$ ), entropy change ( $\Delta S$ ) and enthalpy change ( $\Delta H$ ) for deprotonation of the ligands and formation of complexes were determined.

## RESULTS AND DISCUSSION

### Nature of potentiometric curves

The curves for the binary complexes and ternary complexes of L-tyrosine, L-histidine and Thiobarbituric acid in 40% ethanol-water at 27°C are presented figure 1 for the purpose of brevity.




Figure 1. Potentiometric titration curves of Thiobarbituric acid(A) and L-Histidine(B) and their binary and ternary complexes in 40% ethanol-water at 27°C ( $I=0.02M NaNO_3$ )

It is observed from the figure above that b curve shifts from acid curve to higher and lower pH while c curve deviates to lower pH region. The divergences of the curves to higher pH and lower pH in relation to the acid curve for the same volume of alkali indicate deprotonation of the ligands [11]. Addition of metal ions to each of the ligand solutions results in deviations of curves, d,e,f,g,h and k from curves b&c to a lower pH value region. This observation suggests coordination of the ligand to the metal ions occurs through displacement of proton(s) from the ligands. The i, j and l curves are found to lie below metal - ligand curves of Thio ligand and this suggests the formation of the mixed-ligand complexes. It is also observed from the figure that the ternary complexes curves do not coincide with any of binary complex curve and this indicate simultaneous coordination of the two ligands to the metal ions in the complexes [12].

### Proton-ligand, Metal-ligand stability constants and effect of Temperature

From curves a, b&c in figure 1, the values of  $\overline{nA}$  (the average number of protons associated with the ligand) were determined by employing the relationship derived by Irving and Rossotti [12]. These  $\overline{nA}$  values were used for determination of proton-ligand stability constants of the ligands by pointwise method. Formation constants of the binary complexes were determined from ligands curves and metal-ligand curves. The average number of ligand molecules coordinated to the metal ion( $\overline{n}$ ) and free ligand concentration(pL) were evaluated and from obtained values, metal-ligand stability constants were obtained by pointwise method in ethanol-water 40% medium[12]. Proton-ligands stability constants of the ligands and stability constants of binary complexes and their thermodynamic parameters are presented in tables 1& 2 respectively.

**Table 1. Proton-ligand stability constants of the ligands at 27°C and 35°C(I=0.02MNaNO<sub>3</sub>) and thermodynamic parameters**

Lligand	27°C		35°C		ΔG(KJ/mol)		ΔH(KJ/mol)	ΔS(J/molK)
	LogK <sub>1</sub> <sup>H</sup>	LogK <sub>2</sub> <sup>H</sup>	LogK <sub>1</sub> <sup>H</sup>	LogK <sub>2</sub> <sup>H</sup>	300K	308K		
L-His	4.5944	11.0651	3.3314	10.7158	-90	-83	-357	-889
L-Tyr	3.6802	11.3279	-	11.1145	-86	-66	-861	-2582
Thio	-	11.9198	-	11.4935	-685	-68	-94	-86

**Table 2. Stability constants of binary complexes at 27°C and 35°C ( $I=0.002MNaNO_3$ ) and thermodynamic parameters.**

Complex	LogK	300K	308K	$\Delta G(KJ/mol)$	$\Delta H(KJ/mol)$	$\Delta S(J/molK)$
L-HisCu(II)	$LogK_{CuHis}^{Cu}$	7.6744	6.5231	-44	-254	-702
L-His-Co(II)	$LogK_{CoHis}^{Co}$	6.5724	6.4519	-38	-27	-137
L-His-Pb(II)	$LogK_{PbHis}^{Pb}$	7.8574	6.5585	-45	-287	-807
L-TyrCu(II)	$LogK_{CuTyr}^{Cu}$	7.5232	7.4981	-43	-6	+125
L-Tyr-Co(II)	$LogK_{CoTyr}^{Co}$	6.7419	5.9816	-39	-168	-431
L-Tyr-Pb(II)	$LogK_{PbTyr}^{Pb}$	7.2704	6.7085	-42	-124	-275
ThioCu(II)	$LogK_{CuThio}^{Cu}$	9.0645	7.5790	-52	-329	-921
Thio-Co(II)	$LogK_{CoThio}^{Co}$	7.2557	6.4497	-42	-178	-455
Thio-Pb(II)	$LogK_{PbThio}^{Pb}$	8.3048	7.4314	-48	-193	-485

Table 1 shows that dissociation of two protons and one occurs in secondary ligands(L-His and L-tyr) and primary ligand(Thio) respectively at 27°C while at 35°C, two protons are dissociated from L-His ligand and one from both Thio and L-Tyr ligands. The mean  $LogK_2^H$  value of 11.9198 at 27°C and  $LogK_2^H$  of 11.4935 at 35° C of Thio ligand are attributed to deprotonation of protons at  $N_3H$  group present in the ligand [13]. The mean values of  $LogK_1^H$  equals to 3.6802 and  $LogK_2^H$  equals to 11.3279 of L-Tyr at 27°C are attributed to deprotonation at COOH group and amine ( $NH_2$ ) group of the ligand respectively. L-His also shows two dissociation constants at mean  $LogK_1^H$  value of 4.5944 and  $LogK_2^H$  value of 11.0651 of L-His at 27°C which are attributed to deprotonation at COOH group and amine ( $NH_2$ ) group of the ligand respectively [14,15]. Only on proton is observed to be deprotonated by L-Tyro at 35°C with deprotonation constant of  $LogK_2^H$  equals to 11.1145 which is ascribed to deprotonation of amine ( $NH_2$ ) group of the ligand[14]. The proton-ligand stability constant values decrease with increase in temperature showing increase in acid strength with temperature. The negative values of  $\Delta G$  and  $\Delta H$  respectively reveal spontaneous and exothermic deprotonation processes undergone by the ligands. The negative values of  $\Delta S$  obtained are attributed to binding of greater amount of

solvent molecules to the deprotonated ligand than the original amount associated with undepronated form[16].

All the ligands formed 1:1 binary complexes and this is confirmed by the inflection points and  $\bar{n}$  evaluated average number of ligand bound per metal in range of 0.1-0.5(17). From table 2, Thio and L-Tyr interact with Cu(II), Co(II) and Pb(II) at 27°C and 35°C with stabilities in order of  $\text{LogK}_{\text{CuTyr}}^{\text{Cu}} > \text{LogK}_{\text{PbTyr}}^{\text{Pb}} > \text{LogK}_{\text{CoTyr}}^{\text{Co}}$  and  $\text{LogK}_{\text{CuThio}}^{\text{Cu}} > \text{LogK}_{\text{PbThio}}^{\text{Pb}} > \text{LogK}_{\text{CoThio}}^{\text{Co}}$ . This order is assigned to polarizability and covalent-bonding effect of the metal ions. Extra stability exhibited by Cu(II) might be due to the Jahn Teller distortion due unique configuration of the metal ion(18,19). L-His ligand interacts with the metal ions at both temperatures to give order of  $\text{LogK}_{\text{PbHis}}^{\text{Pb}} > \text{LogK}_{\text{CuHis}}^{\text{Cu}} > \text{LogK}_{\text{CoHis}}^{\text{Co}}$ . The order could be attributed to covalent-bonding index and polarizability of the metals. The ligands give the order of  $\text{LogK}_{\text{CuThio}}^{\text{Cu}} > \text{LogK}_{\text{CuHis}}^{\text{Cu}} > \text{LogK}_{\text{CuTyr}}^{\text{Cu}}$  at 27°C and 35°C which could be ascribed to the nature of donor atoms of the ligands [20,21]. The order of  $\text{LogK}_{\text{CoThio}}^{\text{Co}} > \text{LogK}_{\text{CoTyr}}^{\text{Co}} > \text{LogK}_{\text{CoHis}}^{\text{Co}}$  is observed and attributed to type of donor atoms involved in the coordination, polizability and basicity of the ligand [20,22] while the order of  $\text{LogK}_{\text{CoHis}}^{\text{Co}} > \text{LogK}_{\text{CoThio}}^{\text{Co}} > \text{LogK}_{\text{CoTyr}}^{\text{Co}}$  at 35°C is attributed to basicity of the ligands. The order of  $\text{LogK}_{\text{PbThio}}^{\text{Pb}} > \text{LogK}_{\text{PbHis}}^{\text{Pb}} > \text{LogK}_{\text{PbTyr}}^{\text{Pb}}$  at 27°C is attributed to basicity of ligands and involvement of S-donor atom in Thio ligand while the order of  $\text{LogK}_{\text{PbThio}}^{\text{Pb}} > \text{LogK}_{\text{PbTyr}}^{\text{Pb}} > \text{LogK}_{\text{PbHis}}^{\text{Pb}}$  at 35°C is assigned to coordinating atoms and polarizability of the ligands[23,24].

In all the complexes, decrease in metal-ligand stability constant (logK) was observed with increase in temperature. The negative  $\Delta G$  values indicate spontaneous complexation of the ligands to the metal ions. It is reported that the divalent metal ions exist in solution as octahedrally hydrated species [16] and the values of  $\Delta H$  and  $\Delta S$  can be determined by considering the combined effects of (a) dissociation of water molecules and (b) metal-ligand bond formation. Negative enthalpy change ( $\Delta H$ ) obtained for the interactions between the ligands and the metals implying exothermic complexation reactions and the reactions are favourable at lower temperature. **The metal-ligand binding process is enthalpy driven and metal-ligand association is fairly strong [25].** Negative values of  $\Delta S$  observed in all the binary complexes indicate that complexation reactions are entrophy unfavourable due to the large amount of coordinating solvent around the ligands while positive  $\Delta S$  observed in L-His-Pb(II) complex implies entropy favoured association between the ligand and the metal[26].

### Stability constants of ternary complexes and thermodynamic parameters

Stability constants of ternary complexes of Thiobarbituric acid as primary ligand and L-Histidine and L-Tyrosine as secondary ligands with Cu(II), Co(II) and Pb(II) ions in 40%(v/v) ethanol-water at 27°C and 35°C are evaluated from the formation curves. The stabilities of the ternary complexes with respects to the binary complexes are quantified by  $\Delta \log K$  and RS%. The stability constants and the thermodynamic parameters are presented in table 3-7.

**Table 3: Stability constants of ternary complexes of L-tyrosine and Thiobarbituric acid in 40% (v/v) ethanol-water with I = 0.02 M (NaNO<sub>3</sub>) at 27°C**

Complex	$\text{Log}K_{\text{MAB}}^{\text{M}}$	$\text{Log}K_{\text{MA}}^{\text{M}}$	$\text{Log}K_{\text{MAB}}^{\text{MA}}$	$\text{Log}K_{\text{MB}}^{\text{M}}$	$\Delta \log K$	RS%
[CuThioTyr]	15.6108	9.0645	6.5463	7.5232	-0.9797	-0.2778
[CoThioTyr]	15.2378	7.2557	7.9821	6.7419	1.2402	10.0114
[PbThioTyr]	15.9948	8.3048	7.6901	7.2704	0.4196	-7.4017

**Table 4: Stability constants of ternary complexes of L-tyrosine and Thiobarbituric acid in ethanol-water 40-60% with I = 0.02 M (NaNO<sub>3</sub>) at 35°C**

Complex	$\text{Log}K_{\text{MAB}}^{\text{MA}}$	$\text{Log}K_{\text{MAB}}^{\text{M}}$	$\text{Log}K_{\text{MA}}^{\text{M}}$	$\text{Log}K_{\text{MB}}^{\text{M}}$	$\Delta \log K$	RS%
[CuThioTyr]	15.1953	7.5790	7.6163	7.4981	0.1183	0.4921
[CoThioTyr]	14.0300	6.4497	7.5803	5.9816	1.5987	17.4434
[PbThioTyr]	15.4782	7.4314	8.0468	6.7034	1.3434	8.2743

**Table 5: Stability constants of ternary complexes of L-histidine and Thiobarbituric acid in ethanol-water40-60% with I = 0.02 M (NaNO<sub>3</sub>) at 27°C**

Complex	$\text{Log}K_{\text{MAB}}^{\text{M}}$	$\text{Log}K_{\text{MA}}^{\text{M}}$	$\text{Log}K_{\text{MAB}}^{\text{MA}}$	$\text{Log}K_{\text{MB}}^{\text{M}}$	$\Delta\text{log}K$	RS%
[CuThioHis]	17.6925	9.0645	8.6280	7.6744	0.9536	-4.8154
[CoThioHis]	15.2314	7.2557	7.9757	6.5724	0.6833	10.2247
[PbThioHis]	16.2401	8.3048	7.9353	7.8574	1.7723	-4.4492

**Table 6: Stability constants of ternary complexes of L-histidine and Thiobarbituric acid in ethanol-water40-60% with I = 0.02 M (NaNO<sub>3</sub>) at 35°C**

Complex	$\text{Log}K_{\text{MAB}}^{\text{M}}$	$\text{Log}K_{\text{MA}}^{\text{M}}$	$\text{Log}K_{\text{MAB}}^{\text{MA}}$	$\text{Log}K_{\text{MB}}^{\text{M}}$	$\Delta\text{log}K$	RS%
[CuThioHis]	16.1824	7.5790	8.6034	6.5231	2.0803	13.5163
[CoThioHis]	14.8441	6.4497	8.3944	6.4519	1.9445	30.1518
[PbThioHis]	15.7759	7.4314	8.3445	6.5585	1.7860	12.2871

**Table 7: Thermodynamic parameters of ternary complexes of L-histidine and Thiobarbituric acid**

Complex	$\Delta G(\text{KJ/mol})$		$\Delta H(\text{KJ/mol})$	$\Delta S(\text{J/mol})$	
	300K	308K		300K	308K
[CuThioHis]	-102	-95	-334	-774	-774
[CoThioHis]	-87	-88	-86	6	6
[PbThioHis]	-93	-93	-103	-31	-31
[CuThioTyr]	-90	-90	-92	-7	-7
[CoThioTyr]	-88	-82	-267	-599	-599
[PbThioTyr]	-92	-91	-114	-75	-75

In the tables  $\Delta \log K$  was found to be positive for all the complexes except Cu(II) ternary complex of L-tyrosine and Thiobarbituric acid at 27°C. The positive  $\Delta \log K$  reveals more stability of the complexes than their corresponding binary complexes. The positive value of RS% affirmed the existence of enhanced stability. The enhanced stability could be attributed to intramolecular aromatic-ring stacking and hydrogen bond while negative  $\Delta \log K$  implies less stability of the ternary complexes than the binary complex of secondary ligand[27]. The stability is also attributed to increase in polarity as a result of redistribution of the electron densities of the metal-ligand bond in ternary therefore mixed ligand complexes are not easily hydrolysed[28,29]. Co(II) ternary complexes exhibited more stability than the corresponding binary complexes among the ternary complexes and this is established by the values of  $\Delta \log K$  and RS%. The decrease in stability values of the complexes with increase in temperature as shown in the tables indicates that complexation of the ligands is favourable at lower temperature. A negative  $\Delta G$  values for all the complexes suggest spontaneous complexation reaction. Negative values of  $\Delta H$  show that the chelation processes are exothermic and favorable at very low temperature[11]. In all the ternary complexes of L-histidine, the order of stability is found to be Cu(II)>Pb(II)>Co(II). This order could be attributed to tetragonal distortion of Cu(II) geometry which gives extra stability to the metal complexes[30]. In all the ternary complexes of L-tyrosine, the stability is in order of Pb(II) > Cu(II) > Co(II) and the extra stability exhibited by Pb(II) complexes could be attributed to increase in covalent –bonding index, polarizability of the metal ions and polarizability of ligand. At both temperatures, all the ternary complexes of L-histidine, except Co(II) ternary complex at 27°C, are more stable than the ternary complexes of L-tyrosine. The more stability of the ternary complexes is attributed to coordination of the ligand through nitrogen donor atoms of imidazole and amino groups[24].

## Conclusion

Interaction of L-histidine, L-tyrosine and Thiobarbituric acid with Cu(II), Co(II) and Pb(II) ions in % (v/v) ethanol–water mixture at different temperatures have been determined by potentiometric method. The metal ions formed binary (1:1) and (1:1:1) ternary complexes with the ligands. The stability of the complexes is affected by basicity of the ligand, nature of donor

atoms, polarizability of ligands, covalent-bonding index and polarizability of the metal ions. The stability of the ternary complexes with respect to binary complexes was determined by  $\Delta \log K$  and RS%. The positive values of  $\Delta \log K$  and high values of RS% affirmed more stability of the ternary complex than the corresponding binary complex while negative value of  $\Delta \log K$  established less stability of ternary complex than the binary complex. The deprotonation constants of the ligands and metal-ligand stability constants decrease with increase in temperature. Thermodynamic evaluation revealed spontaneous, exothermic and entropically favoured and unfavoured complexation process. Thiobarbituric acid and L-histidine ligands showed high affinity for Pb(II) ion while L-tyrosine exhibited high affinity for Cu(II) ion. The mixed-ligand of L-histidine and Thiobarbituric acid showed high affinity for Pb(II) ion while mixed ligand of L-tyrosine and Thiobarbituric acid showed high affinity for Cu(II) ion. The individual ligand and their mixture can be used for extraction, transportation and detoxification of metals.

## References.

- [1]. Martell, A. E. and Hancock, R. D. *Metal Complexes in Aqueous Solutions*, New York: Plenum Press, 1996.
- [2]. Martell, A. E. and Motekaitis, R. J. *The Determination and Use of Stability Constants*, New York: VCH Publishers. 1988.
- [3]. Hartley, F. R., Burgess, C. and Alcock, R. *Solution Equilibria*, New York: Ellis Horwood, 1980.
- [4]. Berthon, G. Ed., *Handbook of Metal-Ligand Interaction in Biological fluids*, 1995, 2, p.1206.
- [5]. Khalil, M. M.; Radalla, A. M. Binary and Ternary Complexes of Inosine, *Talanta*. 1998; (46), 53-61.
- [6]. Khalil, M. M. (2000). Complexation Equilibria and Determination of Stability Constants of Binary and Ternary Complexes with Ribonucleotides (AMP, ADP, and ATP) and Salicylhydroxamic Acid as Ligands, *J. Chem. Eng. Data*. 2000; 45(1):70-74
- [7]. Ondrej, Z., Marketa, R., Jaromir, H., Tomas, E., Vojtech, A. and Rene, K. From Amino Acids to Proteins as Targets for Metal-based Drugs, *Current Drug Metabolism*. 2012; 13, 306-320.
- [8]. Lisa, M., Jacqueline, L. S. and James, A. H. Immobilized peptides/amino acids on solid supports for metal remediation, *Pure Appl. Chem*. 2004; 76(4):777-787
- [9]. Narendra, P., Sudarshan, R. D., Madhava, R. P. and Satyanarayana, B. Quantitative equilibrium studies on binary complexes of 3-hydroxy quinazolin-4-one with inner transition metal ions, *Res J Chem Envir.* 2003; 7(2): 43-47.
- [10]. Janrao, D. M., Jamil P., Kayandeb, D. D. and Jabber, J. M. An overview of potentiometric determination of stability constants of metal complexes, *Sci. Revs. Cchem. Commun*. 2014; 4(1):11-24
- [11]. Jeyaprakash, D., Paramasivam, S., Thirugnanasamy, E., Sutha, S. and Saravanan, R. Complexation Equilibria and Coordination Aspect of Zn(II) Complexes Contain 2 Aminobenzamide and some Bioactive Amino acid Mixed Ligands pH-metric, Spectroscopic and Thermodynamic Studies, *Acta Chimica Slovica*. 2014; 61, 803-812.
- [12]. Abdulbaset, A. Z., Mazahar, F. B. and Janrao, D. M. Study of stability constant of biological active molecules (drug) using potentiometric titration technique, *Journal of Chemical, Biological and Physical Sciences*. 2011; 2(1):67-80.
- [13]. Naciye, T. and Suat, A. M. Complex Formation of Nickel(II) and Copper(II) with Barbituric Acid, *International Journal of Analytical chemistry*. 2014; 2014. 1-5.

- [14]. Siham, A. L. and Reda, A. A. Mixed-ligand complex formation of cu(ii) with 1,2diphenylethylenediamine as primary ligand and amino acids as secondary ligands, *International Journal of Electrochemical Science*.2012;7, 7699 – 7711.
- [15]. Jahangir, M. A., Ullah, M.R. and Mousume, A. (2018). Determination of stability constant of mixed ligand complexes of Ni(II) and Cu(II) metal ions with Diethylenetriamine (DET) and Ethylenediamine(en), Glycine (gly), alpha-Alanine (alpha-ala), Phenylalanine (ph-ala). Oxalic acid (ox), Tyrosine (tyr), *Journal of Ultra Scientist of Physical sciences*.2018;30(2):12-19.
- [16]. Ahmed A. E., Mohamed R. S., Mohamed, M. S. and Mohammad, H. Bar. Thermodynamic Investigation and Mixed Ligand Complex Formation of 1,4-Bis-(3-aminopropyl)-piperazine and Biorelevant Ligands, *Bioinorganic Chemistry and Application*, 2012, 1-10.
- [17]. Nieboer, E. and Richardson, D.H.S.. The replacement of the nondescript term 'heavy metals' by a biologically and chemically significant classification of metal ions, *Environmental Pollution Series B, Chemical and Physical*. 1980;1(1): 3-26.
- [18]. Atkins, P W and Shriver, D.F. Inorganic Chemistry, 5th ed. Oxford; Oxford University Press. 2010.
- [19]. Grisser, R. and Sigel, H. Ternary complexes in solution-VII. *Inorg. chem*. 1979; 9, 1238-1246
- [20]. Jain, A. K. and Khan, F.. Polarographic study of mixed-ligand complexes of cadmium(II) with l-amino-acid and vitamin b-5. *Indian chem. soc.* 1998;75, 31-32.
- [21]. Martell, A. E. and Sillen, I. G. *Stability constants of metal ion complexes*. London: The Chemical society, 1971.
- [22] Salvatore, M., Andrea, A., Arcangelo, M. and Angelo, V. Electronic and vibrational polarizabilities of the twenty naturally occurring amino acids, *Biophysical Chemistry, Elsevier*. 2007; 132,(2-3):139-145.
- [23]. Amal, A.M. and Asma, A.A. Characterization by Potentiometric Procedures of the Stability Constants of the Binary and Ternary Complexes of Cu(II) and Duloxetine Drug with Amino Acids, *Journal of Chemistry*, 2019;1-13.
- [24]. Singh, S.V., Thakur, M. F. and Naikwad, S.D. Formation of Transition Metal Complexes Carrying Medicinal Drugs: Thermodynamic Study. *J. Chem. Bio and Phy. Sci.*, 2014;4(1): 1-7.
- [25]. Promila, D.T. and Lonibala, R.. Potentiometric Studies on the Complexation Reactions of N-(2-[1 (3-aminophenyl)ethylidene] hydrazino-2 oxoethyl) benzamide with Ni<sup>2+</sup>, Cu<sup>2+</sup>, and Cd<sup>2+</sup> ions in Aqueous-dioxane and Micellar media, *Journal of Chemical and Engineering data*. 2010;55, 1166-1172.

[26]. Usama, E., Kenawy, I.M. and Abu, E.Y.G. Synthesis, thermal and spectral studies of first-row transition metal complexes with Girard-T reagent-based ligand, *Journal of Molecular Structure*.2007;871, (3):14-23

[27]. El-Haty, M.T., Amrallah, A.H., Mahmoud, R.A. and Ibrahim, A.A.. PH-metric studies on ternary metal complexes of some amino acids and benzimidazole, *Talanta*.1995;42(11):1711-1717

[28]. Jyotsna, G., Renu, N. and Dwivedi, K. (2015). Ternary Complexes of Cd(II) and Gd (III) involving biologically important ligands, *International Journal of Theoretical & Applied Science*.2015;7(1):22-34.

[29]. Ayser, A.A., Mohammed, R.S., Mohammed, S. and Saied, H. Coordination properties of dehydroacetic acid - Binary and ternary complexes, *Journal of coordination chemistry*. 2008;61(12):1906-1916.

UNDER PEER REVIEW