

1 **PRELIMINARY PHYTOCHEMICAL ANALYSIS AND *IN VITRO* ANTIMICROBIAL**  
2 **STUDY OF THE ROOT AND STEM BARK EXTRACTS OF *FICUS SYCOMORUS***  
3 ***LINN.***

4 I.Toma<sup>1\*</sup>, D. Dahiru<sup>2</sup>, and M.A. Madusolumou<sup>2</sup>

5 <sup>1</sup>Department of Biochemistry, Adamawa State University Mubi, Nigeria.

6 <sup>2</sup>Department of Biochemistry, Modibbo Adama University of Technology, Yola, Nigeria

7  
8  
9 \* Correspondence Author: dalitoma2014@gmail.com

10  
11 **Abstract**

12 This study was conducted to carryout preliminary phytochemical analysis and *in vitro*  
13 antimicrobial activities of aqueous and ethanolic root and stem bark extracts of *Ficus sycomorus*.  
14 Qualitative phytochemical analysis for tannins, saponin, terpenoids, flavonoids, alkaloids,  
15 glycosides, steroids, phenols, and reducing sugar was done using standard methods. The  
16 antimicrobial activities of the extracts were tested against four micro- organisms; *Escherichia*  
17 *coli*, *Staphylococcus aureus*, *Shigella dysentrae*, and *Salmonella typhi*. Agar well diffusion  
18 method was used for the antimicrobial studies. Phytochemical screening of both root and stem  
19 bark aqueous extracts showed the presence of tannin, saponin, terpenoid, flavonoid , alkaloids,  
20 glycoside, steroid, reducing sugar, and phenol. Glycoside was not detected in both the aqueous  
21 and ethanolic extracts of the root bark. The result of the antimicrobial studies showed that the  
22 aqueous root extract have higher antimicrobial activity ranging from (2-12 mm) on the tested  
23 microorganisms than aqueous stem bark extract (3-9 mm), while for ethanol extract both stem  
24 and root bark extract has almost the same effect or antimicrobial activity on the tested pathogens  
25 ranging from (2-15 mm) which is having higher activity compared to the aqueous extracts. The  
26 Minimum inhibitory concentration (MIC) and Minimum bactericidal concentration (MBC) of  
27 both the extracts were found to be 50 mg/mL and 100 mg/mL respectively. From this study, it  
28 can therefore be concluded that, the root and stem bark extract is a potential antimicrobial agents  
29 which support the claim of the traditional users of this plant in herbal medicine for the treatment  
30 of diseases that are of microbial origin.

31 **Key words:** *Ficus sycomorus*, *Escherichia coli*, *Staphylococcus aureus*, *Shigella dysentrae*,  
32 and *Salmonella typhi*, phytochemical screening, antimicrobial activity.

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63

## INTRODUCTION

Nature has been a source of medicinal agents for thousands of years and an impressive number of modern drugs have been isolated from natural sources [1]. At least 12000 of such compounds have been isolated so far, a number estimated to be less than 10% of the total[2,3]. Chemical compounds in plants mediate their effects on human body through processes identical to those we already understand for their chemical compound in conventional drugs in terms of how they work. This enables herbal medicines to have beneficial pharmacology, but also gives them the same potential as pharmaceutical drugs to cause side effects [2, 3].

Plant lives longer than every other living thing due to their ability to synthesize phytochemicals in their cells which serve as strong antifungal, antibacterial and antimicrobial agents, as a result of this, their susceptibility to diseases attack to some extent is low as compared to other living things (personal contact, 15<sup>th</sup>, January 2016).

In Nigeria like many African countries, several plants are still being used for the treatment of various ailments. Nigeria is naturally blessed with both savannah and tropical rainforest vegetation and these offer a wide distribution of plants believed to possess secondary metabolites which are responsible for treating or curing various diseases [4]. Quite a number of plants are used as medicines virtually in all cultures of the world. A good number of these medicinal plants are in common use in African traditional medicine. Most of the plants grow near houses and are easily overlooked, especially by urban dwellers [5].

This research work was carried out to study the Phytochemical screening and *in vitro* antimicrobial activities of root and stem bark extract of *Ficus sycomorus* on some selected microorganisms. *Ficus sycomorus* is a common savannah tree that grows or can be found almost everywhere. It is called in English Language as “Wild fig” “sycamore fig”, or common cluster fig. Spanish call it “sicomoro”. The Sukur people call it “Dashakwai”, Tiv people called it “Tur”, in Hausa it is known as “Baure”, Kilba and Marghi people called it “Kamda”, in Fali Language is called “Boduven” and Gude call it “Bodeva” (personal contact, 15<sup>th</sup>, January 2016). It grows in high water table areas, it can be found along water courses such as streams, rocky places, swamps and water holes[6]. The sycamore fig is sensitive to frost but can withstand some

64 cold. The relevance of this plant in traditional medicine is as a result of the secondary  
65 metabolites such as glycosides, reducing sugar, phenols, saponins, steroids, tannins, alkaloids,  
66 terpenoids and flavonoids which they have been screened to contain. Also referred to as  
67 phytochemicals, they are reported to possess inhibitory activities against the growth and disease  
68 inducing activities of some pathogenic microorganisms [7,8,9,10,11].

69 The root and stem-bark of *Ficus sycomorus* are said to be used as herb in Northern  
70 Nigeria for treatment of diseases like diarrhea, dysentery, cough, sore throat, chest diseases, and  
71 infertility and as antidote for snake. Therefore, this study was conducted to carry out the  
72 phytochemical screening and to evaluate antimicrobial activity of root and stem-bark of *Ficus*  
73 *sycomorus* in order to validate the claims of the traditional users of this plant.

## 74 MATERIALS AND METHODS

### 75 Sample Collection and identification of plant material.

76 Plant roots and stem-barks of the plant *Ficus Sycomorus* were collected from Sukur  
77 Kingdom in Madagali Local Government Area, Adamawa State, Nigeria. It was identified and  
78 authenticated by a Botanist from the Department of Biological Sciences, Adamawa State  
79 University, Mubi. A voucher number AD170023 was assigned. Sampling was carried out in the  
80 month of May from the tree.

### 81 Sample preparation

82 The root and Stem-barks (cut into small pieces) washed with water and rinsed with  
83 distilled water and then dried in the shade for two weeks. The dried samples was grinded by  
84 wooden mortar and pestle and sieve using clean Kitchen sieve to obtain a fine powder and was  
85 stored in a tight container until required for use.

### 86 Extraction

#### 87 Aqueous Extract

88 For the water extraction was done by cold maceration method according to the procedure  
89 described by [12, 13] with little modification. Two hundred grams (200 g) of each of the stem and  
90 root barks powder was weighed and soaked in 1000 mL of distilled water in a beaker for 48 h to  
91 obtain aqueous extracts. The aqueous extracts were filtered using sterile filter paper (Whatman  
92 No.1) into a clean conical flask. The filtrate was concentrated with a rotary evaporator. The  
93 extracts were then stored in a refrigerator.

94 Percentage yield was calculated as:  $\text{weight of extract} / \text{weight of dried powdered sample} \times 100$

95

## 96 **Preparation of ethanol extracts**

97 Maceration method of extraction as described by [12, 13] was adopted in this study. Two  
98 hundred grams (200g) each of the root and stem bark powdered material was weighed and  
99 soaked in 1000 mL of 70% ethanol and left for 24 h .Thereafter, it was decanted. The procedure  
100 was repeated with another 1000 mL to ensure complete extraction of the active ingredient .The  
101 extract was filtered and evaporated to dryness wit rotary evaporator. The dried extract was then  
102 weighed and stored in tightly closed bottle in a refrigerator until required.

103 Percentage yield was calculated as: weight of extract/ weight of dried powdered sample  $\times$  100

## 104 **Qualitative Phytochemical analysis.**

105 The qualitative phytochemical screening of the samples was carried out as described by [14, 15,  
106 16] ith slight modification. The root or stem bark extracts was screened for carbohydrates,  
107 alkaloids, flavonoids, steroids, phenols and tannins, saponin, glycosides, and proteins.

### 108 **Preparation of stock solution**

109 Two grams (2g) each of root or stem bark extracts were dissolved in 10 mL of water or ethanol  
110 to make a concentration of 200 mg/mL

### 111 **Test for Tannins**

112 One milliliter (1 mL) of the extracts was taken in a test tube and 2 mL of 5 % ferric chloride was  
113 added. Formation of blue –black, green or blue – green precipitate was taken as evidence for the  
114 presence of tannins.

### 115 **Test for Saponins**

116 One milliliter (1 mL) of the extracts was shaken with 5 mL of distilled water in a test tube for 5  
117 min. Frothing which persists on warming was taken as evidence for the presence of Saponins.

### 118 **Test for Terpenoids**

119 Five milliliters (5mL) of aqueous extract of each plant sample was mixed with 2mL of  $\text{CHCl}_3$  in  
120 a test tube and then 3mL of concentrated  $\text{H}_2\text{SO}_4$  was carefully added to the mixture to form a  
121 layer. An interface with a reddish brown coloration was considered as indication for the presence  
122 of terpenoids.

123

124 **Test for Flavonoids**

125 A little amount of magnesium powder and a few drops of concentrated hydrochloric acid were  
126 added to 3 mL of the extracts. A red or intense coloration indicated the presence of flavonoids.

127 **Test for Alkaloids**

128 To 2 mL of plant extracts, 2 mL of concentrated hydrochloric acid was added. The mixture was  
129 filtered and then 3 drops of Mayer's reagent was added. Presence of green colour or white  
130 precipitate indicated the presence of alkaloids.

131 **Test for glycosides**

132 Two milliliter (2 mL) of the extracts was hydrolyzed with HCl solution and neutralized with  
133 NaOH solution. A few drops of Fehling's solution A and B were added. Presence of red  
134 precipitate indicates the presence of glycosides.

135 **Test for Steroids (Salkowski's test)**

136 To 1 mL of plant extract, equal volume of chloroform and 3 drops of concentrated sulphuric acid  
137 was carefully added to form a lower layer. Formation of brown ring indicates the presence of  
138 steroids.

139 **Test for phenols**

140 Five drops of 10% ferric chloride was added to 1 mL of the extracts in a test tube. Formation of  
141 green or dirty green precipitate indicated the presence of phenols.

142 **Test for reducing Sugar**

143 To 2 mL of plant extract, 1 mL of Molisch reagent and 4 drops of concentrated sulphuric acid  
144 was added. Formation of purple or reddish ring indicates the presence of carbohydrates.

145

146 **Antimicrobial Analysis**

147 *Escherichia coli*, *Shigella dysenteriae*, *Salmonella typhi* and *Staphylococcus aureus* were used in  
148 this study. The microorganisms were obtained at the Microbiology Laboratory of Modibbo  
149 Adama University of Technology, (MAUTECH) Yola, Nigeria.

150 **Standardization of Isolates:**

151  
152 Test organisms were sub-cultured onto fresh plates of MacConkey agar and incubated  
153 aerobically at 37°C for 24 h. Colonies from these plates were suspended in Mueller- Hinton  
154 broth to a turbidity matching 0.5 McFarland standard (108cfu/ml). Mueller-Hinton agar was then  
155 used for antimicrobial assay. All the broth cultures were incubated at 37°C.

156

157

## 158 **Preparation of the Extract for Antimicrobial Study**

159

160 Two grams (2g) each of aqueous and ethanol root or stem bark extracts were separately  
161 dissolved in 10 mL of dimethylsulfoxide (DMSO) to obtain a concentration of 200mg/mL.

162 This was the initial concentration of each of the extracts used.

163

### 164 **Antimicrobial Test:**

165

166 The method described by the National committee for Clinical Laboratory Standard [17] was  
167 used.

168 Suspensions of the bacteria obtained contained approximately  $1 \times 10^8$ cfu/mL. Each labeled  
169 plate was uniformly seeded with a test organism by means of sterile swab stick rolled in the  
170 culture medium. Five wells, 4mm each in diameter were created using cork borer. Aliquots were  
171 dropped in each well to fullness at various concentrations of 100, 50, 25 and 12.5 mg/mL for  
172 both the root and stem bark extracts on different plates. Each plate was kept in the refrigerator  
173 for 1 hour to allow the extracts to diffuse into the culture medium while the immediate growth of  
174 the organism was stopped from taking place. These plates were then incubated at 37°C for 24 h.  
175 The zones of inhibition around the wells were measured in millimeter (mm). Control antibiotic  
176 (tetracycline capsule 100 µg/mL) was placed in a well on each plate along with the test extracts  
177 as control.

### 178 **Determination of the Minimum Inhibitory Concentration (MIC)**

179 The minimum inhibitory concentration of the extract was evaluated by the method described by  
180 [18].

181 The extract concentration were serially diluted with distilled water to various concentrations of  
182 100, 50, 25and 12.5mg/mL. The extract and the nutrient agar broth were mixed in the sterile test  
183 tube; the cultured medium was added to each test tube and incubated for 24hrs at 37°C .The  
184 lowest zones of inhibition for all the tested organisms showing no visible growth of **bacteria** was  
185 taken as the MIC.

### 186 **Minimum Bactericidal Concentration (MBC).**

187 The minimum bactericidal concentration (MBC) was determined after the minimum inhibitory  
188 concentration (MIC) was obtained. This was **carried out** by selecting the test tube that shows no  
189 growth during the MIC determination. A loopful from the test tube containing the media and the  
190 extract were inoculated into a sterile nutrient broth media. This was further incubated for another

191 24-48 hrs at 37°C for bacteria, after which was examined for bacteria for any microbial growth.  
192 The lowest concentration at which no growth was observed on the plate was taken as the MBC  
193 [18].

## 194 **RESULTS AND DISCUSSION**

195 This study was undertaken to investigate the antimicrobial activity and phytochemical  
196 screening the aqueous and ethanolic root and stem bark extracts of *Ficus sycomorus* Linn. Due  
197 to the side effects of the current drugs and the resistance that pathogenic microorganisms build  
198 against antibiotics, much attention has led to the study of biologically active compounds isolated  
199 from plant species used in herbal medicine [19]. Different scientific studies provided evidence  
200 that medicinal plants might indeed be potential sources of new antibacterial agents even against  
201 some antibiotic-resistant strains [20].

202 The yield of the plant extracts is presented in Table 2. It was observed that Ethanol stem bark  
203 extract (ESBE) gave the highest yield 16.00g (8.0%) followed by Ethanol root bark extract (ERBE)  
204 14.14 g (7.07 %) then Aqueous root bark extract (ARBE) 12.23 g (6.12%) and the lowest is Aqueous  
205 stem bark extract (ASBE) 11.16 g (5.58 %). In general the solvent, ethanol gave higher yield  
206 irrespective of the plant part than the aqueous solvent.

207 The result of this study shows the presence of phytochemicals considered as active  
208 medicinal chemical constituents as shown in table 2. Phytochemicals such as tannins, saponin,  
209 terpenoids, flavonoids, alkaloids, glycosides, steroids, phenols and reducing sugars were all  
210 found to be present in both the ethanol extracts of roots and stem bark of *Ficus sycomorus*.  
211 However, glycosides was the only constituent not detected in Aqueous extracts of the root and  
212 stem bark. The result is contrary to the findings of [21] who reported the presence glycoside in  
213 the methanolic stem bark extract of *Ficus sycomorus* obtained from Zaria city of Kaduna State,  
214 Nigeria. This difference could be attributed to geographical location of the samples.

215 The various phytochemical compounds detected are known to have beneficial  
216 importance in industrial and medicinal sciences. These secondary metabolites exert antimicrobial  
217 activity through different mechanisms. Plant phenolic compounds especially flavonoids are  
218 currently of growing interest owing to their supposed properties in promoting health (anti-  
219 oxidants) [22]. Flavonoids have been demonstrated to have antiinflammatory, anti-allergenic,  
220 anti-viral, anti-aging, and anti-carcinogenic activity. The broad therapeutic effects of flavonoids  
221 can be largely attributed to their antioxidant properties. In addition to an antioxidant effect,

222 flavonoid compounds may exert protection against heart disease through the inhibition of  
223 cyclooxygenase and lipoxygenase activities in platelets and macrophages[23].Tannins are  
224 reported to possess physiological astringent and haemostatic properties, which hasten wound  
225 healing and ameliorate inflamed mucus membrane and also inhibit the growth of microorganisms  
226 by precipitating microbial proteins and making nutritional proteins unavailable for them; they  
227 form irreversible complexes with proline rich proteins, resulting in the inhibition of the cell  
228 protein synthesis. They have important roles such as stable and potent antioxidants [23, 24].  
229 They act as binders and for treatment of diarrhea and dysentery [25] Tannins also reported to  
230 exhibit antiviral, antibacterial, anti-tumor activities. It was also reported that certain tannins are  
231 able to inhibit HIV replication selectivity and is also used as diuretic [23]

232 The results of the zones of inhibition of the different extracts (ARBE, ASBE, ESBE  
233 and ERBE) against the tested pathogens are exhibited in Tables 3 – 6. It showed that the extracts  
234 have dose dependent antimicrobial activities against the pathogens at various concentrations used  
235 in this study. It was noticed that the extract was more effective at concentration of 100 mg/mL,  
236 but the effectiveness increases as the concentration increases. The highest activity was shown by  
237 the ESBE and ERBE at 100 mg/mL (15mm) against *E. coli*. Although most of the extracts at the  
238 various concentrations used showed activity against the pathogens, it was observed on the  
239 general that the extracts are more effective at 100 mg/mL on *E. coli*, which showed similar  
240 activity with the standard drug (Tetracycline at 100µg/mL) used. At lower concentrations, the  
241 extracts seem to show more activity against shigella *dysenteriae* as seen in tables 3 - 6.

242 From table 3, it is revealed that the zones of inhibitions of the extract (ARBE)  
243 against the tested pathogens showed that the extract has antimicrobial activities against the  
244 pathogens at various concentrations respectively. It was noticed that the extract was very  
245 effective at a concentration of 100 mg/mL, the effectiveness increases as the concentration  
246 increases. The control was more effective on *E. coli* with zone of inhibition up to 20 mm. Table  
247 4 shows the zones of inhibitions of the aqueous stem bark extract (ASBE) on the  
248 microorganisms. The result shows that the extract was effective at different concentrations with  
249 various zones of inhibitions as the concentration increases. However, *E. coli* was resistant  
250 against the extract at higher concentration of 100 mg/mL and 50 mg/mL but effective at lower  
251 concentration 25 mg/mL and also the control which has the highest zone of inhibition (11mm) on  
252 *E. coli*. From table 5, the ethanol stem bark extract (ESBE) also showed considerable  
253 antimicrobial activities on the tested clinical isolates at various concentrations used. The result

254 shows that at a higher concentration the extract was active against the clinical isolates or  
255 pathogens but more effective on *Shigella* at lower concentration (25 mg/mL) with zone of  
256 inhibition 10 mm, also the control was more effective with the highest zone of inhibition 16 mm.  
257 This extract show more activity against *E. coli* than the control drug at 100 mg/mL with 15 mm  
258 zone of inhibition. From table 6 the results of ethanol root extract (ERBE) against the pathogens  
259 also shows that the antimicrobial potential of the extract increases considerably as the  
260 concentration increases.

261 The result of the antimicrobial activity of root and stem bark extracts in this study is  
262 similar to that of [26,27,28] who asserted that many plants have been reported for therapeutic  
263 purposes because of the chemical compounds synthesized in these plants. The antibacterial  
264 activities of the ethanolic extracts of the leaves and stem bark of *F. sycomorus* have been previously  
265 reported [27]. The present study suggests that *F. sycomorus* may serve as a potential source of  
266 antibacterial and/or antimicrobial agents of plants origin. Hence, the observed antimicrobial  
267 activity of the root and stem bark extracts against the test organisms in this study may be due to  
268 the presence of phytochemical components. The findings demonstrated that the stem and root  
269 bark extract were sensitive to all the tested organisms and thus showed that the extract contained  
270 potential antimicrobial agents such as tannin, saponin, alkaloid, glycosides as secondary  
271 metabolite responsible for curing various sicknesses .The presence of tannin in all the extract  
272 could be probably responsible for the observed antimicrobial activity. The claim of literature  
273 that *F. sycomorus* has antimicrobial activity is hereby verified. The anti-microbial activity of the  
274 extracts, both the ethanol and aqueous of root and stem have shown a reasonable zone of  
275 inhibition to the concentration from 12.5 – 100 mg/mL and the control drug (Tetracycline) at 100  
276 µg/mL concentration. However, the ASBE extracts of *F. sycomorus* was observed to be less  
277 potent against the tested clinical isolate respectively.

278  
279 **The Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration**  
280 **(MBC) of the extracts are shown in Tables 7 and 8.** The result has shown that the MIC for all  
281 extracts of root and stem bark was 50 mg/mL. At this concentration, the extract was able to  
282 inhibit the growth of microorganisms. The result also revealed that the MBC was at 100 mg/mL  
283 these means that at this concentration the extract was able to kill the bacteria completely. This  
284 result is similar to the work of [26] who reported that the Minimum Inhibitory Concentration  
285 (MIC) and Minimum Bactericidal Concentration (MBC) of the aqueous and ethanolic root and

286 stem bark extracts of *Ficus sycomorus* extracts ranged from 3.125 mg/mL to 100 mg/mL. [21].  
 287 also reported that the minimum inhibitory concentration (MIC) of methanol root bark extract of *F.*  
 288 *sycomorus* was observed within the range of 2.5– 5.0mg/ml against *E. faecalis*, *E. coli*, *S. typhi*, *S.*  
 289 *dysenteriae* and *C. albicans*. This result therefore suggests that the extracts are more of  
 290 bacteriostatic.

291

292 **Table 1:** Percentage yield of the root and stem extracts

Extract	Initial weight	YIELD(g)	%
ERBE	200.00g	14.14g	7.07
ESBE	200.00g	16.00g	8.00
ARBE	200.00g	12.23g	6.12
ASBE	200.00g	11.16g	5.58

293 Key:

294 ERBE-----Ethanol Root Extract

295 ESBE-----Ethanol Stem Bark Extract

296 ARBE-----Aqueous Root Extract

297 ASBE-----Aqueous Stem Bark Extract

298  
 299 **Table 2: Qualitative Phytochemical analysis of the root and stem bark extract of**  
 300 *Ficus sycomorus*

301

TEST	Aqueous extract		Ethanol extract	
	Root	Stem bark	Root	Stem bark
Tannins	+	+	+	+
Saponin	+	+	+	+
Terpenoid	+	+	+	+
Flavonoids	+	+	+	+
Alkaloids	+	+	+	+
Glycosides	-	-	+	+
Steroids	+	+	+	+

310

311	Phenols	+	+	+	+
312	Reducing sugar	+	+	+	+

313

314           + = Present           - = Absent

315

316 **Table 3: Zone of Inhibition in (mm) Aqueous Root bark Extract (ARBE) Against**  
 317 **Opportunistic Pathogens.**

318

S/No.	Name of Organism	Concentration mg/mL				
		100	50	25	12.5	Tetracycline(Control)
	<i>S. aureus</i>	7	6	5	2	13
	<i>Escherichia coli</i>	10	8	7	4	20
	<i>Salmonella spp</i>	12	7	5	R	13
	<i>Shigella spp</i>	10	9	7	4	13

319 Key:

320 **Resistant---- R**  
 321 **Aqueous Root bark Extract----- ARBE**

322

323 **Table 4: Zone of inhibition in (mm) of Aqueous stem bark extract (ASBE) against**  
 324 **Opportunistic Pathogen**

Name of Organism	Concentration mg/mL				
	100	50	25	12.5	Tetracycline(Control)
<i>S. aureus</i>	9	6	4	3	7
<i>Escherichia coli</i>	R	R	10	4	11
<i>Salmonella spp</i>	7	5	4	3	8
<i>Shigella spp</i>	9	6	5	4	10

325

326 **Key: Resistant----- R           Aqueous stem bark extract ----ASBE**

327 **Table 5: Zone of Inhibition in (mm) of Ethanol stem bark extract (ESBE) against**  
 328 **Opportunistic Pathogens**

Name of Organism	Concentration mg/mL				
	100	50	25	12.5	Tetracycline(Control)
<i>S. aureus</i>	6	5	4	2	10

<i>Escherichia coli</i>	15	9	3	2	12
<i>Salmonella spp</i>	10	6	5	3	11

S/No.	Name of Organism	Concentration mg/MI				
		100	50	25	12.5	Tetracycline(Control)
	<i>S. aureus</i>	6	5	4	2	10
	<i>Escherichia coli</i>	15	9	3	2	12
	<i>Salmonella spp</i>	10	5	6	3	11
	<i>Shigella spp</i>	10	5	5	4	16
	<i>Shigella spp</i>	5	4	10	5	16

329 **Key: Ethanol stems bark extract----- ESBE**

330

331 **Table 6: Zone of Inhibition (mm) of Ethanol root bark Extract (ERBE) against**  
 332 **Opportunistic Pathogens.**

333

	100	50	25	12.5
<i>Staphylococcus aureus</i>	-	-	+	+
<i>Escherichia coli</i>	-	-	+	+
<i>Salmonella spp</i>	-	-	+	+
<i>Shigella spp</i>	-	-	+	+

334 **Key: Ethanolic root bark extract----- ERBE**

335

336 **Table 7: The Result of Minimum Inhibitory Concentration (MIC) of both aqueous and**  
 337 **ethanol extracts of root and stem bark of *Ficus sycomorus***

338

339 **Microorganism MIC (mg/mL)**

340 **+= Growth ; - = No growth**

341

342

343

344

345  
346  
347  
348

---

	100	50	25	12.5
<i>Staphylococcus aureus</i>	-	+	+	+
<i>Escherichia coli</i>	-	+	+	+
<i>Salmonella spp</i>	-	+	+	+
<i>Shigella spp</i>	-	+	+	+

---

349  
350  
351  
352  
353  
354  
355  
356

**Table 8: The Result of Minimum Bactericidal Concentration (MBC) of both aqueous and ethanol extracts of root and stem bark of *Ficus sycomorus***

---

Microorganism	MBC (mg/mL)
---------------	-------------

357 + = Growth ; - = No growth

358 **CONCLUSION**

359 Phytochemicals such as tannins, saponin, terpenoids, flavonoids, alkaloids, glycosides, steroids,  
360 phenols and reducing sugars were all found to be present in both the aqueous extracts of roots  
361 and stem bark of *Ficus sycomorus*.

362 From the studies of the antimicrobial activities, the research revealed that, for aqueous stem and  
363 root bark, ARBE had more antimicrobial potentials against the selected pathogens than the  
364 ASBE, but for ethanol stem and root bark both have almost the same inhibitory activities on the  
365 tested pathogens.

366 From the research, it was noticed that both the root and stem bark have antimicrobial potentials  
367 on the tested pathogens. This validates the claim of the traditional users who used it to treat  
368 diseases of microbial origin. Therefore, it can be used for therapeutic purposes.

369

370 **ACKNOWLEDGEMENT**

- 371 The authors are grateful to Laboratory Technologists, in person of Mr. Umar, Mallam Sani,  
372 Esther Dauda, all from Microbiology Department, Modibbo Adama University of Technology  
373 Yola, for their contributions for the success of the research.  
374  
375  
376  
377
- 378 **REFERENCES**
- 379 1. Cragg, G.M, and Newman D.J., (2001). Natural product drug discovery in the next  
380 millennium. *Pharmaceutical Biology*; Vol. 39, Supplement, pp. 8–17  
381
- 382 2. Tapsell LC, Hemphill I, Cobiac L.,(2006). "Health benefits of herbs And spices: the past, th  
383 resent, the future". *Med. J. Aust.* **18** (4 *Suppl*): 4–24.  
384
- 385 3. Lai P.K. and Roy J., (2004). Antimicrobial and chemopreventive properties of herbs and  
386 spices. *Current Medicinal Chemistry*, 11(11): 1451-1460
- 387 4. Rabiou, M.K., Safiyya, A., Sani, A. K., and Gambo, C., (2018). Phytochemical Compositions  
388 In Some Nigerian Medicinal Plants and Their Pharmacological Properties: A Review;  
389 *Journal of Anesthesiology*; 6(1): 15-25.  
390
- 391 5. Sofowora, A. (2008). *Medicinal Plants and Traditional Medicine in Africa*, 3rd Edition,  
392 Spectrum Books Ltd., Ibadan, Nigeria, page: 23-25  
393
- 394 6. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A (2009). *Agroforestry Database:a tree*  
395 *reference and selection guide version 4.0* [Online] Availableat: [http:// www World](http://www.worldagroforestry.org/treedb2/speciesprofile.php?Spid=620#)  
396 [agroforestry.org/treedb2/speciesprofile.php?Spid=620#](http://www.worldagroforestry.org/treedb2/speciesprofile.php?Spid=620#) [Accessed 19 August, 2017].  
397
- 398 7. Hassan, S.W., Lawal M., Muhammad, B.Y., Umar, R.A., (2007). Antifungal activity and  
399 phytochemical analysis of Column Chromatographic fractions of stem bark extracts of  
400 *Ficus sycomorus* L (Muraceae). *Journal of Plant Sciences.* 2(2): 209-215.  
401
- 402 8. Oyeleke, S. B., Dauda, B. E. N., Boye, O. A. (2008): Antibacterial activity of *Ficus capensis*.  
403 *Afr. J. Biotechnol.* 7: 1414 - 1417.  
404
- 405 9. Sandabe, U. K.; Onyeyili, P. A. & Chibuzo, G. A.(2006). Phytochemical screening and effect  
406 of aqueous extractof *Ficus sycomorus* L (moraceae) stem bark on muscularActivity in  
407 laboratory animals. *Journal of Ethnopharmacology*, **104**: 203 - 285  
408
- 409 10. Solomon-Wisdom G. O., Shittu G. A., Agboola Y. A. (2011): Antimicrobial And  
410 Phytochemical Screening Activities Of *Ficus Sur* (Forssk). *New York Science Journal*  
411 4(1):15-18.  
412
- 413 11. Udobi C. E., Onaolapo J.A., Agunu A. (2008): Antibacterial activities and bioactive  
414 Components of the aqueous fraction of the stem bark of *Parkia biglobosa* (JACQ)

- 415 (Mimosaceae). Nigerian Journal of Pharmaceutical Sciences; 7(1): 49-55.  
416
- 417 12. Fatope, M.O., Ibrahim H. and Takada Y. (1993). Screening for higher plants reported as  
418 Pesticides using the brine shrimp lethally Assay. *International Journal of Pharmacology*  
419 11(6): 250-254.
- 420 13. Nguta, J.M., Mbaria, J.M., Gakuya, D.W., Gathumbi, P.K., Kabasa, J.D., Kiama, S.G.  
421 (2011). Biological Screening of Kenya Medicinal Plants using *Artemia salina* L. (Artemiidae),  
422 *Pharmacology online*, 2, 458-478  
423  
424
- 425 14. Trease, G.E. and Evans, W.C. (2002). *Pharmacognosy*. 15th Ed. Saunders Publishers,  
426 London. pp. 42-44, 221-229, 246-249, 304-306, 331-332, 391-393.  
427
- 428 15. Nweze E.T., Okafor J.I. and Njoku O., (2004). Antimicrobial activities of methanolic extract  
429 of *Trumeguineesis* (Schumm and Thorn) and *Morinda lucinda* Benth used in Nigerian  
430 herb medicinal practice. *Journal of Biological Research and Biotechnology*, 2(1): 34-46.  
431
- 432 16. Senthilkumar P.K. and Reetha D., (2009). Screening of antimicrobial properties of certain  
433 Indian medicinal plants. *J. Phytol.*, 1(3): 193-198.  
434
- 435 17. National Committee for Clinical Standards. Reference method for both dilution antifungal  
436 susceptibility testing of yeast Approved Standard M27-APA: National Committee for  
437 Clinical Laboratory Standards Wayne, 2000.  
438
- 439 18. Ibekwe, V. I., Nnanyere, N. F. and Akujobi, C. O. (2001). Studies of Antibacterial Activity  
440 and Phytochemical Qualities of Extracts of Orange Peels. *International Journal of*  
441 *Environment and Human Health*, 2(1):41-46.  
442
- 443 19. Essawi, T., Srour, M. (2000). Screening of some Palestinian medicinal plants for antibacterial  
444 activity, *Journal of Ethnopharmacology*, 70, 343-349  
445
- 446 20. Kone, W. M., K. K. Atindehou, C. Terreaux, K. Hostettmann, D. Traore, and M. Dosso.  
447 (2004). Traditional medicine in North Cote-d'Ivoire: screening of 50 medicinal plants for  
448 antibacterial activity. *Journal of Ethnopharmacology*, 93, 43-49.
- 449 21. Abubakar, U. S., Danmalam, U. H., Musa, K. Y., Banni, Z., Yahaya, I., Abba, A. and Sani,  
450 A., (2015). Phytochemical and antimicrobial screening of methanol root bark extract of *Ficus*  
451 *sycomorus* linn. (moraceae). Nigerian Journal of Pharmaceutical Sciences;  
452 Vol. 14, No.2, pp1 -6  
453
- 454 22. Rauha JP, Remes S, Herinonen W, Hopia M, Kgjala T, Pitinlaja K ,(2000). Antimicrobial  
455 effects of finished plant extract containing flavanoids and other phenolic compounds. *Int. J*  
456 *Food Microbiol* ; 56: 3-12  
457
- 458 23. Fateh, A.L., Rahman F. Magbool, Elamin I. E., Shayoub, M. E., Salah, E. O. H., (2017).  
459 Phytochemical and Antimicrobial Screening of Stem Bark and Leaves Extracts from *Ficus*  
460 *sycomorus*. *World Journal of Pharmaceutical and Medical Research*; 3(11), 234-239

- 461  
462 24. Ogunleye, D.S., Ibitoye,S.F., (2003). Studies of antimicrobial activity and chemical  
463 constituents of *Ximenia Americana*. Trop. J Pharm Res.; 2: 239-241. 35.  
464  
465 25. Dharmananda S.(2003).Gallnuts and the uses of tannins in Chinese medicine. A paper  
466 Delivered at the Institute for Traditional Medicine, Portland, Oregon, 2003.  
467  
468 26. Abdullahi M. (2014). Susceptibility Profiles of Some Bacteria Isolated From Stool of  
469 Diarrhoeal Patients to the Stem and Root Barks Extracts of *Ficus Sycomorus* Linn  
470 ( Moraceae). M.Sc. Thesis Submitted to the School of Postgraduate Studies  
471 (unpublished), Ahmadu Bello, University, Zaria.  
472  
473 27. Adeshina, G.O., Okeke C. L. E., Osuagwu N.O. and Ehinmidu, J.O. (2010). Preliminary *in*  
474 *vitro* antibacterial activities of ethanolic extracts of *Ficus sycomorus* Linn and *Ficus*  
475 *platyphylla* (Moraceae). *African Journal of Microbiology Research*; **Vol. 4(8)** pp 598-601.  
476  
477 28. Bello, O.M., Ojediran, O.J., Dada, O.A., Olatunya, A.M. and Awakan, O.J., (2015): *In Vivo*  
478 Toxicity Studies and Phytochemical Screening of Stem Bark of *Ficus sycomorus* Linn  
479 (Moraceae). *Journal of Environmental Science, Toxicology and Food Technology* 9(3):72-74  
480

481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500

