

Original Research Article

Phenolic and mineral characteristics of seed coats and kernels from 24 species from Raipur area, India

ABSTRACT

Aims: The objective of the present work is the investigation of the physicochemical characteristics of seed coats and kernels from 24 species with medicinal and food applications.

Methodology: Seeds from 24 species (2 herbs, 11 vines and 11 trees), belonging to 13 families, were sampled in Raipur (India) in 2017. The collected seeds were dried and weighed, after which seed coats were manually peeled and separately weighed **separately**. Phenolic and mineral contents in the seed coats and kernels were analyzed by spectrophotometric and X-ray fluorescence (XRF) techniques, respectively.

Results: The seed coat fraction represented from 12% to 95% of the seed mass, depending on the species. The concentrations of total polyphenols, flavonoids and minerals in the seed coats varied from 1800 to 32300 mg/kg, from 1200 to 26900 mg/kg, and from 5876 to 36499 mg/kg, respectively. **Whereas** in the seed kernels, TPh, Fla and minerals ranged from 780 to 31760 mg/kg, from 300 to 31300 mg/kg, and from 12595 to 40810 mg/kg, respectively. **Elements:** P, S, K, Mg, Ca and Fe were found to be the main macro- and micro-elements. Seed coats from Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae families featured the highest total polyphenol contents, and those from Lauraceae and Rutaceae families showed the highest flavonoid concentrations. The highest total mineral contents corresponded to seed coats from Lauraceae, Rutaceae and Euphorbiaceae families.

24 **Conclusion:** Indian-laurel and curry tree stand out as promising phytochemical and nutrient
25 sources.

26 **Keywords:** Seed coat, Seed Kernel, Total polyphenol, Flavonoid, Mineral.

27 INTRODUCTION

28 The seed coat protects the internal parts of the seed from fungi, bacteria and insects, and prevents
29 water loss. It is composed of cellulose, fibre, polyphenols, starch, wax, etc. Its outer layer, called
30 testa, is generally hard and thick, while its inner layer, known as the tegmen, is softer [1].
31 Enrichment of various compounds (viz. minerals, cellulose, fibre, polyphenols, starch, wax, etc.)
32 in seed coats have been reported in the literature [2-7]. Among these phytochemicals:
33 polyphenols have become the subject of increasing research efforts owing to their potential
34 beneficial effects on human health [8,9].

35 Among the plants found in the Raipur area, Black Siris (*Albizia odoratissima* (L.f.) Benth.),
36 Malabar spinach (*Basella rubra* L., syn. *Basella alba* L.), wax gourd (*Benincasa hispida*
37 (Thunb.) Cogn.), squash (*Cucurbita maxima* (Duchesne) Duchesne ex Poir.), watermelon
38 (*Citrullus lanatus* (Thunb.) var. *lanatus*), Persian melon (*Cucumis melo* var. *cantalupo* Ser.),
39 Liane Cacorne (*Entada gigas* (L.) Fawc. & Rendle), tree cotton (*Gossypium arboreum* L.),
40 physic nut (*Jatropha curcas* L.), Persian walnut (*Juglans regia* L.), hyacinth bean (*Lablab*
41 *purpureus* (L.) Sweet), calabash (*Lagenaria siceraria* Standl.), Chinese-okra (*Luffa acutangula*
42 Roxb.), sponge gourd (*Luffa aegyptiaca* Mill.), Indian-laurel (*Litsea glutinosa* (Lour.) C.B.Rob.),
43 Indian-lilac (*Melia azadirachta* L., syn. *Azadirachta indica* A.Juss.), bitter melon (*Momordica*
44 *charantia* L.), curry tree (*Murraya koenigii* Spreng.), emblic (*Phyllanthus emblica* L.), East
45 Indian kino (*Pterocarpus marsupium* Roxb.), Indian sandalwood (*Santalum album* L.), Ceylon-
46 oak (*Schleichera oleosa* (Lour.) Oken), clearing-nut-tree (*Strychnos potatorum* L.f.), and Indian

47 tuliptree (*Thespesia populnea* Sol. ex Corrêa) **is are** widely used as medicine, food and fodder for
 48 animals [10-23].

49 Accumulation of the nutrients and polyphenols in some seed coats **were have been** reported **in**
 50 **the literature** [6, 24-28]. In this work, the physical and chemical characteristics of the seed coats
 51 and kernels from **these aforementioned** 24 species (2 herbs, 11 vines and 11 trees) are analyzed,
 52 with emphasis on their polyphenol contents.

53 **METHODS AND MATERIALS**

54 **Sample collection and handling**

55 Seeds from **aforementioned the selected** twenty-four species were collected in Raipur area
 56 (21.25°N 81.63°E), Chhattisgarh, India, during their maturation period in 2017. The seeds were
 57 manually separated and sun-dried in a glass room for one week, after which they were further
 58 dried in a hot air oven at 50 °C for 24 h. The mass of the seeds was measured using an AG245
 59 (Mettler Toledo, Columbus, OH, USA) electronic balance. The seed coats were then carefully
 60 peeled with the aid of a surgical blade and their mass was measured. The separated seed coats
 61 and kernels were crushed into a fine powder, and particles of mesh size ≤ 0.1 mm were sieved
 62 out. The samples were preserved in a deep freezer at -4 °C until the analyses were conducted.

63 **Analyses**

64 Sigma-Aldrich AR grade reagents were used for the analysis of polyphenols. 0.1 g of powdered
 65 seed coat were extracted with **an** acetone: water mixture (7:3, v/v), as recommended by Bertaud
 66 et al. [29]. An appropriate fraction was allowed to react with Folin-Ciocalteu reagent for colour
 67 development, and absorbance was measured at $\lambda=740$ nm with a UV-1800 (Shimadzu, Kyoto,
 68 Japan) UV-Vis spectrophotometer [30]. Three replicates for each solvent extract were performed

to determine the total phenolic content (TPh), which was expressed in terms of tannic acid equivalents by using a standard calibration curve. For flavonoid (Fla) analysis, a fraction of the extract was reacted with an aluminium chloride solution to develop a yellow coloured complex, measuring the absorbance at $\lambda=410$ nm [31]. The Fla concentration was determined with the aid of a standard quercetin calibration curve and indicated in terms of quercetin equivalents. Three replicates for each solvent extract were performed, and results are presented as average values across the three replicates.

A III Tracer-SD portable XRF (Bruker, Billerica, MA, USA) spectrophotometer was used for the quantification of 15 elements: K, Rb, Mg, Ca, Sr, Al, P, S, Cl, Ti, Mn, Fe, Cu, Zn and Pb. Standard brown and white cowpea (*Vigna unguiculata* (L.) Walp.) seeds were used as reference material to standardize the analyte concentration [32].

Statistical analyses

Cluster analysis was used to assess similarities in the micro- and macro-elements content in the seed coats. IBM (Armonk, NY, USA) SPSS v.25 software was used.

RESULTS AND DISCUSSION

Physical characteristics

The physical characteristics of the seeds and seed coats under study (shown in **Fig. 1**) are summarized in **Table 1**. Large differences in seed mass were found, with average values ranging from 25 to 23623 mg per seed, with the highest weights for the seeds from *Entada gigas* (23623 mg), followed by those from *Juglans regia* (12200 mg). The seed coat mass represented from 12 to 95% of the total seed weight.




Fig. 1. Seed samples from: (1) *P. embilica*, (2) *P. marsupium*, (3) *L. aegyptiaca*, (4) *B. hispida*, (5) *C. lanatus*, (6) *C. melo*, (7) *C. maxima*, (8) *L. acutangula*, (9) *L. siceraria*, (10) *M. charantia*, (11) *L. glutinosa*, (12) *G. arboreum*, (13) *T. populnea*, (14) *S. album*, (15) *M. koenigii*, (16) *S. oleosa*, (17) *M. azadirachta*, (18) *S. potatorum*, (19) *A. odoratissima*, (20) *B. rubra*, (21) *L. purpureus*, (22) *J. curcas*, (23) *J. regia*, and (24) *E. gigas*.

Table 1. Physico-chemical characteristics of seeds and seed coats. Total phenolic contents and flavonoid contents correspond to the seed coat and kernel samples

Species	Family	Type	Seed mass (mg)	Seed Colour	Seed coat (%)	Seed coat TPh (mg/kg)	Seed coat Fla (mg/kg)	Seed kernel TPh (mg/kg)	Seed kernel Fla (mg/kg)
<i>B. rubra</i>	Basellaceae	V	38	BrB	47±2	11400	10500	3457	1650
<i>C. maxima</i>	Cucurbitaceae	V	132	YeW	18±1	3100	1900	4931	1100
<i>L. siceraria</i>	Cucurbitaceae	V	216	WhBr	42±2	14400	8900	1956	1400
<i>C. lanatus</i>	Cucurbitaceae	V	38	ReBr	49±2	18500	13100	2278	1280
<i>L. aegyptiaca</i>	Cucurbitaceae	V	105	B	43±2	3100	2500	780	620
<i>C. melo</i>	Cucurbitaceae	V	25	LY	28±1	2900	2100	965	300
<i>L. acutangula</i>	Cucurbitaceae	V	122	B	47±2	8300	7200	2144	1380
<i>B. hispida</i>	Cucurbitaceae	V	64	YeW	47±	2900	2600	4074	2280
<i>M. charantia</i>	Cucurbitaceae	V	189	YeBr	35±1	30847	1700	1769	1180
<i>J. curcas</i>	Euphorbiaceae	H	758	Br	47±2	14700	4000	1501	4260
<i>L. purpureus</i>	Fabaceae	V	293	DBr	34±1	22000	3700	1260	2550
<i>A. odoratissima</i>	Fabaceae	T	159	LBr	42±	27000	4100	2492	4300

<i>E. gigas</i>	Fabaceae	V	23623	DBr	40±1	26900	3900	18840	2650
<i>P. marsupium</i>	Fabaceae	T	933	LY	93±3	25800	3800	31760	5800
<i>J. regia</i>	Juglandaceae	T	12200	PY	32±1	9600	1900	1045	1520
<i>L. glutinosa</i>	Lauraceae	T	248	DBr	43±2	29200	26900	4931	3880
<i>S. potatorum</i>	Loganiaceae	T	280	B	24±1	26000	15000	2707	1640
<i>G. arboreum</i>	Malvaceae	H	82	Br	48±2	4000	3500	7263	7160
<i>T. populnea</i>	Malvaceae	T	162	LBr	47±2	16800	8000	15839	12020
<i>M. azadirachta</i>	Meliaceae	T	972	DBr	65±2	1800	1200	1822	1300
<i>P. embilica</i>	Phyllanthaceae	T	920	PW	95±3	27000	3500	4476	3750
<i>M. koenigii</i>	Rutaceae	T	155	B	12±1	32300	25300	3457	3650
<i>S. album</i>	Santalaceae	T	180	DBr	40±2	10900	6200	7075	2750
<i>S. oleosa</i>	Sapindaceae	T	352	DBr	49±2	8500	5400	2198	1950

V = Vien, H = Herb, T = Tree, BrB = Brownish black, YeW = Yellowish white, WhBr = Whitish brown, ReBr = Reddish brown, B = Black, YeBr = Yellowish brown, DBr = Dark brown, LuB = Luster black, LY = Light Yellow, PY = Pale yellow, DB = Dark black, PW = Pale white


Polyphenol contents

The concentration of TPh and Fla in the seed coats and kernels varied from 1800 to 32300 mg/kg, from 780 to 31760 mg/kg, from 1200 to 26900 mg/kg and from 300 to 12020 mg/kg, respectively **with a mean value of 15748, 5376, 6954 and 2932 mg/kg (Table 1)**. The [TPh]/[Fla] ratio in the studied seed coats and kernels ranged from 1.1 to 18.1 and 0.4 to 7.1 **with a mean value of 3.5 and 1.9**, respectively. Higher contents of TPh and Fla were observed in the seed coats than in the seed kernels **was marked**.

Large variations in the polyphenol content were observed from one species to another, with noticeably higher TPh and Fla values in seed coats and kernels from tree species (**Fig. 2**).

Similarly, remarkable differences in the polyphenol content of the seed coat and kernel samples were detected as a function of the family (**Fig. 3**). **In the case of seed coats, Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae** four families showed the highest TPh contents in the case of seed coats.: **Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae** while. Whereas in the case of kernels, three families exhibited the highest TPh contents in the case of kernels

117 | were exhibited by: Fabaceae, Malvaceae and Santalaceae families. As regards Fla contents, the
 118 | highest concentrations corresponded to seed coats from Lauraceae and Rutaceae families.



119 | **Fig. 2.** Polyphenol concentration variation in (a) seed coats or and (b) seed kernels with respect to plant
 120 | types. TPh and Fla stand for total phenolic content and flavonoid content, respectively.
 121 |
 122 |




Fig. 3. Polyphenol concentration variation in (a) seed coats **or and** (b) seed kernels with respect to the family. TPh and Fla stand for total phenolic content and flavonoid content, respectively.

Mineral contents

The mineral contents of 15 elements (viz. K, Rb, Mg, Ca, Sr, Al, P, S, Cl, Ti, Mn, Fe, Cu, Zn and Pb) in the seed coats are summarized in **Table 2**. The total concentrations (Σ_{M15}) ranged from 5876 to 36499 mg/kg, with the highest values for seed coats from *Jatropha J. curcas*. Remarkably high mineral contents were observed in the seed coats from three families: Lauraceae, Rutaceae and Euphorbiaceae (**Fig. 4**).

P and K nutrients were abundant in the seed coats, ranging from 99 to 4983 mg/kg and from 1714 to 21982 mg/kg, respectively. The highest P contents were observed in seed coats from

Cucurbitaceae family, while the highest K contents (>15000 mg/kg) were detected in seed coats from *P. marsupium*, *L. glutinosa*, *T. populnea* and *M. koenigii*.

Table 2. Mineral contents in the seed coats from the 24 species under study, expressed in mg/kg.

Species	Mg	Al	P	S	Cl	K	Ca	Rb	Sr	Ti	Mn	Fe	Cu	Zn	Pb
<i>B. rubra</i>	1012	78	99	194	71	6305	3409	12	4	39	71	389	4	33	3
<i>C. maxima</i>	1762	67	4220	1307	1178	10591	992	13	3	7	85	583	2	4	2
<i>L. siceraria</i>	2020	55	2799	966	88	11324	3833	9	2	9	47	295	2	5	1
<i>C. lanatus</i>	1913	41	3474	1865	78	3247	2965	10	32	8	57	524	11	10	1
<i>L. aegyptiaca</i>	3344	67	2273	909	55	6791	4080	9	1	11	31	142	3	9	1
<i>C. melo</i>	1638	44	4983	1719	101	3913	338	21	1	12	37	364	8	49	1
<i>L. acutangula</i>	1754	81	3486	1302	142	7134	845	9	3	7	20	125	12	10	1
<i>B. hispida</i>	561	98	1878	1063	131	5859	1604	14	3	8	70	313	1	13	2
<i>M. charantia</i>	2642	432	2441	1444	88	5470	2763	8	2	9	77	308	1	3	1
<i>J. curcas</i>	4002	47	1991	1264	91	14636	14210	15	27	11	40	147	13	4	1
<i>L. purpureus</i>	1382	38	2344	1156	48	8541	3176	4	3	7	32	84	6	9	1
<i>A. odoratissima</i>	1738	46	1745	3140	71	8049	6256	15	9	8	132	125	3	5	1
<i>E. gigas</i>	1096	35	104	195	65	6278	3405	16	12	11	58	65	17	27	1
<i>P. marsupium</i>	2098	61	985	1897	59	15236	6685	18	29	38	79	491	914	3	3
<i>J. regia</i>	105	55	254	116	66	7297	2292	3	9	12	17	68	4	4	1
<i>L. glutinosa</i>	542	43	1559	1424	121	21982	3403	23	12	27	94	181	35	27	1
<i>S. potatorum</i>	5808	68	375	975	105	1714	7734	5	2	7	127	166	6	4	1
<i>G. arboreum</i>	2067	77	4001	1399	91	9312	1387	1	5	9	14	137	5.5	40	1
<i>T. populnea</i>	662	87	1631	1062	4859	16894	3256	17	4	11	33	419	402	1	1
<i>M. azadirachta</i>	323	844	826	725	132	7515	2205	9	3	30	28	685	5.5	12	1
<i>P. embilica</i>	175	59	465	243	111	3405	1008	6	20	9	18	348	4	4	1
<i>M. koenigii</i>	1395	49	1311	480	81	20233	6401	7	13	11	21	271	5	11	1
<i>S. album</i>	911	54	886	1100	77	12405	5280	26	29	23	70	327	7	16	1
<i>S. oleosa</i>	1635	65	1814	1227	68	2568	4982	3	13	12	34	334	14	6	1





Fig. 4. Total mineral contents in seed coats as a function of the family to which the plant species belongs.

S and Cl concentrations in seed coats were in the 116–3140 mg/kg and in the 48–4859 mg/kg range, respectively. The highest values for S and Cl corresponded to *A. odoratissima* and *T. populnea*, respectively.

Mg and Ca elements, probably present as silicates, ranged from 105 to 5808 mg/kg and from 338 to 14210 mg/kg, respectively. In this case, the highest concentrations of Mg and Ca were observed in seed coats from *S. potatorum* and *J. curcas*.

The concentrations of other elements in the seed coats, expressed in mg/kg, were in the following ranges: 1–26 (Rb), 1–32 (Sr), 35–844 (Al), 7–39 (Ti), 14–132 (Mn), 65–685 (Fe), 1–914 (Cu), 1–49 (Zn) and 1–3 (Pb). The highest concentrations of Rb, Sr, Al, Ti, Pb, Mn, Fe, Cu and Zn were found in the seed coats from *S. album*, *C. lantus*, *M. azadirachta*, *B. rubra*, *P. marsupium*, *A. odoratissima*, *M. azadirachta*, *P. marsupium* and *C. melo*, respectively. It is worth noting that seed coats from *T. populnea* featured high contents of Cl, K, Fe and Cu.

Noticeable differences in the mineral contents were also found depending on plant type. Total mineral contents were at least 29 and 55% higher in seed coats from trees and herbs, respectively, than in vine samples (**Fig. 5A**). Higher concentrations of major elements (P, S, Mg, Ca and Al) were observed in the herb samples (**Fig. 5B**), while those of Cl, K, Mn, Cu, Ti and Sr were higher in the tree samples (**Fig 5C**). As regards samples from vines, high contents of Rb, Fe and Zn were detected (**Fig 5D**).



160

161 **Fig. 5.** Variation of mineral contents in seed coats with respect to plant type: (A) total mineral content;
 162 (B) major elements; (C) Al, Cl and Fe; (D) trace elements. V, T and H stand for vine, tree and herb,
 163 respectively.

164

165 On the basis of their mineral contents, the seed coats from the 24 species under study were
 166 categorized into two groups by using cluster analysis (**Fig. 6**). Group-I consisted of 19 species,
 167 and the other 5 species were included in group-II, in such a way that the mean concentration
 168 value of Σ_{MIS} in the seed coats that belonged to group-II was at least twice that of group-I ones.

TPh = Total polyphenol content, Fla = Flavonoid content

The minerals content **of minerals** in the seed coats and kernels **of the prominent**from four of the species with the highest mineral contents. **seeds** are shown in **Table 4**. **The nutrients are well abundant in the seed coats than the kernels**. Fifteen elements were detected in the coats, while. **However**, in the kernels, only ten elements were identified. Generally, **a** higher concentration **of nutrients** were detected in the seed coats than in the seed kernels **was marked**, **(Fig. 7)**.

Species	Mg	P	S	K	Ca	Mn	Fe	Cu	Zn	Rb	Sr
<i>C. maxima</i>	1130	5505	1614	3920	102	63	157	14	78	11	2
<i>J. curcas</i>	1905	5513	1596	8032	2509	26	96	10	42	16	2
<i>P. marsupium</i>	5555	6743	5993	17119	5095	41	145	39	50	20	11
<i>M. koenigii</i>	551	1436	525	13130	2891	10	110	8	15	4	5

Species	Mg	P	S	K	Ca	Mn	Fe	Cu	Zn	Rb	Sr
<i>C. maxima</i>	1130	5505	1614	3920	102	63	157	14	78	11	2
<i>J. curcas</i>	1905	5513	1596	8032	2509	26	96	10	42	16	2
<i>P. marsupium</i>	5555	6743	5993	17119	5095	41	145	39	50	20	11
<i>M. koenigii</i>	551	1436	525	13130	2891	10	110	8	15	4	5




Fig. 7. Distribution of total elements in seed coats and kernels.

CONCLUSIONS

Seed coats are major sources of polyphenols and minerals, with concentrations at least twice those found in o folds higher nutrients in seed coats than the seed kernels are available.

Remarkably high total polyphenol contents (of up to 32300 mg/kg) were detected in the seed coats from tree species of the Loganiaceae, Phyllanthaceae, Lauraceae and Rutaceae families, while the highest flavonoid concentrations (of up to 26900 mg/kg) corresponded to seed coats from the latter two families. As regards mineral contents, the highest total values were observed in the seed coats from three families: Lauraceae, Rutaceae and Euphorbiaceae. The highest concentrations of major elements (P, S, Mg, Ca and Al) were observed in seed coats from herb species, while those of Cl, K, Mn, Cu, Ti and Sr were higher in the tree samples. In turn, samples from vines featured high contents of Rb, Fe and Zn. Seed coats from Indian-laurel and curry tree stand out as particularly promising phytochemical and nutrient sources.

CONSENT

Not applicable.

ETHICS APPROVAL

211 Not applicable.

212 CONFLICT OF INTEREST

213 The authors declare no conflict of interest.

214

215 REFERENCES

- 216 1. Moïse JA, Han S, Gudynaite-Savitch L, Johnson DA, Miki BLA. Seed coats: Structure,
217 development, composition, and biotechnology. In Vitro Cellular & Developmental Biology
218 - Plant. 2005;41(5):620-644. <https://doi.org/DOI: 10.1079/IVP2005686>
- 219 2. Attree R, Du B, Xu B. Distribution of phenolic compounds in seed coat and cotyledon, and
220 their contribution to antioxidant capacities of red and black seed coat peanuts (*Arachis*
221 *hypogaea* L.). Industrial Crops and Products. 2015;67:448-456. <https://doi.org/DOI: 10.1016/j.indcrop.2015.01.080>
- 222 3. Zhang RF, Zhang FX, Zhang MW, Wei ZC, Yang CY, Zhang Y, Tang XJ, Deng YY,
223 Chi JW. Phenolic composition and antioxidant activity in seed coats of 60 Chinese black
224 soybean (*Glycine max* L. Merr.) varieties. Journal of Agricultural and Food Chemistry.
225 2011;59(11):5935-5944. <https://doi.org/DOI: 10.1021/jf201593n>
- 226 4. Abutheraa R, Hettiarachchy N, Phillips GK, Horax R, Chen P, Morawicki R, Kwon YM.
227 Antimicrobial activities of phenolic extracts derived from seed coats of selected soybean
228 varieties. Journal of Food Science. 2017;82(3):731-737. [https://doi.org/10.1111/1750-](https://doi.org/10.1111/1750-3841.13644)
229 [3841.13644DOI: 10.1111/1750-3841.13644](https://doi.org/10.1111/1750-3841.13644)
- 230 5. Phommallath S, Teraishi M, Yoshikawa T, Saito H, Tsukiyama T, Nakazaki T, Tanisaka T,
231 Okumoto Y. Wide genetic variation in phenolic compound content of seed coats among
232

- black soybean cultivars. Breeding Science. 2014;64(4):409-415. [doiDOI: 10.1270/jsbbs.64.409](https://doi.org/10.1270/jsbbs.64.409)
6. Ribeiro ND, Maziero SM, Prigol M, Nogueira CW, Rosa DP, Possobom MTF. Mineral concentrations in the embryo and seed coat of common bean cultivars. Journal of Food Composition and Analysis. 2012;26(1-2):89-95. <https://doi.org/10.1016/j.jfca.2012.03.003> DOI: 10.1016/j.jfca.2012.03.003
7. Wu XJ, James R, Anderson AK, Mineral contents in seed coat and canning quality of selected cultivars of dark red kidney beans (*Phaseolus vulgaris* L.). Journal of Food Processing and Preservation. 2005;29(1):63-74. <https://doi.org/10.1111/j.1745-4549.2005.00013.x> DOI: 10.1111/j.1745-4549.2005.00013.x
8. Ganesan K, Xu B. A critical review on polyphenols and health benefits of black soybeans. Nutrients. 2017;9(5):455. [doiDOI: 10.3390/nu9050455](https://doi.org/10.3390/nu9050455)
9. Sevindik M, Akgul H, Pehlivan M, Selamoglu Z. Determination of therapeutic potential of *Mentha longifolia* ssp. *longifolia*. Fresen Environ Bull, 2017; 26: 4757-4763.
10. Sivasankar V, Moorthi A, Sarathi Kannan D, Suganya devi P. Anthocyanin, and its antioxidant properties in selected fruits. Journal of Pharmacy Research. 2011;4(3):800-806.
11. Prasad DMR, Izam A, Khan MMR. *Jatropha curcas*: Plant of medical benefits. Journal of Medicinal Plants Research. 2012;6(14):2691-2699. [https://doi.org/DOI: 10.5897/JMPR10.977](https://doi.org/10.5897/JMPR10.977)
12. Saboo SS, Thorat PK, Tapadiya GG, Khadabadi SS. Ancient and recent medicinal uses of *Cucurbitaceae* family. International Journal of Therapeutic Applications. 2013;9:11-19.

- 254 13. Hassan GA, Bilal T, Ahmad BT, Sameena W, Irshad AN. Economic and ethno-medicinal
 255 uses of *Juglans regia* L. in Kashmir Himalaya. Unique Journal of Ayurvedic and Herbal
 256 Medicines. 2013;1(3):64-67.
- 257 14. Ramana KV, Raju AJS. Traditional and commercial uses of *Litsea glutinosa* (Lour.) C.B.
 258 Robinson (Lauraceae). Journal of Medicinal Plants Studies. 2017;5(3):89-91.
- 259 15. Yadav KN, Kadam PV, Patel JA, Patil MJ. *Strychnos potatorum*: Phytochemical and
 260 pharmacological review. Pharmacognosy Review. 2014;8(15):61-66. doi: 10.4103/0973-
 261 7847.125533
- 262 16. Singh E, Sharma S, Pareek A, Dwivedi J, Yadav S, Sharma S. Phytochemistry, traditional
 263 uses and cancer chemopreventive activity of Amla (*Phyllanthus emblica*): The Sustainer.
 264 Journal of Applied Pharmaceutical Science, 2011;2(1):176-183.
- 265 17. Kumar SR, Loveleena D, Godwin S. Medicinal property of *Murraya koenigii* - A review.
 266 International Research Journal of Biological Sciences. 2013;2(9):80-83.
- 267 18. Kumar R, Anjum N, Tripathi YC. Phytochemistry and pharmacology of *Santalum album* L.:
 268 A review. World Journal of Pharmaceutical Research. 2015;4(10):1842-1876.
- 269 19. Bhatia H, Kaur J, Nandi S, Gurnani V, Chowdhury A, Reddy PH, Vashishtha A, Rath B. A
 270 review on *Schleichera oleosa*: Pharmacological and environmental aspects. Journal of
 271 Pharmacy Research. 2013;6(1):224-229. <https://doi.org/10.1016/j.jopr.2012.11.003>
 272 [DOI: 10.1016/j.jopr.2012.11.003](https://doi.org/10.1016/j.jopr.2012.11.003)
- 273 20. Sathyanarayana T, Sarita T, Balaji M, Ramesh A, Boini MK. Antihyperglycemic and
 274 hypoglycemic effect of *Thespesia populnea* fruit in normal and alloxan-induced diabetes in
 275 rabbits. Saudi Pharmaceutical Journal. 2004;12(2):107-111.

- 276 | 21. Al-Snafi AE. The pharmacology and medical importance of *Lathyrus* *dolichos* *lablab* (*Lablab*
 277 | *purpureus*)-A review. IOSR Journal of Pharmacy. 2017;7(2):22-30. DOI: 10.9790/3013-
 278 | 0702012230
- 279 | 22. Van Andel TR, Croft S, Van Loon EE, Quiroz D, Towns AM, Raes N. Prioritizing West
 280 | African medicinal plants for conservation and sustainable extraction studies based on market
 281 | surveys and species distribution models. Biological Conservation. 2015;181:173-181.
 282 | <https://doi.org/10.1016/j.biocon.2014.11.015> DOI: 10.1016/j.biocon.2014.11.015
- 283 | 23. Moraghan JT, Grafton K. Genetic diversity and mineral composition of common bean seed.
 284 | Journal of the Science of Food and Agriculture. 2001;81(4):404-408. [https://doi.org/DOI:](https://doi.org/DOI:10.1002/1097-0010(200103)81:4<404)
 285 | 10.1002/1097-0010(200103)81:4<404
- 286 | 24. Moraghan JT, Grafton K. Distribution of selected elements between the seed coat and
 287 | embryo of two black bean cultivars. Journal of Plant Nutrition. 2002;25(1):169-176.
 288 | [https://doi.org/DOI: 10.1081/PLN-100108788](https://doi.org/DOI:10.1081/PLN-100108788)
- 289 | 25. Tajoddin M, Shinde M, Lalitha J. Polyphenols of mung bean (*Phaseolus aureus* L.) cultivars
 290 | differing in seed coat color: Effect of dehulling. Journal of New Seeds. 2010;11(4): 369-379.
 291 | [https://doi.org/DOI: 10.1080/1522886X.2010.520146](https://doi.org/DOI:10.1080/1522886X.2010.520146)
- 292 | 26. Yang QQ, Gan RY, Ge YY, Zhang D, Corke H. Polyphenols in common beans (*Phaseolus*
 293 | *vulgaris* L.): Chemistry, analysis, and factors affecting composition. Comprehensive
 294 | Reviews in Food Science and Food Safety. 2018;17(6):1518-1539. [https://doi.org/DOI:](https://doi.org/DOI:10.1111/1541-4337.12391)
 295 | 10.1111/1541-4337.12391
- 296 | 27. Marles MA, Vandenberg A, Bett KE. Polyphenol oxidase activity and differential
 297 | accumulation of polyphenolics in seed coats of pinto bean (*Phaseolus vulgaris* L.)

- characterize postharvest color changes. Journal of Agricultural and Food Chemistry. 2008;56(16):7049-7056. doiDOI: 10.1021/jf8004367.
28. Zhong L, Wu G, Fang Z, Wahlqvist ML, Hodgson JM, Clarke MW, Junaldi E, Johnson SK. Characterization of polyphenols in Australian sweet lupin (*Lupinus angustifolius*) seed coat by HPLC-DAD-ESI-MS/MS. Food Research International. 2019;116:1153-1162. <https://doi.org/DOI:10.1016/j.foodres.2018.09.061>
29. Bertaud F, Tapin-Lingua S, Pizzi A, Navarrete P, Petit-Conil M. Characterisation of industrial barks for their tannin contents for further green-wood based adhesives applications. InTech Fibre. COST FP0901-Hamburg. 2010.
30. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods in Enzymology. 1999;299:152-178. [https://doi.org/DOI:10.1016/S0076-6879\(99\)99017-1](https://doi.org/DOI:10.1016/S0076-6879(99)99017-1)
31. Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. Journal of Food and Drug Analysis. 2002;10(3):178-182.
32. Towett, E. K., Shepherd, K. D., Drake, B. L. (2016) Plant elemental composition and portable X-ray fluorescence (pXRF) spectroscopy: quantification under different analytical parameters, X-Ray Spectrom. 45, 117–124. DOI: 10.1002/xrs.2678

Formatted: Font: Italic