

Case study

Nutritional, spectral and thermal characteristic of Lamiaceae seeds

Formatted: Font: Not Italic

ABSTRACT

Aims: Species of the family Lamiaceae possess a rich tradition of use for flavoring and medicinal purposes. This paper focusses on the nutritional and thermal characteristics of the seeds from eight Lamiaceae species belonging to this family: *Gmelina arborea* Roxb. ex Sm., *Hyptis suaveolens* (L.) Poit., *Leonotis nepetifolia* (L.) R.Br., *Ocimum americanum* L., *Ocimum sanctum* L. (Rama Tulsi), *Ocimum tenuiflorum* L. (Krishna Tulsi), *Origanum vulgare* L. and *Tectona grandis* L.f.).

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Methodology: The oil, starch, total polyphenol, flavonoid and mineral contents for aforementioned seeds were determined. The Fourier-transform infrared (FTIR) spectroscopy was used to assess the phytoconstituents. The thermogravimetric/derivative thermogravimetric analyses (TG/DTG) and differential scanning calorimetry (DSC) analyses were performed to analyze the decomposition patterns.

Formatted: Font: Not Italic

Results: The concentrations of oil, starch, total polyphenol, flavonoids and minerals for eight the seeds from the eight plants under study were ranged from 11.8 to 50.4%, from 0.22 to 1.84%, from 295 to 5842 mg/kg, from 1660 to 12680 mg/kg and from 11756 to 33927 mg/kg, respectively. Unsaturated oils, polyphenols and lignin were recognized by vibrational spectroscopy. The sequence of thermal effects in the seed pyrolysis process above 100 °C have been put in relation to seed protein crystallization (endotherm at 200 °C), oxidation reactions and

degradation of hemicellulose and other fiber components (at around 300 °C), and decomposition of polyunsaturated (at 357 °C) and mono-unsaturated (at 391 °C) triglycerides.

Conclusion: Lamiaceae seeds are **medicinally** potential food alternatives to the cereals.

Formatted: Font: Not Italic

Keywords: Lamiaceae, seeds, FTIR Fourier-Transform Infrared Spectroscopy, Thermal analysis, chemical compounds, Oil, Starch, Polyphenol, Mineral composition.

Formatted: Font: Not Italic

INTRODUCTION

The Lamiaceae or Labiatae (Clade: Angiosperms / Eudicots / Asterids. Order: Lamiales) are a family of flowering plants comprising about 200 genera and 3,200 species, commonly with aromatic, herbage, quadrangular stems, and verticillate inflorescences. They are widely cultivated for medicinal, perfumery, culinary and ornamental purposes. [1]. Members of this family are a source of **essential oils for flavoring and perfumes include the** strong aromatic essential oils, tannins, saponins and organic acids. [2, 3]. Eight Lamiaceae plants **with a widespread distribution in central India**, viz. *Gmelina arborea* Roxb., *Tectona grandis* L.f., *Hyptis suaveolens* (L.) Poit, *Leonotis nepetifolia* (L.) R.Br., *Ocimum americanum* L., *Ocimum sanctum* L., *Ocimum tenuiflorum* L., and *Origanum vulgare* L., and *Tectona grandis* L.f., **with a widespread distribution in central India** are studied herein. *Gmelina arborea* (Malay bush-beech) and *T. grandis* (Bangkok teak) are large deciduous trees harvested for local use as a wood, food and medicine purposes and as a source of oils. [4, 5]. *Hyptis suaveolens* (pignut) is a strong-scented herb considered to be stimulant, carminative, **endorific** and lactagogue [6], which grows as a weed over large areas in barrel land in the rainy season. *Leonotis nepetifolia* (Christmas candlestick) is an annual short-lived perennial plant, often found **at in roadsides**, canal and riversides in the rainy season. **Other species:** *Ocimum americanum* L., *Ocimum sanctum* L. (Rama Tulsi, light holy basil), *Ocimum tenuiflorum* L. (Krishna Tulsi, dark holy

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Comment [AM1]: Is it correct??

Comment [PMR2]: Yes, it is (see, for instance, <http://www.efloraofgandhinagar.in/herb/hyptis-suaveolens>)

Comment [AM3]: This references is about Ocimum. Does include Hyptis in the text?? OR put it in Ocimum

Comment [PMR4]: Relocated

basil), and *Origanum vulgare* are perennial and aromatic plants, used in treatments of various diseases. [6][7]. These plants have shown promising properties as functional foods, in pain therapy and as bactericides and fungicides [8, 9, 10, 11, 12, 13]. The chemistry and uses of *G. arborea* and *H. suaveolens* seed oils have been reported [14, 15], but most of the characteristics of Lamiaceae seeds from Indian origin remain undescribed. The purpose of this paper is to report the nutritional, and mineral contents, and the thermal features behaviour of the seeds from the eight selected species.

Comment [AM5]: This references is about Ocimum. Does include Hyptis in the text?? OR put it in Ocimum

Comment [PMR6]: Thanks for bringing this to our attention. It has been relocated.

Formatted: Font: Not Italic

MATERIALS AND METHODS

Sample Collection

Seeds from the eight Lamiaceae plants (viz. *G. arborea* (GA), *T. grandis* (TG), *H. suaveolens* (HS), *L. nepetifolia* (LN), *O. americanum* (OA), *O. sanctum* (OS), *O. tenuiflorum* (OT) and *O. vulgare* (OV) under study were collected in the Raipur city area, India (21.25°N 81.63°E), and were authenticated by a plant taxonomist and by using a standard monograph. [16]. The ripening periods of TG, HS and LN; OA, OS, OT and OV; and GA were October-November, December and May, respectively. Their leaves and fruits were collected in the relevant period together with near-surface soil samples in year, 2017.

Formatted: Font: Not Italic

Sample Preparation

The seeds were manually separated from their carpels/fruits manually. All samples were sun-dried for one week in a glass room, and further dried in an oven at 50 °C overnight. Subsequently, they were crushed into fine powder and sieved out particles of mesh size, ≤100 µm. They were stored in the glass bottle and preserved in the refrigerator at -4 °C.

Comment [AM7]: Are you sure?? Because I believe, you are considering as seed, the carpel enclosing the real seed, i. e., a diaspore (seed + fruit or floral parts). I believe that you have been separating diaspores from persistent floral parts or perhaps acresent calyx.

Comment [PMR8]: Thank you for bringing this to our attention. The wording was misleading (we made a mistake in the translation into English language). We actually studied only the seeds. Figure 1 has been updated (see comment below)

Analyses

The pH value of the soil was determined by keeping a 5 g sample in a 100 mL conical flask with deionized water (15 mL) overnight. The pH value of the decanted solution was measured with a Hanna Instruments (Woonsocket, RI, USA) pH meter. The moisture content of the seeds was determined by drying the seeds at 105 °C in an air oven for 6 hr period to the analysis, and mean values were determined. All characterization results are presented on a dry weight (dw) basis.

The oil content in the seeds was determined by extraction from a 5 g powdered sample (kernel of GA and TG, and whole seed of HS, LN, OA, OS, OT or OV) in n-hexane (25 mL), as described in [the](#) literature. [17]. The oil content was presented as a percentage on the basis of the dry weight (dw) of the seeds.

[Analytical reagent \(AR\)](#) grade sodium maleate (CAS 371-47-1) buffer, sodium acetate (CAS 127-09-3) buffer, potassium hydroxide (CAS 1310-58-3), amyl glucosidase (CAS 9032-08-0), pancreatic- α -amylase (MDL MFCD00081319) [were purchased from Sigma-Aldrich](#), and glucose oxidase–peroxidase was purchased from Megazyme International Ireland Ltd. The starch content of seeds was determined by the enzymatic method. [18].

Analytical grade Folin-Ciocalteu reagent (MDL MFCD00132625), aluminum chloride (CAS 7446-70-0), tannic acid (CAS 1401-55-4), gallic acid (149-91-7) and quercetin (CAS 117-39-5) for the analysis of the phenols were purchased from Sigma–Aldrich. For the analysis of [the](#) total polyphenol content (TPC), 100 mg of sample (whole seed of GA, TG, HS, LN, OA, OS, OT or OV) in powder form [was were](#) mixed with 5 mL of an acetone: water mixture (70:30, v/v), and subjected to sonication for 20 minutes at 20 °C in an ultrasonic bath, according to the procedure described by [Bertaud et al.](#) [19]. The TPC of each extract was determined by use of the Folin-Ciocalteu reagent and expressed as tannic acid equivalents (TAE). [20]. The flavonoid content

Comment [AM9]: Beginning the phrase you must put the complete word, not use abbreviations.

Comment [PMR10]: Corrected

Formatted: Font: Not Italic

Formatted: Font: Italic

(Fla) was analyzed by the aluminum chloride method and expressed as quercetin equivalents (QE). [21].

A Bruker Tracer 5i portable X-ray fluorescence (pXRF) spectrometer (Serial Number 900F4473), equipped with a 4W rhodium anode and Xflash Silicon Drift Detector (SDSD) with a typical resolution of 2028 channels, was employed for the elemental analysis of the seed samples. Four standard reference materials, brown and white cowpea [*Vigna unguiculata* (L.) Walp.] seeds, cowpea and mango (*Mangifera indica* L.), leaves with reference values from ICP-OES and MS (As, Mo and Se in mg/kg) after Aqua Regia (HCl: HNO₃, 4:1) digestion were used for validation of the pXRF results. **Whereas,** a standard soil sample (NCS DC 73382 CRM) was used for as a reference for the soil **analysisanalyses to generate the precise data base.**

The vibrational spectrum in the 400 to 4000 cm⁻¹ **spectral** range was characterized using a Thermo Scientific (Waltham, MA, USA) Nicolet iS50 **Fourier-Transform Infrared (FTIR)** spectrometer, equipped with an in-built diamond attenuated total reflection (ATR) system, with a 1 cm⁻¹ spectral resolution and by averaging 64 scans.

Thermogravimetric/derivative thermogravimetric analyses (TG/DTG) and differential scanning calorimetry (DSC) analyses were conducted with a Perkin-Elmer (Waltham, MA, USA) STA6000 simultaneous thermal analyzer by heating the samples in a slow stream of N₂ (20 mL/min) from room temperature up to 800 °C, at a heating rate of 20 °C /min. Pyris v.11 software was used for data analysis.

RESULTS AND DISCUSSION

Seeds Physical Characteristics

Comment [AM11]: Is it correct??

Comment [PMR12]: Corrected

Formatted: Font: Italic

The seeds of the eight Lamiaceae under study (GA, TG, HS, LN, OA, OS, OT and OV) were colored from yellow to black and featured different shapes, viz. circular, elliptic, lanceolate, ovate and sagittate (**Fig. 1** and **Table 1**). Apropos of seed weights, *Ocimum* seeds were the smallest, with weights ranging from 0.3 to 1.0 mg per seed, whereas LN, HS, GA and TG seeds ranged from moderate to (very) big size, featuring weights from 2.6 to 481 mg per seed. The moisture content of the seeds varied from 4.2% to 8.1% and **had showed a fare fair** correlation with seed mass ($r = 0.75$).

Oil and Starch Concentration

As noted above, the oil from Lamiaceae seeds has wide medicinal uses. In the seeds from the eight species discussed herein, oil contents ranged from 11.8% to 50.4% (**Table 1**). Among them, GA and TG seeds were highly oily, with lipid fractions in the 41.2 to 50.4% range. Similar **oil composition of oilspercentage** in some *Ocimum*, GA and TG seeds **were have been** reported elsewhere. [22, 23].

The concentration of total starch in the eight seeds ranged from 0.22% to 1.84%. Four seeds –GA, LN, OM and TG– showed low starch contents (1.24 to 1.84%), while in the other seeds it was contained at trace levels (0.22 to 0.64%). The concentration of resistant starch in the seeds ranged from 0.03% to 1.13%, with a high fraction of insoluble starch in seven of the seeds (except for GA), varying from 17% to 78%.

(a) (b) (c) (d)

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted Table




Fig. 1: Seed image: (Aa) *Gmelina arborea*, (Bb) *Hyptis suaveolens*, (Cc) *Leonotis nepetifolia*, (Dd) *Ocimum americanum*, (Ee) *Ocimum sanctum*, (Ff) *Ocimum tenuiflorum*, (Gg) *Origanum vulgare* L., and (Hh) *Tectona grandis*.

Formatted: Font: 14 pt

Formatted: Font: 14 pt

Comment [PMR13]: Two of the photographs were confusing: e.g., for *T. grandis* the entire fruit, with the corky layer, was shown. The photographs of the seeds from *G. arborea* and *T. grandis* have been updated

Phenol Concentration

The TPC and Fla contents in the Lamiaceae seeds varied from 295 to 5842 mg/kg and from 1660 to 12680 mg/kg (in TAE and QE, respectively). These contents were noticeably lower than the TPC and Fla concentration in their leaves, which varied from 15406 to 29900 mg/kg and from 9804 to 34800 mg/kg, respectively. **The basil (*O. cimumsanctum*) and *T. grandis* species** were the richest in flavonoids.

Formatted: Font: Not Italic

Comment [AM14]: All Ocimum?? So you must put *Ocimum* spp.

Comment [PMR15]: Clarified

Mineral Concentration

The sum of the concentrations of the 19 elements under analysis (P, S, Cl, K, Rb, Mg, Ca, Sr, Ba, Ti, V, Cr, Mn, Fe, Co, Cu, Zn, Mo and Pb) in the GA, HS, LN, OA, OS, OT, OV and TG seeds was found to be 11756, 22629, 30977, 23586, 27133, 18641, 33927 and 22077 mg/kg, respectively. Their reduced concentration in the GA seeds (whole seed, **including the carpels, with seed coat**) was **noticednoteworthy**.

Comment [AM16]: Verify if you are speaking of seed or diaspore, and what is the difference respect to other species??.

Comment [PMR17]: Clarified

P, S, K, Rb, Mg, Ca, Sr, Mn, Fe, Cu and Zn were detected in all eight seeds, and were in the 2954 to 6829, 884 to 2543, 3304 to 16284, 4 to 17, 1047 to 2862, 567 to 11368, 6 to 94, 10 to 68, 86 to 1239, 7 to 49 and 11 to 120 mg/kg range, respectively (**Table 1**). Cobalt was identified in all seeds (except GA and LN) at traces level, 1 mg/kg. Barium was detected in HS, OA and OS seeds at low levels (ranging from 13 to 25 mg/kg). Other toxic elements, such as Ti, V, Cr, Mo and Pb, were detected in **the** OS seeds also at low levels, in the range of 2 - 70 mg/kg. Clusters of mineral elements at high concentrations were identified for the seeds of three species: P-Mg-Ca-Mn, for LN; Rb-Sr-Ti-V-Cr-Fe-Mo-Pb, for OS; and S-Cu, for HS. The maximum K and Zn concentrations were detected in **the** OV and GA seeds.

For comparative purposes, the concentration of trace elements in the leaves of HS was tested. Their total concentrations were remarkably higher in the leaves, 42241 mg/kg, probably due to coordination with polyphenols. Elements i.e. S, Cl, K, Mg, Ca and Fe were the dominated

elements in the leaves. The concentration of elements i.e. P, S, Cl, K, Rb, Mg, Ca, Sr, Ti, Cr, Mn, Fe, Mo and Pb in the dried HS leaves was found to be 820, 1925, 6733, 24376, 6, 1979, 5028, 50, 114, 11, 175, 1020, 4 and 4 mg/kg.

Table 1. Physico-chemical characteristics of Lamiaceae seeds and leaves (in brackets)

Parameter	<i>Gmelina arborea</i>	<i>Hyptis suaveolens</i>	<i>Leonotis nepetifolia</i>	<i>Ocimum americanum</i>	<i>Ocimum sanctum</i>	<i>Ocimum tenuiflorum</i>	<i>Origanum vulgare</i>	<i>Tectona grandis</i>
Color	light yellow	dark brown	salty black	black	dark brown	dark brown	black	light brown
Shape	obovate	sagittate	lanceolate	narrow ovate	broad ovate	broad ovate	elliptic	circular
Mass, mg	94.7	5.0	2.6	1.1	0.3	0.3	1.0	481
Moisture, %	7.5	6.4	4.2	4.9	5.2	5.5	4.2	8.1
Oil, %	50.4	17.2	30.3	13.4	12.7	11.8	14.8	41.2
Total starch, %	1.42	0.22	1.84	1.82	0.45	0.62	0.64	1.24
Resistant starch, %	0.03	0.08	0.31	1.13	0.31	0.34	0.50	0.41
Resistant/total starch, %	2.1	36	17	62	69	55	78	33
TPh, mg/kg	1447 (21476)	4181 (23875)	3538 (16042)	2117 (15406)	295 (28900)	5842 (25898)	750 (29900)	4904 (25580)
Fla, mg/kg	2075 (34800)	3025 (9804)	2825 (16538)	5100 (9966)	15850 (18900)	5350 (24295)	3600 (28900)	7375 (21548)
P, mg/kg	4880	3634 (820)	6829	4631	3983	3630	4738	2954
S, mg/kg	1507	2543 (1925)	2235	1940	1678	1537	2096	884
Cl, mg/kg	ND	0 (6733)	ND	ND	ND	ND	183	ND
K, mg/kg	3304	4355 (24376)	7165	7775	6950	5544	16284	11816
Rb, mg/kg	10	6 (6)	11	9	17	14	7	4
Mg, mg/kg	1047	2070 (1979)	2862	1804	2775	1189	2339	1120
Ca, mg/kg	567	9736 (5028)	11368	7201	10186	6363	7917	5060
Sr, mg/kg	6	21 (50)	29	59	94	18	29	26
Ba, mg/kg	ND	14	ND	13	25	ND	ND	ND
Ti, mg/kg	ND	0 (114)	ND	ND	70	ND	ND	ND
V, mg/kg	ND	ND	ND	ND	2	ND	ND	ND
Cr, mg/kg	ND	ND	ND	ND	5	ND	ND	ND
Mn, mg/kg	66	44 (175)	68	10	43	21	25	34
Fe, mg/kg	219	145 (1020)	346	86	1239	255	235	160
Co, mg/kg	ND	1	ND	1	1	ND	1	1

Comment [AM18]: It is an important table, perhaps other format would be better for to have a stable and clear data visualization.

Comment [PMR19]: We have used this format in other papers previously published in this journal and we would prefer to keep it consistent, in case the readers would like to compare results with those of other families.

Cu, mg/kg	30	49	36	12	13	13	30	7
Zn, mg/kg	120	11	28	45	48	57	43	11
Mo, mg/kg	ND	0 (4)	ND	ND	2	ND	ND	ND
Pb, mg/kg	ND	0 (4)	ND	ND	2	ND	ND	ND

Bioaccumulation

The pH value of soil solutions (n = 8) **was** ranged from 7.5 to 8.9, with a mean value (p = 0.05) of 8.1 ± 0.4 . The concentrations of Cl, P, S, K, Rb, Mg, Ca, Sr, Mn, Fe, Cu and Zn elements were in the following ranges: 109 **to** 161, 118 **to** 185, 190 **to** 280, 1101 **to** 1170, 3 **to** 8, 1180 **to** 1740, 4480 **to** 7852, 40 **to** 59, 1000 **to** 1480, 13400 **to** 19700, 30 **to** 58 and 17 **to** 28 mg/kg, respectively (with mean values of 138 ± 13 , 154 ± 16 , 243 ± 22 , 1504 ± 137 , 6 ± 1 , 1504 ± 137 , 6402 ± 800 , 51 ± 5 , 1278 ± 117 , 17055 ± 1549 , 48 ± 6 and 24 ± 3 mg/kg, respectively). Among them, K, Mg, Ca, Mn and Fe were the **dominated main** elements in the near-surface soil.

Three nutrients (P, S ad K) were found to be hyperaccumulated in all seeds with respect to their mean soil values, with bioaccumulation factors in the **range of** 19 to 44, 4 to 10 and 2 to 11 **range**, respectively (with mean values of 29, 7 and 6, respectively). On the other hand, in HS leaves, Cl and K were the elements that were strongly hyperaccumulated (**with** bioaccumulation factors of 49 and 17, respectively).

Correlation Coefficients

The taxonom~~iesy~~ of the eight Lamiaceae plants under study are different, and they can be grouped into three classes: (i) GA and TG; (ii) HS and LN; and (iii) OA, OS, OT and OV. The correlation coefficient values for the different parameters under analysis were computed for group (iii) and are shown in **Table 2**. A high correlation among oil, S, P, K and Cl nutrients was

Formatted: Font: Not Italic

found. Several elements (Mg, Ca, Sr, Ba, Ti, V, Cr, Mn and Fe) featured high correlations, suggesting their role as cofactor elements in the bioaccumulation of each other. Rb co-factor element showed a good correlation with heavy metals (Ti, V, Cr, Fe, Cu, Zn, Mo and Pb), while Co cofactor element had a high correlation with P, S, Mg, Ca and Sr. Elements i.e. Rb, Mg, Ca, Sr, Ba, V, Cr, Mn, Mo and Pb had an excellent correlation in their accumulation pattern.

Table 2. Correlation coefficients of chemical variables for the four *Ocimum* seeds under study.

	P	S	K	Cl	Rb	Mg	Ca	Sr	Ba	Ti	V	Cr	Mn	Fe	Co	Cu	Zn	Mo	Pb
P	1																		
S	0.98	1.00																	
Cl	0.62	0.75	1.00																
K	0.75	0.85	0.98	1.00															
Rb	-0.85	-0.87	-0.69	-0.75	1.00														
Mg	0.36	0.37	0.30	0.39	0.12	1.00													
Ca	0.03	0.03	0.00	0.07	0.46	0.94	1.00												
Sr	0.02	-0.07	-0.41	-0.29	0.50	0.73	0.85	1.00											
Ba	-0.08	-0.19	-0.53	-0.41	0.58	0.64	0.80	0.99	1.00										
Ti	-0.33	-0.36	-0.33	-0.30	0.77	0.73	0.92	0.86	0.86	1.00									
V	-0.33	-0.36	-0.33	-0.30	0.77	0.73	0.92	0.86	0.86	1.00	1.00								
Cr	-0.33	-0.36	-0.33	-0.30	0.77	0.73	0.92	0.86	0.86	1.00	1.00	1.00							
Mn	-0.16	-0.06	0.32	0.28	0.42	0.77	0.81	0.40	0.34	0.74	0.74	0.74	1.00						
Fe	-0.43	-0.44	-0.34	-0.33	0.82	0.67	0.88	0.79	0.80	0.99	0.99	0.99	0.76	1.00					
Co	0.78	0.73	0.33	0.49	-0.33	0.81	0.63	0.63	0.53	0.33	0.33	0.33	0.28	0.22	1.00				
Cu	-0.73	-0.84	-0.99	-0.99	0.79	-0.28	0.05	0.40	0.52	0.41	0.41	0.41	-0.20	0.43	-0.41	1.00			
Zn	-0.78	-0.88	-0.97	-1.00	0.74	-0.44	-0.12	0.22	0.35	0.26	0.26	0.26	-0.30	0.29	-0.55	0.98	1.00		
Mo	-0.33	-0.36	-0.33	-0.30	0.77	0.73	0.92	0.86	0.86	1.00	1.00	1.00	0.74	0.99	0.33	0.41	0.26	1.00	
Pb	-0.33	-0.36	-0.33	-0.30	0.77	0.73	0.92	0.86	0.86	1.00	1.00	1.00	0.74	0.99	0.33	0.41	0.26	1.00	1.00

Vibrational Characterization

The ATR-FTIR spectra of the seeds samples under study are depicted in **Fig. 2**. The main bands and their assignments are summarized in **Table 3** [24]. Peaks at around 3300 (ν O-H) are characteristic absorption bands from cellulose_-[25]. Other prominent bands attributed to cellulose are those at 1317-1305 cm^{-1} (C-H vibration), 1159-1154 cm^{-1} (ν C-O-C in bridge, asymmetric), 1059-1057 cm^{-1} ($>$ CH-O-CH₂), 1035-1027 cm^{-1} (ν C-O), 921-896 cm^{-1} (ν C-O-C in bridge, symmetric) and 668-665 cm^{-1} (β -glycosidic linkage). Peaks indicative of the presence of hemicellulose are those **which that** appear at 1644-1637 cm^{-1} (C=O stretching), 1378-1377 cm^{-1}

(-CH₃ symmetric deformation) and 1240 - 1236 cm⁻¹ (C-C-O asymmetrical stretching from acetylated glucomannan). Peaks at 2924 - 2922 cm⁻¹ (-CH₂ stretching) and 2854 - 2853 cm⁻¹ (-CH stretching) indicate the presence of cutine and wax. The peak that appears at 1651 cm⁻¹ can be attributed to C=C absorption of cellulose when it is cross-linked and dehydrated, but it **also** may also appear when the samples are rich in unsaturated oils. This band can be put in relation with the band at 1738 cm⁻¹ (C=O stretching), typical of hemicellulose, but which may also be associated with the stretching vibration of the ester carbonyl functional groups of triglycerides. The presence of this band, typical of the vinyl group, could justify the quantitative presence of unsaturated oils in the seeds under study. The sharp, intense C-H wags at 1000 - 997 cm⁻¹ are also indicative of vinyl. The bands that appear in *T. grandis* sample at 1606 cm⁻¹ (aromatic ring stretching) and at 896 cm⁻¹ are typical of lignin. The bands at 1520 - 1505 cm⁻¹ (aromatic skeletal vibration) are also typical of lignin. The presence of pectin is indicated by peaks at 1457 - 1455 cm⁻¹ (associated with O-CH₃ stretching) for pectic ester and at 1417 - 1411 cm⁻¹ (COO⁻ symmetric stretching vibration) for calcium pectate. [26]. Bands near 921 - 916 cm⁻¹ probably correspond to α -glycosidic linkage. The bands at 1710 cm⁻¹ (conjugated C=O), 1436 cm⁻¹ (CH₂ scissoring, known as the marker of crystallinity) and 814 cm⁻¹ (aromatic C-H out-of-plane bending or to C-O-C deformation) are not assigned to a particular plant fraction or component. The band at 721 cm⁻¹ (due to O-C=O in-plane deformation or a CH₂ rocking deformation) is attributed to phenolic components.




Figure 2. ATR-FTIR spectra of seed samples from species of the *Lamiaceae* family. Some offset has been added to the y axis for clarity purposes.

An interesting feature for the *Lamiaceae* is that they show vibrational spectra similar to those exhibited by other families of the Asterid clade (*Sapotaceae* and *Asteraceae*). Bands at 1651 cm^{-1} and $1644 - 1637\text{ cm}^{-1}$ are in good agreement with those exhibited in *Sapotaceae* for their kernel and coat fractions, respectively [27]. The absence of the bands at 1015 cm^{-1} and $\sim 878\text{ cm}^{-1}$ (apart from that of 795 cm^{-1}) is also a common feature with the *Asteraceae* (unpublished results). The absence of bands at 1574 , 1563 , 1336 , 1138 , 1076 , 857 , 778 and 757 cm^{-1} has been previously observed for *Salvia hispanica* L. [28].

Formatted: Font: Not Italic

Table 3. Main absorption bands in the ATR-FTIR spectra of seed samples from eight species of the *Lamiaceae* family and their assignments (all wavenumbers are expressed in cm^{-1}) [24,29,30].

<i>Gmelina arborea</i> kernel	<i>Hyptis suaveolens</i> whole seed	<i>Leonotis nepetifolia</i> whole seed	<i>Ocimum americanum</i> whole seed	<i>Ocimum sanctum</i> whole seed	<i>Ocimum tenuiflorum</i> whole seed	<i>Origanum vulgare</i> whole seed	<i>Tectona grandis</i> whole seed	Assignments
3284	3294	3289	3289	3282	3288	3289	3345	O-H stretching (cellulose)
2922	2924	2922	2924	2924	2924	2924	2923	-CH ₂ stretching (cutine, wax and pectin)
2853	2954	2853	2854	2853	2853	2853		-CH ₂ stretching (cutine and wax)
1745	1744	1744	1743		1743	1743	1738	C=O stretch (uronic ester groups)

								in hemicellulose)
	1710	1709	1710					C=O stretching of aldehyde/ketone
	1651			1651	1651	1651		C=C (cellulose) / COO ⁻ symmetric stretch
1637	1644	1640	1644	1639	1645	1640	1600	C=O stretching (hemicellulose) / C=C-C=C (cellulose) / aromatic ring stretch (lignin)
1535	1538	1540	1537	1539	1538	1539	1538	COO ⁻ symmetric stretching
			1519		1520	1520	1505	aromatic skeletal
1457	1456	1456	1455	1455	1456	1455	1455	O-CH ₃ stretching
	1417	1417	1417	1436	1411	1417	1417	CH ₂ scissoring
								COO ⁻ stretching / typical of pyranoside
	1397		1393				1393	CH rocking
1377	1377	1378	1378	1378	1378		1372	-CH ₃ symmetric deformation (hemicellulose) stretching of C-H bending in the CH ₂
	1315		1305		1307	1312	1317	C-H (cellulose)
1235	1238	1236	1237	1240	1236	1236	1236	C-C-O asymm stretching, acetylated glucomannan
1158	1158	1143	1159		1158	1156	1154	C-O-C in bridge, asymmetric (cellulose)
1117								C-O (cellulose)
1096	1096	1094	1097		1097	1097	1097	C-O-C stretch in pyranose
			1057		1059	1058		>CH-O-CH ₂ (cellulose) / -C-O-H (fructose)
	1035			1027			1027	C-O stretching (cellulose) / -C-O-H (glucose)
997					1001			C-H wags, vinyl
	916				921		896 814	C-O-C symmetric / glycosidic linkages (cellulose); aromatic C-H out-of-plane binding or C-O-C deformation
721		721						O-C=O in-plane def. or a CH ₂ rocking def
698		691	695	694	699	695		cis C=C
	665	668		665		668	666	β -glycosidic linkage (cellulose)
525	510	518	506		525	518		saccharide moieties

Thermal Characterization

Differences in weight loss for the different seeds of the *Laminaceae* family can be observed in the TG curves depicted in **Fig. 3** [Error! Reference source not found.](#) Additional DTG peaks and DSC thermal effects are shown in **Fig. 4**. The main endotherm at around 110 °C can be related both to dehydration and to gelatinization of starch (an order-disorder transition for the starch/moisture system). The small endotherm at around 200 °C indicates seed protein

crystallization to β -crystals accompanied by the random-coil \rightarrow β -form conformational transition. The chain of thermal events above 240 °C began with those related to oxidation reactions, followed by those attributed to the degradation of hemicellulose and other fiber components (at around 300 °C), and by those associated with the decomposition of the polyunsaturated (at 357 °C) and mono-unsaturated (at 391 °C) triglycerides. The very slow weight loss above 450 °C can be attributed to remaining lignin mass loss.





Figure 3. Comparison of TG curves for the seeds from eight species of the *Lamiaceae* family.






Figure 4. DSC (dotted line, y-axis on the left side of the graph), TG (solid line, first y-axis on the right side of the graph) and DTG (dashed line, second (rightmost) y-axis on the right side of the graph) curves for the seeds from eight species of the *Lamiaceae* family under study.

CONCLUSIONS

The nutritional potential of the seeds from eight *Lamiaceae* species was assessed by determining [their](#) proximate and phytochemical composition. Results indicated that the oil contents ranged from 11.8 to 50.4 mg/kg, with the highest levels for *G. arborea* and *T. grandis*. Total polyphenols varied from 295 to 5842 mg/kg (lower than in leaves) and mineral elements from 11756 to 33927 mg/kg (with concentration following the Ca>K>P>S>Mg>Fe>Zn sequence). Three main nutrients (P, S and K) were found to be hyperaccumulated in all seeds with respect to their mean soil values. Some toxic elements as Ba and Pb were found, albeit at

Formatted: Font: Not Italic

low levels. These results suggest that the seeds of the Lamiaceae studied species can be nutritive despite the presence of some toxic components.

Formatted: Font: Not Italic

The thermal profiles displayed by basil species (*O. americanum*, *O. tenuiflorum*, *O. sanctum*) and *O. vulgare* differed from those of *H. suaveolens*, *L. nepetifolia* and *T. grandis*, and all of them were clearly distinguishable from that of *G. arborea*. Differences should be related to lipids (tryglycerides) and lignin contents.

CONSENT

Not applicable.

ETHICS APPROVAL

Not applicable.

REFERENCES

1. Raja RR. Medicinally potential plants of Labiatae (Lamiaceae) family: an overview. Research Journal of Medicinal Plants. 2012;6(3):203-13. DOI: 10.3923/rjmp.2012203.213
2. Mechergui K, Jaouadi W, Coelho JA, Serra MC, Khouja ML. Biological activities and oil properties of *Origanum glandulosum* Desf: a review. Phytothérapie. 2016;14(2):102-108. DOI 10.1007/s10298-015-0963-x
3. Klimankova E, Holadova K, Hajslova J, Cajka T, Poustka J, Koudela M. Aroma profile of five basil (*Ocimum basilicum* L.) cultivars grown under conventional and organic conditions. Food Chemistry. 2008;107:464-472. DOI:10.1016/j.foodchem.2007.07.062

Formatted: Portuguese (Portugal)

4. Pathala D, Harini A, Hegde PL. A review on Gambhari (*Gmelina arborea* Roxb.). Journal of Pharmacognosy and Phytochemistry. 2015;4(2):127-132.
5. Dhawan NG, Khan AS, Srivastava P. A general appraisal of *Leonotis nepetifolia* (L.) R.Br: an essential medicinal plant. Bulletin of Environment, Pharmacology and Life Sciences. 2013;2(8):118-121.
6. Kalita J, Khan ML. Commercial potentialities of essential oil of *Ocimum* members growing in North East India. International Journal of Pharmacy & Life Science. 2013;4(4):2559-2567.
7. Uritu CM, Mihai CT, Stanciu GD, Dodi G, Alexa-Stratulat T, Luca A, Leon-Constantin, MM, Stefanescu R, Bild V, Melnic S, Tamba BI. Medicinal plants of the family Lamiaceae in pain therapy: a review. Pain Research Management. 2018;8:7801543. DOI: 10.1155/2018/7801543. **eCollection 2018**
8. Goswami DV, Nirmal SA, Patil MJ, Dighe NS, Laware RB, Pattan SR. An overview of *Tectona grandis*: chemistry and pharmacological profile. Pharmacognosy Review. 2009;3(5):170-174.
9. Jesus NZT, Falcão HS, Lima GRM, Caldas Filho MRD, Sales IRP, Gomes IF, Santos SG, Tavares JF, Barbosa-Filho JM, Batista LM. *Hyptis suaveolens* (L.) Poit. (Lamiaceae), a medicinal plant protects the stomach against several gastric ulcer models. Journal of Ethnopharmacology. 2013;150(3):982-988. <https://doi.org/DOI:10.1016/j.jep.2013.10.010>
10. Carović-Stanko K, Petek M, Grdiša M, Pintar J, Bedeković D, Čustić MH, Šatović Z. Medicinal plants of the family Lamiaceae as functional foods – a review. Czech Journal of Food Science. 2016;34(5):377-390. DOI: 10.17221/504/2015-CJFS

Formatted: Font: Italic

11. Krishna MS, Jayakumaran NA. Antibacterial, cytotoxic and antioxidant potential of different extracts from leaf, bark and wood of *Tectona grandis*. International Journal of Pharmaceutical Science and Drug Research. 2010;2(2):155-158.
12. Bozin B, Mimica-Dukic N, Simin N, Anackov G. Characterization of the volatile composition of essential oils of some Lamiaceae spices and the antimicrobial and antioxidant activities of the entire oils. Journal of Agricultural and Food Chemistry. 2006;54(5):1822-1828. <https://doi.org/DOI:10.1021/jf051922u>
13. Marques JL, Volção LM, Funk GD, Kroning IS, Silva WP, Fiorentini AM, Ribeiro GA. Antimicrobial activity of essential oils of *Origanum vulgare* L. and *Origanum majorana* L. against *Staphylococcus aureus* isolated from poultry meat. Industrial Crops and Products. 2015;77:444-450.
14. Basumatary S, Deka DC, Deka DC. Composition of biodiesel from *Gmelina arborea* seed oil. Advances in Applied Science Research. 2012;3(5):2745-2753.
15. Bachheti RK, Rai I, Joshi A, Satyan RS. Chemical composition and antimicrobial activity of *Hyptis suaveolens* Poit. seed oil from Uttarakhand State, India. Oriental Pharmacy Experimental Medicine. 2015;15(2):141-46. DOI: 10.1007/s13596-015-0184-8
16. Joshi MC. Handbook of Indian Medicinal Plants. Scientific Publishers, India. 2007. ISBN-10: 8172334508
17. Górnaś P, Rudzińska M, Segliņa D. Lipophilic composition of eleven apple seed oils: a promising source of unconventional oil from industry by-products. Industrial Crops and Products. 2014;60:86-91. <https://doi.org/DOI:10.1016/j.indcrop.2014.06.003>

Formatted: Portuguese (Portugal)

18. AOAC Official Method. Resistant sStarch in sStarch and pPlant mMaterials eEnzymatic dDigestion fFirst aAction 2002.
19. Bertaud F, Tapin-Lingua S, Pizzi A, Navarrete P, Petit-Conil M. Characterisation of industrial barks for their tannin contents for further green-wood based adhesives applications. InTech Fibres COST FP0901-Hamburg 2010.
20. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. 1999;299:152-178.
21. Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis*. 2002;10(3):178-182.
22. Angers P, Morales MR, Simon JE. Fatty acid variation in seed oil among *Ocimum* species. *Journal of the American Oil Chemists Society*. 1996;73(3):393-395.
23. Bachheti RK, Sharma A, Rai I, Joshi A, Mamgain R. Fatty acid composition and elemental analysis of seed oil of *Tectona grandis* collected from Dehradun, Uttarakhand, India. *International Journal of ChemTech Research*. 2012;4(3):1119-1123.
24. Stuart BH. *Infrared spectroscopy: fundamentals and applications*. *Methods Analytical Techniques in the Sciences*, Ando, DJ (Ed), England, Wiley 2004;8.
25. Wang XQ, Zhou RW, Groot G, Bazaka K, Murphy AB, Ostrikov K. Spectral characteristics of cotton seeds treated by a dielectric barrier discharge plasma. *Scientific Reports*. 2017;7(1):5601. DOI: 10.1038/s41598-017-04963-4

Formatted: Portuguese (Portugal)

Formatted: Font: Italic

Comment [AM20]: I did not find cited in the text

Comment [PMR21]: Fixed. Now it is cited in the text.

26. Fengel D, Ludwig M. Möglichkeiten und Grenzen der FTIR-Spektroskopie bei der Charakterisierung von Cellulose. I: Vergleich von Verschiedenen Cellulosefasern und Bakterien-Cellulose. Das Papier 1991;45(2):45-51.

27. Chakradhari S, Deb MK, Patel KS, Martín-Gil J, Towett EK, Martín-Ramos P. Evaluation of the phytochemical and mineral characteristics of some selected Sapotaceae plants. European Journal of Medicinal Plants. 2019;27(4):1-16. DOI: 10.9734/EJMP/2019/v27i430122

28. Carrión-Prieto P, Martín-Ramos P. et al. Vibrational analysis and thermal behaviour of *Salvia hispanica*, *Nigella sativa* and *Papaver somniferum* seeds. Pharmacognosy Journal. 2017;9(2):157-162.

29. Chakradhari S, Rajhans KP, Patel KS, Towett EK, Martín-Gil J, Martín-Ramos P. Nutritional and spectral characteristics of Terminalia plants. European Journal of Medicinal Plants. 2019; 27(4):1-13. DOI: 10.9734/ejmp/2019/v27i430120.

30. Chakradhari S, Patel KS, Towett E, Martín-Gil J, Martín-Ramos P. Chemical composition of Caesalpinioideae seeds. European Journal of Medicinal Plants. 2019;28(1):1-12. DOI: 10.9734/ejmp/2019/v28i130123

Formatted: Spanish (Spain, International Sort)