

13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

**Effect of Sorghum-Tigernut *Ibyer* (A Traditional Gruel) on the Fasting Blood Glucose Levels of Alloxan-Induced Diabetic Rats**

---

**ABSTRACT**

**Background:** There is growing interest in the use of natural foods in the management of chronic diseases like diabetes. *Ibyer* is a fibre rich gruel consumed amongst the Tiv people of Benue State made from whole sorghum or millet flours.

**Aim:** The aim of the study was to evaluate the effect of sorghum-tigernut *ibyer* on the fasting blood glucose levels and body weight of alloxan monohydrate-induced diabetic rats.

**Methods:** Sorghum flour (SF) and tigernut flour (TNF) were blended at different proportions (100:00; 90:10; 80:20; 70:30) for the purpose of *ibyer* production. The flour samples were subjected to proximate analysis using standard analytical procedures, the sensory attributes of *ibyer* produced from the different flour samples was evaluated on a 9-point hedonic scale. Thirty (30) male Wistar rats (100–180 g body weight) were grouped into five (1-5) each group containing six rats. They were induced with diabetes by injecting them with 150ml/kg of body weight with alloxan monohydrate dissolved in saline water (0.9% NaCl) except for group 1. Blood samples were collected from the tail of the rats, prior to induction, 48hrs after induction and 72hrs after three days of continuous feeding with test diet. Fasting blood glucose was measured using a standard glucometer and test strips.

**RESULTS:** The sensory attributes indicated that *ibyer* produced from the flour samples were generally acceptable. Fasting blood glucose levels after 72hrs of feeding were found to be lowered more in groups giving flours with a higher proportion of Tigernut.

**CONCLUSION:** The results indicated that sorghum-tigernut *ibyer* exerted hypoglycaemic effect on the experimental animals.

13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

*Keywords: Sorghum, Tigernut, Ibyer, Diabetic rats, Blood glucose level*

**1. INTRODUCTION**

Diabetes is a chronic disease in which food (particularly the nutrient – carbohydrate) is not properly absorbed in the body to be used as energy [1]. Diabetes and hypertension are global health disorders afflicting millions of people worldwide with an ever-increasing incidence and prevalence the upsurge of diabetes in Africa has been linked with rapid urbanization and changing dietary habits [2, 3]. Use of indigenous foods has been advocated to reduce the incidence of chronic, diet-related non-communicable diseases such as obesity, diabetes, cardiovascular diseases and stroke [4]. The development of diabetes mellitus, obesity, cancer and cardiovascular disease (CVD) has been reported to be linked to the intake of high glycaemic index (GI) foods while intake of low GI foods has been shown to play a positive role in the management of these diseases [5].

27 *Ibyer* is an indigenous non-alcoholic gruel made from cereals (maize, sorghum and millet),  
28 consumed in Nigeria especially in Benue state by Tiv people [6]. It is prepared by cooking  
29 reconstituted whole cereal flour or paste in water and is often served with beans products  
30 such as *akpukpa* (native bread) and beans cake. Sour type (*ibyer-i-angen*) arises as result of  
31 the fermentation step that is undertaken before the porridge is prepared. The flour or wet  
32 milled paste is usually reconstituted with water to form slurry or paste after which it is left for  
33 a given time to enable fermentative microbial organisms to act on the product, producing the  
34 characteristic sourness associated with the product. Traditionally, the fermentation time lasts  
35 overnight.

36 Sorghum (*Sorghum Bicolor L. Moench*) is the fifth most important cereal after wheat, rice,  
37 maize and barley in terms of production and utilization. The total world annual sorghum  
38 production is over 60 million tons from cultivated area of 45 mile for food, alcoholic  
39 beverages and the grain is one of the staple foods for poor and rural people. Sorghum is  
40 gluten-free thus can be consumed by people with celiac diseases [7]. Other important  
41 nutrients of sorghum include; dietary fibre, fat-soluble and B-vitamins and minerals [8].  
42 Sorghum flour is used for flours, porridge and side dishes, malted and distilled beverages  
43 and special food such as popped grain, its protein content is higher than many grains.  
44 Sorghum is rich in antioxidant which is believed to help lower the risk of cancer, diabetes,  
45 heart diseases and other neurological diseases.

46 Tigernut has been used extensively mainly for human consumption in Spain [9]. It was an  
47 important food in ancient Egypt [10]. Nowadays tigernuts are cultivated in Northern Nigeria,  
48 Mali, Senegal, Ahana and Togo where they are used primarily uncooked as a side dish [11].  
49 The flour is a good alternative for other flour like wheat flour, as it is gluten free and good for  
50 people who cannot take gluten in their diets. It is also used to make cakes and biscuits and  
51 the oil is used for cooking [12]. The dietary fibre content of tigernut is effective in the  
52 treatment and prevention of diseases such as colon cancer, coronary heart diseases,  
53 obesity, diabetes and gastro-intestinal disorders [13]. This research was aimed at  
54 determining the effect of sorghum-tigernut '*Ibyer*' on fasting blood glucose levels of alloxan  
55 monohydrate induced diabetic rats.

56  
57

## 58 **2.0 MATERIAL AND METHODS**

59  
60

### 61 **2.1 Sample Preparation**

62 The red sorghum grain variety and dried tigernuts were purchased from Wadata Market in  
63 Makurdi, Benue State, Nigeria. Alloxan monohydrate was purchased from Sigma Chemical  
64 Co. (St. Louis, MO, USA).

### 65 **2.2 Sample Preparation**


66 **2.2.1 Sorghum Flour:** The preparation of sorghum flour was done according to the  
67 method described by [14]. Sorghum grains were dry-cleaned by handpicking, washed in a  
68 large volume of water to remove impurities and sun-dried. The dried sorghum grains were  
69 milled into sorghum flour using hammer mill.

70 **2.2.2 Tigernut Flour:** Tigernut flour was prepared according to the method described by  
71 [15]. Dry tigernuts (brown variety) were sorted to remove unwanted materials like stones,  
72 pebbles and other foreign seeds before washing with tap water. The cleaned nuts were sun  
73 dried to a moisture content of about 13% then milled and sieved.

74  
75

76 **2.3 Formulation of Flour Blends:** the two flours (SF and TNF) were blended in a ratio  
77 of 100: 0, 90:10, 80:20 and 70:30 respectively.

78 **2.4 Preparation of *Ibyer-i-angen*:** After the preparation of the sorghum-tigernut flour  
79 blends, the flour samples were mixed with 200ml of water in five different containers with a  
80 cover and kept at ambient temperature ( $30\pm 2^{\circ}\text{C}$ ) for 24hrs to allow natural fermentation take  
81 place. The fermented paste from the two flour blends was diluted with water and cooked in a  
82 tower aluminium pot containing about 80cl of water on a gas cooker. The mixture was stirred  
83 for about 7-8min to achieve desired consistency. For feeding to the animals, the  
84 prepared/cooked *ibyer* was oven dried and milled.  
85 Flow chat for production of *ibyer-i-angen* is shown in figure.1  
86



96 **Fig.1: Traditional method of production of *Ibyer-i-angen***

97 **2.5 Determination of Proximate Composition**

98 The samples were separately analysed for moisture, ash, fat, and fibre contents respectively  
99 using standard methods of the Association of Official Analytical Chemists [16]. Crude protein  
100 contents ( $N \times 6.25$ ) were estimated from the crude nitrogen contents of the samples, as  
101 determined using Micro Kjeldahl method [17]. Carbohydrate content of the sample was  
102 determined by difference as described by [18].  
103

104 **2.6 Sensory Analysis**

105 Sensory characteristics of *ibyer-i-angen* prepared from sorghum and tigernut was evaluated  
106 and assessed by 20 semi-trained panellists of Department of Food Science and Technology,  
107 University of Agriculture, Makurdi who are familiar with *ibyer-i-angen*. Fresh samples of  
108 cooked porridge/gruel were assessed for appearance, taste, mouth feel, flavour and general  
109 acceptability. The judges recorded the quality characteristics of each sample using nine-  
110 point hedonic scale, where 1 = dislike extremely, 5 = neither like nor dislike, and 9 = like  
111 extremely as described by [19].  
112

113 **2.7 Animal Treatments**

114 Thirty (30) male Wistar rats (100–180 g body weight) were obtained from the small Animal  
115 Experimental Unit of the National Veterinary Research Institute Vom, Plateau State. The rats  
116 were housed under standard hygienic conditions in metal cages with wood shavings as  
117 bedding. Rats were also kept under natural thermal environmental conditions with ambient  
118 temperature of 24 C–26 C and relative humidity of 70%–80%, and approximately alternating  
119 12hr light/dark cycles. They were given access to a standard pelletized rat chow and water

120 *ad libitum*. The animal experiments adhered to the Guide for the Care and Use of Laboratory  
121 Animals.

122

## 123 **2.8 Induction of Experimental Diabetes**

124 Diabetes mellitus (DM) was induced in overnight-fasted rats by a single i.p. injection of  
125 freshly-prepared alloxan monohydrate, dissolved in a cold physiological saline (0.9% NaCl)  
126 solution at the dose rate of 150 mg/kg body weight. The animals were given free access to  
127 5% glucose solution in order to overcome the alloxan-induced hypoglycaemia for the first  
128 one-hour post-treatment with alloxan monohydrate. Blood glucose concentration of the rats  
129 was estimated 48hours after alloxan administration and DM was confirmed by analysis of  
130 blood samples, collected from the vein at the tip of the tail, using a portable blood  
131 glucometer and glucose test strips (On Call®Plus, Hannover, Germany). Animals with blood  
132 glucose concentration equal or more than 14 mmol/L were considered diabetic and used in  
133 the entire experimental group. Animals were weighed and randomly assigned to six groups  
134 and treated as follows:

135 Group 1, Control group: (normal saline only);

136 Group 2, Diabetic + Treated with Sample A (alloxan 150 mg/kg, i.p);

137 Group 3, Diabetic + Treated with Sample B (alloxan 150 mg/kg, i.p);

138 Group 4, Diabetic + Sample C (alloxan 150 mg/kg, i.p);

139 Group 5, Diabetic + Sample D (alloxan 150 mg/kg, i.p);

140 Experimental animals were fed thirty grams (30g) each of the test diet, after 48hrs of  
141 verification of diabetes, daily for 3 days.

142

## 143 **2.9 Measurement of Blood Glucose**

144 Experimental animals were rearranged according to the blood glucose concentration, except  
145 the control group, before commencement of treatment. Blood glucose concentration in all  
146 experimental groups were recorded following overnight fasting 72h after commencing the  
147 feed trials, using a portable glucometer (On Call®Plus, Hannover, Germany) and glucose  
148 test strips.

149

## 150 **2.10 Measurements of Body Weight**

151 Rats were weighed individually at Day 0 (before induction of DM), day 2 (post induction of  
152 DM) and 72 h (post-treatment with feed samples) using a digital precision weighing balance,  
153 and the body weights were recorded to calculate the body weight gains.

154

## 155 **2.11 Statistical Analysis**

156 Data collected were subjected to Analysis of Variance (ANOVA). Means were separated with  
157 Fisher's LSD using SPSS software (2009 model) and judged significantly different at 95%  
158 confidence level ( $p < 0.05$ ).

159

# 160 **3. RESULTS AND DISCUSSION**

## 161 **3.1 Results**

### 162 **3.1.1 Proximate composition of blends of sorghum and tigernut flours**

163 Table 1 shows the results obtained for the proximate composition of blends of sorghum and  
164 tigernut flours. Moisture, protein and carbohydrate showed a decreasing trend with  
165 increasing incorporation levels of Tigernut flour. For the rest of the parameters the trend was  
166 opposite.

167

### 168 **3.1.2 Sensory properties of lbyer produced from blends of sorghum and tigernut flours.**

169

170 The results of the sensory evaluation of lbyer produced from blends of sorghum and tigernut  
171 flours are shown in table 2. The sensory qualities measured were appearance, mouth feel,

172 taste, flavour and general acceptability. All the samples were generally acceptable though  
 173 sample B was most preferred.

174 **Table 1: Proximate composition of Blends of sorghum and tigernut flours**

| Parameters (%) | Samples                 |                         |                         |                         | LSD  |
|----------------|-------------------------|-------------------------|-------------------------|-------------------------|------|
|                | A                       | B                       | C                       | D                       |      |
| Moisture       | 10.1 <sup>a</sup> ±0.02 | 8.2±0.01 <sup>b</sup>   | 6.2 <sup>c</sup> ±0.01  | 5.5 <sup>d</sup> ±0.01  | 0.02 |
| Protein        | 7.5 <sup>a</sup> ±0.00  | 7.2 <sup>b</sup> ±0.00  | 7.0 <sup>c</sup> ±0.00  | 6.3 <sup>d</sup> ±0.01  | 0.02 |
| Fat            | 3.2 <sup>e</sup> ±0.01  | 4.8 <sup>d</sup> ±0.00  | 8.4 <sup>c</sup> ±0.00  | 10.2 <sup>b</sup> ±0.00 | 0.01 |
| Fibre          | 1.0 <sup>d</sup> ±0.03  | 2.1 <sup>c</sup> ±0.03  | 2.7 <sup>b</sup> ±0.01  | 3.2 <sup>a</sup> ±0.02  | 0.04 |
| Ash            | 2.1 <sup>d</sup> ±0.01  | 2.4 <sup>c</sup> ±0.00  | 2.7 <sup>b</sup> ±0.01  | 2.8 <sup>a</sup> ±0.00  | 0.01 |
| Carbohydrate   | 76.1 <sup>a</sup> ±0.02 | 75.3 <sup>b</sup> ±0.01 | 73.0 <sup>c</sup> ±0.02 | 72.1 <sup>d</sup> ±0.01 | 0.03 |

175  
 176 *Mean with different superscript within the same column are significantly different at (P<0.05).*

177 LSD: Least Significant Difference SF: Sorghum Flour; TNF: Tigernut flour.

178 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30%  
 179 TNF

180 **Table 2: Results of the Sensory scores of “lbyer” produced from blends of sorghum**  
 181 **and tigernuts flour.**

| Sample                | A                 | B                 | C                  | D                  | LSD  |
|-----------------------|-------------------|-------------------|--------------------|--------------------|------|
| Appearance            | 7.15 <sup>a</sup> | 7.05 <sup>a</sup> | 6.85 <sup>a</sup>  | 6.20 <sup>ab</sup> | 0.94 |
| Flavour               | 6.65 <sup>a</sup> | 6.75 <sup>a</sup> | 6.75 <sup>a</sup>  | 6.05 <sup>a</sup>  | 1.17 |
| Mouth feel            | 7.15 <sup>a</sup> | 6.95 <sup>a</sup> | 6.30 <sup>ab</sup> | 5.60 <sup>bc</sup> | 0.96 |
| Taste                 | 6.90 <sup>a</sup> | 6.75 <sup>a</sup> | 6.65 <sup>a</sup>  | 6.50 <sup>a</sup>  | 1.17 |
| General acceptability | 7.40 <sup>a</sup> | 7.15 <sup>a</sup> | 6.75 <sup>a</sup>  | 6.45 <sup>a</sup>  | 1.02 |

182 *Mean with different superscript within the same column are significantly different at (P<0.05).*

183 LSD: Least Significant Difference; SF: Sorghum Flour; TNF: Tigernut flour.

184 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30%  
 185 TNF

186

### 187 **3.1.3 Effect of Sorghum-tigernut lbyer on the Fasting Blood glucose levels of alloxan** 188 **induced diabetic rats.**

189 The results of the fasting blood glucose levels of alloxan induced diabetic rats before and  
 190 after feeding with test diets are presented in figure 2. The results showed that the average  
 191 fasting glucose levels of the diabetic rats before feeding with the test diets ranged from  
 192 22.30mm/L to 32.00mm/L and 3.15mm/L to 6.05mm/L in the non-diabetic rats (control  
 193 group). However, there was considerable reduction in average fasting blood glucose levels  
 194 of the diabetic rats 0hrs through to 72hrs post treatment. The results ranged from 5.52mm/L  
 195 from group 1 (control group) through to 5.45mm/L in group 5 with rats' group 2 having the  
 196 highest average fasting blood glucose levels of 17.60mm/L

197


### 198 **3.1.4 Body Weight of Rats before and after Feeding with Test Diet**

199 Figure 3 presents results of body weight changes during the experimental period. Results  
 200 showed a weight increase in the non-diabetic rat group from 98 to 118g from 0hrs to 72hrs

201 post feeding. However, the diabetic rat groups showed slight variations in body weight  
 202 changes.

203  
 204  
 205

**Fig. 2. Fasting blood glucose levels of alloxan induced diabetic rats before and after feeding with test diet.**



206  
 207  
 208  
 209  
 210  
 211  
 212  
 213


Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample B at 90%SF and 10%TNF) Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)

key:

SF; Sorghum flour

TNF: Tigernut flour

**Fig. 3. The average body weight of the rats (grams).**



214  
 215  
 216  
 217  
 218  
 219

Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample A at 90%SF and 10%TNF) Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)

Key: SF; Sorghum flour

TNF: Tigernut flour

220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272

## **3.2 Discussion**

### **3.2.1 Proximate composition of sorghum and tigernut flour blends.**

The moisture content of the sample flour decreased with increase in the tigernut flour substitution, which is due to the low moisture content found in tigernuts. The result is in agreement with those of [20] who reported a decrease in moisture content in flour. Low moisture content is an indication of good keeping quality of any food product; it also reduces microbial growth thereby extending the shelf life of the product.

The protein content decreased with increase in tigernut flour, which is attributed to the low protein content of tigernut as reported by [21]. The protein content of the four samples meets the recommended dietary allowance (RDA) of 0.8g of protein per kilogram of body weight per day [22].

The fat content of the samples increased steadily from samples A to D with increase in tigernut substitution, as tigernut has been reported to be a rich source of fats [21, 24, 25, 26]. Sorghum flour has been reported to contain fats of up to 4.25%, [18, 23]. The increase in the fat content could be attributed to the increase in the tigernut substitution. The fibre content of samples also increased with increase in tigernut substitution. These results were in accordance with [21] who reported an increase in the fibre content of tigernut-wheat flour with increasing tigernut flour substitution. There was a significant ( $P<.05$ ) difference in the ash content of all the samples as the level of tigernut substitution increased. The ash content result is in agreement with the reports of [27] and comparable with those of [28]. Ash content is an indication of the mineral content in flour.

There was however, a decrease in the Carbohydrate content of Samples with a corresponding increase in tigernut substitution. There was a significant ( $P<.05$ ) difference in carbohydrate content of all the samples. The decrease in carbohydrate content is attributed to an increase in tigernut flour.

### **3.2.2 Quality characteristics on the sensory attributes of lbyer produced from blends of sorghum and tigernut flour.**

The significant ( $P<.05$ ) difference in sample with the highest proportion of Tigernut could be attributed to the fact that tigernut has a colour that is distinct from that of sorghum. It was however noted that the sensory scores for appearance were considerable high even though with slight differences. This means all products were appealing to the eyes.

The significant difference in mouth feel in sample C and D could be attributed to higher amounts of fibre in samples C and D. *lbyer* is a gruel made from whole cereals, particularly sorghum and millet, due to this it has a coarse feel when consumed.

There was also no significant ( $P<.05$ ) difference among the samples in terms of taste, flavour and general acceptability. All the samples were generally acceptable indicating that *lbyer* of acceptable eating qualities can be produced from flour blends and sorghum and tigernuts. These results are in agreement with the reports of [6] who reported general acceptability of *lbyer* produced from sorghum and soy addition meaning addition of other legumes to the product may not affect the eating qualities of the product.

### **3.2.3 Effect of sorghum-tigernut lbyer on the Fasting Blood Glucose Levels Alloxan Induced diabetic rats**

The average fasting blood glucose of the rats in the various groups differed significantly. After three days of consuming approximately 30g of the test diets daily, their fasting blood glucose levels dropped significantly, showing that rat groups fed with samples containing higher amounts of tigernut had lower fasting blood glucose levels than those of the control group. This could be attributed to the fact that tigernut is a rich source of fibre [29, 30]. Fibre has been reported to exert some hypoglycaemic effects in subjects with type II diabetes. It is

273 also reported that arginine an amino acid (not determined) found in tigernut has the potential  
274 of stimulating the release of insulin thereby ameliorating the effect of diabetes [31]. This  
275 could be one of the reasons for the lowered blood glucose levels after three days of  
276 consecutive feeding with the test diet. Several studies have shown that sorghum extracts  
277 and sorghum rich diets exert hypoglycaemic effects in either human subjects or lab animals,  
278 thus agreeing with the findings of this research [32, 33].  
279  
280

#### 281 4. CONCLUSION

282 Results of the sensory evaluation of *ibyer* produced from sorghum and tiger flour blends  
283 showed that all the samples were generally acceptable. **The test diet was observed to have**  
284 **some hypoglycaemic effect on the experimental animals, however, mechanism of action of**  
285 **achieving these results was not studied and so remains unclear. Therefore, it would be**  
286 **worthwhile, to further investigate the effect of this diet rich in Tigernut on blood glucose**  
287 **levels.**

288 From the results, it is evident that nutritious diets can be formulated by complementing  
289 unexploited tuber and cereals like tigernut and sorghum respectively, such blends could be  
290 used to diversify their uses to develop new products.  
291

#### 292 REFERENCES

- 293 1. Bellows L and Nichols K (2012). Position Statement of the American Diabetes  
294 Association. Nutrition Recommendation and intervention for diabetes: Diabetes  
295 Care, vol.31,suppl.1.
- 296 2. Animaw W, Seyoum Y. Increasing prevalence of diabetes mellitus in a developing  
297 country and its related factors. PLoS One [Internet]. 2017 Nov 7;12(11):e0187670–  
298 e0187670. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29112962>
- 299 3. WHO. WHO | Double burden of malnutrition: Infographics. WHO [Internet]. 2017  
300 [cited 2019 Feb 7]; Available from: [https://www.who.int/nutrition/double-burden-](https://www.who.int/nutrition/double-burden-malnutrition/en/)  
301 [malnutrition/en/](https://www.who.int/nutrition/double-burden-malnutrition/en/)
- 302 4. Checkley W, Ghannem H, Irazola V, Kimaiyo S, Levitt NS, Miranda JJ, et al.  
303 Management of NCD in low- and middle-income countries. Glob Heart  
304 [Internet]. 2014 Dec;9(4):431–43. Available from:  
305 <https://www.ncbi.nlm.nih.gov/pubmed/25592798>
- 306 5. Mcculloch DK, Nathan DM, Lipman TO. Nutritional considerations in type 2 diabetes  
307 mellitus. 2016 [cited 2019 May 7];1–26. Available from:  
308 [https://www.uptodate.com/contents/nutritional-considerations-in-type-2-diabetes-](https://www.uptodate.com/contents/nutritional-considerations-in-type-2-diabetes-mellitus)  
309 [mellitus](https://www.uptodate.com/contents/nutritional-considerations-in-type-2-diabetes-mellitus)
- 310 6. Kure OA, Wyasu G. Influence of natural fermentation, malt addition and soya-  
311 fortification on the sensory and physico-chemical characteristics of Ibyer-Sorghum  
312 gruel. Adv Appl Sci Res [Internet]. 2013 [cited 2019 May 7];04(1):345–9. Available  
313 from: [http://www.imedpub.com/abstract/influence-of-natural-fermentation-malt-](http://www.imedpub.com/abstract/influence-of-natural-fermentation-malt-addition-and-soyafortification-on-thernsensory-and-physicochemical-characteristics-of-ibyersorghum-gruel-14775.html)  
314 [addition-and-soyafortification-on-thernsensory-and-physicochemical-characteristics-](http://www.imedpub.com/abstract/influence-of-natural-fermentation-malt-addition-and-soyafortification-on-thernsensory-and-physicochemical-characteristics-of-ibyersorghum-gruel-14775.html)  
315 [of-ibyersorghum-gruel-14775.html](http://www.imedpub.com/abstract/influence-of-natural-fermentation-malt-addition-and-soyafortification-on-thernsensory-and-physicochemical-characteristics-of-ibyersorghum-gruel-14775.html)
- 316 7. Taylor JRN, Achoch TJ and Bean SR (2006). Novel Food and Non-food uses for  
317 Sorghum and Millets. Articles in press *Journal of Cereal Science*, Vol.44: Pg.252-  
318 271.
- 319 8. Waniska RO, Poe JH and Bandy O.R. (2004). Effects of growth conditions on grain  
320 molding and phenols in sorghum Caryopsis, *Journal of cereal Science*, Vol. 10, Pg.  
321 217-225.

- 322 9. Gambo, A. (2012). Sensory evaluation of yoghurt produced from blend of Cowmilk,  
323 Soymilk and Tigernut milk. In: Proceedings of 36th Annual Conference/AGM of  
324 Nigerian Institute of Food Science and Technology, Lagos October. Pp.440- 441.
- 325 10. Neigbi OA (1992): Quality characteristics of Candies Produced from Tigernuts  
326 Tubers (*Cyperusesculentus*) and Melon Seeds (*Colocynthiscitrulus*) Milk Blends.  
327 *Global journal of Science Frontier Research: Agriculture and veterinary*. Vol.15 Issue  
328 2 Version
- 329 11. Omode MO, Ibiom OF and Okoi A. (2009): Studies on the fungi and phytochemical  
330 and proximate composition of dry and fresh tigernuts (*Cyperuseculentus*. L).  
331 *International Research journal of Biotechnology*. Vol.4 (1) pp. 11-14.
- 332
- 333 12. Adebayo-Oyetero AO, Ogundipe OO, Lofinmakin FK, Akinwande FF, Aina  
334 DO, Adeyeye SAO. Production and acceptability of chinchin snack made  
335 from wheat and tigernut (*Cyperus esculentus*) flour. *Cogent Food Agric*  
336 [Internet]. 2017 Jan 1;3(1):1282185. Available from:  
337 <https://www.tandfonline.com/doi/abs/10.1080/23311932.2017.1282185>
- 338 13. Forouhi NG, Misra A, Mohan V, Taylor R, Yancy W. Dietary and nutritional  
339 approaches for prevention and management of type 2 diabetes. *BMJ* [Internet]. 2018  
340 Jun 13;361:k2234. Available from:  
341 <http://www.bmj.com/content/361/bmj.k2234.abstract>
- 342 14. Sergio Serna-Saldivar (2010). *Cereal Grains: Properties, Processing, and Nutritional*  
343 *Attributes* CRC Press, ISBN 97814398156012010
- 344 15. Adejuyitan JA, Otunolam ET, Akandem ET, Bolarinwa IF and Oladokun FM (2009):  
345 Some physicochemical properties of flour obtained from fermentation of tigernut  
346 (*Cyperuseculentus*) sourced from a market u Ogbomoso, Nigeria. *African Journal of*  
347 *Food Science*. Vol.3(2) pp.051-055.
- 348 16. AOAC (2010): *Official methods of Analysis*, Association of Official Analytical  
349 Chemist Washington, D. C.
- 350 17. AOAC (2005): *Official methods of Analysis*, Association of Official Analytical  
351 Chemist Washington, D. C.
- 352 18. Ihekoronye and Ngoddy (1985): *Integrated Food Science and Technology for the*  
353 *Tropics*.
- 354 19. Iwe, M.O (2002). *Hand book of sensory Evaluation methods and analysis* Rojoint  
355 communication services Ltd. Enugu. Pp 72-75.
- 356 20. Abiose SH and Ikujenlola AV (2014): Comparison of chemical composition,  
357 functional properties and amino acids composition of quality protein maize and  
358 common maize (*Zea may* L.)
- 359 21. Ade-Omomowaye BIO, Akinwande BA, Bolarinwa IF and Adebisi AO (2008).  
360 Evaluation of tigernut (*cyperuseculentus*) -wheat composite flour and bread. *African*  
361 *Journal of Food Science* Vol.(2) pp.087-091.
- 362 22. Yemi B. (2010): *Malnutriton; the case of Death In Nutrition*, Pg. 785.
- 363 23. Ibrahim, F.S., E.E. Babiker, N.E. Yousif and A.H. El-Tinay, (2005). Effect of whey  
364 protein supplementation and/or fermentation on biochemical and sensory  
365 characteristics of sorghum flour. *J. Food Technol.*, 3: 118-125
- 366 24. Estoshola E and Oraedu AC (2012). Fatty acid composition of Tigernut tubers,  
367 Baobab seeds (*Adansoniadigitata* L.) and their mixtures. *American Journal of*  
368 *Chemical Sciences*. Vol.73: Pg. 255-267.
- 369 25. Zahras A and Ahmed MS (2014). Explaining the suitability of incorporating tigernut  
370 flour as novel ingredient in gluten-free biscuit. *Journal of Food and Nutritional*  
371 *Sciences*. Vol. 64, No1, pp 27-36.
- 372 26. Nwaignikpe R (2010). The phytochemical proximate and amino acid composition of  
373 the extracts of two varieties of tigernut (*Cyperuseculentus*) and their effects on  
374 sickle cell hemoglobin polymerization. *J.med. Med.sci*, 2010.1, 543-549.

- 375 27. Adebowale, A.A., Adegoke, M.T., Sanni, S.A., Adegunwa M.O., and Fetuga,G.O.  
376 (2012). Functional Properties and Biscuit Making Potentials of Sorghum-wheat Flour  
377 Composite. *American Journal of Food Technology*, 7: 372-379.
- 378 28. Aremu M. O, Bamidele TO, Agere H, Ibrahim H and Aremu SO (2015): Proximate  
379 Composition and Amino Acid Profile of Raw and Cooked Black Variety of Tigernut  
380 (*Cyperusesculentus* L.) Grown in Northern Nigeria. *Journal of Biology Agriculture*  
381 *and Health care*. Vol.5. No.7.
- 382 29. Sánchez ZE, Fernández LJ and Angel Pérez AJ (2012). Tigernut (*Cyperus*  
383 *esculentus*) Commercialization: Health Aspects, Composition, Properties and Food  
384 Applications. *Comprehensive Reviews in Food Science and Food Safety*. 11(4),  
385 pp.366–377.
- 386 30. Gambo A and Da' (2014). Tigernut (*cyperus esculentus*): composition, products,  
387 uses and health benefits – a review. *Bayero journal of pure and applied sciences*.  
388 Vol. 7(71), Pg.56– 61.
- 389 31. Wu GY and Meininger CJ (2000). Arginine nutrition and cardiovascular function.  
390 *Journal of Nutrition*. Vol. 130(11), Pg. .2626–2629.
- 391 32. Chung IM, Kim EH, Yeo MA, Kim SJ, Seo M and Moon HI (2011). Antidiabetic  
392 effects of three Korean sorghum phenolic extracts in normal and streptozotocin-  
393 induced diabetic rats. *Food Research International*. Vol. 44(1), pp.127–132.
- 394 33. Park, J. H., Lee, S. H., Chung, I. M., & Park, Y. (2012). Sorghum extract exerts an  
395 anti-diabetic effect by improving insulin sensitivity via PPAR-?? in mice fed a high-fat  
396 diet. *Nutrition Research and Practice*, 6(4), 322–327.  
397 <https://doi.org/10.4162/nrp.2012.6.4.322>

398  
399  
400