

Contribution of eggs and other cholesterol-containing foods to total dietary cholesterol intake, and their influence on serum lipid profile of adults in Calabar, Nigeria.

ABSTRACT

Aim: Eggs have sometimes been regarded as unhealthy foods due to their relatively high cholesterol content. The aim of this study is to determine contribution of eggs and other cholesterol containing food to total dietary cholesterol and their influence on serum lipid profile of adults.

Study design: Cross sectional and experimental.

Place and Duration of Study: Department of Biochemistry, University of Calabar, Calabar. February to July 2017.

Methodology: A cross-sectional survey to determine consumption pattern was carried out on 400 respondents using food frequency questionnaire (FFQ) and 24hour dietary recall. The respondents were further grouped into four based on their reported weekly egg consumption. A detailed follow-up study was conducted on 50 participants selected from across the 4 groups, using a 3-day repeated 24hour dietary recall to determine their consumption of egg and other cholesterol-containing foods. Serum blood lipid profile of these 50 participants was also determined using Randox cholesterol test kits. Food composition tables were used to calculate dietary cholesterol intake (DCI). The mean DCI of the 4 groups were cross-tabulated with mean serum cholesterol levels. Percentage contribution of eggs and other frequently consumed cholesterol-containing foods (such as milk and fish) to total DCI was calculated. Statistical significance was accepted at $p = .05$.

Results: For the follow-up participants, it was observed that results of correlations between DCI and the lipid profile parameters showed negative correlation (at $p = .01$) in both males and females, except slight positive correlations between cholesterol intake and HDL-c ($r=0.191$) among the males, and cholesterol with TC ($r=0.265$) among the females. Apart from this, no association was observed between DCI and the lipid profile parameters. Furthermore, the > 5eggs/week group had the lowest TC and LDL-c (4.23 ± 0.19 mmol/L and 2.38 ± 0.10 mmol/L). Based on the respondents' consumption patterns, eggs (boiled and fried) contributed the highest- 34.8% to total DCI, followed by milk (15.9%); salad cream contributed lowest (0.3%) to total DCI.

Conclusion: Increased DCI from cholesterol-containing foods (such as eggs), did not cause an adverse increase in serum cholesterol levels of normocholesterolemic people.

Keywords: blood lipid profile, eggs, cholesterol, foods

1. INTRODUCTION

With the increase in the prevalence of malnutrition and non-communicable diseases (NCDs) worldwide, it has become necessary to study the aetiology of growing number of diet-related diseases which populations are being faced with, in a bid to proffer solutions [1]. Dietary adjustments/modifications have also become quite popular and effective in the treatment and management of non-communicable diseases [2][3].

Healthy foods/diets are essential for maintaining good health and preventing diseases. The recent increase in the incidence of many NCDs worldwide has brought about a lot of research on the effect of various foods on people's health and wellbeing. Some foods are considered healthy depending on their

23 nutrient content while others are considered unhealthy. The nutrient composition of various foods
24 depends on several factors which include species, breeds, cultivars, ecological factors, post-harvest
25 handling, preservation and storage techniques [4].

26 Foods are either of plant or animal sources and comprise of various species/breeds. These foods may be
27 grouped broadly into: cereal grains, legumes, nuts and seeds, fruits, vegetables, milk and milk products,
28 meat and poultry, fish, eggs, fats and oils, fat replacers, roots and tubers, herbs and spices, sweeteners
29 [5]. Each of these groups consist of a wide variety of foods rich in the various essential nutrients. Animal
30 foods contain more complete proteins compared to plant foods which usually have what are called limiting
31 amino acids, but most animal-source foods have also been found to contain cholesterol – a compound
32 which has recently attracted quite some attention in the medical field. This is because increased serum
33 cholesterol levels (hypercholesterolaemia) is reported to be a major risk factor for cardiovascular diseases
34 such as hypertension and stroke [6].

35 Cholesterol, one of the most important and abundant steroids in the body, is found in the liver, bile salts
36 and skin where it forms vitamin D. Cholesterol in the body is obtained from animal food sources like eggs,
37 milk and meat; it is also synthesized in the liver from fats, carbohydrates and proteins. There is no
38 cholesterol in vegetables and plant foods [7]. There was also a fad that 'eggs are bad for your health'
39 because an egg yolk contains about 250 mg of cholesterol. In 2000, the American Heart Association
40 (AHA) revised its dietary guidelines and declared eggs to be nutritionally fit for healthy adults. The AHA
41 guidelines now allow an egg a day for healthy adults while still advising a total of daily cholesterol limit of
42 300 mg [5].

43 An article by the Cancer Care Ontario [8] reports that a healthy food/diet is one that helps maintain or
44 improve overall health. It provides the body with the essential nutrients in their right proportions. On the
45 other hand, an unhealthy diet is a major risk factor for many disease conditions like obesity, hypertension,
46 cardiovascular disease and cancer. Globally, unhealthy diets are estimated to cause about 19% of
47 gastrointestinal cancer, 31% of ischemic heart disease, and 11% of strokes [9,10], thus making it one of
48 the leading preventable causes of death worldwide [11]. According to WHO [12], NCDs such as heart
49 disease, stroke, cancer, chronic respiratory diseases and diabetes are the leading cause of mortality in
50 the world. About 38 million of the 56 million global deaths in 2012 were due to NCDs; and 48% of NCD
51 deaths in low- and middle- income countries in 2012 occurred before the age of 70. In Nigeria, NCDs are
52 estimated to account for 24% of total deaths and the probability of dying between the ages of 30 and 70
53 years from the four main NCDs (which are cancer, diabetes, cardiovascular diseases, and chronic
54 respiratory disease) is 20% [13].

55 Hypercholesterolaemia is a major risk factor for hypertension. Some research has also shown that the
56 effects of cholesterol-rich foods on serum cholesterol are small and clinically insignificant when compared
57 with the much greater effects of dietary saturated fatty acids on serum cholesterol [14]. Others suggest
58 that dietary cholesterol increases the ratio of total cholesterol (TC) to HDL-cholesterol (HDL-c) and hence
59 adversely affects an individual's lipid profile [15]; while other studies show that moderate consumption of
60 eggs, up to an egg a day, does not appear to increase the risk of heart disease in healthy individuals [16].

61 This study therefore, seeks to ascertain the influence of dietary consumption of cholesterol on the serum
62 profile of adults. It also aims at evaluating the contribution of some frequently consumed foods to the total
63 dietary cholesterol intake of a population.

64
65

66 2. METHODOLOGY

67
68

68 2.1 Consumption survey and Dietary assessment

69 2.1.1 Area of study

70 The study was carried out in the University of Calabar, Calabar in Cross River State. Cross River State
71 has a population of about 4million people with Calabar the State capital having a population of about
72 330,000 people (National Population Commission, 2006). The University of Calabar was established in
73 October, 1975. It is located in Calabar Municipal of Cross River State. This Federal University has one
74 Graduate School, 10 Faculties and 3 Institutes. Currently, from the records available at the University's
75 Registry, the student population stands at about 40,000, while the staff are about 3,000 bringing the total
76 population to about 43,000. The University community is comprised of people from different ethnic groups
77 in Nigeria and other nationalities like Cameroun, Ghana and Liberia; but the predominant tribes are the
78 Efiks, Ibibios and Ibos.

79 2.1.2 Population of the study

80 The population for the cross-sectional study consisted of the 3,000 staff- men and women within the age
81 range of 25 to 65 years, working at the University of Calabar, Calabar.

82 2.1.3 Sample size determination

83 This was calculated using Cochran's formula [17] as shown below:

$$84 \quad n = \frac{t^2 \times p(1-p)}{m^2}$$

85 n = required sample size

86 t = confidence level at 95% (standard value of 1.96)

87 p = estimated prevalence of hypercholesterolaemia in the area. According to a recent study by Akpa *et al.*
88 [18] carried out in Port Harcourt (South-South, Nigeria), the prevalence of hypercholesterolaemia was
89 31.5%.

90 m = margin of error at 5% (standard value of 0.05)

91
92 The calculated sample size of 332 was increased by 20% (66) to make room for contingencies like
93 dropouts, non-responses or incorrectly-filled questionnaires. The total sample size was rounded up to 400
94 adults
95

96 2.1.4 Sampling procedure

97 A two-stage sampling technique was employed for selecting the sample of the study. In the first stage,
98 University of Calabar was stratified into the 10 Faculties, 3 Institutes, Bursary, Registry and Vice
99 Chancellor's office (16 sample clusters in all). A list of staff in each of the 16 sample clusters was
100 obtained (sampling frame). In the case of faculties, the staff list was obtained from the various
101 departments. In the second stage, a number of participants proportional to the size of each cluster was
102 randomly selected for the study.

103 2.1.4.1 Exclusion criteria: Participants who did not meet the desired sample criteria- those who were
104 chronically ill, diabetic, hypertensive patients, pregnant and lactating mothers, were dropped from the
105 study (particularly the detailed dietary assessment) and replaced by others in the same sample cluster.
106 The health status of the participants was determined by observation and interaction, during which medical
107 history was taken.

108 2.1.4.2 Ethical approval: Appropriate ethical approval was obtained from the University of Calabar
109 Teaching Hospital (UCTH) for this research work.

110
111 2.1.4.3 Informed consent: An informed consent form was designed containing information on this
112 research.

113 The participants were made to read and then sign the informed consent form to formally indicate their
114 consent to participate in this study.

115

116 2.1.4.4 Questionnaire design and administration: A semi-structured questionnaire was designed to gather
117 information from the 400 participants who had read and signed the consent form. The questionnaire was
118 structured to gather socio-economic data, medical history, information on dietary intake (including egg
119 consumption pattern) and lifestyle of the participants. A food frequency questionnaire and 24 hour dietary
120 recall form was also attached. The questionnaires were filled mostly by interviewer-administered pattern
121 (in order to minimize errors) except in some cases where the respondents were literate enough to
122 complete them.

123
124 2.1.5 3-day 24 hour dietary intake study: Based on the responses from the questionnaire interviews
125 respondent were grouped according to their reported egg as found below.

- 126 1. Those who consumed more than 5 eggs per week (frequent egg consumers)
- 127 2. Those who consumed 3 to 5 eggs per week (moderate egg consumers)
- 128 3. Those who consumed less than 3 eggs per week (occasional egg consumers)
- 129 4. Those who do not consume eggs at all.

130 A detailed dietary assessment was conducted on 50 randomly-selected participants, using Gibson [19] 24
131 hour multi-pass dietary recall method. The 24 hour dietary recall was conducted on three different days of
132 the week (2 work days and 1 weekend day). Different sizes of solid materials and pictures from food
133 model materials (representing foods which have already been weighed), were used in order to increase
134 the accuracy of meat, fish, fruits and other foods quantification [19]. Standard measures and weights
135 were used to calculate the foods consumed by the participants based on the descriptions and
136 quantifications they gave. The weight of the foods consumed were converted into nutrients and calories
137 by the use of the West African Food Composition Table [20] and USDA National Nutrient Database [21]
138 was used in calculating the cholesterol content of the foods consumed and the dietary intake of
139 cholesterol by the participants. The percentage contribution of eggs and other foods to total dietary
140 cholesterol intake was also calculated and recorded. The mean individual's daily dietary cholesterol
141 intake was compared with the recommended dietary allowance (RDA). After dietary intake assessment,
142 the 50 participants were further grouped into 3 based on their DCI:

- 143 1. 0 – 150 mg/day
- 144 2. 151 – 300 mg/day
- 145 3. > 300 mg/day

146 **The mean serum cholesterol levels of these groups of participants were then calculated.**

147 2.2.1 Lipid profile analysis

148 On the last day of the dietary intake assessment carried out on the 50 participants, their blood samples (5
149 ml) were collected by venous puncture for fasting lipid profile tests, after a 12 hour overnight fast. Whole
150 blood samples were stored in clean sample bottles in the refrigerator for 1 to 2 days before analysis.
151 Serum lipid profile was determined using Randox Rx Monza analyser.

152 2.2.2 Statistical analysis

153 Responses from the questionnaire, were coded, and entered into the computer and analysed using
154 Microsoft Excel 2013 spreadsheets and SPSS version 20.0. Descriptive statistics such as frequencies,
155 percentages, mean and standard deviation from the means were used to present the results. Chi-square
156 analysis was used to determine the association between mean cholesterol intakes of the groups of
157 participants (based on reported egg consumption and DCI) and their mean serum cholesterol levels.
158 Linear correlation was also used to check the relationship between contribution of dietary cholesterol
159 intake from different foods and serum lipid profile results of the 50 participants. Significance was accepted
160 at $p = .05$.

161

162 **3. RESULTS AND DISCUSSION**

163

164 **3.1 Food Frequency analysis and Egg consumption pattern**

165 A summary of the personal/socio-demographic data of the respondents (as seen on Table 1a) showed
166 that all the respondents had a reasonable source of income and over 60% of them were earning
167 adequate income. About 87% of the respondents had obtained a degree from a tertiary institution. This
168 shows that most of them were educationally enlightened people. Analysis of the food frequency
169 questionnaire showed that egg (both boiled and fried forms) was the most frequently consumed food item
170 containing a significant amount of cholesterol. Pastries (such as cakes, pies and egg rolls) were also
171 frequently consumed and most of them are usually prepared using eggs. The percentage frequency
172 consumption of eggs by respondents, per week is shown in figure 1 below. Out of the 92.5% respondents
173 who consumed eggs, majority (73%) of them consumed < 3 eggs/week; 21% of them consumed 3 - 5
174 eggs/week while only 6% consumed > 5 eggs/week.

175 Table 1b shows the egg consumption pattern of the respondents. Out of the 400 staff surveyed, 92.5%
176 consumed eggs; 37.5% consumed only boiled eggs, 13.2% consumed only fried eggs while 34.3%
177 consumed eggs in both boiled and fried forms. Only 7.4% of the respondents who consumed eggs drank
178 them raw. The most frequently consumed species of eggs was the exotic chicken egg (83.5%). None of
179 the respondents was found to consume guinea fowl eggs and turkey eggs. The least consumed egg
180 species was quail egg which was consumed by only seven respondents (< 2%); out of which five persons
181 (71.4%) drank raw quail eggs. Most of the people who drank raw eggs did so occasionally (88.5%), and
182 not habitually (11.5%): only 4.2% drank up 3 -5 raw eggs in a week.

183 The result of the questionnaire survey showed that 91% of the respondents ate snacks and the most
184 frequently consumed snacks were pastries such as cakes, meat pies, and egg rolls. These pastry snacks
185 are usually made with eggs (as seen in the recipes), and this also contributed to the consumer's daily
186 dietary cholesterol intake. Egg was also widely consumed by people in the study population (93% of the
187 respondents either ate or drank eggs). Out of the five egg species, the exotic chicken egg was the most
188 consumed (both among the raw and boiled forms), obviously as a result of its availability and accessibility
189 (being relatively inexpensive). Scientific research has shown that exotic chicken egg is just as healthy as
190 the other eggs; only that some species like the quail and guinea fowl eggs have significantly higher
191 protein and micronutrient contents. Cholesterol content of the exotic chicken egg is lower than that of
192 turkey and guinea fowl eggs but higher than that of local chicken and quail eggs [22]. Most of the
193 respondents who consumed quail eggs took it in the raw form- probably for perceived health benefits.
194 Also, over half of the number of people who consumed eggs, ate < 3 eggs/week while almost all those
195 who drank raw eggs, did so occasionally (< 3 eggs/week). This was in line with the AHA [5] dietary
196 recommendations which says that an egg a day is safe for healthy adults, except in conditions where the
197 individual is genetically predisposed to hypercholesterolaemia. It was generally observed that the study
198 population comprised of relatively healthy individuals. Education and awareness goes a long way in
199 informing people of the need for consuming healthy diets and for healthy feeding practices, especially as
200 a person ages. This enables people to make enlightened food choices. Adequate income also ensures
201 increased purchasing power, access to healthy food and proper health care.

202

203

204

205

206


207

208

209

210

211 **Figure 1. Percentage frequency consumption of eggs**



212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230
231
232
233
234
235
236

Table 1a. Personal/Socio-demographic data of respondents

Variable	Sub-variable	Frequency	Percentage
Age group (years)	25 - 35yrs	182	46.1
	36 - 45yrs	114	28.8
	46 - 55yrs	62	15.7
	56 - 65yrs	37	9.4
	Total	395	100
Sex	Male	191	47.6
	Female	209	52.4
	Total	399	100
Educational attainment	No formal education	4	1.0
	Primary	4	1.0
	Secondary	44	11.0
	Tertiary	348	87.0
	Total	400	100
Marital status	Married	243	60.8
	Single	137	34.3
	Divorced	5	1.3
	Widowed	15	3.6
	Total	400	100
Income per month (in Naira)	<50,000	97	24.5
	50,000 to 100,000	128	32.3
	100,000 to 300,000	150	37.9
	>300,000	21	5.3
	Total	396	100
Occupation	Academic	95	24.1
	Administrative	221	55.9
	Medical Practitioner	65	16.5
	Technologist	14	3.5
	Total	395	100
Household size (persons)	<4	128	32.4
	4 – 6	164	41.5
	>6	103	26.1
	Total	395	100
Hypertensive	Response	Frequency	%
	Yes	21	5.3
	No	372	94.7
	Total	393	100
Diabetic	Yes	6	1.6
	No	380	98.4
	Total	386	100

History of BCDs	Yes	17	4.3
	No	313	79.4
	Don't know	64	16.3
	Total	394	100

*BCDs means blood cholesterol disorders

237
238
239

Table 1b. Frequency distribution of egg consumption by respondents

Variable	Egg consumption	n	%
Consume eggs	Consume	368	92.5
	Do not consume	30	7.5
	Total	398	100
Form of consumption	Raw	28	7.4
	Boiled	142	37.5
	Fried	50	13.2
	Scrambled	4	1.1
	Boiled & Fried	130	34.3
	Boiled, Fried and Scrambled	25	6.6
	Total	379	100
Type of eggs eaten	Exotic chicken	319	83.5
	Local chicken	21	5.5
	Turkey	0	0
	Quail	7	1.8
	Guinea fowl	0	0
	Chicken (both)	35	9.2
	Total	382	100
Drinking of raw egg	Drink	29	7.4
	Do not drink	362	92.6
	Total	391	100
Manner of consumption of raw eggs	Habitually	3	11.5
	Occasionally	23	88.5
	Total	26	100
Type of egg drank	Exotic chicken	16	61.6
	Local chicken	5	19.2
	Turkey	0	0
	Quail	5	19.2
	Guinea fowl	0	0
	Total	26	100
Number of eggs drank weekly	< 3 eggs	23	95.8
	3 to 5 eggs	1	4.2
	Total	24	100

240
241
242
243
244

245 3.2 Dietary cholesterol intake versus serum lipid profile.

246

247 Table 2 shows the total and mean nutrient intakes of the male and female respondents, and the
248 recommended nutrient intakes (RNI). The mean energy intake of the male respondents (10,175 KJ)
249 exceeded the RNI (8400 KJ) while that of the females (6,840 KJ) was not up to the RNI (8400 KJ). Also,
250 the mean cholesterol intake of the males (150 mg) and females (153 mg), did not exceed the RDA for
251 cholesterol (300 mg). Only 3 males and 4 females exceeded the RDA for cholesterol intake; and a female
252 participant had the highest cholesterol intake while 3 participants had the lowest cholesterol intake. Forty
253 eight percent of the participants had fat intake that exceeded the RNI.

254

255 Table 3 shows the average results of the lipid profile parameters for the four groups of participants who
256 were tested. The mean TC values for the four groups were all within normal range. Similarly, mean values
257 of Low Density Lipoprotein cholesterol (LDL-c), Very Low Density Lipoprotein cholesterol (VLDL-c) and
258 Triglycerides (TG) were within normal, only HDL values were slightly borderline high for all the groups
259 except the > 5 eggs a week group. All the lipid profile mean values for the groups were statistically
260 different ($p = .05$) from each other, despite having approximately similar values. The 3 - 5 eggs a week
261 group had the highest mean TC (4.83 ± 0.34 mmol/L) but the < 3 eggs a week group had significantly ($p =$
262 $.05$) higher LDL-c (2.78 ± 0.17 mmol/L) than the other three groups. The > 5 eggs a week group had the
263 lowest TC and LDL (4.23 ± 0.19 mmol/L and 2.38 ± 0.10 mmol/L) and this was significantly ($p = .05$)
264 different from the values of the other three groups. Table 4 showed similar result as table 3 but here the
265 participants were grouped according to their level of dietary cholesterol intake: those who consumed 150
266 – 300 mg of cholesterol per day showed slightly higher mean serum cholesterol values. The 3 groups had
267 their mean lipid profile parameters also within normal range. Statistical correlations of these 2 groupings
268 and their mean serum cholesterol levels showed no association between DCI and the serum lipid profile
269 parameters (at $p = .05$) and showed mostly negative correlation (at $p = .01$) in both males and females,
270 except slight positive correlations between DCI and HDL-c ($r = 0.191$) among the males, and DCI with TC
271 ($r = 0.265$) among the females. Apart from this, no association was observed between DCI and the lipid
272 profile parameters (at $p = .05$).

273

274 The mean nutrient intakes were compared with RNIs and RDA in the case of cholesterol and the mean
275 energy intakes for males exceeded their RNI but that of the females was not up to the RNI. This may be
276 as a result of people's (in this case, particularly women) awareness of the need to reduce caloric intake in
277 order to stay physically fit and prevent the occurrence of NCDs such as obesity and diabetes. This may
278 be attributed to the fact that the study population consisted of mostly literate and enlightened people.
279 Most African women tend to put on weight after childbirth and so they become more conscious of their
280 need to watch their weight by reducing caloric intake in order not to become obese. This is not the case
281 with men who do a lot of physical exercise and take a lot of calories to meet their bodies' energy
282 requirement. The fact that the men were also income-earners and had access to enough food, may also
283 help in explaining their meeting their RNI for energy. Also, the males and females exceeded the RNI for
284 carbohydrates and fats, while the females did not meet up with the RNI for protein. It should also be noted
285 that the participants (both males and females), did not meet up the RNI for dietary fibre which has been
286 found to help in lowering fat and cholesterol levels in the body [23]. Some micronutrients have been found
287 to play roles in cholesterol metabolism and these include micronutrients include vitamin A and C. Strong
288 clinical and experimental evidence suggests that chronic vitamin C deficiency results in
289 hypercholesterolaemia [24]. In a research carried out using rat models, Olivoros *et al.* [25] reported that
290 vitamin A deficiency induced a hypolipidemic effect by reducing serum cholesterol levels. In the case of
291 dietary cholesterol, most of the participants' intakes were within safe limits, with only seven people
292 exceeding the RDA of 300 mg. The contribution of eggs to each person's dietary cholesterol varied
293 widely. In the case of a certain respondent where egg contributed 98% of the dietary cholesterol, the
294 serum TC and LDL-c were within normal range. Similarly, in another respondent in whose case egg only
295 made up 38% of the dietary cholesterol intake, the lipid profile was also very normal; hence, eggs do not
296 appear to raise serum TC and LDL-c in normocholesterolemic individuals. In the overall picture, eggs
297 contributed most to the total dietary cholesterol intake of the 400 respondents. This was followed by milk
298 and meat pies.

299

300 As a result of the rising prevalence of NCDs such as hypertension and obesity, a lot of research is
 301 ongoing in the area of risk factors causing these diseases. The risk factors include hyperglycaemia,
 302 hypercholesterolaemia and unhealthy lifestyles. The results of this research showed that increase in
 303 dietary cholesterol intake did not cause a corresponding (unhealthy) increase in serum cholesterol levels.
 304 It is important to note here that this study focused mainly on healthy individuals hence an exclusion
 305 criteria was used (Table 1a showed that less than 5%. This agreed with the results of most similar
 306 researches carried out in different parts of the world such as that of Hu *et. al.* [16] and Natoli *et. al.* [26]. In
 307 studying the group of participants who consumed more than five eggs per week, it was observed that
 308 their TC and LDL-c levels were still within normal range. This implied that increased dietary cholesterol
 309 intake may only lead to hypercholesterolaemia in individuals with genetic or already-existing problems of
 310 dyslipidaemia. In order to maintain a healthy serum lipid profile, such individuals with dyslipidemias have
 311 to restrict their dietary intake of not only cholesterol but also that of fat (especially saturated and trans
 312 fats); not only from eggs but also from other dietary sources which have significant cholesterol levels such
 313 as meat and certain seafoods like prawns and shrimps (Note that since these foods are rich in essential
 314 nutrients, they may be consumed in smaller portions to reduce the risks of elevated serum cholesterol
 315 levels in such individuals). On the other hand, the slight positive correlation (among the males) between
 316 dietary cholesterol intake and HDL-c indicates that certain cholesterol-containing foods (including eggs)
 317 may actually cause an increase in the HDL-c (good cholesterol) which helps to reduce the LDL-c (bad
 318 cholesterol) levels in the blood thereby reducing the risk of some NCDs. McNamara [27] reported a
 319 similar finding which indicated that increase in dietary cholesterol also increased HDL-c thereby reducing
 320 the LDL/HDL ratio and risk of CHD. This should go a long way to dispel the myth that 'eggs are bad for
 321 your blood cholesterol' thereby allowing a lot of people (not children only), to benefit from the exceptional
 322 nutritional value of various species of eggs.

323
324
325
326
327
328 **Table 2. Mean nutrient intakes of participants.**
329

	Protein (g)	Fat (g)	Carb (g)	Fibre (g)	Ash (g)	Energy (KJ)	Calcium (mg)	Vit A (mcg)	Vit C (mg)	Cholesterol (mg)
Males										
Mean	56.9	85.6	270.2	16.9	26.5	10176.0	421.9	1241.8	99.8	149.7
RNI	46.0	65.0	130.0	25.0	21.0	8400.0	1000.0	700.0	75.0	300
% met	123.7	131.7	207.8	67.6	126.2	121.1	42.2	177.4	133.1	49.9
Females										
Mean	53.3	43.7	203.9	16.4	18.8	6840.0	331.2	1165.5	89.5	153.3
RNI	46.0	65.0	130.0	25.0	21.0	8400.0	1000.0	700.0	75.0	300
% met	115.9	67.2	156.8	65.6	89.5	81.4	33.1	166.5	119.3	51.1

330 *% met refers to the percentage of RNI/RDA that is met by the mean nutrient intake; Source for RNIs and RDA (for cholesterol):
 331 FAO/WHO (2002); FNRI (2002).
 332
333
334
335
336
337
338
339
340
341

342
343
344
345
346
347
348
349
350
351
352

Table 3. Lipid profile of subjects according to number of eggs consumed per week.

	N	TC (mmol/L)	TG (mmol/L)	HDL-c (mmol/L)	LDL-c (mmol/L)	VLDL-c (mmol/L)
Normal range		3.6 - 5.2	0.6 - 1.7	0.9 - 1.5	1.9 - 3.5	<0.8
No eggs	10	4.78 ± 0.28 ^b	1.21 ± 0.12 ^d	1.57 ± 0.10 ^d	2.65 ± 0.23 ^c	0.56 ± 0.06 ^d
< 3 eggs	20	4.83 ± 0.24 ^d	1.12 ± 0.09 ^b	1.54 ± 0.07 ^b	2.78 ± 0.17 ^d	0.52 ± 0.04 ^b
3 – 5 eggs	8	4.83 ± 0.34 ^c	1.20 ± 0.19 ^c	1.56 ± 0.10 ^c	2.59 ± 0.23 ^b	0.55 ± 0.09 ^c
>5 eggs	12	4.23 ± 0.19 ^a	1.04 ± 0.08 ^a	1.36 ± 0.11 ^a	2.38 ± 0.10 ^a	0.49 ± 0.04 ^a

Values are expressed as mean ± SEM

Values with different superscript in the same column are heterogeneous at p < 0.05

N represents number of persons in each group and differences in the size of N was responsible for similar numeric means being statistically different

TC means Total Cholesterol, TG means Triglyceride, HDL-c means High Density Lipoprotein cholesterol, LDL-c means Low Density Lipoprotein cholesterol, VLDL-c means Very Low Density Lipoprotein cholesterol

353
354
355
356
357
358
359
360
361
362
363
364

Table 4. Lipid profile of subjects according to mean daily dietary cholesterol intake.

Cholesterol Intake (mg/day)	N	TC (mmol/L)	TG (mmol/L)	HDL-c (mmol/L)	LDL-c (mmol/L)	VLDL-c (mmol/L)
		3.6 - 5.2	0.6 - 1.7	0.9 - 1.5	1.9 - 3.5	<0.8
0 – 150 mg	37	4.69 ± 0.16 ^a	1.11 ± 0.06 ^a	1.51 ± 0.06 ^a	2.67 ± 0.12 ^a	0.51 ± 0.03 ^a
151 – 300 mg	7	4.84 ± 0.40 ^a	1.29 ± 0.19 ^a	1.60 ± 0.11 ^a	2.64 ± 0.26 ^a	0.60 ± 0.10 ^a
> 300 mg	6	4.22 ± 0.17 ^a	1.05 ± 0.11 ^a	1.38 ± 0.10 ^a	2.33 ± 0.09 ^a	0.50 ± 0.04 ^a

Values are expressed as mean ± SEM

Values with similar superscript in the same column are homogeneous at p < 0.05

N represents number of persons in each group and differences in the size of N was responsible for varied numeric means being statistically similar

TC means Total Cholesterol, TG means Triglyceride, HDL-c means High Density Lipoprotein cholesterol, LDL-c means Low Density Lipoprotein cholesterol, VLDL-c means Very Low Density Lipoprotein cholesterol

365
366
367
368
369
370

371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404


3.3 Percentage contribution of egg and other cholesterol-containing foods to total dietary cholesterol.

Figure 2 shows the percentage contribution of various cholesterol-containing foods to the overall dietary cholesterol intake of all the respondents at the end of the dietary assessment period. Eggs (both boiled and fried added together) contributed the highest- 34.8%. This was followed by milk which had a total contribution of about 15.9%; next to this was meat pies (7.2%) and beef (6.7%). Periwinkle and salad cream had the lowest percentage contributions of 0.4% and 0.3%, respectively to the total dietary cholesterol intake. When the different forms of eggs consumed were calculated separately, boiled eggs contributed slightly higher than fried eggs to the overall dietary cholesterol intake.

These contributions are as a result of the cholesterol content of each food item and also the quantity of the food item consumed by the participants. Apart from eggs which have been reported to contain about 500mg/100 g, other foods liver and the sea foods- crayfish, prawns, shrimps also have a relatively high content of cholesterol per 100 g [28]. The reason why their percentage contributions may not be as high as that of egg is due to the fact that in this part of the world, such sea foods are not consumed in excessively large amounts nor are they consumed too frequently.

A correlation-regression analysis between the mean cholesterol intake from eggs, milk and fish (which were the three highest cholesterol-contributing foods in the study population), and the mean respondent's serum cholesterol levels, showed no association between DCI (in mg) from each of the foods and the mean serum cholesterol levels. Only a slight negative association ($r = -0.714$) was observed (at $p = .05$) between mean DCI from eggs and serum triglycerides concentration.

Figure 2. Percentage contribution of cholesterol-containing foods to total dietary cholesterol intake.



405
406
407
408
409
410
411
412
413
414
415
416

4. CONCLUSION

The results of the cross-sectional survey, dietary intake assessment and serum lipid profile tests showed no significant association between dietary cholesterol intake and the serum cholesterol levels since increased dietary cholesterol intake from cholesterol-containing foods (such as eggs), did not cause any adverse increase in serum cholesterol levels of normocholesterolemic people. Consequently, consuming an egg a day may not necessarily lead to hypercholesterolaemia (a risk factor for certain NCDs) in healthy individuals. It was also observed that among the study population, eggs contributed most to the total dietary cholesterol intake of the participants.

417

REFERENCES

418
419
420
421
422
423
424

1. FAO, IFAD & WFP (2014). The state of food insecurity in the world. Food and Agriculture Organisation of the United Nations, Rome.
2. Bhatnagar, D., Soran, H. & Durrington, P. N. (2008). Hypercholesterolaemia and its management. *British Medical Journal*, 337: a993.

425
426

3. American Health Association, AHA (2008). Diet and lifestyle recommendations. Available at: <http://www.americanheart.org/presenter.jhtml>

427
428

4. FAO (2013). Combating Micronutrient Deficiencies: Food-based Approaches. The Food and Agriculture Organization of the United Nations. Eds B. Thompson and L. Amoroso).

- 429 5. Mann, J. & Truswell, S. (2002). *Essentials of Human Nutrition*. 2nd Edition. Oxford University Press Inc.,
430 New York.
- 431 6. Li, Y, Zhou, X. & Li, L. (2013). Egg consumption and risk of cardiovascular diseases and diabetes: a
432 meta analysis. *Altherosclerosis*, 229: 524-530.
- 433 7. Timberlake, K.C. (2004). *General Organic & Biological Chemistry*. Pearson, USA.
- 434 8. Cancer Care Ontario (2012). Cancer Fact: Cancer and other chronic diseases share several risk
435 factors. <http://www.cancercare.on.ca/cancerfacts>.
- 436 9. World Health Organization (1990). Diet, nutrition, and the prevention of chronic diseases. Report of a
437 WHO study group. General (WHO Technical Report Series, No. 797).
- 438 10. NCD Alliance (2016). Unhealthy diets and obesity. A WHO attachment: Fact sheet.
- 439 11. Lopez, A. D., Mathers, C. D., Ezzati, M., Mamison, D. T. & Murray, C. J. (2006). "Global and Regional
440 burden of disease and risk factors, 2001: Systematic analysis of population health data". *Lancet*, 367
441 (9524): 1747–1757.
- 442 12. World Health Organization (2015). Global Health Observatory (GHO) data > Reports > World Health
443 Statistics.
- 444 13. World Health Organization (2014). Non-communicable Diseases (NCD) Country Profiles.
- 445 14. Gray, J. & Griffin, B. (2009). Eggs and dietary cholesterol dispelling the myth. British Nutrition
446 Foundation. *Nutrition Bulletin* 34; 66-70.
- 447 15. Weggemans, R. M., Zock, P. L. & Katan, M. B. (2001). Dietary cholesterol from eggs increases the
448 ratio of total cholesterol to high density lipoprotein cholesterol in humans: meta analysis. *American*
449 *Journal of Clinical Nutrition*, 73(5): 885-891.
- 450 16. Hu, F. B., Stampfer, M. J., Rimm, E. B. (1999). A prospective study of egg consumption and risk of
451 cardiovascular disease in men and women. *Journal of the American Medical Association*, 281(15): 1387-
452 1394.
- 453 17. Bartlette, J. E., Kotliik, J. W. & Higgins, C. C. (2001). Organizationl research: determining appropriate
454 sample size in survey research. *Information Technology, Learning, and Performance Journal*, 19(1): 43-
455 50.
- 456 18. Akpa, M. R., Agomouh, D. I. & Alasia, D. D. (2006). Lipid profile of healthy adult Nigerians in Port
457 Harcourt, Nigeria. *Nigerian Journal of Medicine*, 15(2): 137-140.
- 458 19. Gibson, R. S. (2005). *Principles of Nutritional Assessment* (2nded.). New York, New York: Oxford
459 University Press.
- 460 20. FAO (2012). West African Food Composition Table. Food and Agriculture Organisation of the United
461 Nations. Rome, Italy.
- 462 21. USDA (2016). USDA National Nutrient Database for Standard Reference, Release 23, Nutrient Data
463 Laboratory Home Page: <http://www.ars.usda.gov/nutrientdata>
- 464 22. Onyenweaku, E. O., Ene-Obong, H. N., Inyang, M. I., & Williams, I. O. (2018). Cholesterol and fatty
465 acid profiles of some bird egg varieties: Possible health implication. *Asian Food Science Journal*, Vol.
466 3(4): 1-9.
- 467

468 23. Brown, L., Rosner, B., Walter, W. W. & Sacks, F. M. (1999). Cholesterol-lowering effects of dietary
469 fibre: a meta-analysis. *American Journal of Clinical Nutrition*, 69: 30–42.

470 24. Turley, S. D., West, C. E. & Horton, B. J. (1976). The Role of Ascorbic acid in the regulation of
471 cholesterol metabolism and in the pathogenesis of Atherosclerosis. *Atherosclerosis*, 24: 1-18.

472 25. Oliveros, L. B., Domeniconi, M. A., Vega, V. A., Gatica, L. V., Brigada, A, M. & Gimenez, M. S. (2007).
473 Vitamin A deficiency modifies lipid metabolism in rat liver. *British Journal of Nutrition*, 97(2): 263-272.

474 26. Natoli, S., Markovic, T., Lim, D., Noakes, M. & Kotsner, K. (2007). Unscrambling the research: Eggs,
475 serum cholesterol and coronary heart disease. *Nutrition and Dietetics*, 64: 105-111.

476 27. McNamara, D. J. (2000). The impact of egg limitations on coronary heart disease risk: Do the
477 numbers add up? *Journal of American College Nutrition*, 19: 540-548.

478 27. USDA (2010). USDA National Nutrient Database for Standard Reference, Release 23, Nutrient Data
479 Laboratory Home Page: <http://www.ars.usda.gov/nutrientdata>.

480

481

482

UNDER PEER REVIEW