

## ***In vitro* study on anti-Salmonella activities of Boerhaavia diffusa (L. syn) Leaf extract**

### **Abstract**

Various strategies have been employed in the treatment and management of *Salmonella* infection however, *Salmonella* strains have gained resistance to antibiotics. This study was to determine *in vitro* anti-Salmonella activity of *Boerhaavia diffusa* leaf extract against clinical isolate of *Salmonella typhi* and *Salmonella typhi* ATCC 14028. The aqueous and ethanol extracts of *B. diffusa* were studied for their antibacterial activity against pathogenic *Salmonella typhi*. This study was carried out between April and September 2018. The *in vitro* antibacterial activity was performed by agar well diffusion method and broth dilution using spectrophotometric method and the results were expressed as the average diameter of zone of inhibition of bacterial growth around the well and optical density respectively. It was observed that aqueous extract exerted slightly higher activity than ethanolic extract as revealed by the mean diameter of zone of inhibitions at a concentration of 200 mg/ml, the aqueous extract had 35.21±0.47 mm (*Salmonella typhi* ATCC 14028) compared with ethanol extract 26.41±0.32 mm (clinical). However, in the broth dilution method, ethanol extract significantly (p=0.05) reduced the cell, at 48 hours, the optical density of clinical isolate of *S. typhi* treated at concentration of 200 mg/ml of extract was 0.47±0.02 nm while at the same concentration of extract, aqueous extract had an optical density of 0.52±0.11 nm respectively. The phytochemical assay revealed that tannin (5.18±0.02 mg/g) and quinone (8.45±0.13 mg/g) in ethanol extract was significantly (p=0.05) higher than aqueous extract while saponin (14.18±0.06 mg/g) was higher in the aqueous extract. The ethanol and aqueous extracts of leaves of *B. diffusa* whole plant exhibited significant antibacterial activity against both clinical and typed *Salmonella typhi*. Therefore, the plant extract could be used for the treatment of Salmonellosis, however, the *in vivo* studies is needed to ascertain the safety of the extract.

**Keywords:** Anti-*Salmonella* activity, plant extracts, agar well diffusion, broth dilution, *Salmonella* strains

## 32 **Introduction**

33 The bacterium *Salmonella typhi* causes typhoid fever [1, 2]. The bacterium is a gram-negative,  
34 motile, non-sporing, non-capsulated bacillus that can be contracted through contaminated water,  
35 milk, food or fruits and vegetables or via convalescent or chronic carriers [3]. It has also been  
36 linked with zoonotic transmission via reptiles and common domestic pets [4]. *Salmonella*  
37 *enterica*, which is a group of Gram-negative bacterial pathogens capable of infecting humans and  
38 animals, cause significant morbidity and mortality worldwide [5]. Certain serotypes adapted to  
39 humans, such as *Salmonella typhi* (*S. typhi*) and *Salmonella paratyphi* (*S. paratyphi*), usually  
40 cause severe diseases in humans, such as enteric fevers (typhoid and paratyphoid fevers). In most  
41 endemic areas like Africa, Asia, and Latin America, approximately 90% of enteric fever is  
42 typhoid. This disease is an important global health problem with an estimated 16 million cases  
43 and 600 000 deaths each year.

44 Various strategies have been employed in the treatment and management of *Salmonella*  
45 infection. Fluoroquinolones and tetracyclines are most commonly used to treat *Salmonella*  
46 infections. However, *Salmonella* strains resistant to these antibiotics have been reported in Korea  
47 and other countries [6]. One major concern to public health has been the global dissemination of  
48 *S. typhimurium* Definitive Type 104, which is resistant to cotrimoxazole, nalidixic acid and  
49 ampicillin [7]. The rise in antibiotic-resistant strains has led to increased interest in the use of  
50 plant materials to develop new effective drugs [6]. Moreover, conventional antityphoid drugs are  
51 becoming more and more unavailable to the common man in Africa due to increased cost [8].

52 The rise in antibiotic-resistant strains has led to increased interest in the use of plant materials to  
53 develop new effective drugs [6]. It has been reported that 80% of the world population are rural  
54 dwellers and rely on medicinal plants for their daily medications, also, plants have been  
55 reported to have minimal or no side effects compared to antibiotics [9, 10]. *Boerhaavia diffusa*  
56 (Spreading Hogweed in English), belonging to the family of the Nyctaginaceae, is mainly a  
57 diffused perennial herbaceous creeping weed of India (known also under its traditional name as  
58 *Punarnava*). *Boerhaavia diffusa* is traditionally known in Nigeria as *Etiponla* in Yoruba,  
59 *Azeigwe* in Igbo and *Babba-juju* in Hausa. *B. diffusa* is a perennial creeping weed, prostrate or  
60 ascending herb, up to 1 m long or more, having spreading branches [11].

61 The root, leaves, aerial parts and the whole plant of *B. diffusa* (L. syn) are used worldwide for  
62 the treatment of a number of disorders e.g. liver complaints, kidney disorders, rheumatism e.t.c.

63 [12]. The quest to identify and isolate novel phyto-compounds from *B. diffusa* has led many  
64 researchers to discover various compounds such as flavonoids, alkaloids, glycosides, steroids,  
65 triterpenoids, lipids, lignans, carbohydrates, proteins, and glycoproteins from its leaves, stems,  
66 seeds and roots [13]. Sourav [14], explored the Antibacterials from *Boerhaavia diffusa*. In his  
67 study, the chloroform and alcohol extracts of the plant were screened against six bacteria viz  
68 *Staphylococcus aureus*, *Escherichia coli*, *Proteus mirabilis*, *Salmonella typhimurium*,  
69 *Pseudomonas aeruginosa* and *Klebsiella aerogenosa*. Chloroform extract showed activity  
70 against *E. coli*, *S. typhimurium* and *P. aeruginosa* while the alcohol extract was active against *P.*  
71 *mirabilis* and *S. typhimurium*. The present study was undertaken to further investigate the  
72 antibacterial activity of *Boerhaavia diffusa* on typed and clinical strains of *Salmonella typhi*  
73 with the view to provide scientific evidence for its application as a medicinal plant.

#### 74 **Materials and methods**

##### 75 **Collection of leaves of *Boerhaavia***

76 Fresh leaves of *Boerhaavia diffusa* (Plate 1) were collected from the School of Health  
77 Technology, Oda Road, Akure, and identified in the Department of Crop, Soil and Pest  
78 Management, Federal University of Technology, Akure Ondo State.



79  
80 **Plate 1: Photograph of *Boerhaavia diffusa* leaf**

81

##### 82 **Preparation of plant extract**

##### 83 **Aqueous extraction**

84 The aqueous extractions of the water-soluble ingredient were carried out using the filter method.  
85 A 2 g of each of the grounded leaves were extracted by successive soaking for 2 days using 50  
86 ml of distilled water in a 250 ml sterile conical flask. The extracts were concentrated in vacuum  
87 at 60 °C and stored in universal bottles and refrigerated at 4 °C prior to use [15].

#### 88 **Ethanol extraction**

89 The organic solvent leaf extract was prepared by 2 g of plant mixture with ethanol and kept for  
90 two days. The extract was concentrated to one-fifth volume, filter sterilized and stored at 4 °C  
91 [15].

#### 92 **Test organism**

93 The clinical bacterial strains were obtained from the Department of Microbiology, Federal  
94 University of Technology Akure. Clinical *Salmonella typhi* and typed (ATCC 14028) *Salmonella*  
95 *typhi* were used. The isolates were confirmed based on cultural, morphological and biochemical  
96 characteristics following standard methods of identifying *Salmonella typhi* [16]. The bacterial  
97 strain was grown in nutrient broth for 12-18 hours at 37 °C on a rotary shaker. Cells were grown  
98 at 37 °C for 18 hours and cultures were kept at 4 °C.

#### 99 **Antimicrobial susceptibility tests**

##### 100 **Standardization of the inoculum**

101 The inoculum was prepared by inoculating colonies of fresh test cultures into sterile  
102 distilled water. The turbidity was compared to 0.5McFarland standard prepared according to the  
103 method of Cheesbrough [16]

##### 104 **Antibacterial susceptibility assay**

105 The extracts were dissolved and diluted using 50 % v/v dimethylsulphoxide (DMSO) to  
106 obtain different concentrations (50, 100 and 200 mg) in 1 mL. The 50 mg/ml, 100 mg/ml and  
107 200 mg/ml of the extracts of *B. diffusa* leaves were introduced into the wells of Muller Hinton  
108 agar plate. The plates were incubated aerobically at 37 °C and examined after 24 hours. The  
109 plates were examined for microbial growth inhibition and the Inhibition Zone Diameter (IZD)  
110 was measured to the nearest millimetre and compared with those produced by the commercial  
111 antibiotic ciprofloxacin which was used as control. Effect of extract on anti-*Salmonella* efficacy  
112 of the extract in the broth was also assayed using the spectrophotometric method, the  
113 absorbance of the tube was read at 620 nm [16, 17].

##### 114 **Antibiotics sensitivity test using commercial**

115 Antibiotics sensitivity test of the bacterial isolates were determined by the disc diffusion  
116 method as described by Cheesbrough [16]. Standard inoculum of 18 hours broth was spread on  
117 Muller Hinton agar using sterile swab in triplicate. The antibiotic discs were placed on the plate  
118 at equidistance. The plates were then incubated for 24 hours at 37°C and the diameter of zone of  
119 inhibition was measured and recorded. The commercial antibiotics discs (Fondoz Laboratories  
120 Ltd, Nigeria) used were; Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg, Ciprofloxacin  
121 (CPX) 10µg, Amoxicillin (AM) 25µg, Augmentin (AU) 30µg, Gentamycin (CN) 10µg,  
122 Pefloxacin (PEF) 5µg, Ofloxacin (OFX) 5µg, Streptomycin (S) 10 µg and Septra (SXT) 30µg.

### 123 **Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration** 124 **(MBC) of *Boerhaavia diffusa* Extracts**

125 The Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)  
126 of the extracts were determined using the broth (tube) dilution technique [18]. Dilutions of the  
127 extract in Mueller Hinton broth were prepared in tubes. The concentration of inoculum was also  
128 standardized to 0.5 McFarland's turbidity, The Mueller Hinton broth in tubes containing the  
129 different concentration of plant extract, 50, 100, and 200 mg/ml was then inoculated with 0.5 ml  
130 of the standardized culture. The tubes were then incubated at 37 °C for 24 hours. MIC and MBC  
131 values were recorded.

### 132 **Screening of phytochemical compounds**

133 The various solvent extracts of the powder of leaves of *Boerhaavia diffusa* were subjected to  
134 phytochemical tests for the identification of various action constituents using the method of  
135 Marcelin *et al.* [17]. The following major pharmaceutical valuable phytochemical compounds  
136 were analyzed qualitatively and quantitatively; alkaloids, phenols, tannins, flavonoids, quinones,  
137 saponins, terpenoids, sterols and cardiac glycosides.

### 138 **Statistical analysis of data**

139 Data obtained were subjected to analysis of variance and means were compared using Duncan's  
140 New Multiple Range Test (DNMRT) with the aid of SPSS software at  $p \leq 0.05$  level of  
141 significance.

### 142 **Results and Discussion**

143 *Salmonellosis* and enteric fever are always a public health concern in most developing countries,  
144 which are mostly low or middle-income countries with inadequate sanitation and hygiene,  
145 particularly, regarding food, water and disposal of human excreta [17]. Different plants and their

146 parts (flowers, buds, leaves, stem, bark, fruits, skin, pulp and root) have been used for thousands  
147 of years to enhance the flavour and aroma of food. In addition, plants are rich in a wide variety  
148 of second metabolites such as Alkaloids, Flavonoids, Phenols, which were found in vitro to have  
149 antimicrobial properties [17, 19].

150 In this study, extracts of *Boerhaavia diffusa* leaves were investigated for antibacterial activity  
151 against *Salmonella typhi*. Plant extracts were used to investigate antibacterial activity against  
152 two bacterial strains (Clinical *Salmonella typhi* and *Salmonella typhi* ATCC 14028). In this  
153 study, the antibacterial activity of *B. diffusa* leaf extracts was compared against the test bacteria  
154 with activities of model antibiotics.

155 The test organisms used for this study were identified based on biochemical characteristics  
156 common to *Salmonella typhi*. The result is presented in Table 1. The antibiotic sensitivity  
157 patterns of commercial antibiotics on the two strains of *S. typhi* are presented in Figure 1. The  
158 result revealed that the zones of inhibition of antibiotics against typed isolates were higher than  
159 that of clinical isolates however, chloramphenicol had the highest inhibition against the isolates  
160 (STC=24.30±0.42 mm, STT=24.36±0.07 mm). The higher antibacterial activity of model  
161 antibiotics is not surprising since the antibiotics are in a refined state. The standard antibiotics  
162 (ampicillin, amoxicillin, ciprofloxacin, ofloxacin, chloramphenicol) used in this study are the  
163 first line drugs employed in the treatment of typhoid fever [1].

164 The results of antibacterial activity of both water and ethanol crude extracts of *B. diffusa*  
165 showed anti*Samonella* activity on the two strains of *S. typhi* tested at different concentrations,  
166 with aqueous extract exerting slightly higher activity than ethanolic extract as revealed by mean  
167 diameter of zone of inhibitions, 200 mg/ml of aqueous extract had the highest (35.21±0.47)  
168 zone of inhibition (Figure 2). Minimum inhibitory concentration (MIC) and minimum  
169 bactericidal concentrations (MBC) of the extracts is shown in Tables 2. The ethanol and  
170 aqueous extract had the same MIC (100 mg/ml) on the typed isolate, also, there was no  
171 difference in the MIC and MBC of ethanol and aqueous extract on typed isolate. The aqueous  
172 and ethanol extracts exhibited different zones of inhibition against the isolates, however, the  
173 aqueous extract had higher zones of inhibition than ethanol extract. Antimicrobial action may be  
174 due to the synergistic action of different chemical constituents, some of which probably are lost  
175 upon extraction with solvent [15, 17, 20]. Water could be a better extraction solvent than  
176 ethanol for *B. diffusa* leaf, also, the demonstration of higher activity by the aqueous solvent may

177 be an indication that the phytoconstituents in the plant leaves are more soluble in water than the  
178 organic solvent [21]. The antimicrobial potential of *B. diffusa* and other plants sourced from  
179 traditional healers through an ethnobotanical survey of anti-infective plants in Egbado South in  
180 Ogun State, Nigeria was previously reported by Abo and Ashidi [22]. This study also  
181 corroborates the findings of Madani and Jain [23] who reported higher anti-*Salmonella* activity  
182 in aqueous extract of *Terminalia bellerica* than chloroform and acetone extracts. It has been  
183 reported that different phytoconstituents have different degrees of solubility in different types of  
184 solvents depending on their polarity. In a traditional setting, water is the solvent largely used to  
185 prepare these concoctions.

186 The anti-*Salmonella* efficacy of *Boerhaavia diffusa* extracts in the broth was assayed and was  
187 shown in Figure 3, 4, 5 and 6. The result presented in Figure 3 revealed the effect of ethanol  
188 extract on clinical isolate of *S. typhi*, it was noted that the extract significantly ( $p < 0.05$ ) reduced  
189 the cell, at 48 hours, the optical density of clinical isolate of *S. typhi* treated with 50, 100, 200  
190 mg/ml of extract were  $0.52 \pm 0.03$ ,  $0.50 \pm 0.10$ ,  $0.47 \pm 0.02$  nm respectively while at the same  
191 concentration of extract, aqueous extract had optical density of  $0.64 \pm 0.21$ ,  $0.54 \pm 0.03$ ,  $0.52 \pm 0.11$   
192 nm respectively (Figure 4). Also, the anti-*Salmonella* efficacy of *B. diffusa* ethanol extracts on a  
193 typed isolate of *S. typhi* is shown in Figure 5. It was observed that the extract significantly  
194 ( $p < 0.05$ ) reduced the cell, at 48 hours, the optical density was of typed isolate of *S. typhi* treated  
195 with 50, 100, 200 mg/ml of extract were  $0.49 \pm 0.00$ ,  $0.48 \pm 0.01$  and  $0.37 \pm 0.12$  while at the same  
196 concentration, aqueous extract had an optical density of  $0.62 \pm 0.03$ ,  $0.53 \pm 0.11$  and  $0.49 \pm 0.21$  nm  
197 respectively (Figure 6). It was noted from this study that plant extracts tested by microdilution  
198 technique and the optical density was measured after 48 hours showed that ethanol extract had  
199 higher anti-*Salmonella* activity compared to aqueous extract which was higher in values obtained  
200 from agar well diffusion technique. It could be that the bioactive components in ethanol extract  
201 did not diffuse into agar in agar well but was able to inhibit microbial cells directly in broth. This  
202 was previously reported by other findings that the active components of the extract do not diffuse  
203 into Muller Hinton agar, however, they were able to cause inhibition of microbial cells in broth  
204 microdilution [6, 24].

205 Both plant extracts (ethanolic and aqueous) were subjected to preliminary qualitative  
206 phytochemical evaluation. The phytochemical profiles of the two solvent extracts from the plant  
207 used in this study are presented in Table 2. The analysis revealed the presence of alkaloids,

208 phenol, glycosides, steroids, carboxylic acid, reducing sugar, flavonoids, saponins, tannins,  
 209 proteins, triterpenoids, quinines, carbohydrates and sterols. Also, tannin ( $5.18 \pm 0.02$  mg/g) and  
 210 quinone ( $8.45 \pm 0.13$  mg/g) in ethanol extract were significantly ( $p < 0.05$ ) higher than aqueous  
 211 extract while saponin ( $14.18 \pm 0.06$  mg/g) was higher in the aqueous extract. The preliminary  
 212 qualitative phytochemical screening carried out showed that the leaf extracts of *B. diffusa*  
 213 contain vital secondary metabolites such as alkaloids, saponins, tannins and glycosides. The  
 214 bioactive compounds in medicinal plants have been reported to be the active principles  
 215 responsible for the pharmacological potentials of medicinal plants [25]. The presence of these  
 216 chemicals in the leaves and root of these plants justify the local uses of these plants for the  
 217 treatment of various ill conditions. Phytoconstituents such as saponins, phenolic compounds and  
 218 glycosides have been reported to inhibit bacterial growth and to be protective to plants against  
 219 bacterial and fungal infections [6, 26]. Ethanol extract of *Boerhavia diffusa* leaves possess some  
 220 phytochemicals like Alkaloids, Anthraquinone, Glycoside, Flavanoids and Tannins. Saponins  
 221 are natural glycosides that act as hypoglycemic, antifungal and serum cholesterol lowering  
 222 agents in animals [27]. Saponins are essential elements in ensuring hormonal balance and  
 223 synthesis of sex hormones [28]. Tannins are bitter polyphenolic compounds that hasten the  
 224 healing of wounds. They also possess anti-diuretic and anti-diarrhoea properties [28].  
 225 Terpenoids were present in both ethanolic extract of and aqueous extracts of AOU and AFU.  
 226 Terpenoids have been found to be useful in the prevention and therapy of several diseases,  
 227 including cancer. Terpenoids are also known to possess antimicrobial, antifungal, antiparasitic,  
 228 antiviral, anti-allergenic, antispasmodic, antihyperglycemic, anti-inflammatory and  
 229 immunomodulatory properties [29]. The presence of these compounds promises its potential  
 230 application in the treatment of microbial ailment. However, tannins were present in aqueous  
 231 extract of but not in the ethanolic extract. Saponin and flavanoid are higher in the aqueous  
 232 extract of the leaf ( $14.18$  and  $11.26$  mg/g) than the ethanolic extract ( $6.36$  and  $9.98$  mg/g).


233 **Table 1: Biochemical characteristics of *Salmonella* strains**

Biochemical characteristics	<i>Salmonella typhi</i> (Clinical isolate)	<i>Salmonella typhi</i> (ATCC 14028)
Gram reaction	-ve	-ve
Shape	Rod	Rod
Motility	+ve	+ve
Catalase	+ve	+ve
Coagulase	-ve	-ve

Citrate	+ve	+ve
H <sub>2</sub> S	+ve	+ve
Lactose	-ve	-ve
Glucose	+ve	+ve
Fructose	+ve	+ve
Sucrose	-ve	-ve
Galactose	+ve	+ve
Indole	-ve	-ve
Methyl red	+ve	+ve
Voges-Proskauer	-ve	-ve
Oxidase	-ve	-ve

234 Key: -ve = negative +ve = positive

235




236

237 **Figure 1: Antibiotic sensitivity pattern of commercial antibiotic discs on *S. typhi* strains**

238 Key: STC – *Salmonella typhi* (clinical isolate), STT – *Salmonella typhi* (typed isolate),  
 239 Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg, Ciprofloxacin (CPX) 10µg, Amoxicillin  
 240 (AM) 25µg, Augmentin (AU) 30µg, Gentamycin (CN) 10µg, Pefloxacin (PEF) 5µg, Ofloxacin  
 241 (OFX) 5µg, Streptomycin (S) 10 µg and Septra (SXT) 30µg.

242



244

245 **Figure 2: Anti-Salmonella activity of *Boerhaavia diffusa* extracts**

246 **Key:** AQ = Aqueous extracts of *Boerhaavia diffusa*, ET= Ethanolic extracts of *Boerhaavia*  
 247 *diffusa*, Cpx= Ciprofloxacin

248


249

250 **Table 2: Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal**  
 251 **Concentration (MBC) of *Boerhaavia diffusa* extracts on *Salmonella***

<i>Boerhaavia diffusa</i> extracts	Ethanolic extract		Aqueous extract	
	S1	S2	S1	S2
MIC (mg/ml)	100	100	50	100
MBC (mg/ml)	50	100	50	100


252 Key: S1 - *S. typhi* (Clinical isolate), S2 – *S. typhi* (ATCC 14028)

253




254

255 **Figure 3: Effect of *Boerhaavia diffusa* ethanol extract on Clinical isolate**




256

257 **Figure 4: Effect of *Boerhaavia diffusa* aqueous extract on clinical isolate**



258

259 **Figure 5: Effect of *Boerhaavia diffusa* ethanol extract on typed isolate**



260

261 **Figure 6: Effect of *Boerhaavia diffusa* aqueous extract on typed isolate**

262 **Table 3: Qualitative analysis of phytochemicals in *Boerhavia diffusa* leaf extracts**

263	Phytochemical	Ethanolic extract	Aqueous extract
264	Alkaloids	-	+
265	Tannins	-	+
266	Flavonoids	+	+

267	Quinones	+	-
268	Saponins	+	+
269	Terpenoids	+	-
270	Sterols	+	-
271	Cardiac Glycosides	-	+
272	Phenols	+	+
273	Key : + = Present, - = Absent		

274  
 275 **Table 4: Quantitative phytochemical screening of aqueous and ethanol extracts of *B.***  
 276 ***diffusa***

Phytochemicals	Ethanol extract	Aqueous extract
Tannins (mg/g)	5.18±0.02 <sup>a</sup>	3.90±0.22 <sup>a</sup>
Quinones (mg/g)	8.45±0.13 <sup>b</sup>	6.60±0.31 <sup>a</sup>
Saponins (mg/g)	6.36±0.24 <sup>a</sup>	14.18±0.06 <sup>b</sup>
Triterpenoids (mg/g)	8.56±0.08 <sup>a</sup>	8.89±0.31 <sup>a</sup>
Steroid (mg/g)	9.03±0.11 <sup>a</sup>	6.73±0.14 <sup>a</sup>
Glycosides (mg/g)	30.39±0.06 <sup>b</sup>	28.29± 0.03 <sup>a</sup>
Flavonoids (mg/g)	9.98±0.61 <sup>a</sup>	11.26±0.33 <sup>a</sup>

277

278

279 **Conclusion**

280 Most of the antibiotics used nowadays have lost their effectiveness due to the development of  
 281 resistant genes in microbes. The antibiotics are sometimes associated with side effects such as  
 282 hypersensitivity, immune suppression and allergic reaction.

283 More interest is being shown in developing alternative antimicrobial drugs for the treatment of  
 284 infectious diseases without side effects. The results of our present study demonstrates anti-  
 285 *Salmonella* activity of aqueous and ethanol extract of *Boerhavia diffusa*, tannin and quinone  
 286 were higher in ethanol extract while saponin was higher in aqueous extract, using agar well  
 287 diffusion, the aqueous extract showed higher anti-*Salmonella* efficacy while the broth  
 288 microdilution examined by spectrophotometer revealed that ethanol extract had higher anti-  
 289 *Salmonella* efficacy. In the present study, the anti-salmonella activity of *Boerhaavia diffusa*  
 290 may be attributed to an individual or synergistic effect of phytoconstituents present in it. The  
 291 ethanol and aqueous extracts of leaves of *B. diffusa* whole plant exhibited significant  
 292 antibacterial activity against both clinical and typed *Salmonella typhi*. Therefore, the plant

293 extract could be used for the treatment of Salmonellosis, however, the in vivo studies is needed  
294 to ascertain the safety of the extract.

### 295 **Recommendation**

296 Based on our findings, it is therefore recommended that both agar well diffusion and broth  
297 dilution method should be used to affirm the antimicrobial efficacy of the plant extracts.

### 298 **References**

- 299 1. Prescott LM, Harley JP, Klein DA (2005). Microbiology, McGraw-Hill, Paris. pp. 937-  
300 960.
- 301 2. Doughari J.H (2005). A comparative study on effects of crude extracts of some local  
302 medicinal plants and some selected antibiotics on *Salmonella typhi* pp 1-15. M. Sc.  
303 Thesis Federal University of Technology, Yola; Adamawa State, Nigeria. Yola. pp. 1-4.
- 304 3. Doughari J.H, Pukuma M.S., De N. (2007). Antibacterial effects of *Balanites aegyptiaca*  
305 L. Drel. and *Moringa oleifera* Lam. on *Salmonella typhi*. *African Journal of*  
306 *Biotechnology*, 6:2212-2215.
- 307 4. Birgitta J., Yvonne A., Carl E (2005). Effect of regulation and education and reptile  
308 associated salmonellosis. *Emerging Infectious Diseases*, 11:398-403.
- 309 5. Christenson JC. *Salmonella* infections. *Pediatr Rev* 2013; **34**(9): 375-83.
- 310 6. Wangari D. U. (2017). *In vitro* study of anti-Salmonella activities of medicinal plants.  
311 *Global Journal of Medicinal Plants Research*. **3** (1): 301-308
- 312 7. Kariuki S, Revathi G, Kiiru J, Mengo DM, Mwituria J, Muyodi J, Munyalo A, Yik YT,  
313 Kathryn EH, Robert AK, Gordon D (2010). Typhoid in Kenya is associated with  
314 dominant multidrug-resistant *Salmonella enterica* Serovar Typhi haplotype that is also  
315 widespread in Southern Asia. *Journal of Clinical Microbiology* **48**(6):2171-2176.
- 316 8. Donald S. T., Donatien G., Siméon P. C. F., Charles F., Fabrice K. and Merline N. D.  
317 (2015). *In vivo* anti-salmonella activity of aqueous extract of *Euphorbia prostrata* Aiton  
318 (Euphorbiaceae) and its toxicological evaluation. *Asian Pacific Journal of Tropical*  
319 *Biomedicine*. **5**(4): 310-318
- 320 9. Bibitha B., Jisha V.K., Salitha C.V., Mohan S., Valsa A.K. (2002). Antibacterial activity  
321 of different plant extracts. *Indian Journal of Microbiology*, 42:361-363.
- 322 10. Maghrani M, Zeggwagh N, Michel J, Eddouks M (2005). Antihypertensive effect of  
323 *Lepidium sativum* L. in spontaneously hypertensive rats. *J. Ethnopharm.* **100**(102): 193-  
324 197.
- 325 11. Nayak P. and Thirunavoukkarasu M. (2016). A review of the plant *Boerhaavia diffusa*:  
326 its chemistry, pharmacology and therapeutical potential. *The Journal o*  
327 *Phytopharmacology*, **5**(2):83-92
- 328 12. Katampe I., Uwem M., and Uwem J. (2018). Antioxidant and Antimicrobial Effects of  
329 Aqueous and Ethanolic Extracts of Selected Medicinal Plants from Southern Nigeria.  
330 *Journal o Advances in Medical and Pharmaceutical Sciences*, **17**(2): 1-12.
- 331 13. Girach R.D. and Singh V.K. (2006). Notable plants in ethnomedicine of Jajpur District,  
332 Orissa, India. In: Recent Progresses in Medicinal Plants. *Biopharmaceuticals*,  
333 14:495-501.

- 334 14. Sourav Das (2012). Antimicrobial activity study of ethanolic extract of *Boerhaavia*  
335 *diffusa* whole plant. *International Journal Of Pharmacy & Life Sciences*, **3**(10): 2006-  
336 2009
- 337 15. Ogoti P., Magiri E., Magoma G., Kariuki D. and Bii C. (2015). In vitro anti-Salmonella  
338 activity of extracts from selected Kenyan medicinal plants. *Journal of Medicinal Plants*  
339 *Research*, **9**(8): 254-261
- 340 16. Cheesbrough, M. (2000). Medical Laboratory Manual for tropical countries. Part II:  
341 Microbiology. Cambridge University. USA. pp. 62-70.
- 342 17. Marcelin U., Bharathidasan R. and Prince L. (2016). Antisalmonella Activity of Selected  
343 Medicinal Plants. *World Journal of Pharmacy and Pharmaceutical Sciences*, **5**(4): 2512-  
344 2519
- 345 18. Anibijuwon I.I and Udeze O.A (2009). Antimicrobial activity of *Carica papaya* (Pawpaw  
346 Leaf) on some pathogenic organisms of clinical origin from South-Western Nigeria.  
347 *Ethnobotanical Leaflets*, **13**(7): 850-864.
- 348 19. Abbas A. M. Noor M. A. and Sarmina Y. (2007). Antimicrobial screening of different  
349 extracts of *Piper longum* Linn. *Res. J Agri and Biol sci.*, **3**(6): 852-857.
- 350 20. Shahina N., Samia A. and Sheikh A. R (2007). In vitro antimicrobial activity of the  
351 extracts derived from *Terminalia catappa*. *Res. J Microbial.*, **2**(2): 180-184.
- 352 21. Marjorie M.C. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*  
353 **12**(4):564-582.
- 354 22. Abo K.A and Ashidi J.S. (1999). Antimicrobial screening of *Bridelia micrantha*,  
355 *Alchornea cordifolia* and *Boerhavia difussa*. *African Journal of Medicine and Medical*  
356 *Sciences* **28**(34):167-169.
- 357 23. Madani A. and Jain S. K. (2008). Anti-Salmonella activity of *Terminalia bellerica*: In  
358 vitro and in vivo studies. *Indian Journal of Experimental Biology*, **46**: 817-821
- 359 24. Olila D, Olwa-Odyek A, Opuda-Asibo J (2001). Antibacterial and antifungal activities of  
360 extracts of *Zanthoxylum chabeum* and *Wurbugia ugandensis*, Uganda medicinal plants.  
361 *Afr. Health Sci.* **1**(2):66-72.
- 362 25. Edeoga H.O, Okwu D.E, Mbaebie O. (2005). Nutritional values of some known  
363 conventional leafy vegetables of Nigeria. *African Journal of Biotechnology*, **4**:685-  
364 688.
- 365 26. Okwute S.K. (1992). Plant derived pesticidal and antimicrobial agents for use in  
366 Agriculture. A review of phytochemical and biological studies on some Nigerian plants.  
367 *J. Agric. Sci. Technol.* **2**(1):62-70.
- 368 27. Desai S.D., Desai D.G., Kaur H. (2009). Saponins and their Biological Activities.  
369 *Pharma Times*, **41**(3):88-92
- 370 28. Okwu D.E. (2004). Phytochemical and vitamin content of indigenous species of  
371 southeastern Nigeria. *Journal of Sustainable Agriculture and the Environment*, **6**:30-37.
- 372 29. Shah A. S., Wakade A. S. and Juvekar A. R. (2008). Immunomodulatory activity of  
373 methanolic extract of *Murraya koenigii* spreng leaves. *Indian J Exp Biol* **46**: 505-9.