

## **Distribution of Primary Nutrients (NPK) to crop production in Profiles of Soils derived from Coastal Plain Sand in Ikot Ekpo, Calabar, Cross River State, Nigeria**

**Eyong, M. O<sup>\*1</sup>, Akpa, E. A<sup>\*2</sup> and <sup>\*3</sup>Peter, P. G.**

<sup>\*</sup>Department of Soil Science, University of Calabar, P.M.B. 1115, Calabar, Cross River State, Nigeria

<sup>\*1</sup> *Corresponding author: [michaeleyong87@yahoo.com](mailto:michaeleyong87@yahoo.com)*

<sup>\*2</sup> *Corresponding Co-author: [enyaanari@gmail.com](mailto:enyaanari@gmail.com)*

### **ABSTRACT**

This study was conducted to evaluate the distribution of primary nutrients (NPK) in profiles of the coastal plain soils of Ikot Ekpo, Calabar, as well as evaluate other soil properties critical to agricultural productivity, in a bid to generate data that will serve as a guide to effective land use and management of the soils for arable crop production. Three (3) Profile pits were dug on the crest, middle slope and valley bottom, and soil samples were collected from their pedogenetic horizons for analysis. Analytical results showed the three profiles of coastal plain soils studied had predominantly sandy particle sizes (ranged from 770 - 910  $\mu\text{m}$  sand across the three profiles) and mostly loamy sand in texture; especially at the topsoil level. The soils were also acidic (pH 4.7 to 5.1) and low in organic matter (1.0  $\text{mg kg}^{-1}$  to 16.0  $\text{mg kg}^{-1}$ ) as expected. Generally, the soils were found to be low in total nitrogen content (0.1 to 1.3  $\text{mg kg}^{-1}$ ) and exchangeable potassium (0.08 to 0.10  $\text{cmolc kg}^{-1}$ ); however, they were high in available phosphorus (17.20 to 29.75  $\text{mg kg}^{-1}$ ). NPK distribution charts showed that N and P decreased consecutively with increasing depth for the crest profile. The middle-slope and valley bottom profile showed no definite pattern of distribution. However, the concentration of NPK was highest at topsoil level across most profiles. N had the shallowest intra-profile distribution with significantly higher levels of topsoil concentration indicated by the high percentages of intra-profile CV (94 %, 85 % & 97 % for CUF, MUF & VUF respectively). P showed a shallow intra-profile distribution across the three profiles but did not vary significantly from the intra-profile mean (12.5%, 12.0% & 2.6 % for CUF, MUF & VUF respectively). On the other hand K was more evenly distributed within all three profiles ( CV of 9.4 %, 5.3 % & 8.6 % for CUF, MUF & VUF respective) compared to N and P. Inter-profile distribution of NPK showed that N and P had higher concentrations at crest level, with P showing consecutive decrease in concentration down the slope. This study therefore recommends adoption of different NPK fertilizer recommendations for different soil depths and topographic locations for optimal productivity.

**Keywords:** Primary nutrients, coastal plain soils, inherent limitations, productivity.

## INTRODUCTION

The challenge for agriculture over the coming decades is to meet the world's increasing demand for food and sustaining optimal production. Declining soil fertility and mismanagement of available soil nutrients have made this task more difficult. As long as agriculture remains a soil-based industry, major increases in productivity are unlikely to be attained without ensuring that plants have an adequate and balanced supply of nutrients required for optimum productivity. This calls for efficient analysis of agricultural soils to determine soil quality in terms of nutrient composition, and to maintain or enhance soil quality when necessary in order to effectively manage plant nutrient and soil fertility [14].

Coastal plain soils are mainly unconsolidated sediments defined by unique geological substrates which consist of wind worked quaternary sand [32]. They are characterized by acidic conditions, low cation exchange capacity, and multiple nutrient deficiencies due to factors such as intensive weathering, leaching and inappropriate agricultural activities [5; 18]. The clay fraction of the soil is rich in kaolinitic clay minerals [18; 37]. Coastal plain sands cover an area of 480 km<sup>2</sup> in Cross River State, 3,470.32 km<sup>2</sup> in Benue State, 42.20 km<sup>2</sup> in Lagos State, 213.16 km<sup>2</sup> in Akwa Ibom State, 12.18 km<sup>2</sup> in Ogun State, 40.62 km<sup>2</sup> in Ondo State; and 5,435.92 km<sup>2</sup> in River State. In Cross River State, coastal plain soils are found mostly in Akpabuyo, Bakassi, Calabar and Odukpani Local Government Areas [16].

Coastal plain soils have good agricultural potentials because they have moderate inherent fertility and availability of water during the dry season. In crop production and land use, the evaluation of soils' chemical properties is important because properties such as pH, organic matter, nitrogen, phosphorus exchangeable bases, cation exchange capacity and base saturation affect plant growth and development [9]. To increase crop performance, there is always the need to establish relationships between soil chemical properties and soil capacity

to produce food crops, that indeed the basis of yield predictions and could be considered as the most useful expression of soil productivity [9].

Soil characterization in relation to evaluation of fertility status of the soils of an area or region is an important aspect in context of sustainable agricultural production. Because of imbalanced and inadequate fertilizer use coupled with low efficiency of other inputs, the production efficiency of chemical fertilizer nutrients has declined tremendously under intensive agriculture in recent years. Introduction of high yielding varieties to agriculture in the mid-sixties compelled the farmers to use high doses of NPK (Nitrogen, Phosphorus and Potassium) fertilizers along with micronutrient fertilizers. NPK are among the nutrients required by plants because they are primarily required for plants to complete their life cycle. They are the most critical plant nutrients in agriculture because they are mobile and not easily lost by leaching due to its adsorption to the clays. The deficiencies of these nutrients have become major constraints to productivity, stability and sustainability of soils. Soils with finer particles and with higher organic matter can generally provide a greater reserve of nutrients whereas, coarse textured soils such as sand have fewer reserves and tend to get depleted rather quickly.

Plants take up nitrogen from the soil solution mainly as nitrate ( $\text{NO}_3^-$ ) and ammonium ions ( $\text{NH}_4^+$ ). Nitrogen is responsible for rapid growth and its content in surface soils ranges from 0.02 to 0.5 %, which is below the critical level of plant nutrient of 2.0 % as reported by Espinoza *et al.* [10]. The mineral forms of nitrogen are soluble in water and are easily lost from soils through leaching and volatilization. Minimization of environmentally damaging impact from soil plant system and regulation of soluble forms of nitrogen through split application and slow release fertilizer is therefore imperative in order to maintain adequate supply in the soil as reported by Espinoza *et.al.* [10].

Phosphorus has been identified as one of the most limited nutrient elements in tropical soils. The total phosphorus level of soils is low and is mostly unavailable for plant uptake because they are highly insoluble. When soluble forms are added to the soil, they are also fixed and in time form unavailable forms, leaving a very small percentage for plant utilization. Low availability of phosphorus in tropical soils is attributed to the nature of chemical forms of the soils phosphorus and high content of oxides of iron and aluminum, which are associated with high phosphorus fixation [25]. Phosphorus promotes root formation, affects quality of seeds, fruit and flowers, and increases disease resistance on the plants [20].

Potassium is the third most important plant element after nitrogen and phosphorus. It is essential for plant growth, useful for helping plants overcome drought stress and increases diseases resistance. Potassium is generally high in most mineral soils but very large portion is unavailable to plants, it is also subject to leaching and removal by plants. Problem associated with potassium use by plants is rarely of its total supply but rather of adequate supply of available forms at depths below plough layer as reported by Espinoza *et. al.* [10].

It is well recognized that soils are the storehouse of most of the plant nutrients essential for plant growth and that the way in which soils are managed will have a major impact on soil fertility, plant growth and agricultural sustainability. Since soils are dynamic systems it is expedient to ascertain levels of primary nutrients in soils and to establish management practices for the soils. This is necessary to maintain and improve agricultural productivity and sustainability.


## **MATERIALS AND METHODS**

### **Location of Study Area**

Ikot-Ekpo, Calabar is located between latitude  $008^{\circ}20.030'E$  and longitude  $05^{\circ}38.45'N$  in Cross River State. Cross River State is located in the South Eastern part of the Federal Republic of Nigeria. Geographically, it lies between latitude  $5^{\circ}32'$  and  $4^{\circ}27'$  North of the equator and  $7^{\circ}50'$  and  $0^{\circ}25'$  East of the Greenwich Meridian. It is bounded by Ebonyi and Abia State in the West, Benue State in the North, Akwa Ibom State in the South, and Atlantic Ocean in the South-East and in the East by the Cameroon Republic. The total land mass of Cross River State is about 23,074,245 square kilometer. Cross River State has a population of 4.6 million people living in the state. The state has thick rainforest vegetation and very rich soils [5].

### **Climate**

The climate of the study area is marked by a rainy season and a dry season. The rainy season lasts from March to July with heavy downpours, and strong wing storm, and the dry season last from November to early March which climaxes between December and January during harmattan. The area has rainfall range of 2000 mm to 2739 mm per annum and a temperature of  $25^{\circ}C$  in the rainy season and  $28^{\circ}C$  in the dry season. The area has a relative humidity range of 70 to 80 %.



**Fig.1** Map of Calabar metropolis showing location of Ikot Ekpo  
 Source: Cartographic unit, Department of Geography, University of Calabar, Calabar

## **Geology**

The area is dominated by coastal plain sands belonging to the tertiary deposit and alluvial deposit on low lying swampy areas.

## **Vegetation**

The area is mainly used for arable cropping and rubber plantation. The plantation is more than 15 years and contains other vegetable crops like *Telferia occidentalis*, *Talinum triangulare* and *Manihot esculentus*.

## **Field Studies**

Three (3) Profile pits were dug on the crest, middle slope and valley bottom. A full range of environmental conditions, selected soils morphological properties and soil physico-chemical properties were recorded on soil description sheets. Each profile pit measured 1.5 m x 2 m x 2 m. The profile pits were dug to impenetrable layer or water level depending on whichever is shallower. Each profile pit was described in the moist state for their full ranges of morphological characteristics. Description was done according to guidelines [15].

## **Laboratory Analysis**

In the laboratory, the soil samples were air dried at room temperature for 48 hours and then gently crushed with pestle and mortar and sieved through 2 mm sieve to obtain fine earth fractions for the analysis. The physico-chemical properties of the soils were determined using standard procedures, as outlined by Udo *et al.* [46]. Bulk cores were taken using cylindrical cores, later oven dried to constant weight and then the bulk density was calculated [4]. The particle size distribution was determined by the Hydrometer method using sodium hexametaphosphate (calgon) as dispersant [19]. The soil texture was determined using the textural triangle. The soil pH was determined potentiometrically after equilibration with

water and 0.01 M  $\text{CaCl}_2$  in a 1:2.5 soils to solution ratio using glass electrode pH meter as outlined by Isirimah *et al.* [26]. Organic carbon was determined by the Walkley-Black wet oxidation method and the value multiplied by Van Bemmelen factor of 1.724 to obtain organic matter value [47]. The method involves the digestion of the soil organic matter with potassium dichromate ( $\text{K}_2\text{Cr}_2\text{O}_7$ ) using concentrated sulphuric acid to increase the temperature and hasten the reaction. The total N was determined by the salicylic acid-thiosulphate digestion method followed by the distillation method using modified micro-kjeldhal method [30]. Available P was extracted by the Bray No. 1 method and 2 mLI of the extract was used to determine P in solution colorimetrically by the ascorbic acid method [38], after 2 mLI of the extract is made up to 50 mLI with distilled water; it is then kept for some time for colour development before taking reading in a spectrophotometer. Exchangeable Cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ) were determined with ammonium acetate (pH 7.0) using 1:10 soil-liquid ratio. Calcium (Ca) and Magnesium (Mg) in the filtrate were then determined with atomic absorption spectrophotometer (AAS) while sodium and potassium were determined by flame photometer analyzer [45]. Exchangeable acidity was determined by successive leaching of soils with neutral unbuffered K chloride (KCl) using 1:10 soil to liquid ratio. Exchangeable hydrogen (H) and aluminum (Al) were determined by titration method [29], for this, 1 mLI of KCl was used as an extracting agent, after adding 5 drops of phenolphthalein indicator. It is titrated with 0.01 mLI of NaOH until permanent pink coloration is reached. The solution is titrated against 0.01 mLI HCl until a colourless solution is obtained. Cation Exchange capacity was determined by neutral ammonium acetate (pH 7.0) saturation method described by Udoh *et al.* [46]. Percentage base saturation was determined by the sum of exchangeable cations to effective cation exchange capacity (ECEC) multiply by 100.



## **Statistical Analysis**

Variability in soil properties NPK within the soil profiles were subjected to analysis of variance (ANOVA). Linear regression was used to explore the relationship between NPK and depth. A table of the physical and chemical properties showing relationship of nutrient with depth was made. Also, a table showing variation of NPK with soil depth in the profiles was made.

## **RESULTS AND DISCUSSIONS**

### **Morphological Properties**

The morphological properties of soil derived from coastal plain sand in Ikot Ekpo, Calabar are presented in Table 1.

### **Soil colour**

The hue values were mostly 10YR, with 2.5YR at AP horizon (0 – 30 cm depth) of the crest. The hue value 2.5YR for surface soil (0-30 cm) crest, and 10YR value, with colour range from dark grayish brown to dark yellowish brown conform to [23]. The dark grayish colour may be related to the organic matter content of the soil. However, the yellowish colour may be an indication of the presence of iron (iii) oxide. This in line with [48], who stated that coastal plain soils merge into deep permeable red-earth or yellow earth strata derived from tertiary sediments, an indication of iron (iii) oxide.

### **Soils structure**

Soil structure is the aggregation of primary soil particles (sand, silt and clay) and provide good health to plant growth. One group of structure was observed in the surface soil namely weak fine granular. The subsurface had two groups and were dominated by moderate to medium sub-angular blocky structure. According to [24], soils derived from coastal plain

sands have medium sub-angular blocky subsurface. The structure of these soils is influenced basically by their particle size distribution.

### **Soil texture**

The texture of the surface and subsurface soils varied from loamy sand surface to sandy clay loam subsurface soils and this agrees with Shaw *et al.* [42], who stated that soils derived from coastal plain sands have sand or loamy sand surface and sandy loam or sandy clay loam subsurface. Such textural classes could have serious implications on hydrological processes such as erosion, aeration and water holding capacity as reported by Schoenholtz *et al.* [41].

### **Soil consistency**

Soil consistency indicates the resistance of soil to deformation. The soil consistency was very friable at surface level. The subsurface level were both firm and friable but were largely dominated by firm consistency. [24] described the consistency of coastal plain soils as none or slightly stick or plastic when wet, and loose when dry.

**Table 1:**                    **The morphological properties of soils derived from coastal plain sands under arable crop production in Ikot Ekpo, Calabar, Cross River State**

Profile location	Horizon	Depth (cm)	Munsell colour (wet)	Texture	Structure	Consistence	Boundary	Other features
CUF	AP	0 - 30	2.5YR <sup>4</sup> / <sub>2</sub>	Loamy sand	1 fgr	Vfr	Cs	Many fine roots
E008 <sup>0</sup> 19.91'	B	30 – 70	10YR <sup>4</sup> / <sub>4</sub>	Sandy clay loam	2 msbk	Fi	Cs	Common fine roots
N05 <sup>0</sup> 03.65'	BC	70 – 120	10YR <sup>4</sup> / <sub>4</sub>	Sandy clay	2 msbk	Fi	Cs	Few fine roots
	C	120 – 150	10YR <sup>3</sup> / <sub>6</sub>	Sandy clay	2 msbk	Fi	Cs	Few fine roots
MUF	AP	0 – 26	10YR <sup>4</sup> / <sub>3</sub>	Loamy sand	1 fgr	Vfr	Cg	Few fine roots, many ants
E008 <sup>0</sup> 19.95'	AB	26 - 57	10YR <sup>6</sup> / <sub>6</sub>	Sandy loam	1 sbk	Fr	Cs	Few fine roots
N05 <sup>0</sup> 03.87'	B	57 – 110	10YR <sup>5</sup> / <sub>4</sub>	Sandy clay	2 msbk	Fi	Cs	Few fine roots
	C	110 – 150	10YR <sup>6</sup> / <sub>8</sub>	Sandy clay	2 msbk	Fi	Cs	Few fine roots
VUF	AP	0 – 20	10YR <sup>3</sup> / <sub>2</sub>	Loamy sand	1 fgr	Vfr	Cs	Many roots
E008 <sup>0</sup> 9.09'	B	20 - 60	10YR <sup>4</sup> / <sub>4</sub>	Sandy loam	1 sbk	Fr	Cs	Few medium roots
N05 <sup>0</sup> 03.65'	BC	60 - 115	10YR <sup>4</sup> / <sub>4</sub>	Sandy clay loam	2 msbk	Fi	Sg	Few fine roots
	C	115 – 150	10YR <sup>5</sup> / <sub>6</sub>	Sandy clay loam	2 msbk	Fi	Sg	Few fine roots

Structure: 1 = weak; 2 = moderate; 3 = strong; f = fine; m = medium; gr = granular; cr = crumb; sbk = sub-angular blocky.

Consistence : Vfr = very friable; Fi = firm; Fr = friable.

Boundary : C = clear; S = smooth; g = gradual; w = wavy.

CUF = Crest Unical Farm

MUF = Middle-slope Unical Farm

VUF = Valley-Bottom Unical Farm

## Physical Properties

Table 2 shows the physical properties of soils derived from coastal plain sands under arable crop production in Ikot Ekpo, Calabar

### Particle size distribution

Particle size distribution of the soil showed that sand ranged from 860  $\text{gkg}^{-1}$  to 910  $\text{gkg}^{-1}$ , silt ranged from 60  $\text{gkg}^{-1}$  to 120  $\text{gkg}^{-1}$  while that of clay ranged from 30  $\text{gkg}^{-1}$  to 110  $\text{gkg}^{-1}$ . Surface soils had a mean of 893.3  $\text{gkg}^{-1}$ , 76.6  $\text{gkg}^{-1}$  and 30  $\text{gkg}^{-1}$  for sand, silt and clay respectively while the subsurface soil had a mean of 803  $\text{gkg}^{-1}$ , 101.1  $\text{gkg}^{-1}$  and 970  $\text{gkg}^{-1}$  for sand silt and clay respectively. Sand was the dominant particles size fraction at both surface and subsurface soils, placing the textural classes of all the soils as loamy sand for top soils and sandy clay loam to sandy clay for sub soils. The sandy nature of the soils at both levels could be attributed largely to the parent material. The predominantly sandy nature of the soils was an indication of low fertility arising from physical causes. Sandy soils have loosed particles and hold little organic matter as well as nutrients [34].

### Bulk density

The bulk density of the soils ranged from 1.4  $\text{gcm}^{-3}$  to 1.8  $\text{gcm}^{-3}$  at all depths. The surface soils had a range of 1.6  $\text{gcm}^{-3}$  to 1.8  $\text{gcm}^{-3}$  with a mean value of 1.7  $\text{gcm}^{-3}$ . The subsurface values ranged from 1.4  $\text{gcm}^{-3}$  to 1.8  $\text{gcm}^{-3}$  with a mean value of 1.5  $\text{gcm}^{-3}$ . The mean value for surface bulk density is higher than that of the subsurface, this agrees with Donahue *et al.* [7] who stated that soils derived from coastal plain sands have higher bulk density in the surface than the subsurface. This higher surface bulk density is likely due to higher levels of organic matter in surface profile of the soils. It could also be attributed to soil compaction as a result of continuous and intensive cultivation of the soils. High levels of bulk

density are capable of impeding root penetration and development thereby limiting crop yield.

### **Particle density**

The particle density ranged between 2.30  $\text{gcm}^{-3}$  and 2.81  $\text{gcm}^{-3}$  at all depths. The surface soils for the three locations ranged 2.30  $\text{gcm}^{-3}$  to 2.57  $\text{gcm}^{-3}$  with a mean of 2.49  $\text{gcm}^{-3}$ . The subsurface soils ranged from 2.45  $\text{gcm}^{-3}$  to 2.81  $\text{gcm}^{-3}$  with a mean of 2.67  $\text{gcm}^{-3}$ . This result is closely ranged with that obtained by [13]. The particle density is higher than the bulk density as expected and is considered moderate to high compare with the generally accepted range of 2.6 to 2.7  $\text{mg m}^{-3}$  (Landon 1991). The relatively high values obtained is likely due to the low level of organic matter observe in the soils because high organic matter tend to lower particle density of associated soils.

### **Total porosity**

The total porosity varied between 7 % and 90 % at all depths. The surface soils porosity ranged from 61 % to 90 % with a mean value of 79.3 %. The subsurface had values ranging from 7 % to 90 % with a mean value of 46.6 %. This indicates that the soils have more porous surface than subsurface. According to [1], top soils form from coastal plain sands had more bulk density porous than subsoil. The highly porous surface soil values reflect typical sandy soils; with porosity that ensures even circulation of air and growth of microorganisms.

**Table 2:** The physical properties of soils derived from coastal plain sands under arable crop production in Ikot Ekpo, Calabar, Cross River State

[illegible]

## Chemical Properties

Table 3 shows the chemical properties of soils derived from coastal plain sands under arable crop production in Ikot Ekpo, Calabar

### Soil pH

The pH values ranged from 4.7 to 5.1 at all depths. The surface soils pH ranged from 4.7 to 5.1 with a mean value of 4.9. The subsurface had values ranging from 4.9 to 5.1 with a mean value of 5.0. From the result, the soils at both surface and subsurface depths are strongly acidic. This pH range does not fall within the optimal pH range of 6.0 - 7.5 for the growth of higher plants and microorganisms as reported by Havlin *et al.* [21]. And the texture was due to the high level of rainfall in the region. High level of rainfall causes the basic cations in the soils to leach away from the profile [23]. According to [11], the soil pH of soils in Calabar had values ranging from 4.8 to 5.5, indicating moderate to strong acidity and this agrees with the findings of this research.

### Organic Carbon (O/C)

The organic carbon content of the soils ranged from  $1.0 \text{ mg kg}^{-1}$  to  $16.0 \text{ mg kg}^{-1}$ . The top soil values ranged from  $7.0$  to  $16.0 \text{ mg kg}^{-1}$  with a mean value of  $10.80 \text{ mg kg}^{-1}$ . The subsurface organic carbon values ranged from  $1.0$  to  $5.0 \text{ mg kg}^{-1}$  with a mean value of  $2.0 \text{ mg kg}^{-1}$ . This result shows that the OC level was significantly higher in the soil surface than in the subsurface depth. This is due to accumulation of organic matter at the top soil. However, the organic matter content at both depths was generally low, as all the values were below the critical level of  $40 \text{ mg kg}^{-1}$  [36]. This may be due to intensive use of the land for agricultural activities without return of plant residue to the soils as reported by Bunemann *et al.* [6].

### **Total Nitrogen ( $N_T$ )**

The total nitrogen content of the soils is shown in Table 4. It ranged from 0.1 to 1.3  $\text{mg kg}^{-1}$  at both surface and subsurface soils. The surface soils ranged from 0.5 to 1.3  $\text{mg kg}^{-1}$  with a mean value of 0.8  $\text{mg kg}^{-1}$  while the subsurface soil ranged from 0.1 to 0.4  $\text{mg kg}^{-1}$  with a mean value of 0.2  $\text{mg kg}^{-1}$ . This result compared with the critical value of 0.35  $\text{mg kg}^{-1}$  [36] show that the total nitrogen content of the soil is low as observed by [17] for acid sands of Eastern Nigeria. Such levels of  $N_T$  in soils might have serious negative implications on soil and crop productivity. These low values may be as a result of low organic matter content as well as leaching of nitrates from the soil; as is common in coastal plain soils and the texture.

### **Available Phosphorus ( $P_A$ )**

The available phosphorus content of the soils as shown in Table 4 ranged from 17.20 to 29.75  $\text{mg kg}^{-1}$  at both depths. The values for surface soils ranged from 21.12 to 29.75  $\text{mg kg}^{-1}$  with a mean value of 25.33  $\text{mg kg}^{-1}$  while the subsurface soil ranged from 17.20 to 23.50  $\text{mg kg}^{-1}$  with a mean value of 20.15  $\text{mg kg}^{-1}$ . This result shows that the soils are generally high in available P; as it exceeds the critical value of 15  $\text{mg kg}^{-1}$  [17]. This high in available P might be attributed to fertilizer application used for crop growth in the area. None of the profiles was observed to fall below the critical value. Furthermore, [9] estimated an average value of 20  $\text{mg kg}^{-1}$  of available P for the soils of coastal plain sands and this conforms to these findings.

### **Exchangeable Bases**

#### **Exchangeable $K^+$**

The exchangeable  $K^+$  content of the soils ranged from 0.08  $\text{cmolc kg}^{-1}$  to 0.10  $\text{cmolc kg}^{-1}$  at both depths. The top soil values ranged from 0.08 to 0.10  $\text{cmolc kg}^{-1}$  with a mean value of 0.09  $\text{cmolc kg}^{-1}$  while the subsurface values ranged from 0.08 to 0.10  $\text{cmolc kg}^{-1}$



with a mean value of  $0.09 \text{ cmolc kg}^{-1}$ . This result indicates that level of exchangeable  $K^+$  is generally very low as observed by Holland *et al.* [22]. This contrasts with the mean value of 0.16 obtained by Ewulo *et al.* [12] for the coastal plain soils of Uyo. The low value of  $K^+$  which is below critical level of  $0.2 \text{ cmolc kg}^{-1}$  as reported by Kyuma *et al.* [33] might be attributed to the high rainfall and leaching intensity normally often encountered in coastal plain soil and the sandy texture..

### **Exchangeable $Ca^{2+}$**

The exchangeable  $Ca^{2+}$  content of the soils varied from  $2.2 \text{ cmolc kg}^{-1}$  to  $4.4 \text{ cmolc kg}^{-1}$  at both depths. The top soil values ranged from 2.2 to  $3.0 \text{ cmolc kg}^{-1}$  with a mean value of  $2.7 \text{ cmolc kg}^{-1}$  while the subsurface values ranged from 2.2 to  $4.4 \text{ cmolc kg}^{-1}$  with a mean value of  $3.2 \text{ cmolc kg}^{-1}$ . This result shows that level of exchangeable  $Ca^{2+}$  is generally low to moderate (Holland *et al.* [22]).

### **Exchangeable $Mg^{2+}$**

The exchangeable  $Mg^{2+}$  content of the soils varied from  $0.8 \text{ cmolc kg}^{-1}$  to  $2.8 \text{ cmolc kg}^{-1}$  at both depths. The top soil values ranged from 1.2 to  $2.4 \text{ cmolc kg}^{-1}$  with a mean value of  $1.7 \text{ cmolc kg}^{-1}$  while the subsurface values ranged from 0.8 to  $2.8 \text{ cmolc kg}^{-1}$  with a mean value of  $1.9 \text{ cmolc kg}^{-1}$ . This result shows that level of exchangeable  $Mg^{2+}$  is generally moderate [2] to high as reported by Holland *et al.* [22]

### **Exchangeable $Na^+$**

The exchangeable  $Na^+$  content of the soils varied from  $0.05 \text{ cmolc kg}^{-1}$  to  $0.08 \text{ cmolc kg}^{-1}$  at both depths. The top soil values ranged from 0.05 to  $0.08 \text{ cmolc kg}^{-1}$  with a mean value of  $0.06 \text{ cmolc kg}^{-1}$  while the subsurface values ranged from 0.06 to  $0.08 \text{ cmolc kg}^{-1}$  with a mean value of  $0.06 \text{ cmolc kg}^{-1}$ . This result shows that level of exchangeable Na is rated very low compared to similar soils in the region as reported by Ewulo *et al.* [12].

## Exchangeable Acidity

### Exchangeable $H^+$

The exchangeable  $H^+$  content of the soils varied from 0.2  $\text{cmolc kg}^{-1}$  to 0.88  $\text{cmolc kg}^{-1}$  at both depths. The top soil values ranged from 0.2 to 0.40  $\text{cmolc kg}^{-1}$  with a mean value of 0.28  $\text{cmolc kg}^{-1}$  while the subsurface values ranged from 0.12 to 0.88  $\text{cmolc kg}^{-1}$  with a mean value of 0.39  $\text{cmolc kg}^{-1}$ . This result shows that the subsurface soil had higher exchangeable  $H^+$  values than the surface soil. The soil is generally low in exchangeable  $H^+$  as reported by Holland *et al.* [22]

### Exchangeable $Al^{3+}$

The exchangeable  $Al^{3+}$  content of the soils varied from 0.24  $\text{cmolc kg}^{-1}$  to 1.04  $\text{cmolc kg}^{-1}$  at both depths. The top soil values ranged from 0.32 to 0.96  $\text{cmolc kg}^{-1}$  with a mean value of 0.53  $\text{cmolc kg}^{-1}$  while the subsurface values ranged from 0.24 to 1.04  $\text{cmolc kg}^{-1}$  with a mean value of 0.57  $\text{cmolc kg}^{-1}$ . This result shows that exchangeable  $H^+$  values of the soils is generally low as reported by Holland *et al.* [22]. The low levels of exchangeable acidity is solely attributed to leaching away of the basic cations ( $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Na^+$ ) from the surface soil, which leaves the surface soil acidic.

### Effective Cation Exchange Capacity (ECEC)

The ECEC content of the soils ranged between 4.85  $\text{cmolc kg}^{-1}$  to 7.06  $\text{cmolc kg}^{-1}$  at both depths. The top soil values ranged from 4.85 to 6.30  $\text{cmolc kg}^{-1}$  with a mean value of 5.44  $\text{cmolc kg}^{-1}$  while the subsurface values ranged from 5.24 to 7.06  $\text{cmolc kg}^{-1}$  with a mean value of 6.32  $\text{cmolc kg}^{-1}$ . This result shows that level of ECEC is rated very low and regarded as unsuitable for crop production [16].

**Base Saturation (BS)**

The base saturation of the soils ranged from 76 % to 115 % at both depths. The top soil values ranged from 76 to 115 % with a mean value of 93 % while the subsurface values ranged from 80 to 9 % with a mean value of 83 %. These values indicate the availability of the basic cations and it shows a higher value than that obtained by [11] for the Utisols of Cross River State and so could be regarded as being moderate to high.

**Table 3: The chemical properties of soils of coastal plain sands under arable crop production in Ikot Ekpo, Calabar, Cross River State**

Profile location	Horizon	Depth (cm)	pH	O/C (mgkg <sup>-1</sup> )	N <sub>T</sub> (mgkg <sup>-1</sup> )	P <sub>A</sub> (mgkg <sup>-1</sup> )	Exchangeable Bases (cmolc kg <sup>-1</sup> )				Exchangeable Acidity (cmolc kg <sup>-1</sup> )		ECEC (cmolc kg <sup>-1</sup> )	BS (%)
							Ca <sup>2+</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	Na <sup>+</sup>	H <sup>+</sup>	Al <sup>3+</sup>		
CUF E008 <sup>0</sup> 19.91' N05 <sup>0</sup> 03.65'	AP	0 - 30	5.1	16.0	1.3	29.80	3.0	1.2	0.08	0.05	0.20	0.32	4.85	89
	B	30 - 70	5.1	5.0	0.4	23.50	4.4	0.8	0.10	0.07	0.90	0.24	6.49	82
	BC	70 - 120	5.2	3.0	0.2	19.25	3.2	2.4	0.09	0.06	0.40	0.83	6.99	82
	C	120 - 150	5.2	1.0	0.1	17.20	3.8	2.0	0.08	0.06	0.12	0.56	6.62	89
MUF E008 <sup>0</sup> 19.95' N05 <sup>0</sup> 03.87'	AP	0 - 26	4.9	7.0	0.5	25.10	3.0	2.4	0.10	0.08	0.40	0.32	6.30	88
	AB	26 - 57	5.0	2.0	0.1	19.40	3.5	2.8	0.10	0.08	0.24	0.24	7.06	93
	B	57 - 110	4.9	2.0	0.1	21.50	3.4	2.2	0.09	0.06	0.48	0.40	6.63	86
	C	110 - 150	4.9	2.0	0.1	18.50	2.2	2.6	0.09	0.06	0.85	0.32	6.15	80
VUF E008 <sup>0</sup> 9.09' N05 <sup>0</sup> 03.65	AP	0 - 20	4.7	9.40	0.8	21.10	2.2	1.5	0.10	0.07	0.24	0.96	5.17	76
	B	20 - 60	4.9	0.40	0.1	21.50	2.6	1.2	0.09	0.07	0.24	1.04	5.24	75
	BC	60 - 115	4.9	1.0	0.1	20.10	2.8	1.8	0.08	0.06	0.24	0.72	5.70	83
	C	115 - 150	4.9	3.0	0.2	20.50	3.0	2.0	0.10	0.07	0.80	0.80	6.05	85
Topsoil Range			4.7- 5.1	7.0 - 16	0.5 - 1.3	21.12- 29.75	2.2 -3.0	1.2-2.4	0.08 -0.1	0.05-0.08	0.2-0.40	0.32- 0.96	4.85-6.30	76 -89
Subsoil Range			4.9- 5.2	1.0 -5.0	0.1 - 0.4	17.20 - 23.50	2.2 -4.4	0.8- 2.8	0.08 -0.1	0.06-0.08	0.12-0.90	0.24- 1.04	5.24-7.06	80 - 93
Topsoil Mean			4.9	10.60	0.8	25.33	2.7	1.7	0.09	0.06	0.28	0.53	5.44	93
Subsoil Mean			4.8	2.1	0.2	20.15	3.2	1.9	0.09	0.06	0.39	0.57	6.32	83

## Distribution of NPK in the soils

The intra-profile and inter-profile distribution of total N, available P and exchangeable K were discussed intensively with respect to Table 4 and 5.

### Total nitrogen

Intra-profile distribution of total N as shown in Table 4 indicates a value of  $1.3 \text{ mg kg}^{-1}$  for top soil of crest profile (CUF). The values ranged from 0.1 to  $1.3 \text{ mg kg}^{-1}$  throughout CUF profile depth with a mean of  $0.5 \text{ mg kg}^{-1}$ , SD of  $0.47 \text{ mg kg}^{-1}$  and CV of 94 %. The middle slope profile (MUF) had a topsoil value of  $0.5 \text{ mg kg}^{-1}$  and ranged from 0.1 to  $0.5 \text{ mg kg}^{-1}$  with a mean value of  $0.2 \text{ mg kg}^{-1}$ , SD of 0.17 and CV of 85 %. While the valley bottom profile (VUF) had a top soil value of  $0.8 \text{ mg kg}^{-1}$  and ranged from 0.1 to  $0.8 \text{ mg kg}^{-1}$  with a mean of  $0.3 \text{ mg kg}^{-1}$ , SD of  $0.29 \text{ mg kg}^{-1}$  and CV of 97 %.

The inter-profile distribution of total N as shown in Table 4 ranged from 0.2 to  $0.5 \text{ mg kg}^{-1}$  across the three profiles (using intra-profile means), with a mean value of  $0.33 \text{ mg kg}^{-1}$ , SD of  $0.125 \text{ mg kg}^{-1}$  and CV of 38.0 %.

These values indicate higher concentration of N in top soil and lower concentration in sub soils below critical value [17]. Total N in CUF decreased consistently with increase in profile depth (Fig.2a) while MUF and VUF showed inconsistent patterns (Fig.3a&4a). However, the topsoil had a significantly higher N concentration for all three profiles (Fig.2a, 3a & 4a) and this accounts for the high CV observed within the profiles; 94 %, 85 % and 97 % for CUF, MUF and VUF respectively (Table 5). This agrees with the findings of [28], who stated that N concentration was significantly higher in topsoil. The high levels of total N in the topsoil could be attributed to the high natural organic matter returns and mineralization of plant residue as well as N fertilizer application. According to [44], biological cycling generally moves nutrients upwards because some proportion of the nutrients absorbed by plants are transported aboveground and then recycled to the soil surface.

Generally, inter-profile distribution chart shows that the upper slope (CUF) had the highest nitrogen concentration and the lowest concentration was observed in the middle slope (CUF > VUF > MUF) (Fig.5a). The level of significance of its inter-profile variation is visible in the percentage of its coefficient of variation (38 %). According to [8], soils on lower slope are more prone to nitrogen runoff because they become saturated with excess precipitation.

**Table 4**  
**Intra-Profile distribution of NPK in soils of coastal plain sands under arable**  
**crop production in Ikot Ekpo, Calabar, Cross River State**

Profile location	Horizon	Depth (cm)	N <sub>T</sub> (mgkg <sup>-1</sup> )	Mean	SD	CV (%)	P <sub>A</sub> (mgkg <sup>-1</sup> )	Mean	SD	CV (%)	K <sup>+</sup> (cmolc kg <sup>-1</sup> )	Mean	SD	CV (%)
CUF E008 <sup>0</sup> 19.91' N05 <sup>0</sup> 03.65'	AP	0 - 30	1.3	0.5	0.47	94	29.80	22.44	4.82	21.5	0.08	0.088	0.008	9.4
	B	30 – 70	0.4				23.50				0.10			
	BC	70 – 120	0.2				19.25				0.09			
	C	120 – 150	0.1				17.20				0.08			
MUF E008 <sup>0</sup> 19.95' N05 <sup>0</sup> 03.87'	AP	0 – 26	0.5	0.2	0.17	85	25.10	21.13	2.54	12.0	0.10	0.095	0.005	5.3
	AB	26 - 57	0.1				19.40				0.10			
	B	57 – 110	0.1				21.50				0.09			
	C	110 – 150	0.1				18.50				0.09			
VUF E008 <sup>0</sup> 9.09' N05 <sup>0</sup> 03.65	AP	0 – 20	0.8	0.3	0.29	97	21.10	20.80	0.54	2.6	0.10	0.093	0.008	8.6
	B	20 - 60	0.1				21.50				0.09			
	BC	60 - 115	0.1				20.10				0.08			
	C	115 – 150	0.2				20.50				0.10			

SD = Standard deviation  
CV = Coefficient of variation  
CUF = Crest Unical Farm  
MUF = Middle-slope Unical Farm  
VUF = Valley-Bottom Unical Farm

**Table 5**  
**Inter-Profile distribution of NPK in soils of coastal plain sands under arable**  
**crop production in Ikot Ekpo, Calabar, Cross River State**

Profile location	N <sub>T</sub> (mgkg <sup>-1</sup> )	Mean	SD	CV (%)	P <sub>A</sub> (mgkg <sup>-1</sup> )	Mean	SD	CV (%)	K <sup>+</sup> (cmolc kg <sup>-1</sup> )	Mean	SD	CV (%)
CUF E008 <sup>0</sup> 19.91', N05 <sup>0</sup> 03.65'	0.5				22.44				0.088			
MUF E008 <sup>0</sup> 19.95', N05 <sup>0</sup> 03.87'	0.2	0.33	0.125	38.0	21.13	21.46	0.71	3.3	0.095	0.092	0.0029	3.2
VUF E008 <sup>0</sup> 9.09', N05 <sup>0</sup> 03.65'	0.3				20.80				0.093			

SD = Standard deviation  
CV = Coefficient of variation  
CUF = Crest Unical Farm  
MUF = Middle-slope Unical Farm  
VUF = Valley-Bottom Unical Farm



### Available phosphorus

Intra-profile distribution of available P as shown in Table 4 shows a topsoil value of 29.80 mg kg<sup>-1</sup> for crest profile (CUF) and the values ranged from 17.20 to 29.80 mg kg<sup>-1</sup> throughout the profile depth with a mean of 22.44 mg kg<sup>-1</sup>, SD of 4.82 mg kg<sup>-1</sup> and CV 21.5 %. The middle slope profile (MUF) had a topsoil value of 25.10 mg kg<sup>-1</sup> and ranged from 18.50 to 25.10 mg kg<sup>-1</sup> with a mean value of 21.13 mg kg<sup>-1</sup>, SD of 2.54 and CV of 12 %. While the valley bottom profile (VUF) had a top soil value of 21.10 mg kg<sup>-1</sup> and ranged from 20.10 to 21.50 mg kg<sup>-1</sup> with a mean of 20.80 mg kg<sup>-1</sup>, SD of 0.54 mg kg<sup>-1</sup> and CV of 2.6 %.

The inter-profile distribution of available P as shown in Table 4 ranged from 20.80 to 22.44 mg kg<sup>-1</sup> across the three slopes (using intra-profile means) with a mean value of 21.46 mg kg<sup>-1</sup>, SD of 0.71 mg kg<sup>-1</sup> and CV of 3.3 %.

These values show that available P is generally high across all the profiles. This is indicated by the low inter-profile coefficient of variation (Table 5). However, in CUF profile it has a significantly higher concentration at surface level and decreased consistently with increased profile depth (Fig.2b), with a relatively high intra-profile coefficient of variation (Table 5). This is suggested to be due to vertical and lateral movement of P in the subsurface soil which could be a characteristic of soils of sand stone parent materials as reported by Salminen *et. al.* [40]. MUF and VUF (Fig.3b&4b) showed no definite pattern of distribution which could be due to variation in clay content and organic matter of these soils as reported by Yadav *et. al.* [49]. However, the top soil (0-26 cm) had the highest P concentration in the MUF profile as observed in the CUF profile. This may be a result of biological cycling as stated by [44].

Inter-profile distribution chart showed consistent decrease in P concentration down the slope; CUF had the highest concentration and VUF had the lowest concentration (CUF > MUF > VUF) (Fig.5b). This may be due to higher rate of runoff associated with down slope as stated by [8].

## Exchangeable potassium


The intra-profile distribution of exchangeable K as shown in Table 4 ranged from 0.08 to 0.10  $\text{cmolc kg}^{-1}$  for crest profile (CUF) with a mean of 0.088  $\text{cmolc kg}^{-1}$ , SD of 0.008  $\text{cmolc kg}^{-1}$  and CV 9.4 %. The middle slope profile (MUF) ranged from 0.09 to 0.10  $\text{cmolc kg}^{-1}$  with a mean value of 0.095  $\text{cmolc kg}^{-1}$ , SD of 0.005  $\text{cmolc kg}^{-1}$  and CV 5.3 %. While the valley bottom profile (VUF) ranged from 0.08 to 0.10  $\text{cmolc kg}^{-1}$  with a mean of 0.093  $\text{cmolc kg}^{-1}$ , SD of 0.008  $\text{cmolc kg}^{-1}$  and CV of 8.6 %.

Inter-profile distribution of exchangeable K as shown in Table 5 ranged from 0.088 to 0.095  $\text{cmolc kg}^{-1}$  across the three slopes (using intra-profile means) with a mean value of 0.092  $\text{cmolc kg}^{-1}$ , SD of 0.0029  $\text{cmolc kg}^{-1}$  and CV of 3.2 %.


These values show low exchangeable K at all depths and profiles; as indicated by the low intra and inter profile CV (Table 4&5). Furthermore, exchangeable K showed no definite distribution pattern down the profile depth of CUF and VUF (Fig.2c&4c). However, MUF showed a slightly higher level of K in shallower part of profile depth (Fig.3c). The VUF profile showed that K was higher in the surface though it did not decrease regularly with soil depths; as its concentration increased at lower profile depth of 116-150 cm (Fig.4c). This could be due to the effect of K cycling by the crops from bottom to surface horizons as reported by Singh *et al.* [43] and leaching respectively. According to [31], leaching moves nutrients downward and may increase nutrient concentrations with depth due to the effect of annual water table fluctuations causing leaching of nutrients down the profile, a characteristic of coastal sands. The distribution of K was shallowest in MUF (Fig.3c) where its abundance was highest (fig.5c) this contrasts with the predictions of [28] who stated that a high ratio of plant uptake to soil supply should result in higher rates of upward transport by plants and, hence, shallower vertical distributions.

Inter-profile distribution reveals that K was higher in valley bottom than crest and highest in the middle-slope (MUF > VUF > CUF) (Fig.5c), this may be due to deposition of K at valley bottom from upper slope [31].


a.



b.




c.




N-CUF = Nitrogen in the crest uncial farm  
P-CUF = Phosphorus in crest uncial farm  
K-CUF = Potassium in crest uncial farm

**Fig. 2:** Intra-profile distribution of primary nutrients (NPK) in crest uncial farm (CUF) soil profile


a.



b.




c.




N-MUF = Nitrogen in middle-slope uncultivated farm  
 P-MUF = Phosphorus in middle-slope uncultivated farm  
 K-MUF = Potassium in middle-slope uncultivated farm

**Fig. 3:** Intra-profile distribution of primary nutrients (NPK) in middle-slope uncultivated farm (MUF) soil profile



N-VUF = Nitrogen in valley bottom uncial farm  
P-VUF = Phosphorus in valley bottom uncial farm  
K-VUF = Potassium in valley bottom uncial farm

**Fig. 4:** Intra-profile distribution of primary nutrients (NPK) in valley bottom uncial farm (VUF) soil profile



**Fig. 5:** Inter-profile distribution of primary nutrients (NPK) along the toposequence: a, b & c for N, P & K respectively

## CONCLUSION

This study affirmed that the three profiles studied had predominantly sandy particle size distributions and mostly loamy sand in texture; especially at the topsoil level. The study also determined that the soils had very friable consistency at topsoil levels but were less friable at subsurface depths. They had weak fine granular structure at topsoil level but had mostly moderate sub-angular blocky structure at subsurface depths. The chemical properties analyzed showed that the soils were of low fertility status because their total N, exchangeable K and organic matter contents were below established critical levels. However, N was high at topsoil level but decreased consistently with increased depth especially at the crest profile. These, among other noticeable limitations including low contents of organic matter and high acidity were determined.

Furthermore, the NPK distribution bar charts showed that N and P decreased consecutively with increased depth for the crest profile. The middle-slope and valley bottom profile showed no definite pattern of distribution. However, the concentration of NPK was highest at topsoil level across almost all profiles. N had the shallowest intra-profile distribution with significantly higher level of topsoil concentration indicated by the high percentage of intra-profile coefficient of variation. P which had the highest concentration across all profiles also showed a shallow intra-profile distribution across the three profiles but did not vary significantly from the intra-profile mean as much as N. On the other hand K was more evenly distributed within all three profiles compared to N and P. Inter-profile distribution of NPK showed that N and P had higher concentrations at crest level, with P showing consecutive decrease in concentration down the slope. Furthermore, N had its lowest concentration in middle-slope profile while K was highest in the middle slope profile.

## RECOMMENDATIONS

Based on the findings of the research, the following recommendations are made.

- i. Conventional NPK fertilizer should be used to replenish inadequate contents of primary nutrients on the soils for crop production.
- ii. Slow release fertilizers can be used or split applications should be practiced.
- iii. Adoption of different NPK fertilizer recommendation with due considerations for different soil depths and topographic locations should be practiced to effectively tackle poor nutrient distribution.
- iv. The soils' acidity can be reduced by liming to replenish lost cations, which will increase the soil pH above the acid range for better crop production.
- v. Planting of acid tolerant crops is recommended for easier soil acidity management.
- vi. Artificial and live mulching in the form of cover crops are also recommended to reduce erosion, runoffs and leaching, and provide organic matter content for better supply of macro nutrients in the soil.
- vii. Application of organic manure can improve the soil structure and bring about the formation of soil aggregates that will withstand the high rainfall associated with the region and minimize water erosion.
- viii. Crop rotation is also recommended to enable natural replacement of nutrients removed from the soil by plants.



## REFERENCES

1. Ahn PM. *West African Soils*; Oxford University Press, London. 1970: 28-34.
2. Babalola O. Spatial Variability of soil water properties in Tropical Soils in Nigeria. *Journal of Soil Science*, 1999;126 (5); 269-279.27-54.
3. Babalola O, Obi ME. Physical Properties of Acid Sands in Relation to Land Use. In Udo E.J. and Subulo (Eds)' Acid Sands of Southern Nigeria. *Soil Science Society of Nigeria*. Spectator Publishing Company. 1981; 27-54.
4. Blake GR. Bulk density: In methods of Soil Analysis, In C. A. Black (Ed) Agronomy monograph, *American society of agronomy*, 1965;9; 374-390
5. Bulktrade ICL. *Main Report on Soil and land use Survey of Cross River State*. Calabar Ministry of Agriculture and Natural Resources. 1989; 376.
6. Bunemann EK., Schwenka GD, Van-Zwiten I. Impact of agricultural inputs on soil organism. A review. *Australian Journal of Soil Research*, 2006; 44, 379-408.
7. Danuhue LR. Miller WRS, Kickcluna CJ. *An introduction on Soils and Plant Growth*. 5<sup>th</sup> Edition, John Willey and Sons, London, 1990; 46-64.
8. Easton ZM Petrovic AM Effect of hill slope nutrient runoff from turf. *The Research*, 2005;109-113
9. Enwezor EG. Fertility status and productivity of acid sands. In acid soils of Southern Nigeria, monograph No. 1, *Soil Science Society of Nigeria*, 1981; 56-73.
10. Espinoza L, Norman R, Slaton N, Daniel M. *The nitrogen and phosphorus cycle in soils*. Agriculture and natural resources, University of Arkansas Cooperative Extension Service Printing Services, United States. 2005; 1-4.
11. Esu IE. *Soil Characterization, classification and management problems of the major soil orders in Nigeria*. Idea Investment and Printing Co. Ltd. Calabar. 2005;75

12. Ewulo BS, Ojanuga AG Ojeniyi SO. *Nature and Properties of Soils with Kandic Horizons in Humid South Western Nigeria*. In Management of the Resources of Nigeria for Sustainable Agriculture, 2002; 144-199.
13. Eyong MO, Akpa EA Effect of topo-position on the morphological and physicochemical properties of soils in Odukpani Local Government Area of Cross River State, *Nigeria journal of soil and tillage research*, 2018; 5:74-90.
14. Fairhust T. Handbook for integrated soil fertility management. Africa soil health consortium. Manabí. 2012.
15. FAO. *World Reference Base for Soil Resources*, World Soil Resources, 2014. Report No. 103.Rome.
16. F.A.O. *Soil Map of the World*. Revised Legend World Soil Resource, 1996. Report No.60.Rome.
17. Federal Ministry of Agriculture and Natural Resources, (FMANR) *Fertilizer Use Series*, 1990. No.1191.
18. Fertilizer Procurement and Distribution Division (FPDD). *Literature Review on Soil Fertility Investigation in Nigeria*. Lagos. Federal Ministry of Agriculture and Natural Resources. 1990.
19. Gee AW, Bauder J Particle size analysis. In Method of soil analysis. Part 1 agronomy 9, American society of agronomy Madison W. I. 1986; 383-411.
20. Hall GF. *Pedology and geomorphology*. L.P. Amsterdam Netherlands, 1993; 140.
21. Havlin JL, Beaton JD, Tisdale, SL, Nelson WL *Soil fertility and fertilizers: An introduction to nutrient management*. 2009. 7th Ed. Pearson Education Inc., Upper Saddle River, New Jersey, U.S.A.

22. Holland MD, Allen RKG, Barken D, Murphy ST. Land evaluation and agricultural recommendation for cross river national park, Oban Division. Report by the overseas development resources institute in Collaboration with WWF for the Federal Republic of Nigeria and the Cross River State Government, 1989.
23. Ibanga IJ. *Guidelines of soil survey, classification and land use*, De-Rio Press Nig. Ltd, Calabar. 2003; 67
24. Ibanga IJ, Eyong MO. *Report on land evaluation for pineapple production at Akpabuyo*, Cross River State Pineapple Project (CRSPAP). Governor's Office Calabar, 2002; 43.
25. Ibia TO, Obi JC, Udoh EJ. Phosphorus forms and fixation of representative soils of Akwa Ibom state of Nigeria. *Geoderma*, 1993; 58: 95-112.
26. Isirimah NO, Dickson AA, Igwe CA. *Introductory soil chemistry and biology for agricultural and biotechnology*. Osia Int'l Publisher Ltd, 2003; 270.
27. Jackson ML. *Soil Chemical Analysis*, Prentice Hall. Inc., Eaglewood Cliffs, New Jersey, (6<sup>th</sup> Ed), 1970; 44.
28. Jobbágy EG, & Jackson RB. The distribution of soil nutrients with depth: Global patterns and the imprint of plants, *Biogeochemistry*, Kluwer Academic Publishers, Netherlands, 2001; 53: 51–77.
29. Juo ASR. *Mineralogical Characteristics of Alfisols and Utisols with Variable Charge*. In B. K. G. Theng (Ed). *Soils with Variable Charge*. New Zealand Society of Soil Science. Lower Hutt, 1981.
30. Juo ASR. *Selected methods of soil Analysis*. IITA manual 1979; No.3,58
31. Kirby MJ. (1985). A basis for soil profile modelling in a geomorphic context. *Journal of soil science*. 36:1985; 97–121

32. Kogbe CA *Early Tertiary Bio-stratigraphy of the Southern Nigeria Sedimentary Basin*. Bull De' IFAN, T 37: 1975; 517-537.
33. Kyuma, K., Losaki T, and Juo ASR. *Evaluation of the soils*. In ASR Juo and Loe (ed) the wetland soil and rice. In sub-Sahara Africa. *Proceedings of international conference of wetland soil utilization for rice production in sub-Sahara Africa*. 1986.
34. Lal, R. Soil carbon sequestration impacts on global climate change and food security. *Soil Science*, 304(5677): 2004; 1623-1627.
35. Landon JR. (1991). Booker Tropical Soil Manual A. Hand book for Soil Survey and Agricultural Land Evaluation in the Tropics and Subtropics.
36. Metson AJ, *Methods of Chemical Analysis for Soil Survey samples*. New Zealand DSIR Soil Bur Bull. 12, Guvt Printer Wellington, New Zealand. 1961
37. Mokuwunye U, Bationo A. *Meeting the phosphorus needs of the soils and crops of West Africa: the role of indigenous phosphate rocks*. In Vanlauwe *et al.* (Eds) Integrated Plant Nutrient Management in Sub-sahara Africa. Cabi publishing, USA, 2002:352.
38. Murphy J, Rilley JP. A Modified Simple Solution for the Determination of Phosphorus in Natural Water, *Anai. Chim. Acta*. Vol. 27: 1962; 31-36
39. Ogban PI, Etim EA, Effiong GS. Mulching Practice on a Tropical Soil. *Journal of Soil Science*, 141(1): 1998; 32-45.
40. Salminen R, Batista MJ, Bidovec M, Demetriades A, De Vivo B, De Vos W, Duris M, Gilucis A, Gregorauskiene, V, Halamic J, Heitzmann, P, Lima A, Jordan G, Klaver G, Klein P, Lis J, Locutura J, Marsina K., Mazreku A, O'Connor PJ, Olsson SÅ, Ottesen RT, Petersell V, Plant JA, Reeder S, Salpeteur I, Sandström H, Siewers U, Steenfelt A, Tarvainen T. FOREGS Geochemical Atlas of Europe, Part 1: Background Information, Methodology and Maps. *Geological Survey of Finland*, Espoo. <http://weppi.gtk.fi/publ/foregsatlas/> 2005; 26

41. Schoenholtz SH, Miegroet HV, Burger JA. Physical and chemical properties as indicators of forest soil quality: Challenges and opportunities. *Forest Ecology and Management*, 138: 2000; 335 – 356.
42. Shaw JN, Truman CC, Reeves DW. *Mineralogy of eroded sediments derived from highly weathered soils*. Auburn University Press, USA. 2003; 81-96.
43. Singh K., Malik RVS, Sing V. Distribution of forms of potassium in alluvial soils. *Journal of Potassium Resources*, 7:2002; 116–118.
44. Stark JM. *Causes of soil nutrient heterogeneity at different scales*. In: Caldwell MM, Pearcy RW. (Eds) *Exploitation of Environmental Heterogeneity by Plants*. Academic Press, San Diego, United States 1994.
45. Thomas GW. *Exchangeable cation* in: AL, Page, RN Miller, DR Keoney (ed). *Method of soil analysis*. Part 2 2nd editions, ASA Madison. 1982; 159 - 164.
46. Udo EJ, Ibia O, Ano AO, Esu, IE. *Manual of Soil, Plant and Water Analysis*. Sibon Books. 2009; 183
47. Walkley A, Black IA. An examination of the digital method for determining soil organic matter and proposed modification of the chronic acid titration method. *Soil Science*. 37: 1934; 29-38
48. Vine H. Latisols of Nigeria and Some Related Soils. *Proceedings of 2<sup>nd</sup> Inter-Africa Soil conference*. 1970; 295-308.
49. Yadav NS, Verma R.S, Trivedi SK. Vertical distribution of forms of potassium in some soil series of vertisols Madhya Pradesh. *Journal of Indian Society of Soil Science*; 47:1999; 431–436.