

Effect of Locally Formulated Watermelon and Moringa Syrup Booster on the Growth Performance of *Heterobranchus bidorsalis* Fingerlings

ABSTRACT

Aim: Evaluation of the effectiveness of formulated Watermelon (*Citrullus lanatus*) and *Moringa oleifera* booster on the growth performance of *Heterobranchus bidorsalis*.

Study design: Data were subjected to statistical analysis using Duncan Multiple Range Statistics and interpretation was done using SPSS Version 22.

Place and duration of study: The experiment was conducted in the laboratory at the Department of Applied and Environmental Biology, Rivers State University, Nigeria for a period of 12 weeks (3 months), between November, 2016 – March, 2017.

Methodology: Total of 240 *Heterobranchus bidorsalis* fingerlings with an initial mean weight (1.61 ± 0.23 g) and length (5.13 ± 0.26 cm) were acclimated for one week. After acclimation, the fishes were fed twice daily (9am and 6pm) with 10% of their body weight after coating 2ml/1kg of the commercial feed with their individual growth booster syrup and allowed to air-dry for about 20 minutes. *Heterobranchus bidorsalis* fingerlings were scooped for the measurements exercise at one week interval after which they were returned to their various tanks.

Results: The result of this study recorded the experimental diets with optimum growth and survivability. From the result, it was deduced that, Commercial feed coated with watermelon syrup booster recorded the best performance in the growth variables of mean weight (26.36 ± 3.19 g), mean length (13.61 ± 1.35 cm), mean weight gain (24.64 ± 3.00 g), mean length gain (8.38 ± 1.35 cm), relative weight gain (25.36 ± 3.19) while Commercial feed coated with commercial syrup booster (CbCf) recorded the best performance in growth variables of Daily Growth Rate (15.15 ± 0.55) and Specific Growth Rate ($2.69 \pm 0.03\%$) against *Moringa* growth booster and commercial feed (control). However, there was less mortality as the fishes in different treatment tanks survived above 90%.

Conclusion: It could be summarized that Commercial feed coated with watermelon syrup booster (WbCf) had the best growth performance against the control groups although their values were not significantly impacted ($p > 0.05$) when compared with other experimental diets.

11
12
13
14
15
16
17
18

Keywords: Commercial feed coated with commercial syrup booster (CbCf), Commercial feed only (Cf), Commercial feed coated with watermelon syrup booster (WbCf), Commercial feed coated with *Moringa* syrup booster (MbCf), *Heterobranchus bidorsalis*.

1. INTRODUCTION

19
20
21
22
23

Aquaculture is a practice used all over the world, especially in some African Countries; millions of people practice aquaculture and have used it immensely in ancient times as their means of livelihood [6]. Fish rearing/ Aquaculture practices needs less labor input compared to other agricultural practices and the expected profit margin when properly maintained is usually on the high side. For a small scale fish farmer who wishes to culture *Heterobranchus*

24 *bidorsalis*, the fluctuating level of ingredients contained in commercial feed becomes a barrier.
25 This necessitates an innovative approach in utilizing an available formulated extract from
26 watermelon and *Moringa* growth booster to enhance the nutrients contained in the feed [7].
27 Watermelon and moringa contains vital nutrients which includes vitamins, minerals, essential
28 amino acid, phytochemicals such as carotenoids of which includes lycopene, Beta carotene,
29 lutein. Lycopene and Beta carotene are strong antioxidant that can help to protect the cell
30 against oxidative damage and therefore reduce risk of degenerative diseases caused by free
31 radicals [14, 5]

32 Growth boosters are effective growth promoters, formulated to supplement balance feed and
33 digestion in the target organism. Growth boosters are classified as Acidifiers, probiotics,
34 synbiotics, phytogenics, feed enzymes and immune stimulants. Watermelon and *Moringa*
35 syrup booster are categorized under phytogenics [10, 16].

36 Phytogenics are gotten from herbs, spices, aromatic plant etc. They are agents of microbes,
37 fungi, virus, oxidation etc, aid in digestion as such, increase the palatability of feed and
38 activate endogenous digestive enzymes, they are said to play major roles on the gut
39 microflora [16]. Given the level of production of both crops by local farmers especially in
40 Nigeria, bolstered by the enhanced awareness of the nutritional value of both plant products,
41 there is the opportunity to incorporate this product into preparation of high energy level fish
42 booster for enhanced productivity of catfish.

43 According to [7], *Heterobranchus bidorsalis*, which belongs to the *Clariidea* family, can be
44 reared on formulated and less expensive feed. The *Clariidea* can withstand unfavorable and
45 harsh environmental condition; they thrive in low oxygen and pH environment [8]. The
46 inadequacy of nutritive fish feed ingredients has been a major constraint to the survival of fish
47 culture in the competitive global food production system [2], as such, fish nutrition experts
48 world over have considered the need to review a natural growth promoter especially from
49 plant source to boost growth performance in fish farming [4]. On this note, plant products
50 comprising *Moringa oleifera* leave as well as Watermelon (*Citrullus lanatus*) were been
51 considered.

52 *Moringa oleifera* which belongs to the *Moringaceae* family is said to have originated from
53 southern foothills of the Himalayas in northwestern India. It is generally planted in tropical and
54 subtropical areas where its young seed pods and leaves are eaten as vegetables. Various
55 parts of *Moringa oleifera* contain some important minerals, essential amino acids, such as
56 methionine, cystine, tryptophan [3], it is also a good sources of energy, protein, vitamins,
57 carotene, and phenolics. *Moringa* is said to have peculiar functions which is essential to
58 human, animal, aquatic lives etc. Considering the benefits (nutritional and medicinal) of
59 *Moringa oleifera*, it has been promoted as a "healthful" food, traditionally used to fight some
60 popular ailments. Watermelon (*Citrullus lanatus*) is becoming an everyday fruit like apples,
61 bananas and oranges because of its usefulness and antioxidant properties. Watermelon was
62 also considered because its rind and seed are not usually consumed with the whole fruit, the
63 pulp is the major parts been consumed especially in Nigeria. This fact adds to the increase of

64 solid food waste responsibility. To prevent agricultural health related hazards, some
65 measures like the utilization of watermelon rind and seed as a growth booster were
66 considered.

67

68 **2. MATERIALS AND METHOD/ EXPERIMENTAL DETAILS/ METHODOLOGY**

69 This research was carried out in the laboratory at the Department of Applied and
70 Environmental Biology, Rivers State University located at Longitude 4.799° N and Latitude
71 6.983° S, Rivers State, Nigeria.

72

73 **2.1 Materials**

74 *Heterobranchus bidorsalis* fingerlings, Commercial feed (skretting of different variables, 1mm,
75 1.8mm and 2.5mm) table 1, *Moringa oleifera* leaves, watermelon seed, rind and pulp,
76 commercial booster (leegrow Tab.2), triple beam balance (OHAUS), scoop net, masking tape,
77 meter rule (cm) and twelve plastic tanks (30litre each).

78

79 **2.2 *Moringa oleifera* syrup booster formulation**

80 Fresh *Moringa oleifera* leaves used for this study were harvested from the Departmental
81 garden at Rivers State University, Rivers state Nigeria while other raw materials were
82 purchased from Ugo Resource and Health Farm Limited in Delta State Nigeria. The leaves
83 were sorted out, washed thoroughly, spread on a tray with evenly spaced openings. The
84 leaves were air-dried under shade at the laboratory for 3 days. After drying, the leaves were
85 thoroughly blended in a fine powdery form. 20% of the blended *Moringa* leaf was mixed with
86 20% distilled water, 50% sucrose base and enzyme (binder), 10% of vitamin and mineral
87 premix. The proximate content of the aqueous extract was analyzed for Moisture, Ash,
88 Protein, Fats and Carbohydrates (see Table 2).

89

90 **2.3 Watermelon syrup booster formulation**

91

92 Watermelon fruits were bought and brought to the Research Laboratory at the Department of
93 Applied and Environmental Biology, Rivers State University, Rivers State Nigeria. It was
94 properly washed and sliced, seeds were picked out, rind was carefully scraped and chopped,
95 and pulp obtained. 60% of Watermelon (pulp, rind and seed) was thoroughly blended and
96 mixed with 30% sucrose base and feed enzyme (binder) and 10% of vitamin and mineral
97 premix were used in watermelon growth booster formulation. The proximate content of the
98 formulated Booster were analyzed as (see Table 2).

99

100 **2.4 Collection of Fish**

101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140

Total of two hundred and forty (240) *Heterocbranchus bidorsalis* fingerlings (mean weight, $1.65 \pm 0.23\text{g}$; mean length $5.13 \pm 0.26\text{cm}$) were obtained from National Institute of Marine and Oceanographic Research, Sapele branch in Delta State Nigeria (NIOMOR). The fishes were transported in a transparent aquarium to the experimental laboratory, Rivers State University, Rivers State, Nigeria. The fishes were put into a transparent aquarium and taken to the laboratory. *Heterobranchnus bidorsalis* fingerlings were evenly distributed into twelve plastic tanks (30litres each) at a stocking density of 20 fingerlings per tank. They were acclimated for one week during which they were placed on a maintenance diet with a commercial feed (skretting) once daily at 3% of their body weight. Water lettuce was introduced in the setup to enhance acclimation.

2.5 Experimental Procedures

After acclimation, twelve plastic tanks (30litre each) were randomly arranged and labeled into 4 Treatments (T. 1, 2, 3 and 4) with two replicates (R1, 2) and (R1, 2).

T1- Commercial booster with commercial feed (**CbCf**) - positive control.

T2- Commercial feed only (**Cf**) - Negative control.

T3- Watermelom booster with commercial feed (**WbCf**)

T4- *Moringa oleifera* booster with commercial feed (**MbCf**)

Fingerlings were weighed and measured to determine its initial mean weight and length. The fishes were fed twice daily (9am and 6pm) with 10% of their body weight after coating 2ml/1kg of the commercial feed with their individual growth booster syrup and allowed to air-dry for about 20 minutes.

Water was siphoned from each treatment tank daily and refilled from the tap maintaining its original water volume (25 liter). Weekly mean weight of the fishes in each labeled plastic tank were taken (to the nearest 0.01g) with OHAUS Triple Beam Balance (2610 g), weekly mean length from individual standard lengths of the fishes in each labeled plastic tank were also recorded (to the nearest 0.1cm) with a meter rule.

Heterobranchnus bidorsalis fingerlings were scooped for the measurements exercise at one week interval after which they were returned to their various tanks. Growth performance was determined and were recorded followed by the observation period of three months (Nov 2016-Feb 2017).

2.6 Monitoring of Physico-chemical Parameters

Temperature and pH values were measured daily using glass thermometer and pH kit respectively. Other physico-chemical parameters were obtained using Extech instrument (DO 700) from Institute of Pollution Studies (IPS) RSU and the values obtained were recorded in Table 3.

141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

2.7 Growth parameters

Growth performance, condition factor and survivability were calculated as follows;

(a) Initial Mean Weight = (g / fish)

(b) Final Mean Weight = (g / fish)

(c) Initial Mean Length = (cm / fish)

(d) Final Mean Length = (cm / fish)

(e) Mean Weight Gain (g) = $W1 - W0$ (Where $W1$ =Final Weight, $W0$ =Initial Weight)

(f) Mean Length Gain (cm) = $L1 - L0$ (Where $L1$ =Final Length, $L0$ =Initial Length)

(g) Daily Growth Rate (g) = $\frac{\text{Mean Weight Gain (g)}}{\text{Initial Body Weight (g)}}$

(h) Relative Weight Gain (g) = $\frac{W1 - W0}{W0}$ (g) (Where $W1$ =Final Weight, $W0$ =Initial Weight)

(i) Specific Growth Rate = $\frac{\ln(W1) - \ln(W0)}{T} * 100$

(Where \ln = Log e Final Weight- Log e Initial Weight, T= Culture period)

2.8 Statistical Analysis

Data generated were subjected to one way analysis of variance (ANOVA) with Duncan's Multiple Range Descriptive Test (Duncan, 1995). The result computation was done using Statistical Package for Social Science (SPSS) version 22. Differences among mean were separated with Turkey HSD (Honest Significant Difference) at $p < 0.05$.

178 **3. RESULTS AND DISCUSSION**
 179

180 **Table 1: Commercial feed of different size 1mm, 1.8mm and 2.5mm (Company:**
 181 **skretting, proximate composition**
 182

Nutrients	Compositions (%)		
	1.1mm	1.8mm	2.5mm
Crude protein	52	45	45
Crude fibre	0.9	2.6	2.6
Crude ash	15	14	14
Fats	9.5	7.5	7.5

183

184

185

186

187

188

189 Table 2 showed the proximate analysis of *Moringa*, watermelon and commercial syrup
 190 booster. This result showed that the moisture content in *Moringa* was significantly higher
 191 ($P=.05$) than that of the control (commercial syrup booster) and watermelon syrup
 192 booster. Comparing proximate composition variables of Ash and crude protein content of
 193 the growth boosters, the control (commercial syrup booster) showed a significantly higher
 194 ($P=.05$) result against *Moringa* and watermelon booster. Fats content of commercial syrup
 195 booster is significantly lower ($P=.05$) than that of *Moringa* and watermelon syrup growth
 196 booster. Carbohydrate is significantly higher ($P=.05$) in commercial syrup booster
 197 followed by watermelon and *Moringa* has the least.

198

199

200 **Table 2: Proximate Analysis of *Moringa*, Watermelon and Commercial Syrup Booster**
 201
 202

203

Variables	Samples (3) /compositions (%)		
	<i>Moringa</i> Syrup Booster	Watermelon Syrup Booster	Commercial Syrup Booster
Moisture	47.81±0.77 ^a	36.55±0.64 ^b	20.37±0.57 ^c
Ash	0.14±0.13 ^{bc}	0.09±0.04 ^{bc}	0.59±0.00 ^a

Protein	0.19±0.12 ^{bc}	0.18±0.11 ^{bc}	0.59±0.00 ^a
Fats	0.61±0.13 ^{ab}	0.53±0.06 ^{ab}	0.07±0.03 ^c
Carbohydrates	51.24±0.64 ^c	62.66±0.24 ^b	77.31±0.47 ^a


*Mean ±SD of moringa, watermelon and commercial syrup booster in the same column with different superscript are significantly different ($P=0.05$)

Table 3 Showed Physicochemical Parameters of water sample obtained using Extech instrument (DO 700) from Institute of Pollution Studies (IPS) RSU and the values obtained were recorded.

Table 3: Physicochemical Parameters of Water Sample

Parameters	Value mean
Temperature (°C)	27.5
PH	6.5
Dissolved Oxygen (mg/l)	5.97
Conductivity(µS/cm)	116
Salinity (%)	0.05
BOD (mg/l)	3.45
Turbidity (NTU)	1.06
Total Dissolved Solid (mg/l)	60

Figure 1 showed mean weight of *Heterobranchus bidorsalis* fed different experimental. Commercial feed coated with watermelon syrup booster (WbCf) recorded the highest mean weight followed by commercial feed coated with commercial syrup booster (CbCf) positive control. Commercial feed (Cf) negative control also recorded gradual increase in mean weight. Commercial feed coated with *Moringa* syrup booster (WbCf) had the least mean weight. There was gradual increase in mean weight in all the experimental diet from week 1 to 8, week 9 to 12 recorded a significantly higher ($P=.05$) weight increase.



235
236
237


Fig. 1: Mean weight of *Heterobranchus bidorsalis* fed experimental diets.

238 Key: CbCf Commercial feed coated with commercial syrup booster
 239 Cf Commercial feed only
 240 WbCf Commercial feed coated with watermelon syrup booster
 241 MbCf Commercial feed coated with *moringa* syrup booster

242
243
244

245 Figure 2 showed mean length of *Heterobranchus bidorsalis* fed different experimental diet:
 246 Commercial feed coated with commercial syrup booster (CbCf), commercial feed (Cf), commercial
 247 feed coated with watermelon syrup booster (WbCf) and commercial feed coated with *Moringa* syrup
 248 booster (MbCf). Experimental diet WbCf had the highest mean length followed by (Cf) negative
 249 control, CbCf. MbCf recorded the least mean length. The increase in length was not significantly
 250 different ($P=.05$) from week 1 (start value) to the 12th week.

251



252


253 **Fig. 2: Mean Length of *Heterobranchus bidorsalis* experimental diets at different**
 254 **durations**

255

256

257 Figure 3 showed Mean weight gain of *Heterobranchus bidorsalis* fed different experimental diets.
 258 From this figure, Commercial feed coated with watermelon syrup booster (WbCf) recorded the highest
 259 mean weight gain, followed by commercial feed only (negative control). Commercial feed coated with
 260 commercial syrup booster (CbCf) and *Moringa* syrup booster (MbCf) had the least mean weight gain.
 261 The increase in mean weight gain of all the experimental diets from week 1-8 was not significantly
 262 impacted ($P=.05$). There was significant different ($P=.05$) from week 8-12 as such, a sharp increase in
 263 weight gain was recorded.


264



265 **Fig. 3:** Mean weight Gain of *Heterobranchus bidorsalis* fed with experimental
 266 diets at different durations.
 267
 268
 269

270
 271
 272
 273
 274
 275
 276
 277

Figure 4 showed mean length gain of *Heterobranchus bidorsalis* fed different experimental diets. Commercial feed coated with watermelon syrup booster (WbCf) recorded the highest mean length gain, (see Fig.4), followed by commercial feed only (negative control). Commercial feed coated with commercial syrup booster (CbCf) and *Moringa* syrup booster (CbCf and MbCf) had the least mean weight gain (Fig.4)



278
 279 **Fig. 4: Mean Length Gain of *Heterobranchus bidorsalis* fed with experimental diets at**
 280 **different durations**
 281

282
 283
 284 Figure 5 showed the Daily growth rate of *Heterobranchus bidorsalis* fed different experimental diets.
 285 Here, the control: commercial feed coated with commercial syrup booster (CbCf) and commercial feed
 286 only (Cf) had the best daily growth performance while commercial feed coated with watermelon and
 287 *moringa* syrup booster had the least daily growth rate performance (Fig.5) Also, a gradual increase in
 288 daily growth rate of all the experimental diets from week 1-7 were recorded. On the 8th week, there
 289 was sharp increase in daily growth rate which affected from week 9 to the end of the experimental
 290 duration.




Fig. 5: Daily Growth Rate of *Heterobranchus bidorsalis* fed with experimental diets at different durations

291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303

Figure 6 showed Relative weight gain of *Heterobranchus bidorsalis* fed different experimental diets. This figure showed rapid Relative weight gain of all the experimental diets from week 8-12. Experimental diet commercial feed coated with watermelon syrup booster (WbCf) recorded the highest Relative weight gain against the control groups (CbCf and Cf). Commercial feed coated with *Moringa oleifera* syrup booster (MbCf) had the least Relative weight gain.





Fig. 6: Relative Weight Gain of *Heterobranchus bidorsalis* fed with experimental diets at different durations

304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318

Figure 7 showed specific growth rate of *Heterobranchus bidorsalis* fed different experimental diets. The control: commercial feed coated with commercial syrup booster (CbCf) and commercial feed only (Cf) recorded the highest specific growth rate against commercial feed coated with watermelon syrup booster. *Moringa* syrup booster had the least specific growth rate. Specific growth rate was also negatively affected in week 1. Week 2-12 recorded increase in specific growth rate although the values were not significantly different ($P=.05$) from each other.




319 **Fig. 7: Specific Growth Rate of *Heterobranchus bidorsalis* fed with experimental diets at**
 320 **different durations**
 321

322

323

324 Figure 8 showed the survivability of *Heterobranchus bidorsalis* fed different experimental diets.
 325 Commercial feed only (Cf) had 100% survivability from week 1-5, commercial feed coated with
 326 *moringa* and watermelon syrup booster (MbCf and CbCf) also had 100% survivability only in week 1.
 327 Generally, *Heterobranchus bidorsalis* fed all experimental diets survived above 90%.



328
329

330 **Fig. 8: Survivability of *Heterobranchus bidorsalis* fed with experimental diets at different**
331 **duration**

332
333
334

335 Table 4 showed the overall growth and nutrient utilization variables of *Heterobranchus bidorsalis* fed
336 with all experimental diets. The growth pattern observed from growth variables under the experimental
337 diet of commercial feed coated with commercial syrup booster (CbCf) showed no significant different
338 ($P=0.05$) result. WbCf recorded the highest numerical values in growth ($P=0.05$) from other experimental
339 diets

340
341

342 **Table 4: Cumulative variables of *Heterobranchus bidorsalis* fed all experimental**
343 **diets after the trial period (mean±SD)**

VARIABLES	CbCf	Cf	WbCf	MbCf
Initial Mean Weight	1.58±0.04 ^b	1.60±0.51 ^b	1.71±0.20 ^a	1.56±0.19 ^b
Initial Mean Length	4.77±0.15 ^c	5.30±0.40 ^a	5.13±0.22 ^b	5.33±0.30 ^a
Final Mean Weight (g)	25.56±0.76 ^a	25.76±2.27 ^a	26.36±3.19 ^a	23.79±1.64 ^b
Final Mean Length (cm)	12.88±0.33 ^b	13.44±0.89 ^{ab}	13.61±1.35 ^a	12.50±0.28 ^c
Mean Weight Gain (g)	23.98±0.75 ^a	24.16±2.31 ^a	24.64±3.00 ^a	22.22±1.45 ^b
Mean Length Gain (cm)	8.11±0.48 ^a	8.14±1.25 ^b	8.38±1.49 ^a	7.16±0.55 ^c

Daily Growth Rate	15.15±0.55 ^a	15.07±1.8 ^a	14.37±0.76 ^b	14.25±0.87 ^b
Relative Weight Gain (%)	24.56±0.76 ^a	24.76±2.27 ^a	25.36±3.19 ^a	22.79±1.64 ^b
Specific Growth Rate (%)	2.69±0.03 ^a	2.68±0.12 ^a	2.63±0.06 ^b	2.63±0.08 ^b
Survivability (%)	92.80±2.58 ^a	92.80±2.58 ^a	95.20±2.88 ^a	90.40±4.08 ^a

*Mean ±SD in the same row with different superscript are significantly different (P<0.05)

Key: CbCf Commercial feed coated with commercial syrup booster
Cf Commercial feed only
WbCf Commercial feed coated with watermelon syrup booster
MbCf Commercial feed coated with *moringa* syrup booster

The use of formulated syrup booster from watermelon and *Moringa* coated with commercial feed has revealed that, *Heterobranchus bidorsalis* fed with the experimental diet CbCf, Cf recorded no significant difference (P=.05) in growth variables. Experimental diets of commercial feed coated with watermelon syrup booster WbCf had the highest Final Mean Weight (26.36±3.19g) and Length (13.61±1.35cm) when compared with other experimental diets: CbCf (25.26 ± 0.76g, 12.88 ± 0.33cm), Cf (25.76 ± 2.77g, 13.44 ± 0.89cm). MbCf had the lowest value (23.79 ± 1.64g, 12.05 ± 0.28cm) in final mean weight and length. The calculated Digestible Energy level of the three booster used were (256.10 kcal /kg) for WbCf, *Moringa* Growth Booster (211.17 kcal/kg) and (312.23 kcal/kg) for CbCf. In contrast to the present findings, high percentage in carbohydrates (62.66 and 51.24%) from proximate analysis of the growth boosters (watermelon and *Moringa oleifera*) (Table 2) is not comparable with the result obtained from United State Department of Agriculture, National Nutrient Data base USDA (2016) [15] on *Moringa oleifera* and watermelon (8.28 and 7.55g) but justifies Carbohydrates as an important non-protein energy source for fish and should be included in the diet at an appropriate level to maximize the use of dietary protein for growth and to facilitate movement of nutrient at the Gastro Intestinal Tract (GIT) which supports nutrient absorption [10].

Commercial feed coated with watermelon syrup booster showed poor performance on Daily Growth Rate (DGR) (14.37±0.76) while commercial feed coated with commercial syrup booster CbCf had the highest value on variables of DGR (15.15±0.55%), Specific Growth Rate (2.69±0.03%), but the values were not significantly different at P>0.05 when compared to other experimental diets. The poor growth performance from the experimental diets commercial feed coated with *Moringa* syrup booster MbCF is similar with that of [2], who reported that, at more than 10% concentration of *Moringa oleifera* inclusion in feed meal, the bitter taste of saponin and phenol becomes obvious, as such, causing low feed intake which invariably affects growth response.

This crude protein level in CbCf is contrary to the report of [4]. They recorded higher crude protein level in feed additives (Aqua pro) for African catfish. With this results, it can be said that watermelon syrup booster recorded the highest value in most of the growth variables although comparing with other experimental diets, the values were not significantly impacted (P=.05).

CONCLUSION

383 From the overall observations, this research has revealed Watermelon growth booster be a promising
384 fish feed supplement having shown the highest growth performance. According to [4], a lot of
385 literatures exist on other growth booster (Aqua booster, Aqua pro, Leegrow) unlike that of *Citrullus*
386 *lanatus* and *Moringa oleifera* as a fish growth booster. As such, fish farmers should look inward on the
387 utilization of watermelon growth syrup booster for effective growth performance of catfish, also for the
388 more accessible cost and within a policy of green circular economy that aims at food waste recover.

389 **Ethical Approval:**

390 As per international standard or university standard ethical approval has been collected and
391 preserved by the authors.

392

393 Consent:NA

394

395

396 **REFERENCES**

397

- 398 1. Afung, W., Siddhuraju, P and Becker, K). Comparative nutritional evaluation of raw, methanol
399 extracted residues and methanol extracts of Moringa (*Moringa oleifera* Lam.) leaves on growth
400 performance and feed utilization in Nile tilapia. *Aquaculture*. 2003; 34: 1147-1159.
401
- 402 2. Alphonsus, O., Ebere, S. and Joseph, O. Replacement of fish meal with maggot in African catfish
403 (*Clarias gariepinus*) diets. Department of Animal Science, Anambra State University Igbariam.
404 *Bioline International journal*. . 2009; (9): 666 - 671.
405
- 406 3. Becker, K., and Francis, G.). Anti-nutritional factors present in plant – derived alternative fish feed
407 Ingredients and their effects in fish. *Aquaculture*. 2001 (199): 3 - 4.
408
409
- 410 4. Dada, A. and Olugbemi, B. Dietary effects of two commercial feed additives on growth
411 performance and body composition of African catfish, *Clarias gariepinus* fingerlings. Federal
412 University of Technology Akure, Ondo State, Nigeria. *African Journal of Food Science*. 2013; 7(9):
413 325 - 328.
414
- 415 5. Di Mascio, P., Kaiser, S., Sies, H. Lycopene as the most efficient biological carotenoid singlet
416 oxygen quencher. *Archives of Biochemistry and Biophysics*. 1989; 274: 532-538
417
- 418 6. Food and Agricultural Organization (FAO). FAO Technical Guidelines for Responsible Fisheries.
419 Precautionary approach to capture fisheries and species introduction.1996; (2):54
420
- 421 7. Foundation for Partnership in Niger Delta (PIND). Fish Feed Value Chain Analysis in Niger Delta.
422 2017: Update of an earlier study, 2012.
423
- 424 8. Fagbenro, O.A, Olaniran, T.S, Esan, A.O. Some aspect of biology of catfish, *Heterobranchus*
425 *bidorsalis* (*Clariidea*) in River Ogbese, Nigeria *African Journal Zoology* 1991:105, 363-327.
426
- 427 9. Huisman, E. and C.J. Reproduction, growth, health control and Aquacultural potential of the
428 African catfish, *Clarias gariepinus*. *Aquaculture*. 1987; (63): 1 - 14.
429

- 430 10. Manner, K. Effects of phytogenic feed additives on growth performance and ileal nutrient
431 digestibility in broiler chicken. *Poultry Science*. 2011; 90(12): 2811 - 2816. doi: 10.3382/ps.2011-
432 01515. PMID 22080020.
433
- 434 11. Madhuri S, Y.P Sahni and Govind Pandey. Herbal feed supplement as drugs and growth
435 promoter to fishes. Department of Zoology and Biotechnology, Government Model Science
436 College, Jabalpur, India. 2012, 3 (9)
437
- 438 12. Richter, N, Siddhuraraju, A, Becker, K. Evaluation of nutritional quality of *Moringa (Moringa*
439 *oleifera* Lam) leaves as alternative protein source of fish meal. *Aquaculture*. 2003; 217: 599 - 611.
440
- 441 13. Shiau, S. and Lin, Y. Carbohydrates utilization and its protein-sparing effect in diets for grouper
442 (*Epinephelus malabaricus*). *Animal Science*. 2001; 73: 299 - 304.
443
- 444 14. Scalbert A, Manach C, Morand C, Remesy C, Jimenez L. Dietary polyphenols and the prevention
445 of diseases. *Critical Reviews in Food Science and Nutrition*. 2005; 45:287-306.
446
- 447 15. United State Department of Agriculture, National Nutrient Data base USDA. Agricultural Research
448 Service. National Nutrients Database for Standard Reference. 2016; Released 28.
449
- 450 16. Windisch W, Schedle K, Pletzner C, Kroismayr A. Use of phytogenics products as feed additives
451 for swine and poultry. *Journal Animal Science*. 2008; 86 (suppl 14):E140-E148.
452