

Numerical Optics Soliton Solution of the Nonlinear Schrödinger Equation Using the Laplace and the Modified Laplace Decomposition method

ABSTRACT:

In this paper, the Laplace decomposition method (LDM) and some modification, namely the Modified Laplace decomposition method (MLDM), are adopted to numerically investigate the optic soliton solution of the nonlinear complex Schrödinger equation (NLSE). The obtained results demonstrate the reliability and the efficiency of the considered methods to numerically approximate such initial value problems (IVPs).

Keywords: Nonlinear Schrödinger equation, Laplace Transform, Adomian polynomials.

AMS 2010 Classification: 35Q40, 35Q60, 35C08, 44A10, 49M27

1. Introduction

The nonlinear complex Schrödinger equation (NLSE) is an equation which models many physical phenomena such as nonlinear optics, water waves, plasma physics, ... etc. Particularly, the nonlinearity effects in an optic fiber including four-wave mixing, self-phase modulation, second harmonic generation, ... etc. are modeled by the NLSE [1], [2]. Moreover, the evolution of the envelope of modulated nonlinear water wave groups are essentially described by the NLSE. All these mentioned physical phenomena are eventually interpreted by the exact solutions for specified values of the NLSE's parameters. In this paper we consider the Nonlinear Schrödinger equation (NLSE) of the form:

$$i \frac{\partial \Psi}{\partial t} + P \frac{\partial^2 \Psi}{\partial x^2} + Q \Psi |\Psi|^2 = 0 \quad (1)$$

where $\Psi(x, t)$ is a complex-valued function of real variables (x, t) , and P, Q are nonzero real parameters. The NLSE (1) admits the optic soliton solution [3]:

$$\Psi(x, t) = \sqrt{\frac{-2P\alpha^2}{Q}} \tanh(\alpha x - 2k\alpha t + \xi_0) e^{i(kx - P(2\alpha^2 + k^2)t + \eta_0)} \quad (2)$$

in which $PQ < 0$ gives the de-focusing case, α is a soliton velocity, $P(2\alpha^2 + k^2)$ is a soliton wave number, k is a nonlinear frequency shift and ξ_0, η_0 are arbitrary constants.

Last recent decades, the methods of decomposing have emerged as a powerful technique and as a subject of extensive analytical and numerical studies for large and general class of linear and nonlinear ordinary differential equations (ODE's) as well as partial differential equations (PDE's), fractional differential equations, algebraic, integro-differential, differential-delay equations [4]–[17]. More precisely, the Adomian decomposition method is knowingly efficient in solving initial-value or boundary value problems without unphysical restrictive assumptions such as linearization, perturbation and so forth. The method provides the solution in an infinite series that is proven to converge rapidly with elegant computable components [4], [5], [10]. In recent years a large amount of research work concerning the developing of the ADM is investigated see for instance [18]–[23]. In particular, the modification that was proposed by Wazwaz and El-Sayed

[22], suggests that the zeroth component of the decomposition series can be divided into two functions in which the first part is only assigned to the zeroth component whereas the second part is combined with recursive relation. This modified form is adopted along with the LDM to formulate the MLDM [24], [25] and to be implemented in the current numerical study.

Laplace Decomposition Method (LDM) was introduced by Khuri [11], [12] and has been successfully utilized for obtaining solutions of differential equations [6], [7], [9], [14], [17], [26]–[34] and the NLSE of our interest. As, for instance, a recent study by Gaxiola [26] who applied the Laplace-Adomian decomposition method to a NLS-like equation, namely the Kundu-Eckhaus equation, and the accuracy as well as the efficiency of the method is proved via examples, as for the nonlinear Schrodinger equation with harmonic oscillator the method of Laplace-Adomian was utilized in a comparison with another semi-analytical method to obtain approximate analytical solutions by Jaradat et. al. [28]. The Powerfulness of this method is its consistency of Laplace transform and Adomian polynomials which guarantees an accelerative, rapid convergence of series solutions when compared with the ADM itself and therefore provides major progress [11], [35], [36]. The main numerical approach in this article is implementing the Laplace decomposition method (LDM) and the Modified Laplace Decomposition method (MLDM) to the NLSE (1), for this purpose the paper is organized to fully analyze the considered methods in Section 2. Numerical results are obtained and plentifully discussed via tables, illustrations and concluding remarks in Section 3. Finally, in Section 4 a brief conclusion is given.

2. Methodology of the used Methods

2.1 LDM algorithm of the NLSE

In this section we begin with reducing the nonlinear Schrödinger equation (NLSE) (1) into a system of coupled nonlinear equations involving the real and imaginary parts, by introducing the following transformation [11], [37] :

$$\Psi(x, t) = \psi_1(x, t) + i\psi_2(x, t) \quad (3)$$

where $\psi_1(x, t)$ and $\psi_2(x, t)$ are real-valued functions. Substituting (3) into (1) we obtain the following system of coupled real equations with an initial value problem (IVP), to take the following from:

$$\frac{\partial \psi_1}{\partial t} + P \frac{\partial^2 \psi_2}{\partial x^2} + Q(\psi_1^2 + \psi_2^2)\psi_2 = 0 \quad (4)$$

$$\begin{aligned} \frac{\partial \psi_2}{\partial t} - P \frac{\partial^2 \psi_1}{\partial x^2} - Q(\psi_1^2 + \psi_2^2)\psi_1 &= 0 \\ \begin{cases} \psi_1(x, 0) = F(x) \\ \psi_2(x, 0) = G(x) \end{cases} \end{aligned} \quad (5)$$

Rewriting (4) in the following operator form:

$$\begin{aligned} L_t \psi_1 &= -(PL_{xx} \psi_2 + QN(\psi_1, \psi_2)) \\ L_t \psi_2 &= PL_{xx} \psi_1 + QM(\psi_1, \psi_2) \end{aligned} \quad (6)$$

where $L_t \equiv \frac{\partial}{\partial t}$, $L_{xx} \equiv \frac{\partial^2}{\partial x^2}$ are the linear differential operators, and $N(\psi_1, \psi_2) = (\psi_1^2 + \psi_2^2)\psi_2$,

$M(\psi_1, \psi_2) = (\psi_1^2 + \psi_2^2)\psi_1$ symbolize the nonlinear operators.

Applying the Laplace Transform on both sides of the system in (6), and using the Laplace properties with the initial conditions, we get:

$$\begin{aligned}\psi_1(x, s) &= \frac{1}{s} F(x) - \frac{1}{s} [PL [L_{xx} \psi_2] + QL [N(\psi_1, \psi_2)]] \\ \psi_2(x, s) &= \frac{1}{s} G(x) + \frac{1}{s} [PL [L_{xx} \psi_1] + QL [M(\psi_1, \psi_2)]]\end{aligned}\quad (7)$$

The method assumes that the unknown functions $\psi_1(x, s)$, $\psi_2(x, s)$ are expressed as infinite series in the form:

$$\psi_1(x, s) = \sum_{n=0}^{\infty} \psi_{1,n}(x, s), \quad \psi_2(x, s) = \sum_{n=0}^{\infty} \psi_{2,n}(x, s) \quad (8)$$

And the nonlinear operators are expressed in terms of an infinite series of the well-known Adomian polynomials (see for example [4], [38]) given by:

$$\begin{aligned}N(\psi_1, \psi_2) &= \sum_{n=0}^{\infty} A_n = \sum_{n=0}^{\infty} \left(\frac{1}{n!} \frac{d^n}{d\lambda^n} \left[N \left(\sum_{i=0}^{\infty} \lambda^i \psi_{1i}, \sum_{i=0}^{\infty} \lambda^i \psi_{2i} \right) \right] \right) \\ M(\psi_1, \psi_2) &= \sum_{n=0}^{\infty} B_n = \sum_{n=0}^{\infty} \left(\frac{1}{n!} \frac{d^n}{d\lambda^n} \left[M \left(\sum_{i=0}^{\infty} \lambda^i \psi_{1i}, \sum_{i=0}^{\infty} \lambda^i \psi_{2i} \right) \right] \right)\end{aligned}\quad (9)$$

Listing below a few components of Adomian polynomials:

$$\begin{aligned}A_0 &= \psi_{1,0}^2 \psi_{2,0} + \psi_{2,0}^3, \\ A_1 &= 2\psi_{1,0} \psi_{1,1} \psi_{2,0} + \psi_{1,0}^2 \psi_{2,1} + 3\psi_{2,0}^2 \psi_{2,1}, \\ A_2 &= \psi_{1,1}^2 \psi_{2,0} + 2\psi_{1,0} \psi_{1,2} \psi_{2,0} + 2\psi_{1,0} \psi_{1,1} \psi_{2,1} + 3\psi_{2,0} \psi_{2,1}^2 + \psi_{1,0}^2 \psi_{2,2} + 3\psi_{2,0}^2 \psi_{2,2}, \\ &\vdots\end{aligned}\quad (10)$$

$$\begin{aligned}B_0 &= \psi_{1,0}^3 + \psi_{1,0} \psi_{2,0}^2, \\ B_1 &= 3\psi_{1,0}^2 \psi_{1,1} + \psi_{1,1} \psi_{2,0}^2 + 2\psi_{1,0} \psi_{2,0} \psi_{2,1}, \\ B_2 &= 3\psi_{1,0} \psi_{1,1}^2 + 3\psi_{1,0}^2 \psi_{1,2} + \psi_{1,2} \psi_{2,0}^2 + 2\psi_{1,1} \psi_{2,0} \psi_{2,1} + \psi_{1,0} \psi_{2,1}^2 + 2\psi_{1,0} \psi_{2,0} \psi_{2,2}, \\ &\vdots\end{aligned}\quad (11)$$

Using (8) and (9) into (7), we have:

$$\begin{aligned}\sum_{n=0}^{\infty} \psi_{1,n}(x, s) &= \frac{1}{s} F(x) - \frac{1}{s} \left[PL \left[L_{xx} \sum_{n=0}^{\infty} \psi_{2,n} \right] + QL \left[\sum_{n=0}^{\infty} A_n \right] \right] \\ \sum_{n=0}^{\infty} \psi_{2,n}(x, s) &= \frac{1}{s} G(x) + \frac{1}{s} \left[PL \left[L_{xx} \sum_{n=0}^{\infty} \psi_{1,n} \right] + QL \left[\sum_{n=0}^{\infty} B_n \right] \right]\end{aligned}\quad (12)$$

According to (for example [4], [11], [12]), comparing both sides of (12) by applying the inverse Laplace transform, we obtain the subsequent components to take the following recursive relation:

$$\begin{aligned}
\psi_{1,0}(x,0) &= F(x) \\
\psi_{2,0}(x,0) &= G(x) \\
\psi_{1,n+1}(x,t) &= -L^{-1} \left[\frac{1}{s} \left[PL [L_{xx} \psi_{2,n}] + QL [A_n] \right] \right] \\
\psi_{2,n+1}(x,t) &= L^{-1} \left[\frac{1}{s} \left[PL [L_{xx} \psi_{1,n}] + QL [B_n] \right] \right], n \geq 0
\end{aligned} \tag{13}$$

Obviously, the practical solution will be the n -term approximations of the infinite series (8). Thus the solution of (3) is given by:

$$\begin{aligned}
\Psi(x,t) &= (\psi_{1,0} + \psi_{1,1} + \psi_{1,2} + \dots) + i(\psi_{2,0} + \psi_{2,1} + \psi_{2,2} + \dots) \\
&= \left[R \cos(kx + \eta_0) \tanh(\alpha x + \xi_0) - Rt \left(2P\alpha \operatorname{sech}(\alpha x + \xi_0)^2 (k \cos(kx + \eta_0) \right. \right. \\
&\quad \left. \left. - \alpha \sin(kx + \eta_0) \tanh(\alpha x + \xi_0)) \right. \right. \\
&\quad \left. \left. + \sin(kx + \eta_0) \tanh(\alpha x + \xi_0) (-k^2 P + QR^2 \tanh(\alpha x + \xi_0)^2) + \dots \right] \right. \\
&\quad \left. + i \left[R \sin(kx + \eta_0) \tanh(\alpha x + \xi_0) + Rt \left(-2P\alpha \operatorname{sech}(\alpha x + \xi_0)^2 (k \sin(kx + \eta_0) \right. \right. \right. \\
&\quad \left. \left. + \alpha \cos(kx + \eta_0) \tanh(\alpha x + \xi_0)) \right. \right. \\
&\quad \left. \left. + \cos(kx + \eta_0) \tanh(\alpha x + \xi_0) (-k^2 P + QR^2 \tanh(\alpha x + \xi_0)^2) + \dots \right] \right] \tag{14}
\end{aligned}$$

$$\text{where } R = \sqrt{-2 \frac{P\alpha^2}{Q}}.$$

2.2 The Modified Laplace decomposition method (MLDM) algorithm of the NLSE

The methodology of the LDM is implemented to the NLSE itself (1), along with Wazwaz modification [20], [21] in which the zero components are split into two parts. According to it once we rewrite (1) in operator form, we proceed as follows:

$$\Psi(x,s) = \frac{1}{s} \Psi(x,0) + i \left(\frac{P}{s} L [L_{xx} \Psi] + \frac{Q}{s} L [N(\Psi)] \right) \tag{15}$$

Where the nonlinear operator $N(\Psi) = \Psi |\Psi|^2$ is decomposed using Adomian polynomials [38] into infinite series:

$$N(\Psi) = \sum_{n=0}^{\infty} A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} \left[N \left(\sum_{i=0}^{\infty} \lambda^i \Psi_i \right) \right] \tag{16}$$

Here we may view the few first Adomian polynomials as follows:

$$\begin{aligned}
A_0 &= \Psi_0 \bar{\Psi}_0 \\
A_1 &= \Psi_1 \bar{\Psi}_0 + \Psi_0 \bar{\Psi}_1 \\
A_2 &= \Psi_2 \bar{\Psi}_0 + \Psi_1 \bar{\Psi}_1 + \Psi_0 \bar{\Psi}_2 \\
A_3 &= \Psi_3 \bar{\Psi}_0 + \Psi_2 \bar{\Psi}_1 + \Psi_1 \bar{\Psi}_2 + \Psi_0 \bar{\Psi}_3 \\
A_4 &= \Psi_4 \bar{\Psi}_0 + \Psi_3 \bar{\Psi}_1 + \Psi_2 \bar{\Psi}_2 + \Psi_1 \bar{\Psi}_3 + \Psi_0 \bar{\Psi}_4 \\
&\vdots
\end{aligned} \tag{17}$$

Now, using (15), (16) and that the decomposition of the series $\Psi(x, t) = \sum_{n=0}^{\infty} \Psi_n(x, t)$, we obtain:

$$\sum_{n=0}^{\infty} \Psi_n(x, s) = \frac{1}{s} \Psi(x, 0) + i \left(\frac{P}{s} L \left[L_{xx} \sum_{n=0}^{\infty} \Psi_n \right] + \frac{Q}{s} L \left[\sum_{n=0}^{\infty} A_n \right] \right) \tag{18}$$

Identifying the recursive relation by comparing both sides of (18), then applying the inverse Laplace transform with its properties and using the given initial conditions we get:

$$\begin{aligned}
\Psi_0(x, t) &= F_1(x, t) \\
\Psi_1(x, t) &= F_2(x, t) + iL^{-1} \left(\frac{P}{s} L [L_{xx} \Psi_0] + \frac{Q}{s} L [A_0] \right) \\
\Psi_{n+1}(x, t) &= iL^{-1} \left(\frac{P}{s} L [L_{xx} \Psi_n] + \frac{Q}{s} L [A_n] \right), n \geq 1
\end{aligned} \tag{19}$$

The approximation is successfully obtained as the truncated series decomposition is given by:

$$\begin{aligned}
\Psi(x, t) &= \Psi_0(x, t) + \Psi_1(x, t) + \dots \\
&= iR \sin(kx + \eta_0) \tanh(\alpha x + \xi_0) + R \left(2Pt\alpha \operatorname{sech}(\alpha x + \xi_0)^2 (-k \cos(kx + \eta_0) + \right. \\
&\quad \left. \alpha \sin(kx + \eta_0) \tanh(\alpha x + \xi_0)) \right. \\
&\quad \left. + \tanh(\alpha x + \xi_0) (\cos(kx + \eta_0) + k^2 Pt \sin(kx + \eta_0) \right. \\
&\quad \left. - QR^2 t \sin(kx + \eta_0)^3 \tanh(\alpha x + \xi_0)^2) \right) + \dots
\end{aligned} \tag{20}$$

$$\text{where } R = \sqrt{-2 \frac{P\alpha^2}{Q}}.$$

3. Numerical results and discussion

In the present numerical computations and for the numerical study purposes, we will use the 2-term approximation (14), (20), due to the massive components of the series solution. We have assumed the involved parameters are given by $k = P = 1$, $Q = -1$, $\alpha = \sqrt{2}$, $\xi_0 = \eta_0 = 2$, the interval of spatial coordinate x is $[-20, 20]$ and maximum value of time is taken as $t = 0.1$ sec.

3.1 LDM results

The module of the exact solution $|\Psi(x, t)|$ and the corresponding module of the numerical solution $|\Psi_{LDM}(x, t)|$ with the help of two-term approximations of the decomposition series solution are shown in **Figure 1**. Although we have used a low-order approximation which is led to high accuracy without loss of generality, this is totally achieved in **Table 1** which exhibits the absolute errors $|\Psi(x, t) - \Psi_{LDM}(x, t)|$ in constructions of the approximated $\Psi_{LDM}(x, t)$.




Figure 1. The plot of surface: (a) Exact module $|\Psi(x, t)|$ of the equation (2). (b) Numerical module of LDM $|\Psi_{LDM}(x, t)|$ of the equation (14).

Table 1. The Error Module $|\Psi(x, t) - \Psi_{LDM}(x, t)|$

| x | t | | | |
|-----|--------------------------|--------------------------|-------------------------|----------|
| | 0.0001 | 0.001 | 0.01 | 0.1 |
| -20 | 1.5987×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| -15 | 1.5543×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| -10 | 1.5321×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| -5 | 2.1538×10^{-14} | 1.6266×10^{-10} | 1.5691×10^{-6} | 0.015573 |
| 0 | 3.0733×10^{-12} | 2.9236×10^{-9} | 1.4213×10^{-6} | 0.01396 |
| 5 | 1.5321×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| 10 | 1.5765×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| 15 | 1.5765×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |
| 20 | 1.5543×10^{-14} | 1.5625×10^{-10} | 1.5625×10^{-6} | 0.015564 |

The calculated errors in **Table 1** indicate a very good approximation with the actual solution by using two terms only and the error grows higher as the time value increases.

Figure 12 (a) illustrates the 3 dimensional absolute error module for values of time $[0, 0.1]$ which its peak appears at $t = 0.1$. Whereas, **Figure 12(b)**, focuses on the peak of the surface where the absolute error module is seen at $t = 0.1$ sec where it has been magnified when $x \in [-5, 5]$. The exact solution $|\Psi(x, t)|$ of the equation (1), the numerical solution module $|\Psi_{LDM}(x, t)|$ of the equation (14) and the absolute error of the module $|\Psi(x, t) - \Psi_{LDM}(x, t)|$ are compiled in **Table 2**. The results we obtain show an acceptable agreement between the exact solution and the approximate solution, even at time where the peak of the error surface appears.




Figure 2. The plots of : (a) The error module $|\Psi(x,t) - \Psi_{LDM}(x,t)|$. (b) The peak of the Error Module Curve when $t = 0.1$.

Table 2. The numerical results of the exact module (1), approximated module (14) and the module error .

| x | $ \Psi(x,t) $ | $ \Psi_{LDM}(x,t) $ | $ \Psi(x,t) - \Psi_{LDM}(x,t) $ |
|-----|---------------|---------------------|---------------------------------|
| -5 | 1.9999 | 2.0155 | 0.015573 |
| -4 | 1.9985 | 2.0142 | 0.015705 |
| -3 | 1.9746 | 1.9925 | 0.017932 |
| -2 | 1.609 | 1.6589 | 0.049846 |
| -1 | 0.58801 | 0.55563 | 0.032376 |
| 0 | 1.875 | 1.889 | 0.01396 |
| 1 | 1.9924 | 2.0079 | 0.015487 |
| 2 | 1.9995 | 2.0151 | 0.01556 |
| 3 | 2. | 2.0155 | 0.015564 |
| 4 | 2. | 2.0156 | 0.015564 |
| 5 | 2. | 2.0156 | 0.015564 |

3.2 Modified LDM results

In this subsection the achieved approximations using modified LDM will be discussed. The interpretation of **Figure 3** indicates to the accuracy of modified LDM decreases considerably as the time interval extends which is certainly due to the complexity of the split and the massive components of the solution series (20) and the data results in **Table 3** of the absolute error $|\Psi(x,t) - \Psi_{MLDM}(x,t)|$ module prove it.




Figure 3. The plot of surface of: (a) Exact module $|\Psi(x,t)|$ of the equation (2). (b) Numerical module of MLDM $|\Psi_{MLDM}(x,t)|$ of the equation (20).

Table 3. The Error Module $|\Psi(x, t) - \Psi_{MLDM}(x, t)|$

| x | t | | | |
|-----|-------------------------|-------------------------|-------------------------|-----------|
| | 0.0001 | 0.001 | 0.01 | 0.1 |
| -20 | 7.9802×10^{-8} | 7.9869×10^{-6} | 0.00080531 | 0.085765 |
| -15 | 5.6377×10^{-8} | 5.6468×10^{-6} | 0.00057374 | 0.066362 |
| -10 | 6.6538×10^{-8} | 6.6335×10^{-6} | 0.00064101 | 0.020131 |
| -5 | 1.6119×10^{-8} | 1.6089×10^{-6} | 0.00015788 | 0.012966 |
| 0 | 3.9583×10^{-9} | 3.67×10^{-7} | 8.8355×10^{-6} | 0.017336 |
| 5 | 8.6222×10^{-8} | 8.6189×10^{-6} | 0.00085836 | 0.080951 |
| 10 | 7.5549×10^{-8} | 7.5638×10^{-6} | 0.00076522 | 0.084397 |
| 15 | 3.4809×10^{-8} | 3.4498×10^{-6} | 0.00031235 | 0.017082 |
| 20 | 1.0024×10^{-8} | 1.0022×10^{-6} | 0.00010004 | 0.0097977 |

At the end, as a completion of our numerical study a representation of the peak of the error surface at time $t = 0.1$ as a two dimensional graph where the approximation and the exact solution meet and diverse is presented in **Figure 4(a)**. Moreover, the graphical and the tabulated data of the comparison between the exact solution and the approximate MLDM solution with its absolute error in **Figure 4(b)** and **Table 4** reveals an acceptable agreement of the series solution.




Figure 4. The plots of : (a) The error module $|\Psi(x, t) - \Psi_{MLDM}(x, t)|$. (b) The peak of the Error Module Curve when $t = 0.1$.

Table 4. The numerical results of the exact module (1), approximated module (20) and the module error .

| x | $ \Psi(x, t) $ | $ \Psi_{MLDM}(x, t) $ | $ \Psi(x, t) - \Psi_{MLDM}(x, t) $ |
|-----|----------------|-----------------------|------------------------------------|
| -5 | 1.9999 | 2.0129 | 0.012966 |
| -4 | 1.9985 | 2.0526 | 0.05414 |
| -3 | 1.9746 | 2.0165 | 0.04196 |
| -2 | 1.609 | 1.7139 | 0.10489 |
| -1 | 0.58801 | 0.79691 | 0.2089 |
| 0 | 1.875 | 1.8577 | 0.017336 |
| 1 | 1.9924 | 2.0136 | 0.021197 |
| 2 | 1.9995 | 2.0853 | 0.085782 |
| 3 | 2. | 1.9495 | 0.050521 |
| 4 | 2. | 2.04 | 0.039992 |
| 5 | 2. | 2.081 | 0.080951 |

Remark:

Obviously, any good numerical schemes should have satisfactory long time numerical behavior which is mostly accomplished by increasing the number of iterations which may be costly in time or try different split in the modified LDM. Despite some studies (see for example [24], [25]) have proposed different splits, more components of the decomposition series have to be calculated.

3.3 Comparison of the numerical results for the NLSE IVP

In this subsection, a comparison between the numerical approximate solutions obtained using the LDM and the MLDM with the exact optic soliton solution (2) throughout their related absolute errors in **Table 5** demonstrates the efficiency and the accuracy of the considered approaches within only a few terms of the series solution. Additionally, the observed absolute errors prove that the LDM has the advantage of being stabilized accurate approach over the MLDM which is also shown via illustrated three dimensional error module graph **Figure 2 (a)**, **Figure 4(a)** at $t = 0.1$ when the peak of the error appears.

Table 5. Numerical comparison of the absolute errors obtained by the Laplace decomposition method

| $\Psi_{LDM}(x, t)$ and the modified Laplace decomposition method $\Psi_{MLDM}(x, t)$. | | | | | | | |
|--|-----|--------------------------|-------------------------|------|-----|-------------------------|-------------------------|
| t | x | $ \Psi - \Psi_{LDM} $ | $ \Psi - \Psi_{MLDM} $ | t | x | $ \Psi - \Psi_{LDM} $ | $ \Psi - \Psi_{MLDM} $ |
| 0.0001 | -20 | 1.5987×10^{-14} | 7.9802×10^{-8} | 0.01 | -20 | 1.5625×10^{-6} | 0.00080531 |
| | -15 | 1.5543×10^{-14} | 5.6377×10^{-8} | | -15 | 1.5625×10^{-6} | 0.00057374 |
| | -10 | 1.5321×10^{-14} | 6.6538×10^{-8} | | -10 | 1.5625×10^{-6} | 0.00064101 |
| | -5 | 2.1538×10^{-14} | 1.6119×10^{-8} | | -5 | 1.5691×10^{-6} | 0.00015788 |
| | 0 | 3.0733×10^{-12} | 3.9583×10^{-9} | | 0 | 1.4213×10^{-6} | 8.8355×10^{-6} |
| | 5 | 1.5321×10^{-14} | 8.6222×10^{-8} | | 5 | 1.5625×10^{-6} | 0.00085836 |
| | 10 | 1.5765×10^{-14} | 7.5549×10^{-8} | | 10 | 1.5625×10^{-6} | 0.00076522 |
| | 15 | 1.5765×10^{-14} | 3.4809×10^{-8} | | 15 | 1.5625×10^{-6} | 0.00031235 |
| | 20 | 1.5543×10^{-14} | 1.0024×10^{-8} | | 20 | 1.5625×10^{-6} | 0.00010004 |
| 0.001 | -20 | 1.5625×10^{-10} | 7.9869×10^{-6} | 0.1 | -20 | 0.015564 | 0.085765 |
| | -15 | 1.5625×10^{-10} | 5.6468×10^{-6} | | -15 | 0.015564 | 0.066362 |
| | -10 | 1.5625×10^{-10} | 6.6335×10^{-6} | | -10 | 0.015564 | 0.020131 |
| | -5 | 1.6266×10^{-10} | 1.6089×10^{-6} | | -5 | 0.015573 | 0.012966 |
| | 0 | 2.9236×10^{-9} | 3.67×10^{-7} | | 0 | 0.01396 | 0.017336 |
| | 5 | 1.5625×10^{-10} | 8.6189×10^{-6} | | 5 | 0.015564 | 0.080951 |
| | 10 | 1.5625×10^{-10} | 7.5638×10^{-6} | | 10 | 0.015564 | 0.084397 |
| | 15 | 1.5625×10^{-10} | 3.4498×10^{-6} | | 15 | 0.015564 | 0.017082 |
| | 20 | 1.5625×10^{-10} | 1.0022×10^{-6} | | 20 | 0.015564 | 0.0097977 |

4. Conclusion

In this work, the LDM and modified version of it, namely the MLDM, have been successfully implemented to approximate an optic soliton solution of the nonlinear complex Schrödinger equation (NLSE) with an initial value problem (IVP). A transformation has been presented so that a system of coupled real partial differential equations is obtained and to be numerically solved in order to approximate the NLSE solution. In spite of, some studies [26], [28] in which the Laplace transform has been applied directly to the equation of interest. On the other hand, based on Wazwaz's modification [20] the solution of the NLSE is examined. The obtained results are investigated via illustrations and tables. Therefore, it is predictable, that the LDM and the MLDM are effective techniques to investigate numerical solutions of nonlinear complex problems. The LDM has the advantage of being a stabilized accurate method over the MLDM. Additionally, the considered methods are converging very rapidly with fewer terms of the series solution.

5. References

- [1] A. Hasegawa and Y. Kodama, *Solitons in optical communications / Akira Hasegawa and Yuji Kodama*. Oxford University Press, New York, 1995.
- [2] W. X. Ma and M. Chen, "Direct search for exact solutions to the nonlinear Schrödinger equation," *Appl. Math. Comput.*, vol. 215, no. 8, pp. 2835–2842, 2009.
- [3] K. M. Abu-alnaja, "Numerical Study of a Class of Nonlinear Partial Differential Equations," University of Umm Al-Gurra, Kingdom of Saudi Arabia, 2009.
- [4] G. Adomian, "Solution of physical problems by decomposition," *Comput. Math. with Appl.*, vol. 27, no. 9–10, pp. 145–154, 1994.
- [5] G. Adomian and R. Rach, "Linear and Nonlinear Schrodinger equations," *Found. Phys.*, vol. 21, pp. 983–991, 1991.
- [6] Ö. F. Gözükızıll and H. Gündoğdu, "Applications of the decomposition methods to some nonlinear partial differential equations," *New Trends Math. Sci.*, vol. 3, no. 6, pp. 57–66, Jul. 2018.
- [7] S. Sirisubtawee and S. Kaewta, "New Modified Adomian Decomposition Recursion Schemes for Solving Certain Types of Nonlinear Fractional Two-Point Boundary Value Problems," *Int. J. Math. Math. Sci.*, vol. 2017, pp. 1–20, 2017.
- [8] D. Ziane, D. Baleanu, K. Belghaba, and M. Hamdi Cherif, "Local fractional Sumudu decomposition method for linear partial differential equations with local fractional derivative," *J. King Saud Univ. - Sci.*, vol. 31, no. 1, pp. 83–88, Jan. 2019.
- [9] Y. Lin, K. H. Chang, and C. K. Chen, "Modified Laplace Adomian decomposition method for analysing the entropy generation of Sakiadis flow with radiation effect," *Int. J. Exergy*, vol. 24, no. 1, p. 21, 2017.
- [10] G. Adomian, "A review of the decomposition method and some recent results for nonlinear equations," *Comput. Math. with Appl.*, vol. 21, no. 5, pp. 101–127, 1991.
- [11] S. A. Khuri, "A Laplace decomposition algorithm applied to a class of nonlinear differential equations," *J. Appl. Math.*, vol. 1, no. 4, pp. 141–155, 2001.
- [12] S. A. Kuri, "A new approach to Bartu's problem," *Appl. Math. Comput.*, vol. 14, pp. 131–136, 2004.
- [13] H. M. Yousef and A. I. B. M. Md Ismail, "Application of the Laplace Adomian decomposition method for solution system of delay differential equations with initial value problem," in *AIP Conference Proceedings*, 2018, vol. 1974.
- [14] A. Hamoud and K. Ghadle, "The Approximate Solutions of Fractional Volterra-Fredholm Integro-Differential Equations by using Analytical Techniques," *Issues Anal.*, vol. 25, no. 1, pp. 41–58, 2018.
- [15] A. A. Hamoud and K. P. Ghadle, "Modified laplace decomposition method for fractional volterra-fredholm integro-differential equations," *J. Math. Model.*, vol. 6, no. 1, 2018.
- [16] A. K. Jaradat, A. A. Obeidat, M. A. Gharaibeh, and M. K. H. Qaseer, "Adomian Decomposition Approach to Solve the Simple Harmonic Quantum Oscillator," *Int. J. Appl. Eng. Res.*, vol. 13, pp. 1056–1059, 2018.
- [17] R. Shah, H. Khan, M. Arif, and P. Kumam, "Application of Laplace-Adomian decomposition method for the analytical solution of third-order dispersive fractional

partial differential equations,” *Entropy*, vol. 21, no. 4, 2019.

- [18] T. A. Abassy, “Improved Adomian decomposition method,” *Comput. Math. with Appl.*, vol. 59, no. 1, pp. 42–54, 2010.
- [19] T. A. Abassy, “Improved Adomian decomposition method (solving nonlinear non-homogenous initial value problem),” *J. Franklin Inst.*, vol. 348, no. 6, pp. 1035–1051, 2011.
- [20] A. M. Wazwaz, “A reliable modification of Adomian decomposition method,” *Appl. Math. Comput.*, 1999.
- [21] A. M. Wazwaz, “The modified decomposition method and Padé approximants for solving the Thomas-Fermi equation,” *Appl. Math. Comput.*, 1999.
- [22] A. M. Wazwaz and S. M. El-Sayed, “A new modification of the Adomian decomposition method for linear and nonlinear operators,” *Appl. Math. Comput.*, vol. 122, no. 3, pp. 393–405, 2001.
- [23] A. M. Wazwaz, “A reliable Technique for solving linear and nonlinear Schrodinger equations by the decompostion method,” *Bull. Inst. Math. Acad.*, vol. 29, pp. 125–134, 2001.
- [24] S. S. Ray, “Solution of the Coupled Klein-Gordon Schrödinger Equation Using the Modified Decomposition Method,” *International Journal of Nonlinear Science*, vol. 4, no. 3, pp. 227–234, 2007.
- [25] R. D. Pankaj, “Laplace Modified Decompostion method to study solitary wave solution of Coupled nonlinear Klien-Gorden Schrodinger equation,” *Int. J. Stat. Math.*, vol. 5, no. 1, pp. 1–5, 2013.
- [26] O. Gonzalaz-Gaxiola, “The Laplace-Adomian Decompostion Method Applied to the Kundu-Eckhaus Equation,” *Int. J. Math. its Appl.*, vol. 5, no. 1-A, p. 2017, 2017.
- [27] N. Manjak, O. O, and E. SULEIMAN, “The Single Soliton Solution of The Nonlinear Schrodinger Equation by Modified Adomian Decomposition Method (ADM),” *Sci. Forum (Journal Pure Appl. Sci.)*, vol. 17, no. 1, p. 41, 2019.
- [28] E. K. Jaradat, O. Alomari, M. Abudayah, and A. M. Al-Faqih, “An Approximate Analytical Solution of the Nonlinear Schrödinger Equation with Harmonic Oscillator Using Homotopy Perturbation Method and Laplace-Adomian Decomposition Method,” *Adv. Math. Phys.*, vol. 2018, pp. 1–11, Dec. 2018.
- [29] O. Gonzalez-Gaxiola, P. Franco, and R. Bernal-Jaquez, “Solution of the Nonlinear Schrodinger Equation with Defocusing Strength Nonlinearities Through the Laplace-Adomian Decomposition Method,” Jan. 2018.
- [30] R. Khalil, M. Al Horani, A. Yousef, and M. Sababheh, “A new definition of fractional derivative,” *J. Comput. Appl. Math.*, vol. 264, pp. 65–70, 2014.
- [31] H. X. Zhu, H. L. An, and Y. Chen, “A laplace decomposition method for nonlinear partial differential equations with nonlinear term of any order,” *Commun. Theor. Phys.*, vol. 61, no. 1, pp. 23–31, 2014.
- [32] M. Khan, M. Hussain, H. Jafari, and Y. Khan, “Application of Laplace Decomposition Method to Solve Nonlinear Coupled Partial Differential Equations,” *Appl. Sci.*, vol. 9, pp. 13–19, 2010.
- [33] A. S. R. Kumar and R. Rangarajan, “Applicability of laplace decompostion method for

solving certain differential difference equation of order 1,2," *Bull. Int. Math. Virtual Inst.*, vol. 3, pp. 103–111, 2013.

- [34] A. Kumar and P. D. Pankaj, "Solitary Wave Solutions of Schrodinger equation by Laplace Adomian decomposition method," *Phys. Rev. Res. Int.*, vol. 3, no. 4, pp. 702–7012, 2013.
- [35] K. Abbaoui and Y. Cherruault, "Convergence of Adomian's method applied to differential equations," *Comput. Math. with Appl.*, vol. 28, no. 5, pp. 103–109, 1994.
- [36] R. Z. Ouedraogo, Y. Cherruault, and K. Abbaoui, "Convergence of Adomian's method applied to algebraic equations," *Kybernetes*, vol. 29, no. 9/10, pp. 1298–1305, 2002.
- [37] H. N. Hassan and M. A. El-Tawil, "Solving cubic and coupled nonlinear Schrödinger equations using the homotopy analysis method," *Int. J. Appl. Math. Mech.*, vol. 7, no. 8, pp. 41–64, 2011.
- [38] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*. 2013.