

Short term outcome of extracorporeal and Intracorporeal anastomosis in laparoscopic colorectal surgeries – A prospective observational study.

Abstract

Aim of study

Laparoscopic colorectal surgery has gained popularity around the Globe. Laparoscopic colectomy significantly improves the short term and long term outcomes of patients such as lower pain scores, less estimated blood loss and shorter hospital stay and reduced rate of incisional hernia. The bowel anastomosis after laparoscopic resection of the tumour can be done in two ways - extracorporeal anastomosis and Intracorporeal anastomosis. Our study observed and evaluated the data of the patients who underwent these two techniques.

Materials and Methods

A prospective observational study was conducted in the department of General Surgery, SMHS hospital of Government Medical College, Srinagar. A total of 32 patients were studied out of which 20 patients had undergone intracorporeal anastomosis and 12 patients had undergone extracorporeal anastomosis. The aims of the study were to assess the operative time, post-operative ileus, length of hospital stay, anastomotic leak and other anastomotic complications, wound infections and extraction site hernias.

Results

The patients in our study were in the range of 30-85 years with a mean age of 59.18 ± 14.92 . and 59.4% of patients were males and the rest 40.6% were females, with male/female ratio of 1.46:1. There was no significant difference in mean operative time between the intracorporeal group and extracorporeal group (188 ± 9.78 minutes vs. 180.3 ± 13.8 minutes). The patients in the intracorporeal group had earlier return of bowel function than extracorporeal group as reflected by earlier appreciation of flatus (median of 2.0 days vs. 3.0 days) and tolerance to orals (median of 3 days vs. 4 days). This had led to the shorter hospital stay of the intracorporeal group than the extracorporeal group of patients (median of 5 days vs. 6.5 days) The wound infection rate was 6.3 percent in our study, 5.0% in the intracorporeal group and 8.3% in the extracorporeal group. Only 1 out of the total 32 patients (3.1%) developed mesenteric twist which belonged to the extracorporeal group (1 out of 12 patients). We did not observe any leak in both the groups of patients in the study. No patient in our study in either group developed extraction site or incisional hernia.

Conclusion

36 Intracorporeal anastomosis in laparoscopic colorectal surgeries leads to earlier return of bowel
37 function, earlier resumption of orals and shorter hospital stay than the extracorporeal
38 anastomosis. There does not exist a significant difference between the two modes of anastomosis
39 in terms of anastomotic and wound related complications.

40 **Keywords;** Laparoscopy, Colonrectum, Anastomosis, Intracorporeal, Extracorporeal

41

42 Introduction

43 Colon cancer is the most common type of gastro intestinal cancer ^[1].Chemotherapy and
44 radiotherapy can improve survival in colorectal cancer patients. However, the only treatment
45 with curative intent is surgical resection of the tumor. Colectomy for cancer can be performed
46 using either open or laparoscopic approach. Laparoscopic colorectal resection has gone through a
47 major evolution since 1991, when the first reports of colorectal resections had been published ^{[2,}
48 ^{3, 4]}.In 1991, Jacobs ^[5] performed the first laparoscopic right hemicolectomy. After right
49 hemicolectomy the ileocolic anastomosis cannot be fashioned “in a natural way” as it normally
50 happens following left hemicolectomy or anterior resection of the rectum; for this reason,
51 different kinds of laparoscopic right colectomy have been proposed ^[6]. The ileocolic anastomosis
52 can be performed using different techniques and devices depending on the intracorporeal and
53 extracorporeal approach.This anastomosis can also be manually performed; despite the lower
54 cost, this requires a greater ability ^[7,8].Since the publication of the first laparoscopic colectomy
55 its use has been increased.^[9] Moreover, randomized trials have demonstrated that laparoscopic
56 surgery for colon cancer achieves good short-term and oncologic outcomes similar to those
57 found in open surgery. ^[10,11] However, laparoscopic surgery for transverse and descending colon
58 cancer requires an advanced technique. Hence, only recently, studies have demonstrated the
59 feasibility and safety of the laparoscopic resection for lesions located in the distal transverse and
60 descending colon. ^[12,13]. Incisional hernias after open surgery occur in 12 to 20% and may lead to
61 significant morbidity. Midline extraction sites have a higher chance of hernias than non-
62 midline. ^[14] Laparoscopic surgery for rectal cancer has been considered technically more
63 demanding when compared with that for colon cancer. However, laparoscopic total mesorectal
64 excision (TME) has been positively employed for the treatment of rectal cancer in Japanese
65 Centres without lateral lymph node metastasis ^[15] or invasion to the adjacent organ, since it has
66 the advantage of providing a good view even in a narrow pelvis and allowing to perform more
67 precise autonomic nerve preservation

68 Rectal transection and anastomosis at the lower rectum is the most challenging part of
69 laparoscopic low anterior resection. Therefore, some have demonstrated that rectal transection
70 should be performed using instruments for open surgery with small laparotomy. In our institute,
71 however, rectal transection using a currently available endo-stapler followed by anastomosis
72 with a double stapling technique is usually performed.However, anastomotic leakage is still a
73 serious problem after sphincter-saving surgery for rectal cancer. ^[16] Diverting stomata are used to

74 reduce leakage-related complications after LAR, but the routine use of diverting stomata is
75 controversial because of reported morbidity associated with their creation and closure.^[17,18]

76 Aims and objectives

77 The aim of this study was to assess the short-term outcome of extracorporeal anastomosis and
78 intracorporeal anastomosis in laparoscopic colorectal surgeries in terms of; operative time,
79 anastomotic leak rates and other complications of anastomosis, Post-operative ileus, Length of
80 hospital stay, Wound infections. Extraction and port site hernias.

81 Material and methods:

82 This prospective observational study was conducted in the department of General and minimal
83 access Surgery, from 2015 to 2018 after obtaining the clearance from the Institutional Ethical
84 Committee. A total of 32 cases were enrolled in the study. This comprised of patients admitted
85 for elective surgery for right colon growth, transverse colon growth, left colon growth, sigmoid
86 colon growth and rectal growth above peritoneal reflection in various surgical wards of hospital.
87 The patients that are included in the study are;

- 88 1. Age >18 years.
- 89 2. Patients who are eligible for curative resection of cancer by means of hemicolectomy.
- 90 3. In case of polyp, a colonoscopic biopsy proven invasive cancer.
- 91 4. For rectosigmoid, patient can be included if the tumour lies above the peritoneal
92 reflection.
- 93 5. Solitary colon carcinoma observed at colonoscopy or barium study.

94 The patients that are excluded from the study are;

- 95 1. Contraindications to general anaesthesia e.g. congestive heart failure, chronic renal
96 failure, chronic obstructive lung disease, un-correctable coagulopathy
- 97 2. General contradictions to laparoscopic surgery.
- 98 3. Metastases in the liver or lungs or pre operative evidence of involvement of adjacent
99 structures as detected by CT, MRI or USG.
- 100 4. Acute intestinal obstruction.
- 101 5. Patients who had conversion to open procedure will be excluded from the analysis.

102 These patients were initially evaluated in the outpatient department (OPD) and then planned for
103 surgery. On admission, a detailed history of the patient was recorded including the presenting
104 complaints, duration of the complaints, past history especially with reference to previous
105 surgery, family history and any other associated condition such as chronic ailment and any drug
106 intake.

107 General physical examination was done with particular consideration of build, height and weight
108 followed by systemic examination. Thorough abdominal examination was done in each patient.
109 The patients were taken for laparoscopic colorectal surgery after proper clinical evaluation and
110 after diagnosing them with the disease on colonoscopy and after confirming malignancy on
111 colonoscopic biopsy. Each patient and his attendants were fully explained about the nature of the
112 procedure and the possible complications inherent to the procedure in the native language and
113 thereafter a written consent was sought from the patient prior to surgery. All base line

114 investigations were performed which includes (Complete blood count, Coagulogram, Liver
115 function test, Kidney function test, Blood sugars, Serum electrolytes (Na⁺/K⁺).Chest X-ray and
116 abdominal USG were also performed to investigate lung and liver metastasis respectively. Pre-
117 operative CECT abdomen was done in all patients as a pre-operative staging. The Serum
118 Carcinoembryonic antigen (CEA) also done in all patients.

119 Pre-operative preparation

120 After completing the routine and specific investigations, patients were assessed for anesthetic
121 fitness to undergo the laparoscopic surgery. All the patients were kept fasting 12 hours before
122 surgery and a proper bowel preparation was done using oral solution of polyethylene glycol.
123 Serum electrolytes were repeated before surgery and necessary corrections were made.
124 Ceftriaxone 1 gm I.V as surgical prophylaxis was given to every patient before surgery.
125 Thromboembolic prophylaxis was only given to high risk patients. Patients were allotted to
126 either extracorporeal or intracorporeal limb based on CT scan and intraoperative findings..

127 Operative technique:

128 The experience gained in basic laparoscopy, like technique of creation of pneumoperitoneum,
129 trocar insertion and dissection techniques forms the foundation for advanced laparoscopic
130 colorectal surgery. Pneumoperitoneum is created either via the percutaneous insertion of a
131 Verres needle or with the open Hassan technique. In LRHC, we used four ports: a 10 mm to 12
132 mm camera port for a 30° laparoscope positioned at the level of umbilicus on left side of
133 abdomen, one 12-mm working port for stapling devices in the left upper abdomen above the
134 camera port and 2 five-mm working ports, one located in the left lower abdomen below the
135 camera port and another in the right lower abdomen. The 5 mm port located in the right lower
136 abdomen is later extended for extraction of the specimen. In LLHC, four port technique was also
137 followed but port positions were mirror images of LRHC. Four ports were also used in cases of
138 sigmoidectomy, anterior resection and low anterior resection, with 10 mm umbilical port for
139 camera, 12 mm working port in right lower abdomen and two 5 mm ports – one to the left of
140 camera port and another in the left lower abdomen. The port located in the left lower abdomen is
141 later extended for extraction of the specimen.

142 Basic common steps in laparoscopic colorectal resection


143 All laparoscopic colon procedures have several steps that are common. These steps include:

- 144 1. Localization of the lesion / tumour.
- 145 2. Mobilization of the lesion (medial-to-lateral approach).
- 146 3. Vessel ligation for devascularization of the specimen.
- 147 4. Bowel division.
- 148 5. Restoring bowel continuity by Anastomosis (extracorporeal or intracorporeal).
- 149 6. Specimen retrieval and protection of the wound during retrieval.


150 Follow up:

151 After discharging from the hospital, the patients were advised to follow in OPD with the
152 histopathology of the resected specimen. Patients with stage 1 disease did not require adjuvant


153 chemotherapy and on further follow up such patients were advised serum CEA levels every
154 three months. Such patients were advised CT scan only if there was rise in serum CEA levels.
155 Patients with stage 2 disease and above were strictly advised to follow the medical oncology for
156 chemotherapy. Serum CEA levels were repeated every three months and an annual CT scan and
157 colonoscopy were advised to look for recurrence of the disease.



158
159 Fig.1 CECT showing Right colonic growth



160 Fig. 2 Mobilization of Right colon



161
162 Fig.3 and 4 Creation of extracorporeal and in tra corporeal anastomosis
163



164
165

Fig 5 specimen of colon with ileum

Fig.6 Port position in right hemicolectomy



166
167
168
169
170
171
172
173
174
175
176
177

Fig.7 port operative scar in right hemicolectomy

Fig.8 Port position in LAR

178 **Statistical Analysis**

179 The recorded data was compiled and entered on a spread sheet (Microsoft excel) and then
 180 exported to data editor of SPSS V.20.0 (SPSS Inc., Chicago, Iclinos, USA).Continuous variables
 181 were summarized in the form of means and standard deviations whereas categorical variables
 182 were summarized as median and percentage. Results were compared using Chi square tests and
 183 Mann-Whitney U-tests. P-values less than 0.05 were considered to be statistically significant.
 184 Graphically the data was presented by Bar Charts and Pie Charts.

185 **Data Analysis:**

186 The study included 32 patients out of which 20 patients had undergone intra-corporeal
 187 anastomosis and 12 patients had undergone extra-corporeal anastomosis after resection of
 188 tumour. The patients in our study were in the range of 30-85 years with a mean age of $59.18 \pm$
 189 14.92 . Maximum number of patients were in the age group of 41-60 years (43.75%) followed by
 190 patients older than 60 years (40.63%). Table 2 shows distribution of patients according to their
 191 gender. 59.4% of patients were male and the rest 40.6% were females with male/female ratio of
 192 1.46: 1. The demographic data is shown in table 1 and 2.

Table1: Age distribution of study patients

Age (years)	Count(Percentage)	Anastomosis		Total
		Intracorporeal	Extracorporeal	
<40	Count(Percentage)	4(12.50 %)	1(3.12 %)	5(15.62 %)
41-60	Count(Percentage)	9(28.12%)	5(15.63 %)	14(43.75 %)
>60	Count(Percentage)	7(21.88%)	6(18.75%)	13(40.63 %)
Total	Count(Percentage)	20(62.50%)	12(37.50 %)	32(100.0 %)

Mean±SD = 59.18 ± 14.92 ,

Table 2: Gender distribution of study patients.

Sex	Count(Percentage)	Anastomosis		Total
		Intracorporeal	Extracorporeal	
M	Count(Percentage)	13(40.62%)	6(18.75%)	19(59.37 %)
F	Count(Percentage)	7(27.88 %)	6(18.75 %)	13(18.75 %)
Total	Count(Percentage)	20(100.0%)	12(100.0%)	32(100.0%)

193 Table 3 shows operation time in the two groups of study patients with the mean operation time of
 194 188.1 ± 9.78 minutes in the intracorporeal group and 180.3 ± 13.8 minutes in the extracorporeal
 195 group. No significant difference in the operation time between the two groups of study
 196 population could be observed as reflected by the p-value of 0.07

Table: 3 Operation time (minutes) in two techniques of study patients.

Anastomosis	Number of Patients	Operation time Mean±SD	p-value
Intracorporeal	20	188.1 ± 9.78	0.075
Extracorporeal	12	180.3 ± 13.8	

197 Table 4 shows absolute and percentage distribution of patients in the two groups of study with
 198 respect to the appreciation of passage of flatus. Patients in the intracorporeal group had the

199 median of 2.0 days where as patients in the extracorporeal group had median of 3.0 days. The
 200 difference in the two groups is statistically significant as shown by p-value of 0.01

Table 4: Time distribution (days) of appreciation of passage of flatus in the two groups of study patients

Flatus	Anastomosis		Total	
	Intracorporeal	Extracorporeal		
1	Count(percentage)	1(5.0%)	1(8.3%)	2(6.3%)
2	Count(percentage)	12(60.0%)	1(8.3%)	13(40.6%)
3	Count(Percentage)	6(30.0%)	5(41.7%)	11(34.4%)
4	Count(Percentage)	1(5.0%)	4(33.3%)	5(15.6%)
5	Count(Percentage)	0(0.0%)	1(8.3%)	1(3.1%)
Total	Count(Percentage)	20(100.0%)	12(100.0%)	32(100.0%)
Median		2.0	3.0	-

p=0.01, Mann-Whitney test

201 Table 5 shows absolute and percentage distribution of patients in the two groups of study with
 202 respect to the first bowel movement. Patients in the intracorporeal group had the median of 4.0
 203 days where as patients in the extracorporeal group had median of 5.0 days. The difference in the
 204 two groups is statistically significant as shown by p-value of 0.01

Table 5: Time distribution (days) of bowel movement in the two groups of patients in study.

Stools	Anastomosis		Total	
	Intracorporeal	Extracorporeal		
2	Count(Percentage)	2(10.0%)	0(0.0%)	2(6.3%)
3	Count(Percentage)	2(10.0%)	1(8.3%)	3(9.4%)
4	Count(Percentage)	12(60.0%)	1(8.3%)	13(40.6%)
5	Count(Percentage)	2(10.0%)	7(58.3%)	9(28.1%)
6	Count(Percentage)	2(10.0%)	3(25.0%)	5(15.6%)
Total	Count(Percentage)	20(100.0%)	12(100.0%)	32(100.0%)
Median		4.0	5.0	-

p=0.01, Mann-Whitney test

205 Table 6 shows absolute and percentage distribution of the two groups of study population with
 206 respect to the day of tolerance of orals. In the intracorporeal group, 45% patients tolerated orals
 207 on day 2, another 45% patients on day 3 and 5% patients on day 4 and day 5 with the median of
 208 3 days. While, in the extracorporeal group 33.3% patients tolerated orals on day 3, 50% patients
 209 on day 4 and 16.7% patients on day 5 with the median of 4 days. There is a statistically
 210 significant difference between the two groups with respect to the day of oral tolerance as
 211 reflected by the p-value of 0.00.

Table 6: Time distribution (days) of oral tolerance in two techniques of anastomosis in study patients.

Orals		Anastomosis		Total
		Intracorporeal	Extracorporeal	
2	Count(Percentage	9(45.0%)	0(0.0%)	9(28.1%)
3	Count(Percentage	9(45.0%)	4(33.3%)	13(40.6%)
4	Count(Percentage	1(5.0%)	6(50.0%)	7(21.9%)
5	Count(Percentage	1(5.0%)	2(16.7%)	3(9.4%)
Total	Count(Percentage	20(100.0%)	12(100.0%)	32(100.0%)
Median		3.0	4.0	-

p=0.00, Mann-Whitney test

212 Table 7 lists different complications of laparoscopic colorectal surgeries reported in literature
213 and their incidence in our study. We did not observe any anastomotic leak or extraction site
214 hernia in our study. Anastomotic twist is described in Table 8 and wound infection in Table 9 in
215 detail.

Table 7: Rate of Complications in two groups of study patients

Complication		Intracorporeal	Extracorporeal	Total
Anastomotic leak	Count(Percentage	0(0%)	0(0%)	0(0%)
Anastomotic twist	Count(Percentage	0(0%)	1(8.3%)	1(3.12%)
Wound infection	Count(Percentage	1(5%)	1(8.3%)	2(6.3%)
Extraction site hernia	Count(Percentage	0(0%)	0(0%)	0(0%)

216 Table 8 shows absolute and percentage distribution of mesenteric twist at site Of anastomosis in
217 the study patients. Only 1 out of the total 32 patients (3.12%) developed mesenteric twist which
218 belonged to the extracorporeal group (1 out of 12 patients). No patient in the intracorporeal
219 group developed this complication. But the difference between the two groups was statistically
220 insignificant as reflected by the p-value of 0.37

Table 8: Rate of mesenteric twist in two techniques of anastomosis in study patients.

Anastomosis	Number of Patients	Mesentric twist	Percentage
Intracorporeal	20	0	0 %
Extracorporeal	12	1	8.3 %
Total	32	1	3.12 %

p>0.37, Chi-square test Exact p

221 Table 9 shows rate of wound infection in the two groups of study population. In the
222 intracorporeal group 1 out of 20 patients (5.0%) developed wound infection while as in the
223 extracorporeal group 1 out of 12 patients (8.3%) developed wound infection. The difference in
224 the rate of wound infection was statistically insignificant (p-value> 0.99). Overall, the wound
225 infection rate was 6.3 percent.

Table 9: Rate of wound infection in two groups of patients in study

Anastomosis	Number of Patients	Wound Infection	Percentage
Intracorporeal	20	1	5 %

Extracorporeal	12	1	8.3 %
Total	32	2	6.3 %

p>0.99, Chi-square test Exact p

226

227 Table 10 shows the absolute and percentage distribution of the patients in the two groups of the
 228 study patients with respect to the number of days of hospital stay. Most of the patients in the
 229 intracorporeal group (70%) stayed in the hospital for 5 – 6 days while as in the extracorporeal
 230 group majority of the patients (75%) stayed in the hospital for 6 – 7 days. Intracorporeal group
 231 had the median hospital stay of 5.0 days where as the extracorporeal group had the median
 232 hospital stay of 6.5 days with the p-value of 0.01 signifying statistically significant difference.

Table 10: length of hospital stay(days) in two groups of study patients

LOHS		Anastomosis		Total
		Intracorporeal	Extracorporeal	
4	Count(Percentage)	2(10.0%)	0(0.0%)	2(6.3%)
5	Count(Percentage)	9(45.0%)	1(8.3%)	10(31.3%)
6	Count(Percentage)	5(25.0%)	5(41.7%)	10(31.3%)
7	Count(Percentage)	3(15.0%)	4(33.3%)	7(21.9%)
8	Count(Percentage)	1(5.0%)	1(8.3%)	2(6.3%)
10	Count(Percentage)	0(0.0%)	1(8.3%)	1(3.1%)
Total	Count(Percentage)	20(100.0%)	12(100.0%)	32(100.0%)
Median		5.0	6.5	-

p=0.01, Mann-Whitney test

233

234 Discussion

235 Application of laparoscopy to colorectal surgery has produced some short- term benefits like
 236 faster return of gut function leading to more rapid discharge from the hospital and a faster return
 237 to normal activity as well as long term benefits like reduced rates of obstruction secondary to
 238 adhesions and reduced incidence of ventral hernia. The creation of bowel anastomosis after
 239 laparoscopic resection of large bowel tumour can be done in two ways – extracorporeal
 240 anastomosis and intracorporeal anastomosis. A number of studies have been carried worldwide
 241 to document the equalities and differences between the two modes of creation of bowel
 242 anastomosis with respect to the short- term outcomes like – operation time, post-operative return
 243 of bowel function, complications of anastomosis, wound infection and length of hospital stay.
 244 So, we also have conducted a study to evaluate these short-term outcomes of extracorporeal and
 245 intracorporeal anastomosis in laparoscopic colorectal surgeries. It was a hospital based
 246 prospective observational study conducted in the department of General and minimal access
 247 Surgery of the hospital of Government Medical College.

248 A total of 32 patients were studied out of which 20 patients had undergone IC anastomosis and
 249 12 patients had undergone EC anastomosis. The results were based on age, gender, operative
 250 time, post-operative return of bowel function in terms of appreciation of flatus, oral tolerance and
 251 first bowel movement, complications such as anastomotic leak, mesenteric twist, wound

252 infections and extraction site hernias and length of hospital stay. The patients in our study were
253 in the range of 30-85 years with a mean age of 59.18 ± 14.92 . Maximum number of patients
254 were in the age group of 41-60 years (43.8%) followed by patients older than 60
255 years(40.6%).Jorge Arredondo Chaves, Carlos Pastor Idoate et al^[19] have reported in their study
256 mean age group of patients as 62.6 ± 13.4 years in the intracorporeal group and 58.9 ± 12.9 years
257 in the extracorporeal group which closely matched with our study.In our study, 59.4% of patients
258 were male and the rest 40.6% were females with male/female ratio of 1.46:1. Jorge Arredondo
259 Chaves, Carlos Pastor Idoate et al have reported male/female ratio of 1.22:1 and Tu Jian-Cheng,
260 BSc, Wang Shu-Sheng, BSc et al^[20] reported male/female ratio of 1.57:1. Both of these studies
261 had comparable gender ratio with our study.

262 In our study there was no significant difference in operative time between two groups.The mean
263 operation time in the intracorporeal group was 188 ± 9.78 minutes and 180.3 ± 13.8 minutes in
264 the extra corporeal group. Comparable results were obtained by Minia Hellen, Casandra
265 Anderson et alwho reported the mean operative time of 190 minutes in the intracorporeal group
266 and 180 minutes in the extracorporeal group.Anania G, Santini M et al^[21] also showed similar
267 results with the mean operative time of 186.8 minutes in the intracorporeal group and 184.8
268 minutes in the extracorporeal group.

269 The appreciation of flatus in the intracorporeal group had the median of 2.0 days where as
270 patients in the extracorporeal group had median of 3.0 days. The difference in the two groups is
271 statistically significant as shown by p-value of 0.007. Comparable results were obtained by
272 Jayleen Grams, Winnie Tong et al^[22],Anania G, Santini M et al who reported mean days of
273 appreciation of flatus in the intracorporeal group as 2.0 days and 2.4 days in the extracorporeal
274 group.Tu Jian-Cheng, BSc, Wang Shu-Sheng, BSc et al reported the mean duration of
275 appreciation of flatus as 2.57 ± 0.08 days in the intracorporeal group and 3.10 ± 0.11 days in the
276 extracorporeal group which are comparable to our study.

277 The first bowel movement in the intracorporeal group had the median of 4.0 days where as
278 patients in the extracorporeal group had median of 5.0 days. The difference in the two groups is
279 statistically significant as shown by p-value of 0.005.Jorge Arredondo Chaves, Carlos Pastor
280 Idoate et al in 2011 reported the median days of the first bowel movement as 3 days in the
281 intracorporeal group 4 days in the extracorporeal group. Anania G Santini Met al in
282 2012reported mean of 3.8 days in the intracorporeal group and 4.9 days in the extracorporeal
283 group for the first bowel movement. In our study patients in the intracorporeal group had the
284 median of 4.0 days where as patients in the extracorporeal group had median of 5.0 days for the
285 first bowel movement. The difference in the two groups is statistically significant as shown by p-
286 value of 0.005. Our results were comparable with the literature.

287 In our study, in the intracorporeal group, 45% patients tolerated orals on day 2, another 45%
288 patients on day 3 and 5% patients on day 4 and day 5 with the median of 3 days. While, in the
289 extracorporeal group 33.3% patients tolerated orals on day 3, 50% patients on day 4 and 16.7%
290 patients on day 5 with the median of 4 days. There is a statistically significant difference
291 between the two groups with respect to the day of oral tolerance as reflected by the p-value of
292 0.001. Our results were comparable to those mentioned in the literature.

293 Anania G, Santini M et al reported the mean of 3.5 days and 4.5 days for the resumption of liquid
294 diet in the intracorporeal and extracorporeal groups respectively. The mean duration for
295 tolerance to solid diet in the two groups was 4.6 days and 5.7 days respectively. Ashley.Vergis,
296 Sarah N. Steigerwaldet al^[23] reported mean of 2.43 days and 3.21 days for tolerance to solid
297 orals in the intracorporeal and extracorporeal groups respectively.

298 In our study we did not observe any leak in both the groups of patients in study. Tu Jian-Cheng,
299 BSc, Wang Shu-Sheng, BSc et al also reported zero leak rates in both the groups. Jayleen Grams,
300 Winnie Tong et al have reported zero leak rate in the intracorporeal group but 1 out of 51 cases
301 (1.96%) in the extracorporeal group had anastomotic leak. However study conducted by Minia
302 Hellen, Casandra Anderson et al^[24] and Milone M, Elmore U et al^[25] showed leak rates of 4.3%
303 and 4.19% in the intracorporeal group respectively and 5.3% in the extracorporeal group each.
304 Significant leak rates in these studies could be due to the larger study design in these studies.
305 In our study, Only 1 out of the total 32 patients (3.1%) developed mesenteric twist which
306 belonged to the extracorporeal group (1 out of 12 patients). No patient in the intracorporeal
307 group developed this complication. But the difference between the two groups was statistically
308 insignificant as reflected by the p-value of 0.375. The patient who developed this complication
309 had hepatic flexure growth and had undergone extended right hemicolectomy. He presented in
310 the postoperative period with features of sub-acute intestinal obstruction (small bowel
311 obstruction) and was re-explored. Minia Hellen, Casandra Anderson et al have reported 1 out of
312 23 patients in the extracorporeal group to develop mesenteric twist. Jorge Arredondo Chaves,
313 Carlos Pastor Idoate et al reported 1 out of 25 patients and Anania G, Santini M et al reported 1
314 out of 33 patients of extracorporeal group to develop this complication.
315 In this study, in the intracorporeal group 1 out of 20 patients (5.0%) developed wound infection
316 while as in the extracorporeal group 1 out of 12 patients (8.3%) developed wound infection. The
317 difference in the rate of wound infection was statistically insignificant (p-value > 0.999).
318 Overall, the wound infection rate was 6.3 percent. Milone M, Elmore U et al. reported overall
319 wound infection rate of 6.83%, 3.84% wound infection rate in intracorporeal anastomosis and
320 10.6% wound infection rate in extracorporeal anastomosis comparable with our study. Jorge
321 Arredondo Chaves, Carlos Pastor Idoate et al. reported rate of wound infection rate of 8% in the
322 extra corporeal group comparable with our study but slightly lower rate in the intracorporeal
323 group (2.86%) with the overall wound infection rate of 5% which was comparable to our study.
324 Ron Shapiro, Uri Keler, et al^[26] also reported wound infection rate of 4.4% in the intracorporeal
325 anastomosis close to our observation.
326 No patient in our study in either group developed extraction site or incisional hernia. Studies
327 carried by Jayleen Grams, et al, Anania et al in 2012, and Milone et al also did not report any
328 extraction site or incisional hernia in their studies. However studies carried by Jorge Arredondo
329 Chaves et al, Shapiro et al reported incisional hernia in few patients. Among all the studies, only
330 Shapiro et al^[27] have recorded a significant percentage of patients in the extracorporeal group to
331 develop incisional hernia. In the study 2.2% patients in the intracorporeal group developed
332 incisional hernia while as 17% in the extracorporeal group developed this complication.
333 In our study, the Intra-corporeal group had the median hospital stay of 5.0 days where as the
334 extracorporeal group had the median hospital stay of 6.5 days with the p-value of 0.010
335 signifying statistically significant difference. Jorge Arredondo Chaves, Carlos Pastor Idoate et al
336 reported average hospital stay of 6 days in the intracorporeal group and 8 days in the
337 extracorporeal group. Roberto Cirocchi, Stefano Trastulli et al^[28] reported average hospital stay
338 of 4 days in the intracorporeal group and 5 days in the extracorporeal group. Ron Shapiro, Uri
339 Keler, et al has reported mean hospital stay of 5.9 ± 2.1 days in the intracorporeal group and $6.9 \pm$
340 3.0 days in the extracorporeal group. The results of all these studies were comparable with our
341 study
342
343

344

345 Conclusion

346

347 Intracorporeal anastomosis in laparoscopic colorectal surgeries leads to earlier return of bowel
348 function, earlier resumption of orals and shorter hospital stay than the extracorporeal
349 anastomosis. There does not exist a significant difference between the two modes of anastomosis
350 in terms of anastomotic and wound related complications.

351

352 References

- 353 1. Canadian cancer society. *Media backgrounder: Colorectal cancer statistics* Toronto: The society; 2002
354 Electronic citation: <http://www.cancer.ca>
- 355 2. Red wine DB, Sharpe DR. Laparoscopic segmental resection of the sigmoid colon for endometriosis. *J*
356 *Laparoendosc Surge* 1991;1(4):217 – 220.
- 357 3. Schlinkert RT. Laparoscopic assisted right hemicolectomy. *Dis Colon Rectum* 1991;34(11):1030 – 1.
- 358 4. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
359 *SurgLaparoscEndosc* 1991;1(3):144 – 50.
- 360 5. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
361 *SurgLaparoscEndosc* 1991 Sep;1(3):144-50.
- 362 6. Slim K. MRC CLASICC trial. *Lancet* 2005 Aug 27-Sep 2;366(9487):712-3. author reply 3e4.
- 363 7. Msika S, Iannelli A, Marano A, Zeitoun G, Deroide G, Kianmanesh R, et al [Hand-sewn intra-abdominal
364 anastomosis performed via video laparoscopy during colorectal surgery]. *Ann Chir* 2000 Jun;125(5):439-43.
- 365 8. Du JJ, Shuang JB, Zheng JY, Li JP, Zhao QC, Hong L, et al [Intracorporeal handsewn technique used in totally
366 laparoscopic colectomy]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2011 Oct;14(10):772-4.
- 367 9. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
368 *SurgLaparoscEndosc*. 1991;1(3):144-50.
- 369 10. Lacy AM, García-Valdecasas JC, Delgado S, Castells A, Taurá P, Piqué JM, et al Laparoscopy-assisted
370 colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. *Lancet*.
371 2002;359(9325):2224-9.
- 372 11. Clinical Outcomes of Surgical Therapy Study Group. A comparison of laparoscopically assisted and open
373 colectomy for colon cancer. *N Engl J Med*. 2004;350(20):2050-9.
- 374 12. Schlachta CM, Mamazza J, Poulin EC. Are transverse colon cancers suitable for laparoscopic resection?
375 *SurgEndosc*. 2007;21(3):396-9.
- 376 13. Yamamoto M, Okuda J, Tanaka K, Kondo K, Tanigawa N, Uchiyama K. Clinical outcomes of laparoscopic
377 surgery for advanced transverse and descending colon cancer: a single-center experience. *SurgEndosc*.
378 2012;26(6):1566-72.
- 379 14. Samia H, Lawrence J, Nobel T, Stein S, Champagne BJ, Delaney CP. Extraction site location and incisional
380 hernias after laparoscopic colorectal surgery: should we be avoiding the midline? *Am J Surg*. 2013;205(3):264-
381 7.
- 382 15. Ueno M, Oya M, Azekura K, Yamaguchi T, Muto T (2005) Incidence and prognostic significance of lateral
383 lymph node metastasis in patients with advanced low rectal cancer. *Br J Surg*, 92:756–763
- 384 16. Hyman N, Manchester TL, Osler T, et al Anastomotic leaks after intestinal anastomosis: it's later than you
385 think. *Ann Surg* 2007; 245:254–8.
- 386 17. Scheidbach H, Benedix F, Hügel O, et al Laparoscopic approach to colorectal procedures in the obese patient:
387 risk factor or benefit? *Obes Surg* 2008;18:66–70.
- 388 18. Tsunoda A, Tsunoda Y, Narita K, et al Quality of life after low anterior resection and temporary loop ileostomy.
389 *Dis Colon Rectum* 2008;51:218–22.
- 390 19. Jorge Arredondo Chaves, Carlos Pastor Idoate, Jorge BaixauliFons, Manuel Bellver Oliver, Nicolás Pedano
391 Rodríguez, Álvaro Bueno Delgado, José Luis Hernández Lizoain. A case-control study of extracorporeal versus
392 intracorporeal anastomosis in patients subjected to right laparoscopic hemicolectomy. *CIR ESP*. 2011;89(1):24–
393 30
- 394 20. Tu Jian-Cheng, BSc, Wang Shu-Sheng, BSc, Zhang Bo, PhD*, Fang Jian, BSc, Zhou Liang, BSc Total
395 laparoscopic right hemicolectomy with 3-step stapled intracorporeal isoperistaltic ileocolic anastomosis for
396 colon cancer. An evaluation of short-term outcomes. *Jian-Cheng et al. Medicine* (2016) 95:48

- 397 21. Anania G, Santini M, Scagliarini L, Marzetti A, Vedana L, Marino S, Gregorio C, Resta G, Cavallesco G. A
398 totally mini – invasive approach for colorectal laparoscopic surgery. *World J Gastroenterol*
399 2012;18(29):3869–3874
- 400 22. Jayleen Grams, Winnie Tong, Alex J. Greenstein, Barry Salky. Comparison of intracorporeal versus
401 extracorporeal anastomosis in laparoscopic-assisted hemicolectomy.. *Surg Endosc* (2010) 24:1886–1891.
- 402 23. Ashley S. vergis, MD, MMed; Sarah n. Steigerwald, MD, MSc; Faizal D, Bhojani, MD; Paul A. Sullivan, MD ;
403 and Krista M. Hardy, MD, MSc. Laparoscopic right hemicolectomy with intra corporeal anastomosis versus
404 extra corporeal anastomosis; a comparison of short-term outcome *Can J Surg*, Vol. 58, No. 1, February 2015:
405 63–68.
- 406 24. Minia Hellan, MD; Casandra Anderson, MD; and Alessio Pigazzi, MD, PhD. Extracorporeal versus
407 intracorporeal anastomosis for laparoscopic right hemi colectomy.. *JSLs, Journal of the Society of*
408 *Laparoendoscopic Surgeons*.(2009)13:312–317
- 409 25. Milone M, Elmore U, Di Salvo E, DelrioP,s Bucci L, Ferulano GP, Napolitano C, Angiolini MR, Bracale U,
410 Clemente M, D’ambra M, Luglio G, Musella M, Pace U, Rosati R, Milone F. Intracorporeal versus
411 extracorporeal anastomosis, results from a multicentre comparative study on 512 right-sided colorectal cancers.
412 *SurgEndosc*. 2015 Aug;29(8):2314–20.
- 413 26. Ron Shapiro, Uri Keler, LiorSegev, Stav Sarna, Kamal Hatib, David Hazzan. *SurgEndosc*. 2015. Laparoscopic
414 right hemicolectomy with intra corporeal anastomosis: short and long term benefits in comparison with
415 extracorporeal anastomosis. *SurgEndosc*. 2016: 3823– 9
- 416 27. Shapiro R, Keler U, Segev L, Sarna S, Hatib K, Hazzan D. Laparoscopic right hemicolectomy with
417 intracorporeal anastomosis: short- and long-term benefits in comparison with extracorporeal anastomosis. *Surg*
418 *Endosc*. 2016 Sep;30(9):3823-9.
- 419 28. Roberto Cirocchi, Stefano Trastulli, EribertoFarinella, Salvatore Guarino, JacopoDesiderio, Carlo Boselli,
420 AmilcareParisi, Giuseppe Noya, Karem Slim. Intracorporeal versus extracorporeal anastomosis during
421 laparoscopic right hemicolectomy - Systematic review and meta-analysis. *Surgical Oncology* 22 (2013) 1–13