

Short term outcome of extracorporeal and Intracorporeal anastomosis in laparoscopic colorectal surgeries – A prospective observational study.

Abstract

Aim of study

Laparoscopic colorectal surgery has gained popularity around the Globe. Laparoscopic colectomy significantly improves the short term and long term outcomes of patients. The bowel anastomosis after laparoscopic resection of the tumour can be done in two ways - extracorporeal anastomosis and Intracorporeal anastomosis. Our study observed and evaluated the data of the patients who underwent these two techniques.

Materials and Methods

A prospective observational study was conducted in the department of General Surgery, SMHS hospital of Government Medical College, Srinagar. A total of 32 patients were studied out of which 20 patients had undergone intracorporeal anastomosis and 12 patients had undergone extracorporeal anastomosis. The aims of the study were to assess the operative time, post-operative ileus, length of hospital stay, anastomotic leak and other anastomotic complications, wound infections and extraction site hernias.

Results

The patients in our study were in the range of 30-85 years with a mean age of 59.18 ± 14.92 . and 59.4% of patients were males and the rest 40.6% were females, with male/female ratio of 1.46:1. There was no significant difference in mean operative time between the intracorporeal group and extracorporeal group (188 ± 9.78 minutes' vs. 180.3 ± 13.8 minutes). The patients in the intracorporeal group had earlier return of bowel function than extracorporeal group as reflected by earlier appreciation of flatus and tolerance to orals. This had led to the shorter hospital stay of the intracorporeal group than the extracorporeal group of patients (median of 5 days vs. 6.5 days) The wound infection rate was 6.3 percent in our study, 5.0% in the intracorporeal group and 8.3% in the extracorporeal group. Only 1 out of the total 32 patients developed mesenteric twist to the extracorporeal group (1 out of 12 patients). We observed no leak or incisional hernia in either groups.

Conclusion

Intracorporeal anastomosis in laparoscopic colorectal surgeries leads to earlier return of bowel function, earlier resumption of orals and shorter hospital stay than the extracorporeal anastomosis. There does not exist a significant difference between the two modes of anastomosis in terms of anastomotic and wound related complications.

37 Keywords; Laparoscopy, Colonrectum, Anastomosis, Intracorporeal, Extracorporeal

38

39 Introduction

40 Colon cancer is the most common type of gastro intestinal cancer ^[1]. Chemotherapy and
41 radiotherapy can improve survival in colorectal cancer patients. However, the only treatment
42 with curative intent is surgical resection of the tumor. Colectomy for cancer can be performed
43 using either open or laparoscopic approach. Laparoscopic colorectal resection has gone through a
44 major evolution since 1991, when the first reports of colorectal resections had been published ^{[2,}
45 ^{3, 4]}. In 1991, Jacobs ^[5] performed the first laparoscopic right hemicolectomy. After right
46 hemicolectomy the ileocolic anastomosis cannot be fashioned “in a natural way” as it normally
47 happens following left hemicolectomy or anterior resection of the rectum; for this reason,
48 different kinds of laparoscopic right colectomy have been proposed ^[6]. The ileocolic anastomosis
49 can be performed using different techniques and devices depending on the intracorporeal and
50 extracorporeal approach. This anastomosis can also be manually performed; despite the lower
51 cost, this requires a greater ability ^[7,8]. Since the publication of the first laparoscopic colectomy
52 its use has been increased. ^[9] Moreover, randomized trials have demonstrated that laparoscopic
53 surgery for colon cancer achieves good short-term and oncologic outcomes similar to those
54 found in open surgery. ^[10,11] However, laparoscopic surgery for transverse and descending colon
55 cancer requires an advanced technique. Hence, only recently, studies have demonstrated the
56 feasibility and safety of the laparoscopic resection for lesions located in the distal transverse and
57 descending colon. ^[12,13] Incisional hernias after open surgery occur in 12 to 20% and may lead to
58 significant morbidity. Midline extraction sites have a higher chance of hernias than non-
59 midline. ^[14] Laparoscopic surgery for rectal cancer has been considered technically more
60 demanding when compared with that for colon cancer. However, laparoscopic total mesorectal
61 excision (TME) has been positively employed for the treatment of rectal cancer in Japanese
62 Centres without lateral lymph node metastasis ^[15] or invasion to the adjacent organ, since it has
63 the advantage of providing a good view even in a narrow pelvis and allowing to perform more
64 precise autonomic nerve preservation

65 Rectal transection and anastomosis at the lower rectum is the most challenging part of
66 laparoscopic low anterior resection. Therefore, some have demonstrated that rectal transection
67 should be performed using instruments for open surgery with small laparotomy. In our institute,
68 however, rectal transection using a currently available endo-stapler followed by anastomosis
69 with a double stapling technique is usually performed. However, anastomotic leakage is still a
70 serious problem after sphincter-saving surgery for rectal cancer. ^[16] Diverting stomata are used to
71 reduce leakage-related complications after LAR, but the routine use of diverting stomata is
72 controversial because of reported morbidity associated with their creation and closure. ^[17,18]

73 Aims and objectives

74 The aim of this study was to assess the short-term outcome of extracorporeal anastomosis and
75 intracorporeal anastomosis in laparoscopic colorectal surgeries in terms of; operative time,
76 anastomotic leak rates and other complications of anastomosis, Post-operative ileus, Length of
77 hospital stay, Wound infections. Extraction and port site hernias.

78 Material and methods:

79 This prospective observational study was conducted in the department of General and minimal
80 access Surgery, from 2015 to 2018 after obtaining the clearance from the Institutional Ethical
81 Committee. A total of 32 cases were enrolled in the study. This comprised of patients admitted
82 for elective surgery for right colon growth, transverse colon growth, left colon growth, sigmoid
83 colon growth and rectal growth above peritoneal reflection in various surgical wards of hospital.
84 The patients that are included in the study are;

- 85 1. Age >18 years.
- 86 2. Patients who are eligible for curative resection of cancer by means of hemicolectomy.
- 87 3. In case of polyp, a colonoscopic biopsy proven invasive cancer.
- 88 4. For rectosigmoid, patient can be included if the tumour lies above the peritoneal
89 reflection.
- 90 5. Solitary colon carcinoma observed at colonoscopy or barium study.

91 The patients that are excluded from the study are;

- 92 1. Contraindications to general anaesthesia e.g. congestive heart failure, chronic renal
93 failure, chronic obstructive lung disease, un-correctable coagulopathy
- 94 2. General contradictions to laparoscopic surgery.
- 95 3. Metastases in the liver or lungs or pre operative evidence of involvement of adjacent
96 structures as detected by CT, MRI or USG.
- 97 4. Acute intestinal obstruction.
- 98 5. Patients who had conversion to open procedure will be excluded from the analysis.

99 These patients were initially evaluated in the outpatient department (OPD) and then planned for
100 surgery. On admission, a detailed history of the patient was recorded including the presenting
101 complaints, duration of the complaints, past history especially with reference to previous
102 surgery, family history and any other associated condition such as chronic ailment and any drug
103 intake.

104 General physical examination was done with particular consideration of build, height and weight
105 followed by systemic examination. Thorough abdominal examination was done in each patient.
106 The patients were taken for laparoscopic colorectal surgery after proper clinical evaluation and
107 after diagnosing them with the disease on colonoscopy and after confirming malignancy on
108 colonoscopic biopsy. Each patient and his attendants were fully explained about the nature of the
109 procedure and the possible complications inherent to the procedure in the native language and
110 thereafter a written consent was sought from the patient prior to surgery. All base line
111 investigations were performed which includes (Complete blood count, Coagulogram, Liver
112 function test, Kidney function test, Blood sugars, Serum electrolytes (Na⁺/K⁺). Chest X-ray and
113 abdominal USG were also performed to investigate lung and liver metastasis respectively. Pre-

114 operative CECT abdomen was done in all patients as a pre-operative staging. The Serum
115 Carcinoembryonic antigen (CEA) also done in all patients.

116 Pre-operative preparation

117 After completing the routine and specific investigations, patients were assessed for anesthetic
118 fitness to undergo the laparoscopic surgery. All the patients were kept fasting 12 hours before
119 surgery and a proper bowel preparation was done using oral solution of polyethylene glycol.
120 Serum electrolytes were repeated before surgery and necessary corrections were made.
121 Ceftriaxone 1 gm I.V as surgical prophylaxis was given to every patient before surgery.
122 Thromboembolic prophylaxis was only given to high risk patients. Patients were allotted to
123 either extracorporeal or intracorporeal limb based on CT scan and intraoperative findings..

124 Operative technique:

125 The experience gained in basic laparoscopy, like technique of creation of pneumoperitoneum,
126 trocar insertion and dissection techniques forms the foundation for advanced laparoscopic
127 colorectal surgery. Pneumoperitoneum is created either via the percutaneous insertion of a
128 Verres needle or with the open Hassan technique. In LRHC, we used four ports: a 10 mm to 12
129 mm camera port for a 30° laparoscope positioned at the level of umbilicus on left side of
130 abdomen, one 12-mm working port for stapling devices in the left upper abdomen above the
131 camera port and 2 five-mm working ports, one located in the left lower abdomen below the
132 camera port and another in the right lower abdomen. The 5 mm port located in the right lower
133 abdomen is later extended for extraction of the specimen. In LLHC, four port technique was also
134 followed but port positions were mirror images of LRHC. Four ports were also used in cases of
135 sigmoidectomy, anterior resection and low anterior resection, with 10 mm umbilical port for
136 camera, 12 mm working port in right lower abdomen and two 5 mm ports – one to the left of
137 camera port and another in the left lower abdomen. The port located in the left lower abdomen is
138 later extended for extraction of the specimen.

139 Basic common steps in laparoscopic colorectal resection


140 All laparoscopic colon procedures have several steps that are common. These steps include:

- 141 1. Localization of the lesion / tumour.
- 142 2. Mobilization of the lesion (medial-to-lateral approach).
- 143 3. Vessel ligation for devascularization of the specimen.
- 144 4. Bowel division.
- 145 5. Restoring bowel continuity by Anastomosis (extracorporeal or intracorporeal).
- 146 6. Specimen retrieval and protection of the wound during retrieval.


147 Follow up:

148 After discharging from the hospital, the patients were advised to follow in OPD with the
149 histopathology of the resected specimen. Patients with stage 1 disease did not require adjuvant
150 chemotherapy and on further follow up such patients were advised serum CEA levels every
151 three months. Such patients were advised CT scan only if there was rise in serum CEA levels.
152 Patients with stage 2 disease and above were strictly advised to follow the medical oncology for

153 chemotherapy. Serum CEA levels were repeated every three months and an annual CT scan and
154 colonoscopy were advised to look for recurrence of the disease.



155
156 Fig.1 CECT showing Right colonic growth



157 Fig. 2 Mobilization of Right colon



158
159 Fig.3 and 4 Creation of extracorporeal and in tra corporeal anastomosis
160



161
162

Fig 5 specimen of colon with ileum

Fig.6 Port position in right hemicolectomy



163
164
165
166
167
168
169
170
171
172
173
174

Fig.7 port operative scar in right hemicolectomy

Fig.8 Port position in LAR

175 **Statistical Analysis**

176 The recorded data was compiled and entered on a spread sheet (Microsoft excel) and then
 177 exported to data editor of SPSS V.20.0 (SPSS Inc., Chicago, Iclinos, USA). P-values less than
 178 0.05 were considered to be statistically significant

179 **Results:**

180 The study included 32 patients out of which 20 patients had undergone intra-corporeal
 181 anastomosis and 12 patients had undergone extra-corporeal anastomosis after resection of
 182 tumour. The patients in our study were in the range of 30-85 years with a mean age of $59.18 \pm$
 183 14.92 . Maximum number of patients were in the age group of 41-60 years (43.75%) followed by
 184 patients older than 60 years (40.63%). Table 2 shows distribution of patients according to their
 185 gender. 59.4% of patients were male and the rest 40.6% were females with male/female ratio of
 186 1.46: 1. The demographic data is shown in table 1 and 2.

Table1: Age distribution of study patients

Age (years)		Anastomosis		Total
		Intracorporeal	Extracorporeal	
<40	Count	4	1	5
	Percentage	12.50 %	3.12 %	15.62 %
41-60	Count	9	5	14
	Percentage	28.12%	15.63 %	43.75 %
>60	Count	7	6	13
	Percentage	21.88%	18.75%	40.63 %
Total	Count	20	12	32
	Percentage	62.50%	37.50 %	100.0 %

Mean±SD = 59.18 ± 14.92 ,

Table 2: Gender distribution of study patients.

Sex		Anastomosis		Total
		Intracorporeal	Extracorporeal	
M	Count	13	6	19
	Percentage	40.62%	18.75%	59.37 %
F	Count	7	6	13
	Percentage	27.88 %	18.75 %	18.75 %
Total	Count	20	12	32
	Percentage	100.0%	100.0%	100.0%

187 Table 3 shows operation time in the two groups of study patients with the mean operation time of
 188 188.1 ± 9.78 minutes in the intracorporeal group and 180.3 ± 13.8 minutes in the extracorporeal
 189 group. No significant difference in the operation time between the two groups of study
 190 population could be observed as reflected by the p-value of 0.07

Table: 3 Operation time (minutes) in two techniques of study patients.

Anastomosis	Number of Patients	Operation time	p-value
		Mean±SD	
Intracorporeal	20	188.1 ± 9.78	0.075

191 Table 4 shows absolute and percentage distribution of patients in the two groups of study with
 192 respect to the appreciation of passage of flatus. Patients in the intracorporeal group had the
 193 median of 2.0 days where as patients in the extracorporeal group had median of 3.0 days. The
 194 difference in the two groups is statistically significant as shown by p-value of 0.01

Table 4: Time distribution (days) of appreciation of passage of flatus in the two groups of study patients

Flatus		Anastomosis		Total
		Intracorporeal	Extracorporeal	
1	Count	1	1	2
	percentage	5.0%	8.3%	6.3%
2	Count	12	1	13
	percentage	60.0%	8.3%	40.6%
3	Count	6	5	11
	Percentage	30.0%	41.7%	34.4%
4	Count	1	4	5
	Percentage	5.0%	33.3%	15.6%
5	Count	0	1	1
	Percentage	0.0%	8.3%	3.1%
Total	Count	20	12	32
	Percentage	100.0%	100.0%	100.0%
Median		2.0	3.0	-

p=0.01, Mann-Whitney test

195 Table 5 shows absolute and percentage distribution of patients in the two groups of study with
 196 respect to the first bowel movement. Patients in the intracorporeal group had the median of 4.0
 197 days where as patients in the extracorporeal group had median of 5.0 days. The difference in the
 198 two groups is statistically significant as shown by p-value of 0.01

Table 5: Time distribution (days) of bowel movement in the two groups of patients in study.

Stools		Anastomosis		Total
		Intracorporeal	Extracorporeal	
2	Count	2	0	2
	Percentage	10.0%	0.0%	6.3%
3	Count	2	1	3
	Percentage	10.0%	8.3%	9.4%
4	Count	12	1	13
	Percentage	60.0%	8.3%	40.6%
5	Count	2	7	9
	Percentage	10.0%	58.3%	28.1%
6	Count	2	3	5

	Percentage	10.0%	25.0%	15.6%
Total	Count	20	12	32
	Percentage	100.0%	100.0%	100.0%
Median		4.0	5.0	-

p=0.01, Mann-Whitney test

199 Table 6 shows absolute and percentage distribution of the two groups of study population with
200 respect to the day of tolerance of orals. In the intracorporeal group, 45% patients tolerated orals
201 on day 2, another 45% patients on day 3 and 5% patients on day 4 and day 5 with the median of
202 3 days. While, in the extracorporeal group 33.3% patients tolerated orals on day 3, 50% patients
203 on day 4 and 16.7% patients on day 5 with the median of 4 days. There is a statistically
204 significant difference between the two groups with respect to the day of oral tolerance as
205 reflected by the p-value of 0.00.

Table 6: Time distribution (days) of oral tolerance in two techniques of anastomosis in study patients.

Orals	Anastomosis		Total
	Intracorporeal	Extracorporeal	
2	Count	9	9
	Percentage	45.0%	28.1%
3	Count	9	13
	Percentage	45.0%	40.6%
4	Count	1	7
	Percentage	5.0%	21.9%
5	Count	1	3
	Percentage	5.0%	9.4%
Total	Count	20	32
	Percentage	100.0%	100.0%
Median		3.0	4.0

p=0.00, Mann-Whitney test

206 Table 7 lists different complications of laparoscopic colorectal surgeries reported in literature
207 and their incidence in our study. We did not observe any anastomotic leak or extraction site
208 hernia in our study. Anastomotic twist is described in Table 8 and wound infection in Table 9 in
209 detail.

Table 7: Rate of Complications in two groups of study patients

Complication		Intracorporeal	Extracorporeal	Total
Anastomotic leak	Count	0	0	0
Anastomotic twist	Percentage	0%	0%	0%
Wound infection	Count	0	1	1
	Percentage	0%	8.3%	3.12%
Wound infection	Count	1	1	2
	Percentage	5%	8.3%	6.3%

Extraction site hernia	Count	0	0	0
	Percentage	0%	0%	0%

210 Table 8 shows absolute and percentage distribution of mesenteric twist at site Of anastomosis in
 211 the study patients. Only 1 out of the total 32 patients (3.12%) developed mesenteric twist which
 212 belonged to the extracorporeal group (1 out of 12 patients). No patient in the intracorporeal
 213 group developed this complication. But the difference between the two groups was statistically
 214 insignificant as reflected by the p-value of 0.37

Table 8: Rate of mesenteric twist in two techniques of anastomosis in study patients.

Anastomosis	Number of Patients	Mesentric twist	Percentage
Intracorporeal	20	0	0 %
Extracorporeal	12	1	8.3 %
Total	32	1	3.12 %

p>0.37, Chi-square test Exact p

215 Table 9 shows rate of wound infection in the two groups of study population. In the
 216 intracorporeal group 1 out of 20 patients (5.0%) developed wound infection while as in the
 217 extracorporeal group 1 out of 12 patients (8.3%) developed wound infection. The difference in
 218 the rate of wound infection was statistically insignificant (p-value> 0.99). Overall, the wound
 219 infection rate was 6.3 percent.

Table 9: Rate of wound infection in two groups of patients in study

Anastomosis	Number of Patients	Wound Infection	Percentage
Intracorporeal	20	1	5 %
Extracorporeal	12	1	8.3 %
Total	32	2	6.3 %

p>0.99, Chi-square test Exact p

220

221 Table 10 shows the absolute and percentage distribution of the patients in the two groups of the
 222 study patients with respect to the number of days of hospital stay. Most of the patients in the
 223 intracorporeal group (70%) stayed in the hospital for 5 – 6 days while as in the extracorporeal
 224 group majority of the patients (75%) stayed in the hospital for 6 – 7 days. Intracorporeal group
 225 had the median hospital stay of 5.0 days where as the extracorporeal group had the median
 226 hospital stay of 6.5 days with the p-value of 0.01 signifying statistically significant difference.

Table 10: length of hospital stay(days) in two groups of study patients

LOHS		Anastomosis		Total
		Intracorporeal	Extracorporeal	
4	Count	2	0	2
	Percentage	10.0%	0.0%	6.3%
5	Count	9	1	10
	Percentage	45.0%	8.3%	31.3%
6	Count	5	5	10
	Percentage	25.0%	41.7%	31.3%

7	Count	3	4	7
	Percentage	15.0%	33.3%	21.9%
8	Count	1	1	2
	Percentage	5.0%	8.3%	6.3%
10	Count	0	1	1
	Percentage	0.0%	8.3%	3.1%
Total	Count	20	12	32
	Percentage	100.0%	100.0%	100.0%
Median		5.0	6.5	-

p=0.01, Mann-Whitney test

227

228 Discussion

229 Application of laparoscopy to colorectal surgery has produced some short- term benefits like
 230 faster return of gut function leading to more rapid discharge from the hospital and a faster return
 231 to normal activity as well as long term benefits like reduced rates of obstruction secondary to
 232 adhesions and reduced incidence of ventral hernia. The creation of bowel anastomosis after
 233 laparoscopic resection of large bowel tumour can be done in two ways – extracorporeal
 234 anastomosis and intracorporeal anastomosis. A number of studies have been carried worldwide
 235 to document the equalities and differences between the two modes of creation of bowel
 236 anastomosis with respect to the short- term outcomes like – operation time, post-operative return
 237 of bowel function, complications of anastomosis, wound infection and length of hospital stay.
 238 So, we also have conducted a study to evaluate these short-term outcomes of extracorporeal and
 239 intracorporeal anastomosis in laparoscopic colorectal surgeries. It was a hospital based
 240 prospective observational study conducted in the department of General and minimal access
 241 Surgery of the hospital of Government Medical College.

242 A total of 32 patients were studied out of which 20 patients had undergone IC anastomosis and
 243 12 patients had undergone EC anastomosis. The results were based on age, gender, operative
 244 time, post-operative return of bowel function in terms of appreciation of flatus, oral tolerance and
 245 first bowel movement, complications such as anastomotic leak, mesenteric twist, wound
 246 infections and extraction site hernias and length of hospital stay. The patients in our study were
 247 in the range of 30-85 years with a mean age of 59.18 ± 14.92 . Maximum number of patients
 248 were in the age group of 41-60 years (43.8%) followed by patients older than 60
 249 years(40.6%).Jorge Arredondo Chaves, Carlos Pastor Idoate et al^[19] have reported in their study
 250 mean age group of patients as 62.6 ± 13.4 years in the intracorporeal group and 58.9 ± 12.9 years
 251 in the extracorporeal group which closely matched with our study.In our study, 59.4% of patients
 252 were male and the rest 40.6% were females with male/female ratio of 1.46:1. Jorge Arredondo
 253 Chaves, Carlos Pastor Idoate et al have reported male/female ratio of 1.22:1 and Tu Jian-Cheng,
 254 BSc, Wang Shu-Sheng, BSc et al^[20] reported male/female ratio of 1.57:1. Both of these studies
 255 had comparable gender ratio with our study.

256 In our study there was no significant difference in operative time between two groups.The mean
 257 operation time in the intracorporeal group was 188 ± 9.78 minutes and 180.3 ± 13.8 minutes in
 258 the extra corporeal group. Comparable results were obtained by Minia Hellen, Casandra
 259 Anderson et alwho reported the mean operative time of 190 minutes in the intracorporeal group
 260 and 180 minutes in the extracorporeal group.Anania G, Santini M et al^[21] also showed similar

261 results with the mean operative time of 186.8 minutes in the intracorporeal group and 184.8
262 minutes in the extracorporeal group.

263 The appreciation of flatus in the intracorporeal group had the median of 2.0 days where as
264 patients in the extracorporeal group had median of 3.0 days. The difference in the two groups is
265 statistically significant as shown by p-value of 0.007. Comparable results were obtained by
266 Jayleen Grams, Winnie Tong et al^[22], Anania G, Santini M et al who reported mean days of
267 appreciation of flatus in the intracorporeal group as 2.0 days and 2.4 days in the extracorporeal
268 group. Tu Jian-Cheng, BSc, Wang Shu-Sheng, BSc et al reported the mean duration of
269 appreciation of flatus as 2.57 ± 0.08 days in the intracorporeal group and 3.10 ± 0.11 days in the
270 extracorporeal group which are comparable to our study.

271 The first bowel movement in the intracorporeal group had the median of 4.0 days where as
272 patients in the extracorporeal group had median of 5.0 days. The difference in the two groups is
273 statistically significant as shown by p-value of 0.005. Jorge Arredondo Chaves, Carlos Pastor
274 Idoate et al in 2011 reported the median days of the first bowel movement as 3 days in the
275 intracorporeal group 4 days in the extracorporeal group. Anania G Santini Met al in
276 2012 reported mean of 3.8 days in the intracorporeal group and 4.9 days in the extracorporeal
277 group for the first bowel movement. In our study patients in the intracorporeal group had the
278 median of 4.0 days where as patients in the extracorporeal group had median of 5.0 days for the
279 first bowel movement. The difference in the two groups is statistically significant as shown by p-
280 value of 0.005. Our results were comparable with the literature.

281 In our study, in the intracorporeal group, 45% patients tolerated orals on day 2, another 45%
282 patients on day 3 and 5% patients on day 4 and day 5 with the median of 3 days. While, in the
283 extracorporeal group 33.3% patients tolerated orals on day 3, 50% patients on day 4 and 16.7%
284 patients on day 5 with the median of 4 days. There is a statistically significant difference
285 between the two groups with respect to the day of oral tolerance as reflected by the p-value of
286 0.001. Our results were comparable to those mentioned in the literature.

287 Anania G, Santini M et al reported the mean of 3.5 days and 4.5 days for the resumption of liquid
288 diet in the intracorporeal and extracorporeal groups respectively. The mean duration for
289 tolerance to solid diet in the two groups was 4.6 days and 5.7 days respectively. Ashley Vergis,
290 Sarah N. Steigerwald et al^[23] reported mean of 2.43 days and 3.21 days for tolerance to solid
291 orals in the intracorporeal and extracorporeal groups respectively.

292 In our study we did not observe any leak in both the groups of patients in study. Tu Jian-Cheng,
293 BSc, Wang Shu-Sheng, BSc et al also reported zero leak rates in both the groups. Jayleen Grams,
294 Winnie Tong et al have reported zero leak rate in the intracorporeal group but 1 out of 51 cases
295 (1.96%) in the extracorporeal group had anastomotic leak. However study conducted by Minia
296 Hellen, Casandra Anderson et al^[24] and Milone M, Elmore U et al^[25] showed leak rates of 4.3%
297 and 4.19% in the intracorporeal group respectively and 5.3% in the extracorporeal group each.
298 Significant leak rates in these studies could be due to the larger study design in these studies.

299 In our study, Only 1 out of the total 32 patients (3.1%) developed mesenteric twist which
300 belonged to the extracorporeal group (1 out of 12 patients). No patient in the intracorporeal
301 group developed this complication. But the difference between the two groups was statistically
302 insignificant as reflected by the p-value of 0.375. The patient who developed this complication
303 had hepatic flexure growth and had undergone extended right hemicolectomy. He presented in
304 the postoperative period with features of sub-acute intestinal obstruction (small bowel
305 obstruction) and was re-explored. Minia Hellen, Casandra Anderson et al have reported 1 out of
306 23 patients in the extracorporeal group to develop mesenteric twist. Jorge Arredondo Chaves,

307 Carlos Pastor Idoate et al reported 1 out of 25 patients and Anania G, Santini M et al reported 1
308 out of 33 patients of extracorporeal group to develop this complication.
309 In this study, in the intracorporeal group 1 out of 20 patients (5.0%) developed wound infection
310 while as in the extracorporeal group 1 out of 12 patients (8.3%) developed wound infection. The
311 difference in the rate of wound infection was statistically insignificant ($p\text{-value} > 0.999$).
312 Overall, the wound infection rate was 6.3 percent. Milone M, Elmore U et al. reported overall
313 wound infection rate of 6.83%, 3.84% wound infection rate in intracorporeal anastomosis and
314 10.6% wound infection rate in extracorporeal anastomosis comparable with our study. Jorge
315 Arredondo Chaves, Carlos Pastor Idoate et al. reported rate of wound infection rate of 8% in the
316 extra corporeal group comparable with our study but slightly lower rate in the intracorporeal
317 group(2.86%) with the overall wound infection rate of 5% which was comparable to our study.
318 Ron Shapiro, Uri Keler, et al^[26] also reported wound infection rate of 4.4% in the intracorporeal
319 anastomosis close to our observation.
320 No patient in our study in either group developed extraction site or incisional hernia. Studies
321 carried by Jayleen Grams, et al, Anania et al in 2012, and Milone et al also did not report any
322 extraction site or incisional hernia in their studies. However studies carried by Jorge Arredondo
323 Chaves et al, Shapiro et al reported incisional hernia in few patients. Among all the studies, only
324 Shapiro et al^[27] have recorded a significant percentage of patients in the extracorporeal group to
325 develop incisional hernia. In the study 2.2% patients in the intracorporeal group developed
326 incisional hernia while as 17% in the extracorporeal group developed this complication.
327 In our study, the Intra-corporeal group had the median hospital stay of 5.0 days where as the
328 extracorporeal group had the median hospital stay of 6.5 days with the p-value of 0.010
329 signifying statistically significant difference. Jorge Arredondo Chaves, Carlos Pastor Idoate et al
330 reported average hospital stay of 6 days in the intracorporeal group and 8 days in the
331 extracorporeal group. Roberto Cirocchi, Stefano Trastulli et al^[28] reported average hospital stay
332 of 4 days in the intracorporeal group and 5 days in the extracorporeal group. Ron Shapiro, Uri
333 Keler, et al has reported mean hospital stay of 5.9 ± 2.1 days in the intracorporeal group and $6.9 \pm$
334 3.0 days in the extracorporeal group. The results of all these studies were comparable with our
335 study
336
337
338

339 Conclusion

340
341 Intracorporeal anastomosis in laparoscopic colorectal surgeries leads to earlier return of bowel
342 function, earlier resumption of orals and shorter hospital stay than the extracorporeal
343 anastomosis. There does not exist a significant difference between the two modes of anastomosis
344 in terms of anastomotic and wound related complications.
345

346 References

- 347 1. Canadian cancer society. *Media backgrounder: Colorectal cancer statistics* Toronto: The society; 2002
348 Electronic citation: <http://www.cancer.ca>
- 349 2. Red wine DB, Sharpe DR. Laparoscopic segmental resection of the sigmoid colon for endometriosis. *J*
350 *Laparoendosc Surge* 1991;1(4):217 – 220.
- 351 3. Schlinkert RT. Laparoscopic assisted right hemicolectomy. *Dis Colon Rectum* 1991;34(11):1030 – 1.

- 352 4. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
353 SurgLaparoscEndosc 1991;1(3):144 – 50.
- 354 5. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
355 SurgLaparoscEndosc 1991 Sep;1(3):144-50.
- 356 6. Slim K. MRC CLASICC trial. Lancet 2005 Aug 27-Sep 2;366(9487):712-3. author reply 3e4.
- 357 7. Msika S, Iannelli A, Marano A, Zeitoun G, Deroide G, Kianmanesh R, et al [Hand-sewn intra-abdominal
358 anastomosis performed via video laparoscopy during colorectal surgery]. Ann Chir 2000 Jun;125(5):439-43.
- 359 8. Du JJ, Shuang JB, Zheng JY, Li JP, Zhao QC, Hong L, et al [Intracorporeal handsewn technique used in totally
360 laparoscopic colectomy]. Zhonghua Wei Chang Wai Ke Za Zhi 2011 Oct;14(10):772-4.
- 361 9. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy).
362 SurgLaparoscEndosc. 1991;1(3):144-50.
- 363 10. Lacy AM, García-Valdecasas JC, Delgado S, Castells A, Taurá P, Piqué JM, et al Laparoscopy-assisted
364 colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. Lancet.
365 2002;359(9325):2224-9.
- 366 11. Clinical Outcomes of Surgical Therapy Study Group. A comparison of laparoscopically assisted and open
367 colectomy for colon cancer. N Engl J Med. 2004;350(20):2050-9.
- 368 12. Schlachta CM, Mamazza J, Poulin EC. Are transverse colon cancers suitable for laparoscopic resection?
369 SurgEndosc. 2007;21(3):396-9.
- 370 13. Yamamoto M, Okuda J, Tanaka K, Kondo K, Tanigawa N, Uchiyama K. Clinical outcomes of laparoscopic
371 surgery for advanced transverse and descending colon cancer: a single-center experience. SurgEndosc.
372 2012;26(6):1566-72.
- 373 14. Samia H, Lawrence J, Nobel T, Stein S, Champagne BJ, Delaney CP. Extraction site location and incisional
374 hernias after laparoscopic colorectal surgery: should we be avoiding the midline? Am J Surg. 2013;205(3):264-
375 7.
- 376 15. Ueno M, Oya M, Azekura K, Yamaguchi T, Muto T (2005) Incidence and prognostic significance of lateral
377 lymph node metastasis in patients with advanced low rectal cancer. Br J Surg, 92:756–763
- 378 16. Hyman N, Manchester TL, Osler T, et al Anastomotic leaks after intestinal anastomosis: it's later than you
379 think. Ann Surg2007; 245:254–8.
- 380 17. Scheidbach H, Benedix F, Hügel O, et al Laparoscopic approach to colorectal procedures in the obese patient:
381 risk factor or benefit? Obes Surg2008;18:66–70.
- 382 18. Tsunoda A, Tsunoda Y, Narita K, et al Quality of life after low anterior resection and temporary loop ileostomy.
383 Dis Colon Rectum 2008;51:218–22.
- 384 19. Jorge Arredondo Chaves, Carlos Pastor Idoate, Jorge BaixauliFons, Manuel Bellver Oliver, Nicolás Pedano
385 Rodríguez, Álvaro Bueno Delgado, José Luis Hernández Lizoain. A case-control study of extracorporeal versus
386 intracorporeal anastomosis in patients subjected to right laparoscopic hemicolectomy. CIR ESP. 2011;89(1):24–
387 30
- 388 20. Tu Jian-Cheng, BSc, Wang Shu-Sheng, BSc, Zhang Bo, PhD*, Fang Jian, BSc, Zhou Liang, BSc Total
389 laparoscopic right hemicolectomy with 3-step stapled intracorporeal isoperistaltic ileocolic anastomosis for
390 colon cancer. An evaluation of short-term outcomes. Jian-Cheng et al. Medicine (2016) 95:48
- 391 21. Anania G, Santini M, Scagliarini L, Marzetti A, Vedana L, Marino S, Gregorio C, Resta G, Cavallesco G. A
392 totally mini – invasive approach for colorectal laparoscopic surgery. World J Gastroenterol
393 2012;18(29):3869–3874
- 394 22. Jayleen Grams, Winnie Tong, Alex J. Greenstein, Barry Salky. Comparison of intracorporeal versus
395 extracorporeal anastomosis in laparoscopic-assisted hemicolectomy.. Surg Endosc (2010) 24:1886–1891.
- 396 23. Ashley S. vergis, MD, MMEd; Sarah n. Steigerwald, MD, MSc; Faizal D, Bhojani, MD; Paul A. Sullivan, MD ;
397 and Krista M. Hardy, MD, MSc. Laparoscopic right hemicolectomy with intra corporeal anastomosis versus
398 extra corporeal anastomosis; a comparison of short-term outcome Can J Surg, Vol. 58, No. 1, February 2015:
399 63–68.
- 400 24. Minia Hellan, MD; Casandra Anderson, MD; and Alessio Pigazzi, MD, PhD. Extracorporeal versus
401 intracorporeal anastomosis for laparoscopic right hemi colectomy.. JSLS, Journal of the Society of
402 Laparoendoscopic Surgeons. (2009)13:312–317
- 403 25. Milone M, Elmore U, Di Salvo E, DelrioP, s Bucci L, Ferulano GP, Napolitano C, Angiolini MR, Bracale U,
404 Clemente M, D'ambra M, Luglio G, Musella M, Pace U, Rosati R, Milone F. Intracorporeal versus
405 extracorporeal anastomosis, results from a multicentre comparative study on 512 right-sided colorectal cancers.
406 SurgEndosc. 2015 Aug;29(8):2314–20.

407 26. Ron Shapiro, Uri Keler, LiorSegev, Stav Sarna, Kamal Hatib, David Hazzan. SurgEndosc. 2015.Laparoscopic
408 right hemicolectomy with intra corporeal anastomosis: short and long term benefits in comparison with
409 extracorporeal anastomosis.SurgEndosc. 2016: 3823– 9
410 27. Shapiro R, Keler U, Segev L, Sarna S, Hatib K, Hazzan D. Laparoscopic right hemicolectomy with
411 intracorporeal anastomosis: short- and long-term benefits in comparison with extracorporeal anastomosis. Surg
412 Endosc. 2016 Sep;30(9):3823-9.
413 28. Roberto Cirocchi, Stefano Trastulli, EribertoFarinella, Salvatore Guarino,JacopoDesiderio, Carlo Boselli,
414 AmilcareParisi, Giuseppe Noya, Karem Slim. Intracorporeal versus extracorporeal anastomosis during
415 laparoscopic right hemicolectomy - Systematic review and meta-analysis.Surgical Oncology 22 (2013) 1–13
416
417
418
419
420
421
422
423
424

UNDER PEER REVIEW