

ESTIMATING OF CORN YIELD BASED ON AGROMETEOROLOGICAL MODELS IN THE STATE OF MATO GROSSO

RESUMO

Aims:The objective of this work was to evaluate the performance of agrometeorological models for estimating the yield potential of corn in municipalities of the State of Mato Grosso, referring to the 2014/2015 harvest period.

Location and duration of the study:Due to their great potential on corn crop cultivation, four cities were chosen within the State of Mato Grosso: Nova Mutum, Lucas do Rio Verde, Sorriso and Sinop.2014/2015 harvest period.

Methodology: The yield data were obtained by the Farming Economy Institute of Mato Grosso-IMEA. The physiological ripening cycle considered was 120 days and the seeding period 02/02/2015 and harvest 01/06/2015.In order to calculate yield and the water stress, it was necessary to gather data from meteorological stations near the cities where the work was produced, these data were available at the National Institute of Meteorology. The estimated potential yield data were compared among the agrometeorological models, as well as the real yield obtained in the selected municipalities.

Results:The estimated potential yield data were compared among the agrometeorological models, as well as the real yield obtained in the selected municipalities. There were variances amongst the agrometeorological models studied, on average the model 1 estimated (5413,68 kg ha⁻¹) and the Model 2 (6766,45 kg ha⁻¹), Table 3.It was observed that Model 1 estimated greater yield for Nova Mutum, followed by Sinop, Sorriso and Lucas do Rio Verde, and yet the Model 2 estimated greater potential for Lucas do Rio Verde, followed by the municipalities of Sinop, Sorriso and Nova Mutum (Picture 1 and Table 3). In this regard, the model 2 has characterized the closest potential yield, from the environment yield reality.

Conclusion:The Model 2 has characterized the potential yield closer to the reality of production environment.The difference of all potential yield of corn from all municipalities studied were all directly related to factor interaction which interferes on its growth and development, consequently the difference among these environments. Taking into account the effects of water conditions, the chosen period for seeding can harm the corn yield in the municipality of Lucas do Rio Verde.

10
11
12
13
14
15
16
17
18
19
20
21
22
23

Keywords: *Agro-climatology, agrometeorological models, productivity, potential yield,, zea mays.*

1. INTRODUCTION

Currently, the corn (Zea Mays) is regarded as second most important crop cultivation in Brazilian Agriculture. Brazil is the third greatest agricultural producer and the second country that exports this cereal [6].

In Brazil, Zea Mays production was approximately 82,1 million tons during the harvest of 2017/2018 [6]. Considering the State of Mato Grosso as the greatest producer of this cereal, obtained during the harvest of 2017/2018it was harvested 27.49 million of tons representing around 43,1% of national production[6].

Differently from other regions in Brazil, the corn produced in Mato Grosso, most of it is characterized as inter-harvest or interim-harvest corn. According to the Conab's research data, 97,94% of cereal in the state is second crop [7].

24
25 The Agriculture is an economic activity which has a great dependency on weather conditions, such as: the rain, the air
26 temperature and solar radiation, the main meteorological variation can interfere the crops' growth, development of its
27 cultivation [8].

28
29 Amongst the factors that could affect the corn yield are the precipitation and temperature.

30
31 In regard to the precipitation regime, the corn can be cultivated in regions where the rain is common from 250 up to
32 5000mm annually, and the plant average consuming on its early cycle remains around 600mm. The daily cereal
33 consuming on its early stages tend not to exceed values beyond than 2,5mm whereas the ripening process, these values
34 could reach 10mm/day at places where the lowest levels of humidity and higher temperatures are presented [9]. Thus, the
35 cultivation could present limitations, especially related to a possible water stress, bearing in mind as the seeding season
36 could occur by the end of the rain season in most regions, it could result in a certain productivity loss due to the water
37 stress problem.

38
39 In addition, higher temperatures allied to the absence of precipitation are responsible for drying the stigma and style and
40 also to produce cob flaws resulting to losses in productivity. In the event of plant blooming, temperatures below than
41 15,5°C can slow down its development. In conclusion, Summer days with the average temperatures below 19°C, and
42 below 12,8°C during the night time are not recommended to corn production [9].

43
44 Turns out the corn yield is highly dependent on its interaction between the weather and phenological periods.

45 In order to attain a better understanding on these interactions, agrometeorological models have been used aiming to
46 distinguish weather variation effects upon grain production. The productivity estimate models based on weather factors
47 allows it to follow the weather effects along the crop cultivating cycle and relate them to its final yield enabling to assess
48 restrictions on agrarian development and to achieve strategies in order to increase production [10].

49 Therefore, the objective of this work was to assess the agrometeorological model's performance estimating the potential
50 corn yield in municipalities in the State of Mato Grosso.

51 52 2. MATERIAL AND METHODS

53 54 2.1 StudyFields

55
56 Nova Mutum, Lucas do Rio Verde, SorrisoandSinop, Municipalities of the State of Mato Grosso, located in the
57 centre-west of Brazil, they were all chosen for this study due to their high yield of corn cultivation.

58 59 2.2 Estimated Potential Yield

60
61 The estimated potential yield calculation has been performed to all four cities, referring to the 2014/2015 harvest period
62 and it was based on two agrometeorological models: Model 1 from [1], with [2] and Model 2 from [3] Adapted [4] and also
63 from [5].

64
65 The estimate yield established by Model 1, [1], equations were used where the plants are divided by four groups regarding
66 to photosynthetic and phenological aspects, the corn being framed on the third group belonging to C4 plants, whereas the
67 optimal photosynthesis temperature situates around 30 and 35°C. The maximum yield is given by the equation below in
68 kg/ha⁻¹:

$$By = (Bn * HI * (100 + U)) / 100$$

69
70
71
72 Where:

- 73 • HI – Represents the harvest index found at table 1, for corn used the inferior value (0,35);
- 74 • U – Product unity on dry base (table 1);
- 75 • O Bn is the dry matter accumulated during all cycle and it is calculated by the equation:

$$Bn = (0,36 * bgm) / (1/N + 0,25 * C) - kg/ha$$

76
77
78
79 **Table 1. Harvesting Index (Cc) and humidity level (U%)**

Crop	Product	Cc	U%
Corn	Grain	0,35 – 0,45	10 – 13

Font: [3] and [4].

Where:

• N is the crops' total cycle and the average photosynthesis gross which is obtained by the following equation:

$$\text{bgm} = F \times \text{bo} \times (0,8 + 0,01 \times \text{Pmax}) + (1-F) \times \text{bc} \times (0,5 + 0,025 \times \text{Pmax})$$

Where:

• F – Fractioned time of cloud covering [1];

• bo- absorption rate during cloudy days (kg/ha/d):

$$\text{bo} = 31,66 + 0,55 \times \text{Ac}$$

• bc- absorption rate during clear days (kg/ha/d):

$$\text{bc} = 107,00 + 0,90 \times \text{Ac}$$

• Pmax – Maximum CO2 Exchange rate [1];

• Ac- This work, regarded that Ac corresponds to 80% of atmospheric transmittance. In this case:

$$\text{Ac} = 0,5 \times 0,8 \times \text{Q0}$$

Where:

• Q0 is the extra-terrestrial short waves of radiation (Table 2).

Table 2. Values Qo (mm/d) for South latitude

Lat	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	14,5	15	15,2	14,7	13,9	13,4	13,5	14,2	14,9	14,9	14,6	14,3
10	15,9	15,7	15	13,8	12,4	11,6	11,9	13	14,4	15,3	15,7	15,7
20	16,7	16	14,5	12,4	10,6	9,6	10	11,5	13,5	15,3	16,2	16,8
30	17,2	15,7	13,5	10,8	8,5	7,4	7,8	9,6	12,2	14,7	16,7	17,6

Font: [11].

The gross photosynthesis was corrected (bgm). The value C for crops with lower protein levels was C30 = 0,0108. Yet, it was considered the adjustment of C with the temperature (T), according with:

$$C = (0,0283 \times (0,044 + (0,00019 \times T) + (0,001 \times T^2)))$$

Where:

$$T = T_d = (3 \times T_{\text{max}} + T_{\text{min}}) / 4$$

• Tmax: Maximum temperature;

• Tmin: minimum temperature.

The model 2 is the method of Agroecological Zone described by [3], where the potential productivity (maximum yield) is given buy the following equation:

$$\text{PP}_R = \text{PPBp} C_{\text{IAF}} C_r C_c \text{ND (kg ha}^{-1}\text{)}$$

Where:

• PPBp: is the production of gross matter in standard crop, which is considered the presence or absence of clouds expressed in kg MS ha⁻¹ d⁻¹ and the extra-terrestrial global solar irradiance (Qo), mm/d (Table 2).

• C_{IAF}: is the correction for the leaf area index giving by the equation 2: 0,0093 + 0,185 IAF – 0,015 IAF², o IAF considered in this work was 4.

• C_R (Correction for the ventilation) it could attain the following values: CR=0,6 for T < 20°C and CR =0,5 T ≥20°C, on T which is the average temperature during the considered period.

• C_c (Correction for the harvested part) was established according to Table 2.

134 • ND: is the considered number of days.
135

136 Then after taken the PPR (Dry matter from the harvest collected) it should be added the residual humidity (U%) which
137 normally is withheld on the product (Table 1), then reaching the final potential yield value (PPF), obtained by the following
138 equation:

$$\text{PPF} = \text{PPR} / (1 - 0,01 \text{ U}\%)$$

143 2.3 Water stress effect upon productivity

144 In order to get acquainted with the water stress, it was necessary to elaborate the water balance sheet according to the
145 methodology adapted by [11]. The materials used to its preparation were: medium temperature, precipitation (obtained at
146 meteorological stations), potential evapotranspiration and available water capacity (CAD) of 100 mm, for annual
147 cultivation.
148

149 The FAO model, proposed by [3], links the relative yield drop (1-PR/PP_F) with the relative deficit of evapotranspiration (1-
150 ETR/ET_c), regarding PR the real yield to be calculated on the following equation below:

$$\text{PR} = (1 - k_y (1 - \text{ETR} / \text{ET}_c)) * \text{PP}_F$$

154 Where:

155 The k_y is the response factor for the crops' initial state; reproductive and ripening, their values were incorporated
156 according to [3].

157 In the calculation of limited yield by the water availability to maximum crop evapotranspiration (ET_c) is linked with the
158 evapotranspiration baseline (ET_o) through a crop coefficient (K_c), as presented in the equation:

$$\text{ET}_c = K_c * \text{ET}_o$$

163 The evapotranspiration baseline was calculated by [12] method which proposed the following calculation
164

$$\text{ETP} = 0,01 Q_o T \text{ ND}$$

167 Whereby:

168 Q_o is the global extra-terrestrial solar irradiance (Table 2), T the average air temperate (oC), at the considered period; and
169 ND the number of days considered.
170

171 The crop coefficient values (K_c) for early stages; reproductive and ripening, were incorporated according to [3] proposal.
172

173 2.4 Yield data

174 The yield data were obtained by the Farming Economy Institute of Mato Grosso-IMEA. The physiological ripening cycle
175 considered was 120 days and the seeding period 02/02/2015 and harvest 01/06/2015.
176

177 The physiological ripening cycle and the seeding season were chosen through agricultural zoning performed by the
178 Department of agriculture, livestock and refuelling, which identifies the capable municipalities and the seeding season for
179 the corn second crop for the state of Mato Grosso.
180

181 In order to calculate yield and the water stress, it was necessary to gather data from meteorological stations near the
182 cities where the work was produced, these data were available at the National Institute of Meteorology [13].
183


184 3. RESULTS AND DISCUSSION

185 The estimated potential yield data were compared among the agrometeorological models, as well as the real yield
186 obtained in the selected municipalities.
187
188
189
190
191
192
193

194
195
196
197
198
199
200
201

There were variances amongst the agrometeorological models studied, on average the model 1 estimated (5413,68 kg ha⁻¹) and the Model 2 (6766,45 kg ha⁻¹), Table 3.

It was observed that Model 1 estimated greater yield for Nova Mutum, followed by Sinop, Sorriso and Lucas do Rio Verde, and yet the Model 2 estimated greater potential for Lucas do Rio Verde, followed by the municipalities of Sinop, Sorriso and Nova Mutum (Picture 1 and Table 3).



202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217

Picture 1. Comparison between estimated yield: models 1 and 2 with the real yield obtained in the municipalities, harvest of 2014/2015

According to [14], the difference of the potential yield amongst the models is associated to the number and quality of input parameters, which constitutes lower or greater level of complexity among themselves, characterizing the determining factor among the agrometeorological models.

[14], still indicates the variation of potential yield on the location (production environment), it could possibly be related to the interaction among the facts that stress culture's growth and development, which diverge from each environment.

In this regard, the model 2 has characterized the closest potential yield, from the environment yield reality.

Table 3. Potential yield (kg ha⁻¹) estimated and real corn yield in the municipalities, harvest of 2014/2015

Municipalities	Model 1	Model 2	Real
Nova Mutum	5476,78	6618,02	6609,50
Lucas do Rio Verde	5371,33	6833,80	6516,90
Sorriso	5380,54	6805,84	6932,48
Sinop	5426,06	6808,12	6231,37
Average	5413,68	6766,45	6572,56

218
219
220
221
222
223

As Model 2 has come closer to real conditions, a comparison was carried out with the real yield obtained in the municipalities, it was noted that Nova Mutum has produced (99,87%), Lucas do Rio Verde (95,36%), Sinop (91,52%) of the average of estimated potential yield (Picture 2). Differently from all other municipalities, Sorriso estimated 1,82% less than real yield.



Picture 2. Comparison amongst estimated yield: Model 2, real yield obtained in the municipalities, harvest of 2014/2015

In terms of water conditions based on real yield (PR), which linked to the relative yield downfall ($1 - PR/PPF$) with the relative deficit of evapotranspiration ($1 - ETR/ETc$). Observing the real yield data (Table 4), It is noted that the municipality of Lucas do Rio Verde was the one which lost the most of its yield in regard of water effect, followed by the municipality of Nova Mutum which was the least to attain the real yield.

Table 4. The estimated potential yield (kg ha⁻¹) of water conditions, real yield and corn yield shortfall from different municipalities, harvest of 2014/2015.


Cities	Estimates	Real	Yield Breakage%
Nova Mutum	5987,12	6609,49	9,53
Lucas do Rio Verde	4161,33	6516,9	39,11
Sorriso	6142,63	6932,47	9,74
Sinop	5942,511	6231,36	12,71

It can be also observed at Picture 3 the water deficit effect upon corn yield for all four phenological periods. There was a major water deficit in all municipalities during the ripening stage. According to [15] the corn presents a critical stage from its pre-flowering stage to its grain filling stage. At this phenological stage, the corn is delicate to water deficit, this sensibility can be noted during the physiological processes connected to zygote development and the grain filling, and by the elevated transpiration which occurs in this period due to its large leaf area.

In Lucas do Rio Verde the water deficit occurred during the flowering stages, ripening and plant development, presenting the major yield shortfall (39,11%) in this municipality due to its water deficit (Table 4).

Probably the major yield shortfall as observed in Lucas do Rio Verde is due to its estimated seeding period for this assignment. According to [16] depending on the seeding season, the risks of yield losses can be reduced by lowering the exposure of cultivation to more plausible periods of low water availability to the plants.

Thus, the chosen period for seeding and harvesting can harm the corn yield in the municipality of Lucas do Rio Verde.



Picture 3. Relation between relative yield and relative evapotranspiration in the four corn phenological stages

4. CONCLUSION

1 The [3] model 2 has characterized the potential yield closer to the reality of production environment.

2 The difference of all potential yield of corn from all municipalities studied were all directly related to factor interaction which interferes on its growth and development, consequently the difference among these environments.

3 Taking into account the effects of water conditions, the chosen period for seeding can harm the corn yield in the municipality of Lucas do Rio Verde.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO. Report on the agro-ecological zones project 48/1-2 Vol. 1 Methodology and results for África. Vol. 2. Results for Southwest Asia. Roma: FAO, 1978. 188 p.
2. Campelo Júnior JH, Caseiro FT, Herbster OF. Zoneamento do potencial de produção de grãos em Mato Grosso. Cuiabá, UFMT, 1990. 30p.
3. Doorenbos J, Kassam AH. Efeito da água no rendimento das culturas. Estudos FAO - Irrigação e Drenagem n.33, 1994. 306p. (Traduzido por Gheyi, H.R. et al. - UFPB).
4. Barbieri V, Tuon RL. Metodologia para estimativa da produção potencial de algumas culturas. DFM/ESALQ/USP, 1992. 17p.

- 286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
5. De Wit, C.T. Photosynthesis of leaf canopies. Wageningen, Pudoc, 1965. 57p (Agricultural Research Report 663).
 6. CONAB –National Supply Company: Perspectives for Agriculture and cattle raising (2018). Accessed 24 May de 2019.
Available: <https://www.conab.gov.br/perspectivas-para-a-agropecuaria>.
 7. IMEA-Instituto Matogrossense de Economia Agropecuária: (2016) Entendendo o Mercado do Milho. Accessed 28 June de 2016.
Available: http://www.imea.com.br/upload/pdf/arquivos/Paper_jornalistas_Milho_AO.pdf.
 8. Hoogenboom G. Contribution of agrometeorology to the simulation of crop production and its application. Agricultural and Forest Meteorology, v.103, p.137-157, 2000.
 9. Cruz CJ. et al. Avaliação de Dez Híbridos Simples de Milho em Altas Densidades de Plantio. In: CONGRESSO NACIONAL DE MILHO E SORGO, 29, 2012, Águas de Lindóia. Anais... Sete lagoas: ABMS, 2012. p 2249-2256.
 10. Tejera NA, Rodés R, Ortega E, Campos R, Lluch C. Comparative analysis of physiological characteristics and yield components in sugarcane cultivars, Field Crops Research, v.102, p.64-72, 2007.
 11. Camargo MBP, Camargo AP. Representação gráfica informatizada do extrato do balanço hídrico de Thornthwaite & Mather. Bragantia, Campinas, v.52, p.169-172, 1993.
 12. Camargo AP. Balanço hídrico no Estado de São Paulo. Boletim Técnico n.116, 1971, IAC. 24p.
 13. INMET - Instituto Nacional de Meteorologia (2016).
Accessed 20 May de 2016.
Available: <http://www.inmet.gov.br/portal/index.php?r=bdmep/bdmep>.
 14. Graeff S, Link J, Binder J, Claupein W. Crop Models as Decision Support Systems in Crop Production. Crop Production Technologies, 1-28p, 2012.
 15. Bergamaschi H, et al. Distribuição hídrica no período crítico do milho e produção de grãos. Pesquisa Agropecuária Brasileira, v.39, p.831-839, 2004.
 16. Wagner MV, Jadoski SO, Marcio F, Saito LR, Lima, Adenilson dos S. Estimativa da produtividade do milho em função da disponibilidade hídrica em Guarapuava, PR, Brasil. Rev. bras. eng. agríc. ambient., Campina Grande, v. 17, n. 2, p. 170-179, Feb. 2013.