

**Fungi associated with major agricultural and forage crops in integrated systems of tropical regions**

**ABSTRACT**

Integrated production systems were developed to preserve productive resources and maintain the profitability of agribusiness. However, the use of seeds of low physiological and sanitary quality and the implantation of agricultural and forage crops in production fields of low sanitary quality may favor the dissemination and proliferation of phytopathogens such as fungi. Therefore, using the scientific literature, this work aimed to identify the fungi associated with the main agricultural and forage crops that cause damage to the integrated production systems of tropical regions and their control measures. This work was based on a literature review in the Scielo, Scopus and Google Scholar databases, with data obtained between 1999 and 2019. The keywords employed were "fungus", "tropical grass"; "agricultural crops"; "ICLS" (Integrated Crop-Livestock); and "ICLF" (Crop-Livestock-Forest) and their respective terms in Portuguese, under different combinations. For the inclusion criteria, publications (papers, books, theses, dissertations, and scientific communiqués) from 1999 to 2019 which fit the study aim were selected, both in the Portuguese and English languages. The publications that did not meet the criteria of this study and were repeated in databases were considered as exclusion criteria. The main fungi associated with forage and agricultural crops and soils of integrated systems of tropical regions are *Bipolaris* sp., *Curvularia* sp., *Exserohilum* syn. *Helminthosporium* sp., *Phoma* sp., *Fusarium* sp., *Macrophomina* sp., *Pythium* sp., *Rhizoctonia* sp. and *Sclerotium* sp. The main methods of fungal control are the use of quality seeds, crop rotation, resistant cultivars, and chemical seed treatment.

*Keywords: Crop rotation, fungicides, resistant cultivars, sustainable agriculture, tropical grass*

**1. INTRODUCTION**

Agribusiness is worldwide in importance, providing nutrients and contributing significantly to the world economy. Due to the widespread essentiality of agribusiness, there is a need to keep the systems that compose it increasingly productive while preserving the land and reducing the need to open new areas. In order to achieve this purpose, sustainable cultivation models have been created and perfected, such as Integrated Crop-Livestock (ICLS) and Crop-Livestock-Forest (ICLF) systems. These systems aim at the maximum use of the land without degradation, improvement in the physical-chemical quality of the soil, in the zootechnical indexes of the animal component and income diversification [1, 2, 3, 4, 5, 6, 7].

Both of the above-mentioned integrated systems (ICLS and ICLF), if poorly managed, might suffer from pest and disease attacks. In addition, the implantation of low sanitary quality

28 crops in production fields, culminating in the use of seeds with low physiological and sanitary  
29 quality may favor the dissemination and proliferation of pests and phytopathogens in  
30 productive areas, reducing the yield of agricultural and forage crops and, consequently,  
31 affecting animal performance [8, 9, 10].

32 Among the phytopathogens that can affect productive areas, fungi and nematodes are the  
33 ones that cause most concern as they can decimate crops when in high incidence, being  
34 difficult to eradicate from the production system [10].

35 In order to adopt fungal control measures, aiming at the maximum yield of integrated  
36 systems in tropical regions, it is necessary to know the fungal incidence in the production  
37 fields of the main crops used in these systems, considering that the cultivation of forages  
38 belonging to the genera *Brachiaria* syn. *Urochloa* and *Panicum*, as well as agricultural crops  
39 such as soybean, maize, sorghum, and millet, are predominant in integrated production  
40 systems [11, 12, 13, 14, 15, 16, 17, 18].

41 Based on this, this study aimed to identify the fungi associated with the main agricultural and  
42 forage crops that damage the integrated production systems of tropical regions, as well as  
43 their control measures.

## 44 **2. METHODOLOGY**

45  
46 This work was based on a literature review in the Scielo, Scopus and Google Scholar  
47 databases, with data between 1999 and 2019.

48 The keywords employed were “fungus”, “tropical grass”; “agricultural crops”; “ICLS”; and  
49 “ICLF”; and their respective terms in Portuguese “fungos”, “capins tropicais”; “culturas  
50 agrícolas”; “iLP”; and “iLPF”, under different combinations.

51 For inclusion criteria, publications (papers, books, theses, dissertations, and scientific  
52 communiqués) from 1999 to 2019 which fit the study aim were selected, both in the  
53 Portuguese and English languages. Publications prior to 1999, that did not meet the criteria  
54 of this study (analyzed by titles and abstract) and were repeated in the databases were  
55 considered as exclusion criteria.

56 After reading the titles of articles and other publications, we selected 100 publications that  
57 met the initially proposed criteria, which were read in full.

## 58 **3. RESULTS AND DISCUSSION**

59 Fungi are phytopathogenic agents at higher rates of association with seeds, presenting  
60 longevity in the productive system through the production of resistance structures [19, 20].  
61 They are among the main causes of diseases in forage plants, causing losses in the yield  
62 and quality of the green mass produced, besides reducing the quality (germination and  
63 vigor) of the seeds [8, 9, 21, 22, 23, 24, 25, 26].

64 Among the fungi present in soils used in integrated systems, and the fungi associated with  
65 *Brachiaria* syn. *Urochloa* sp., *Panicum* sp. and main agricultural crops, there are  
66 phytopathogenic taxa belonging to the genera *Bipolaris* sp., *Curvularia* sp., *Exserohilum* syn.  
67 *Helminthosporium* sp., *Phoma* sp., *Fusarium* sp., *Macrophomina* sp., *Pythium* sp.,  
68 *Rhizoctonia* sp., and *Sclerotium* sp.. There are also secondary fungi such as *Alternaria* sp.,  
69 *Aspergillus* sp., *Cladosporium* sp., *Epicoccum* sp., *Nigrospora* sp., *Penicillium* sp., and  
70 *Trichoderma* sp. [8, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36].

71 ***Bipolaris sp.***

72 *Bipolaris* sp. is a pathogenic fungus with a 100% seed transmission rate to the seedlings. It  
73 is the main fungus that attacks *Panicum* sp., causing leaf spot disease. However, this fungus  
74 can be associated with the integument and interior of the seeds and with crop residues of  
75 susceptible crops, such as tropical grasses, maize, wheat, rice, and coffee. In addition, it is  
76 favored by temperatures between 22 and 30 °C [24, 25, 37, 38, 39, 40, 41, 42, 43, 44, 45].

77 The disease and its symptoms are similar in susceptible cultures, being manifested as brown  
78 spots and coalescence of the lesions in episodes of severe infestation. These symptoms can  
79 be observed 50 days after the sowing of susceptible crops in infected areas and/or using  
80 contaminated seeds [37, 38, 46]. In addition to the symptoms of contamination by  
81 phytopathogens in tropical plants, it causes disturbances in animals and humans, such as  
82 allergies, pulmonary and cutaneous infections [47].

83 Listing as alternatives to control the disease, the use of resistant cultivars and fungicides is  
84 commonly employed. The application of pyraclostrobin associated with epoxyconazole or  
85 tebuconazole alone has been promising for the reduction of the disease intensity in *P.*  
86 *maximum* seed production fields, increasing the speed of germination. In addition, cultural  
87 control, such as crop rotation, is an effective measure for disease control, since it reduces  
88 the initial inoculum [37, 38, 48].

89 Furthermore, the fungal control can be performed by means of seed treatment with  
90 fungicides, aiming to reduce the incidence of fungi in the seeds and in the soil, especially in  
91 the initial stages of development of the susceptible plants, besides avoiding the introduction  
92 or re-inoculation of phytopathogens [49, 50].

93 ***Curvularia sp.***

94 *Curvularia* sp. is a pathogenic fungus found in several regions of the world that has an  
95 important incidence in forage plants of the *P. maximum* species, causing the leaf spot  
96 disease [24, 25, 51]. Furthermore, it can cause rotting, reduction of germination up to seed  
97 unfeasibility (associating to the integument and endosperm), and death of seedlings in  
98 susceptible species, such as tropical grasses and maize, **reproducing asexually by**  
99 **ascomyces** [28, 37, 39, 52, 53, 54, 55].

100 In addition to the diseases that it causes in susceptible plants, the fungus can cause allergic  
101 conditions, endocarditis, pheochycolosis, mycetoma, onychomycosis, keratitis, brain  
102 abscesses, urinary and pulmonary infections, and infectious wounds in animals and humans  
103 [56, 57, 58, 59, 60, 61].

104 For the *Curvularia* sp. control it is possible to employ a seed treatment with the thiram or  
105 thiram-associated carboxin fungicides, which also have efficiency in the control of other  
106 fungi, such as *Alternaria* sp., *Gerlachia* sp. and *Drechslera* sp.; as well as the fungicides  
107 fludioxonil combined with metalaxyl-M, colorless fludioxonil combined with metalaxyl-M,  
108 and thiram alone, which also have fungus control efficiency on *Penicillium* sp., *Alternaria* sp.,  
109 *Drechslera* sp., and so forth. In addition, the association of thiram and thiabendazole  
110 fungicides in seed treatment may promote a greater emergence of seedlings [28, 62, 63].

111 ***Exserohilum syn. Helminthosporium sp.***

112 Such as the fungus *Curvularia* sp., the fungus ***Exserohilum syn. Helminthosporium*** sp.  
113 causes rotting, seed unfeasibility and death of seedlings, as well as leaf and stem spot

114 diseases. The manifested symptoms are necrotic and elliptic lesions with intense sporulation  
115 of the pathogen [9, 37, 64].

116 It survives in crop remains (saprophytic form) and in the soil, forming resistance structures  
117 and associating to seeds (integument and endosperm) and hosts, such as tropical grasses,  
118 sorghum and maize (causing helminthosporiosis). In addition, the fungus is favored by  
119 temperatures between 18 and 27 °C and high moisture. The main control method is through  
120 genetic resistance and crop rotation with resistant cultures [39, 41, 42, 43, 44, 64].

#### 121 ***Phoma* sp.**

122 The pathogenic fungus *Phoma* sp. affects several crops from many continents, such as  
123 soybean and coffee. In the European region, the main etiological agent in soybean is *Phoma*  
124 *pinodella* (= *Phoma sojicola*). The causal agent in Brazil is the fungus *Phyllosticta sojicola*  
125 [35, 65, 66, 67].

126 Furthermore, this fungus also focuses on forages of the genus *P. maximum*. The symptoms  
127 depend on the severity of the incidence, beginning four or five days after the inoculation,  
128 causing leaf spot with elongated, necrotic and irregular characteristics, rotting, seed  
129 unfeasibility and death of seedlings. By possessing fast and aggressive growth, it can even  
130 kill infected seeds before germination [25, 37, 51, 68]. The fungus belongs to the group of  
131 mitosporic fungi and is a Coelomycete, presenting cylindrical, hyaline and small non-septate  
132 conidia [69].

133 An efficient alternative to control this fungus is the treatment of seeds with thiram fungicides,  
134 thiram associated to carboxin, fludioxonil combined with metalaxyl-M, colorless fludioxonil  
135 comminuted with metalaxyl-M, and thiram alone [28, 62, 63].

#### 136 ***Fusarium* sp.**

137 Fungi of the genus *Fusarium* sp. cause damage to agricultural production systems  
138 composed of several agricultural crops, tropical grasses, and animals. This occurs due to the  
139 incidence of fusariosis and production of mycotoxin. The symptom of fusariosis in plants is  
140 dependent on the phytopathogen species and on the interspecific relation between host-  
141 phytopathogen. Mycotoxins can cause symptoms such as false heat, abortion, stillbirths,  
142 infertility, problems in the digestive system, bleeding, anemia due to the destruction of the  
143 bone marrow, vomiting, necrosis of the epidermis and death of the animals [70, 71, 72].

144 The *F. solani* species causes red root rot in the soybean crop, a symptom of which is the  
145 rotting of the root system; *F. moniliforme* causes the fusarium rot disease in crops of cotton,  
146 rice, maize, sorghum, and tropical grasses, in addition to the potential for intoxication of  
147 animals due to the production of mycotoxins (zearalenone, fumonisins, and vomitoxins or  
148 deoxynivalenol). *F. graminearum*, *F. equiseti*, and *F. tricinctum* also produce mycotoxins in  
149 maize, sorghum, soybean, wheat, and oat crops, as well as *F. pallidoroseum*, which can  
150 break the stem and lead to the tipping of the cotton, beans and soybean plants [70, 73].

151 Furthermore, tropical grass seeds susceptible to phytopathogens may increase the inoculum  
152 potential in the area and act as a reservoir for future dissemination in crops that will succeed  
153 in the area, such as pine, cotton, wheat, rice, bean, soybean, maize, sugarcane, and so  
154 forth. The incidence level can be influenced by ideal climatic conditions of high temperature  
155 and soil moisture [24, 25, 34].

156 An effective measure for the control of these phytopathogens is the chemical treatment of  
157 seeds with fungicides. Among the available options in the commercialization, as previously  
158 mentioned for the control of *Curvularia* sp. and *Phoma* sp, the use of thiram or carboxin  
159 associated with thiram is highlighted, besides the fungicides fludioxonil combined with  
160 metalaxyl-M, colorless fludioxonil combined with metalaxyl-M, and thiram alone. In the  
161 absence of resistant cultivars, well-drained and fertilized soils and healthy and certified  
162 seeds can be employed [28, 34, 62, 63].

### 163 ***Macrophomina* sp.**

164 The genus *Macrophomina* sp. inhabits the soils and manages to multiply in vegetal remains  
165 by means of its propagation structures produced by the mycelium (microsclerocios). These  
166 structures are resistant to adverse conditions for long periods in the soil. The ideal conditions  
167 for phytopathogens to develop are high temperatures and moisture. Moreover, after the  
168 insertion into the production area, it attacks roots, stems, leaves, and fruits of susceptible  
169 crops, such as sunflower, cotton, sorghum, maize, soybean, and bean, among others. The  
170 main species, *M. phaseolina*, triggers symptoms corresponding to grayish lesions that may  
171 evolve to rot and tissue destruction. On the other hand, this tissue disruption causes  
172 chlorosis, wilt, drought, and death of susceptible plants [34, 74].

173 The broad spectrum of susceptible species and the absence of resistant cultivars hinder the  
174 control through crop rotation, although the performing of this practice with forage grasses is  
175 recommended. Also, it is always recommended to use healthy and certified seeds [34].

176 **It is observed** that soybean plants produce the phytoalexin gliceolin when infected by *M.*  
177 *phaseolina*, a compound that has the potential to restrict the development of the mentioned  
178 fungus through the rapid biosynthesis of glycerol by the plant during fungal infection. This  
179 feature may promote genetic improvement programs in the search for resistant plants to *M.*  
180 *phaseolina*. As a palliative measure to reduce the incidence of these phytopathogens, it is  
181 possible to use cultivars with higher tolerance to drought and/or high temperatures [75, 76].

### 182 ***Pythium* sp.**

183 This pathogen is an inhabitant of the soil which can infect seeds and seedlings. It can be  
184 associated with plant remains (saprophytes) or susceptible plants, such as soybean,  
185 sorghum, cotton, bean, maize, wheat and tropical grasses. The fungus presents resistance  
186 structures (oospores) that allow its survival in adverse conditions [73, 77].

187 **The most frequent species are *P. graminicola* and *P. debaryanum*, but *P. ultimum* causes**  
188 **the most impact. Due, the lesions develop rapidly causing chlorosis, growth reduction, wilt,**  
189 **root rot and tipping of the plant in ideal conditions of development (high moisture and mild**  
190 **temperature) [73, 77].**

191 Furthermore, it also causes economic losses to the agricultural production system since it  
192 contributes to the onset of sudden death (root rot and plant tipping) of the Marandu grass (*B.*  
193 *syn. Urochloa brizantha* cv. Marandu), along with *Rhizoctonia* sp., *Fusarium* sp. and water  
194 stress [37, 78].

### 195 ***Rhizoctonia* sp.**

196 The fungus *Rhizoctonia* is a saprophytic fungus, being able to exert parasitism on several  
197 crops and animals. It presents a high gene flow, genetic diversity, sexual reproduction and

198 dispersion of clones with high adaptability, as well as producing resistance structures  
199 (sclerotia) which remain in the soil for long periods [73, 79, 80].

200 The *R. solani* species can cause leaf burning, collection rot and death in tropical grasses,  
201 soybean and maize; in soybean, it causes plant tipping and root rot, which may reduce the  
202 vigor and germination rate of the seeds, as well as toxins that inhibit plant growth. In cotton,  
203 the fungus causes the tipping of the plants [73, 79, 80, 81].

204 The methods that have effectiveness in controlling the fungi living in the soil, such as  
205 *Pythium* sp. and *Rhizoctonia* sp., are based on the chemical treatment of seeds, rotation of  
206 crops with resistant species (grasses) and elimination of crop residues. With regard to the  
207 biological control, isolates of the fungus *Trichoderma* spp. have effectiveness in the control  
208 of *R. solani* [34, 82, 83].

#### 209 ***Sclerotium* sp.**

210 The main species, *Sclerotium rolfsii*, lives in the soil and affects crops of soybean, bean,  
211 potato, and tomato, among others. It causes the rotting of roots and colon, wilt and tipping of  
212 infected plants. The symptoms are manifested in the region of the lap of the plant and  
213 correspond to dark spots that originate the cortical rot. This rot can be identified by the  
214 formation of a white mycelium and brown-colored resistance structures (sclerocytes). The  
215 destruction of tissues occurs under these structures and, with that, wilt, drought and death of  
216 the plants. The ideal development conditions occur in regions of tropical climate, with  
217 temperatures within 25 and 35 °C and soil moisture in 70% of the field capacity [34, 73, 84].

218 As a control method, the need to use healthy and certified seeds, the elimination of crop  
219 residues and crop rotation with maize and cotton (resistant plants) are highlighted, as well as  
220 the efficiency of the fungicide tebuconazole in the colony growth and in the germination of  
221 sclerotia of *S. rolfsii* with regard to the chemical control [34, 85, 86].

222 In general, for the fungal control, it is always necessary to employ seeds with high  
223 physiological and sanitary quality, as well as crop rotation, resistant cultivars and chemical  
224 treatment of seeds with fungicides. Seed treatment has a low cost and can improve seed  
225 germination and seedling development. However, for effective seed treatment and fungal  
226 control, effective fungicides are necessary [34, 49, 87, 88].

#### 227 **Occurrence regions and economic impacts**

228 In Brazil, *Bipolaris* sp. is more frequent in the Southern region, although with severity ranging  
229 from low to average. However, *Exserohilum* syn. *Helminthosporium* sp. can reduce up to  
230 50% of the production, primarily in the second harvest of sorghum, in the Brazilian states of  
231 São Paulo, Pernambuco, and Distrito Federal, and of maize and tropical grasses in the  
232 Brazilian states of São Paulo and Minas Gerais, respectively [89, 90, 91]. At world level,  
233 *Exserohilum* syn. *Helminthosporium* sp. can also be found in the Dominican Republic [92].

234 The fungus *Curvularia* sp. in tropical grasses can be found in the Brazilian states of Mato  
235 Grosso, Minas Gerais, and São Paulo, as well as in maize crops in the states of Mato  
236 Grosso, Mato Grosso do Sul and Pernambuco [91]. The worldwide distribution of this fungus  
237 occurs in tropical and subtropical regions [93]. *Phoma* sp. occurs both in the European  
238 region and in Brazil [35, 65, 66, 67]. Reported in Honduras, Costa Rica, Panama, and  
239 Colombia in the 1960s, *Fusarium* sp., has primarily reached the Northeast region of Brazil  
240 [94].

241 The occurrence of *Macrophomina* sp. associated with soybean crops can be found in the  
242 Center-West region of Brazil, besides the states of Paraná, Rio Grande do Sul, São Paulo,  
243 Minas Gerais, and Maranhão. The association with cotton is reported in São Paulo, Paraíba,  
244 Pernambuco, Minas Gerais, and Paraná. The association with bean crops might occur in the  
245 entire Brazilian territory, except for the states of Amazonas, Roraima, Amapá, Rondônia,  
246 and Acre. The association with maize might occur in the South, Southeast and Central-West  
247 regions, besides the states of Pernambuco and Bahia. The association with sunflower is  
248 verified in Mato Grosso, São Paulo, and Paraná. The association with sorghum is reported in  
249 the Brazilian Northeast region and in the state of Rio Grande do Sul [91]. It can also be  
250 verified causing damage in Venezuelan crops [92].

251 The presence of *Rhizoctonia* sp. associated with tropical grasses can be verified in the  
252 Brazilian states of Pará and Mato Grosso; in the soybean crop it presents a wide distribution  
253 throughout the Brazilian territory, except in the states of Goiás, Espírito Santo, and Rio de  
254 Janeiro; as for cotton crops, it only occurs in the state of Paraná [91]. There are also  
255 incidence and damage reports of *Rhizoctonia* sp. and *Pythium* sp. in the state of Florida, in  
256 the United States of America [92]. Furthermore, the fungus *Rhizoctonia* sp. might cause  
257 economic impacts through the reduction of the initial plant population, generating the need  
258 for resowing, which costs 6.44% of the total production cost [81]. The occurrence of  
259 *Sclerotium* sp. has increased in the last few years, and it has been isolated from several  
260 locations in the Mid-North region of Brazil [34].

261 Generally, the severe incidence of fungi might lead to the increase in the production costs  
262 through the need for resowing due to damage in the initial stand of the crops [81]; and  
263 through the need for the application of chemical (via seed and leaves) and biological  
264 treatments. Furthermore, it negatively affects the crop yield since it causes a reduction in the  
265 production of green mass and in the photosynthetic ability, with a reflection of these effects  
266 in the production and quality of the grains. The percentages of cost increase and yield  
267 reduction in the crops are variable and dependent on the level of incidence by the  
268 phytopathogens, interspecific interaction between pathogen and host, disease severity and  
269 abiotic conditions [95, 96, 97, 98, 99, 100].

270 Briefly, based on the scientific findings presented in this paper, the following table was  
271 generated (Table 1).

272 **Table 1.** Most occurring regions, susceptible crops, damages and control of the fungi discussed in this paper.

<b>Fungi</b>	<b>Regions</b>	<b>Susceptible crops</b>	<b>Damages</b>	<b>Controls</b>
<b><i>Bipolaris sp.</i></b>	Regions with a temperatures between 22 and 30 °C; Brazil	Tropical grasses, maize, wheat, rice and coffee	In plants: brown spots and coalescence of the lesions; In animals and humans: allergies, pulmonary and cutaneous infections	Resistant cultivars; Fungicides tebuconazole; pyraclostrobin with epoxyconazole
<b><i>Curvularia sp.</i></b>	Found in several regions of the world; tropical and subtropical regions; Brazil	Tropical grasses and maize	In plants: leaf spot disease; In animals and humans: allergic conditions, urinary and pulmonary infections, among others	Fungicides thiram; thiram with carboxin; fludioxonil with metalaxyl-M and thiabendazole.
<b><i>Exserohilum sp.</i></b>	Regions with a temperatures between 18 and 27 °C and high moisture; Brazil and Dominican Republic	Tropical grasses, sorghum and maize	In plants: leaf and stem spot diseases, rotting, seed unfeasibility and death of seedlings	Genetic resistance; crop rotation
<b><i>Phoma sp.</i></b>	Brazil and European region	Soybean and coffee	In plants: leaf spot, rotting, seed unfeasibility and death of seedlings	Fungicides thiram; thiram with carboxin and fludioxonil with metalaxyl-M
<b><i>Fusarium sp.</i></b>	Regions with high temperature and soil moisture; Honduras, Costa Rica, Panama, Colombia and Brazil	Pine, cotton, wheat, rice, bean, soybean, maize, sugarcane, sorghum, and tropical grasses	In plants: red root rot and fusarium rot disease; In animals: intoxication due mycotoxins	Resistant cultivars; healthy and certified seeds; Fungicides thiram; carboxin with thiram and fludioxonil with metalaxyl-M
<b><i>Macrophomina sp.</i></b>	Regions with high temperatures and moisture; Brazil and Venezuela	Sunflower, cotton, sorghum, maize, soybean and bean	In plants: rot and tissue destruction of roots, stems, leaves, and fruits	Cultivars with higher tolerance to drought and/or high temperatures
<b><i>Pythium sp.</i></b>	Regions with high moisture and mild temperature; Brazil and United States of America	Soybean, sorghum, cotton, bean, maize, wheat and tropical grasses	In plants: chlorosis, growth reduction, wilt, root rot and tipping	Chemical treatment of seeds; rotation of crops; elimination of crop residues; biological control with <i>Trichoderma spp.</i>
<b><i>Rhizoctonia sp.</i></b>	Brazil and United States of America	Tropical grasses, soybean, maize, and cotton	In plants: leaf burning, rot, tipping, inhibit growth and death	Chemical treatment of seeds; rotation of crops; elimination of crop residues; biological control with <i>Trichoderma spp.</i>
<b><i>Sclerotium sp.</i></b>	Regions with temperatures within 25 and 35 °C; Brazil	Soybean, bean, potato, and tomato	In plants: cortical rot, wilt, drought and death	Healthy and certified seeds; elimination of crop residues; crop rotation; fungicide tebuconazole



273 **4. CONCLUSION**

274

275 The main fungi associated with forage and agricultural crops and soils of integrated systems  
276 in tropical regions are *Bipolaris* sp., *Curvularia* sp., *Exserohilum* syn. *Helminthosporium* sp.,  
277 *Phoma* sp., *Fusarium* sp., *Macrophomina* sp., *Pythium* sp., *Rhizoctonia* sp. and *Sclerotium*  
278 sp.

279 The main methods of fungal control involve the employment of high-quality seeds, crop  
280 rotation, resistant cultivars and chemical seed treatment.

281 **COMPETING INTERESTS**

282

283 We declare that no competing interests exist.

284

285 **REFERENCES**

286

287 1. Almeida RG, Ramos AKB, Zimmer AH, Machado LAZ, Kichel NA, Gontijo Neto MM et al.  
288 Performance of tropical forages in crop-livestock and crop-livestock-forest integration  
289 systems. In: Cordeiro LAM, Vilela L, Kluthcouski J, Marchão RL, editors. **Crop-livestock-**  
290 **forest integration: the producer asks, Embrapa responds.** 1st ed. Brasília: Embrapa; 2015.  
291 Portuguese.

292

293 2. Balbino LC, Cordeiro LAM, Porfírio-da-Silva V, Moraes A, Gladys BM, Alvarenga RC et al.  
294 Technological evolution and productive arrangements of crop-livestock-forestry integration  
295 systems in Brazil. *Pesquisa Agropecuária Brasileira*, 2011;46(10):1-12. Portuguese DOI:  
296 10.1590/S0100-204X2011001000001

297

298 **3. Costa** JL, Rava CA. Influence of *Brachiaria* in the management of bean diseases with soil  
299 origin. In: Kluthcouski J, Stone LF, Aidar H, editors. *Crop-livestock integration: 1st ed.* Santo  
300 Antônio de Goiás: Embrapa; 2003. Portuguese.

301

302 4. Leonel FP. Crop-livestock integration: production and forages quality. In: Zervoudakis JT,  
303 Cabral LS. *Nutrition and production of beef cattle.* 1st ed. Cuiabá: UFMT; 2011. Portuguese.

304

305 5. Macedo MCM. Crop and livestock integration: the state of the art and the near future.  
306 *Revista Brasileira de Zootecnia.* 2009;38(spe):133-146. Portuguese. DOI: 10.1590/S1516-  
307 35982009001300015

308

309 6. Mello, LMM. Crop-livestock integration in no-till system. In: *Brazilian Congress of Soil*  
310 *Science. Annals.* Viçosa: Brazilian Society of Soil Science; 2003. Portuguese.

311

312 7. Vilela L, Martha Junior GB, Macedo MCM, Marchão RL, Guimarães Júnior R, Pulrolnik K  
313 et al. Integrated crop-livestock systems in the Cerrado region. *Pesquisa Agropecuária.*  
314 2011;46:1127-1138. **Portuguese.** DOI: 10.1590/S0100-204X2011001000003

315

316 8. Vechiato MH, Aparecido CC. Fungi in forage grasses seeds: phytosanitary restriction and  
317 detection methods. São Paulo: Biological Institute of São Paulo; 2008. Portuguese.

318

319 9. Vechiato MH, Aparecido CC, Fernandes CD. CD. Frequency of fungi in batches of  
320 commercialized *Brachiaria* and *Panicum* seeds. Campo Grande: Embrapa; 2010.  
321 Portuguese

322

- 323 10. Fernandes CD. The sanity of forage seeds and their impact on crop-livestock-forestry  
324 integration systems. In: Brazilian Congress of Seeds, 18. The seed in agricultural  
325 productivity and conservation of genetic resources: abstracts. Florianópolis: Brazilian  
326 Association of Seed Technology; 2013. Portuguese.  
327
- 328 11. Embrapa Sede. Technologic solutions: *Brachiaria brizantha* cv. Marandu. Accessed 12  
329 Mar 2019. Available: [https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/863/brachiaria-brizanthacv-marandu)  
330 [servico/863/brachiaria-brizanthacv-marandu](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/863/brachiaria-brizanthacv-marandu).  
331
- 332 12. Semeata. *Brachiaria ruziziensis*. Accessed 02 Mar 2019. Available:  
333 <http://www.semeata.com.br/?sessao=produto&folder=1&ver&id=7>.  
334
- 335 13. Pedreira CGS, Moura JC, Silva SC, Faria VP. In: Pastures and the environment. 23°  
336 Symposium on Pasture Management. Annals. Piracicaba: 23° Symposium on Pasture  
337 Management; 2006: 165-185. Portuguese  
338
- 339 14. Embrapa Sede. Technologic solutions: *Panicum maximum* cv. Mombaça. Accessed 13  
340 Mar 2019. Available: [https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/882/panicum-maximumcv-mombaca)  
341 [servico/882/panicum-maximumcv-mombaca](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/882/panicum-maximumcv-mombaca).  
342
- 343 15. Embrapa Sede. Technologic solutions: *Panicum maximum* cv. Tanzânia-1. Accessed 13  
344 Mar 2019. Available: [https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/885/panicum-maximumcv-tanzania-1)  
345 [servico/885/panicum-maximumcv-tanzania-1](https://www.embrapa.br/busca-de-solucoes-tecnologicas/-/produto-servico/885/panicum-maximumcv-tanzania-1).  
346
- 347 16. Alvarenga RC, Gontijo Neto MM, Ramalho JH, Garcia JC, Viana MCM, Castro AADN.  
348 Crop-Livestock Integration System: the model implemented in Embrapa Milho e Sorgo. Sete  
349 Lagoas: Embrapa; 2007. Portuguese  
350
- 351 17. Alvarenga RC, Rodrigues JAS, Santos FC, Gontijo Neto MM, Viana MCM. Sorghum  
352 cultivation in integrated crop-livestock or crop-livestock-forestry systems. Sete Lagoas:  
353 Embrapa; 2011.  
354
- 355 18. Rodrigues JAS, Cruz JC. Growing millet. Sete Lagoas: Embrapa; 2009. Portuguese.  
356
- 357 19. Ministério da Agricultura, Pecuária e Abastecimento. Rules for seed analysis. Brasília:  
358 MAPA/ACS; 2009. Portuguese  
359
- 360 20. Kruppa PC, Russomano OMR. Fungi in medicinal, aromatic and condimentary plants -  
361 soil and seed. 2009. Portuguese. Accessed 14 Jun 2019. Available:  
362 [http://www.infobibos.com/Artigos/2009\\_1/Fungos/index.htm](http://www.infobibos.com/Artigos/2009_1/Fungos/index.htm)>. Acesso em 14 de junho 2019.  
363
- 364 21. Lucca-Filho OA, Porto MDM, Maia MS. Fungi in Ryegrass-annual seeds (*Lolium*  
365 *multiflorum* Lam.) e and its effects on the establishment of the pasture. Revista Brasileira de  
366 Sementes. 1999;21(2):142-147. Portuguese. DOI: 10.17801/0101-3122/rbs.v21n2p142-147  
367
- 368 22. Schafer M, Kotanen PM. The influence of soil moisture on losses of buried seeds to  
369 fungi. Acta Oecologica. 2003;24(5-6):255-263. DOI: 10.1016/j.actao.2003.09.001  
370
- 371 23. Mertz LM, Henning FA, Maia MS, Meneghello GE, Henriques A, Madail R. Physiological  
372 and sanitary quality of cawpea seeds processed in gravity table. Revista Brasileira de  
373 Sementes. 2007;29(3):1-7. Portuguese. DOI: 10.1590/S0101-31222007000300001  
374

- 375 24. Marchi CE, Fernandes CD, Bueno ML, Batista MV, Fabris LR. Fungi associated to  
376 comercial seed of braquiária grass. Arquivos do Instituto Biológico. 2010;77(1):65-73.  
377 Portuguese.
- 378
- 379 25. Mallmann G, Verzignassi JR, Fernandes CD, Santos JM, Vechiato MH, Inácio CA et al.  
380 Fungi and nematodes associated with tropical forage seeds. Summa Phytopathologica.  
381 2013;39(3):201-203. Portuguese. DOI: 10.1590/S0100-54052013000300010  
382
- 383 26. Marcos MF, Jank L, Fernandes CD, Verzignassi, JR, Mallmann G, Queiroz CA et al.  
384 Reaction to *Bipolaris maydis*, the causal agent of leaf spot, in apomictic hybrids of *Panicum*  
385 *maximum*. Summa Phytopathologica. 2015;41(3):197-201. Portuguese. DOI: 10.1590/0100-  
386 5405/2078  
387
- 388 27. Vechiato MH. Sanity of grass forages seeds. In: Brazilian Symposium on Seed  
389 Pathology, 8. Annals. Londrina: Abrates; 2004. Portuguese.  
390
- 391 28. Lasca CC, Vechiato MH, Kohara EY. Control of *Brachiaria* spp. seed-borne fungi:  
392 effectiveness of fungicide and influence of the storage period of treated seeds on the action  
393 of these products. Arquivos do Instituto Biológico. 2004;71(4):465-472. Portuguese.  
394
- 395 29. Fernandes CD, Marchi CE, Jerba VF, Borges MF. Pathogens associated with tropical  
396 forage seeds and control strategies. In: Zambolim L. Seeds, sanitary quality. Viçosa: Federal  
397 University of Viçosa; 2005. Portuguese.  
398
- 399 30. Marchi CE, Fernandes CD, Jerba VF, Trentin RA, Bueno ML, Guimarães LR et al.  
400 Tropical forage seeds: associated pathogens and control strategies. In: Brazilian  
401 Symposium on Seed Pathology, 10. Annals. Londrina: Abrates; 2006. Portuguese.  
402
- 403 31. Marchi CE, Fernandes CD, Borges CT, Santos JM, Jerbal VF, Trentin RA et al.  
404 Phytopathogenic nematofauna of commercial seeds of tropical forages. Pesquisa  
405 Agropecuária Brasileira. 2007;42(5):655-660. Portuguese. DOI: 10.1590/S0100-  
406 204X2007000500007  
407
- 408 32. Witt FAP, Oliveira FF, Takeshita V, Ribeiro LFC. Quality of health seeds the *Urochloa*  
409 and *Panicum* sold in matogrossense north. Enciclopédia Biosfera. 2015;11(21):1636-1645.  
410 Portuguese.  
411
- 412 33. Silva AEL, Reis EM, Tonin RFB, Danelli ALD, Avozani A. Identification and quantification  
413 of fungi associated with seeds of ryegrass (*Lolium multiflorum* Lam.). Summa  
414 Phytopathologica. 2014;40(2):156-162. Portuguese. DOI: 10.1590/0100-5405/1935  
415
- 416 34. Athayde Sobrinho C, Belmino CS. Feijão-Caupi: Fungi. Accessed 14 Jun 2019. Available:  
417 [https://www.agencia.cnptia.embrapa.br/gestor/fejao-caupi/arvore/CONTAG01\\_59\\_51020068](https://www.agencia.cnptia.embrapa.br/gestor/fejao-caupi/arvore/CONTAG01_59_510200683537.html)  
418 3537.html.  
419
- 420 35. Vieira Junior JR, Fernandes CDF, Silva DSG, Santana L, Minosso S, Almeida U et al.  
421 Isolated fungi from *Brachiaria* sp. associated with sintoms type "brachiaria disease". In:  
422 Brazilian Congress of Phytopathology, 44. Annals. Brasília: Tropical Plant Pathology; 2011.  
423 Portuguese  
424
- 425 36. Embrapa Soja. Soybean production technologies – Central Region of Brazil 2009. 1st  
426 ed. Londrina: Embrapa; 2008. Portuguese.  
427

- 428 37. Marchi CE, Fernandes CD, Verzignassi JR. Diseases in forage plants. Campo Grande:  
429 Embrapa; 2011. Portuguese.  
430
- 431 38. Costa RV, Silva DD, Cota LV. Corn bipolares leaf spot. Sete Lagoas: Embrapa; 2014.  
432
- 433 39. Santos GR, Tschoeke PH, Silva LG, Silveira MCAC, Reis HB, Brito DR et al. Sanitary  
434 analysis, transmission and pathogenicity of fungi associated with forage plant seeds in  
435 tropical regions of Brazil. Journal of Seed Science. 2014;36(1):54-62. DOI: 10.1590/S2317-  
436 15372014000100007  
437
- 438 40. Tavanti TR, Takada J, Ribeiro LFC, Moraes SRG, Pedreira BC. Occurrence of *Bipolaris*  
439 *maydis* leaf spot on tanzania guineagrass in the north region of the Mato Grosso state.  
440 Revista de Ciências Agroambientais. 2016;14(1):82-85. Portuguese. DOI:  
441
- 442 41. Amorim L, Rezende JAM, Bergamin-Filho A, editors. Phytopathology manual. 4th ed.  
443 Piracicaba: Agronômica Ceres; 2011. Portuguese.  
444
- 445 42. Bueno CJ, Ambrósio MCQ, Souza NL. Production and evaluation of survival of resistance  
446 structures of soilborne phytopathogenic fungi. Summa Phytopathologica. 2007;33(1):47-55,  
447 2007. Portuguese. DOI: 10.1590/S0100-54052007000100007  
448
- 449 43. Danelli AD, Viana E, Fiallos FG. Pathogenic fungi found in wheat seeds and medium  
450 early maturity, produced in three parts of Rio Grande do Sul, Brazil. Scientia Agropecuaria.  
451 2012;3(1):67-74. Portuguese.  
452
- 453 44. Quadros DG, Andrade AP, Oliveira GC, Oliveira EP, Moscon ES. Components of seed  
454 production of two cultivars of *Brachiaria brizantha* (Hochst. ex A. Rich.) Stapf haverst by  
455 manual or mechanical sweeping. Semina Ciências Agrárias. 2012;33(5):2019-2028.  
456 Portuguese. DOI: 10.5433/1679-0359.2012v33n5p2019  
457
- 458 45. Caligiorne RB, Resende MA, Melillo PHC, Peluso CP, Carmo FHS, Azevedo V. In vitro  
459 susceptibility of chromoblastomycosis and phaeohyphomycosis agents to antifungal drugs.  
460 Medical Mycology. 1999;37(6): 405-409. DOI: 10.1111/j.1365-280X.1999.00245.x  
461
- 462 46. Michalski V, Charchar Jd'A, Anjos JRN, Fernandes FD, Silva MS, Silva WAM.  
463 Transmission of seeds fungi to *Panicum maximum* seedlings. In: Faleiro FG, Andrade SEM.  
464 Meeting of young talents from Embrapa Cerrados abstracts presented, 3. Planaltina:  
465 Embrapa; 2007. Portuguese.  
466
- 467 47. Fryen A, Mayser P, Glanz H, Füssle R, Breithaupt H, Hoog GS. Allergic fungal sinusitis  
468 caused by *Bipolaris (Drecheslera) hawaiiensis*. Eur. Arch. Otorhinolaryngol.  
469 1999;256(7):330-334.  
470
- 471 48. Faria AYK, Albuquerque MCF, Neto DC. Physiological quality of cottnseeds submitted to  
472 chemical and biological treatments. Revista Brasileira de Sementes. 2003;25(1):121-127.  
473 Portuguese. DOI: 10.1590/S0101-31222003000100019  
474
- 475 49. Carvalho NM, Nakagawa J. Seeds: science, technology and production. 4th ed.  
476 Jaboticabal: FUNEP; 2000. Portuguese.  
477
- 478 50. Guerra JI, Costa MLN. Evaluation of diseases management in *Brachiaria brizantha* cv  
479 BRS Piatã. In: Meeting of Teaching, Research and Extension - Scientific Education:

480 Scientific Innovations and Regional Development, 3. Annals. Dourados: UEMS; 2012.  
481 Portuguese.  
482  
483 51. Mentem JOM, Ishizuka MS. Health of tropical forage seeds - Upgrade course of tropical  
484 forage seeds production: relevant aspects to seed production. Presidente Prudente:  
485 AEAPP/FATL; 2015. Portuguese.  
486  
487 52. Guimarães GI, Lima MLP. **General** and morphological aspects of the fungus *Curvularia*  
488 sp. General and morphological aspects of *Curvularia* sp. Accessed 12 Mar 2019. Available:  
489 [https://fitopatologia1.blogspot.com/2010/10/aspectos-gerais-e-morfologicos-do-](https://fitopatologia1.blogspot.com/2010/10/aspectos-gerais-e-morfologicos-do-fungo_7393.html)  
490 [fungo\\_7393.html](https://fitopatologia1.blogspot.com/2010/10/aspectos-gerais-e-morfologicos-do-fungo_7393.html)  
491  
492 53. Yago JI, Roh JH, Bae S, Yoon YN, Kim HJ, Nam MH. The effect of seed-borne  
493 mycoflora from sorghum and foxtail millet seeds on germination and disease transmission.  
494 Mycobiology. 2011;39(3):206-218. DOI: 10.5941/MYCO.2011.39.3.206  
495  
496 54. Revankar SG. Dematiaceous fungi. Mycoses. 2007;50(2):91-101. Portuguese.  
497  
498 55. Ferreira LS. Characterization of isolates of *Curvularia* spp corn endophytic by  
499 morphological and molecular parameters. **(Dissertation)**. Federal University of Paraná; 2010.  
500 Portuguese. Accessed 14 Jun 2019. Available:  
501 [https://acervodigital.ufpr.br/bitstream/handle/1884/24194/LISANDRA%20SANTOS%20FER-](https://acervodigital.ufpr.br/bitstream/handle/1884/24194/LISANDRA%20SANTOS%20FERREIRA%20150710versao%20que%20foi%20impressa.pdf?sequence=1&isAllowed=y)  
502 [REIRA%20150710versao%20que%20foi%20impressa.pdf?sequence=1&isAllowed=y](https://acervodigital.ufpr.br/bitstream/handle/1884/24194/LISANDRA%20SANTOS%20FERREIRA%20150710versao%20que%20foi%20impressa.pdf?sequence=1&isAllowed=y)  
503  
504 56. Carter E, Boudreaux C. Fatal cerebral phaeohyphomycosis due to *Curvularia lunata* in  
505 an immunocompetent patient. Journal of Clinical Microbiology. 2004;42(11): 5419-5423. DOI:  
506 10.1128/JCM.42.11.5419-5423.2004  
507  
508 57. Moura RM, Pedregosa EMR, Guimarães LMP. New data on the etiology of the black  
509 bark of yam in northeastern Brazil. Nematologia Brasileira. 2001;25(2):235-237. Portuguese.  
510  
511 58. Herraes P, Rees C, Dunstan R. Invasive phaeohyphomycosis caused by *Curvularia*  
512 species in a dog. Veterinary Pathology. 2001;38(4):456-459. DOI: 10.1354/vp.38-4-456.  
513  
514 59. Dasgupta S, Saha D, Saha A. Levels of common antigens in determining pathogenicity  
515 of *Curvularia eragrostidis* in different tea varieties. Journal of Applied Microbiology.  
516 2005;98(5):1084-1092. DOI: 10.1111/j.1365-2672.2005.02538.x  
517  
518 60. Shan FY, Wei SW, Jie LY, Zhang JC, Gao DJ. Parasitic fitness and RAPD analysis of  
519 *Curvularia* species on corn. Journal of Agricultural University of Hebei. 2008;31(1):37-41.  
520 DOI:  
521  
522 61. Pimentel JD, Mahadevan K, Woodgyer A, Sigler L, Gibas C, Harris OC, Lupino M, Athan  
523 E. Peritonitis due to *Curvularia inaequalis* in an elderly patient. Journal of Clinical  
524 Microbiology. 2005;43(8):4288-4292. DOI: 10.1128/JCM.43.8.4288-4292.2005  
525  
526 62. Schuch UZ, Lucca Filho AO, Peske STP, Dutra LMC, Brancão MF, Rosenthal MD et al.  
527 Physiological and sanitary quality of rice seeds stored with different seed moisture contents  
528 and fungicide treated. Revista Brasileira de Sementes. 2006;28(1):45-53. Portuguese. DOI:  
529 10.1590/S0101-31222006000100007  
530

- 531 63. Pinto JC, Nabinger C, Maraschin GE. Determination of harvest time for seeds of  
532 *Paspalum guenoarum* Arech. f. azulão grass. Revista da Sociedade Brasileira de Zootecnia.  
533 1984;13(1):61-66. Portuguese.  
534
- 535 64. Costa RV, Casela CR, Cota LV. Foliar Diseases. Accessed 12 Mar 2019. Available:  
536 [https://www.agencia.cnptia.embrapa.br/gestor/milho/arvore/CONTAG01\\_32\\_161682005111.](https://www.agencia.cnptia.embrapa.br/gestor/milho/arvore/CONTAG01_32_161682005111.html)  
537 [html](https://www.agencia.cnptia.embrapa.br/gestor/milho/arvore/CONTAG01_32_161682005111.html).  
538
- 539 65. Salgado M, Pfenning LH. Identification and characterisation of *Phoma* species in Brazil.  
540 In: Paulista Congress of Phytopathology. Annals. Campinas: Paulista Association of  
541 Phytopathology; 2000. Portuguese.  
542
- 543 66. Aveskamp MM, Gruyter J, Crous PW. Biology and recent developments in the  
544 systematics of *Phoma*, a complex genus of major quarantine significance. Fungal Diversity.  
545 2008;31:1-18.  
546
- 547 67. Irinyi L, Kövics GJ, Sándor E. Taxonomical re-evaluation of *Phoma*-like soybean  
548 pathogenic fungi. Mycological Research. 2009;113(Pt 2):249-60. DOI:  
549 10.1016/j.mycres.2008.11.003  
550
- 551 68. Anjos JRN, Charchar MJA, Anjos SSN, Teixeira RN. *Phoma* sp. (sect. *Peyronellaea*), as  
552 causal agent of leaf spot on *Paspalum atratum* in Brazil. Fitopatologia Brasileira.  
553 2005;30(1):72-74. Portuguese. DOI: 10.1590/S0100-41582005000100012  
554
- 555 69. Leite GG, Silveira LF, Fernandes FD, Gomes AC. Growth and chemical composition of  
556 the grass *Paspalum atratum* cv. Pojuca. Brasília: Embrapa; 2001. Portuguese.  
557
- 558 70. Silva LC. Mycotoxins in grain and grain products. Revista Grãos Brasil. 2009;8(39):13-  
559 16. Portuguese.  
560
- 561 71. Nor Azliza I, Hafizi R, Nurhazrati M, Salleh B. Production of major mycotoxins by  
562 *Fusarium* species isolated from wild grasses in Peninsular Malaysia. Sains Malaysiana.  
563 2014;43:89-94.  
564
- 565 72. Carmo ALM. *Fusarium* in Pinus seeds in Brazil: distribution, pathogenicity, species  
566 identification and biological control. Thesis (Doctor degree). Federal University of Paraná;  
567 2017. Portuguese. Accessed 14 May 2019. Available:  
568 [https://acervodigital.ufpr.br/bitstream/handle/1884/47122/R%20-%20T%20-](https://acervodigital.ufpr.br/bitstream/handle/1884/47122/R%20-%20T%20-%20ANA%20LIDIA%20MOURA%20DO%20CARMO.pdf?sequence=1&isAllowed=y)  
569 [%20ANA%20LIDIA%20MOURA%20DO%20CARMO.pdf?sequence=1&isAllowed=y](https://acervodigital.ufpr.br/bitstream/handle/1884/47122/R%20-%20T%20-%20ANA%20LIDIA%20MOURA%20DO%20CARMO.pdf?sequence=1&isAllowed=y)  
570
- 571 73. Agrolink. *Fusarium* - dry rot (*Fusarium solani*). Portuguese. Accessed 14 May 2019.  
572 Available: [https://www.agrolink.com.br/problemas/fusariose\\_1258.html](https://www.agrolink.com.br/problemas/fusariose_1258.html).  
573
- 574 74. Batista ICA, Boari ADJ, Quadros AFF. Molecular identification of *Macrophomina*  
575 *phaseolina* in bean-tube. In: Scientific Initiation Seminar, 20 e Postgraduate Seminar of  
576 Embrapa Amazônia Oriental, 4. Annals. Belém: Embrapa; 2016. Portuguese.  
577
- 578 75. Lygin AV, Hill CB, Zernova OV, Crull L, Widholm JM, Hartman GL, Lozovaya VV.  
579 Response of soybean pathogen to glyceollin. Phytopathology. 2010;100(9):897-903. DOI:  
580 10.1094/PHYTO-100-9-0897  
581
- 582 76. Almeida AMR, Seixas CDS, Farias JRB, Oliveira MCN, Franchini JC, Debiasi H.  
583 *Macrophomina phaseolina* in soybean. 1st ed. Londrina: Embrapa; 2014. Portuguese.

584  
585 77. Agrolink. Stretching: Root rot (*Pythium* spp.). Portuguese. Accessed 14 May 2019.  
586 Available:[https://www.agrolink.com.br/problemas/estiolamento\\_1711.html](https://www.agrolink.com.br/problemas/estiolamento_1711.html)  
587  
588 78. Andrade CMS, Valentim JF. Grass brizantão death syndrome in Acre: characteristics,  
589 causes and technological solutions. 1st ed. Rio Branco: Embrapa; 2007. Portuguese.  
590  
591 79. Dias PP, Berbara RLL, Fernandes MCA. *Rhizoctonia solani* and *Fusarium oxysporum*  
592 f.sp. *phaseoli* control by biopreparation with *Trichoderma* spp. isolates Summa  
593 Phytopathologica. 2019;39(4):258-262. Portuguese. DOI: 10.1590/S0100-  
594 54052013000400005  
595  
596 80. Poloni NM, Molina LMR, Mesa EC, Garcia IL, Ceresini PC. Evidence that the fungus  
597 *Rhizoctonia solani* AG-1 IA adapted to Urochloa in Colombia keeps a broad host range  
598 including maize. Summa Phytopathologica. 2016;42(3):228-232. Portuguese. DOI:  
599 10.1590/0100-5405/2176  
600  
601 81. Goulart ACP. Reaction of cotton cultivars to *Rhizoctonia solani* at seedling stage and  
602 benefits of fungicide seed treatment. Summa Phytopathologica. 2016;42(4)308-312.  
603 [Portuguese]. DOI: 10.1590/0100-5405/2116  
604  
605 82. Lucon CMM, Koike CM, Ishikawa AI, Patrício FRA, Harakava R. Bioprospection of  
606 *Trichoderma* spp. isolates to control *Rhizoctonia solani* on cucumber seedling production.  
607 Pesquisa agropecuária brasileira. 2009;44(3):225-232. Portuguese. DOI: 10.1590/S0100-  
608 204X2009000300002  
609  
610 83. Lewis JA, Lumsden RD. Biocontrol of damping-off of greenhouse-grown crops caused  
611 by *Rhizoctonia solani* with a formulation of *Trichoderma* spp. Crop Protection.  
612 2001;20(1):49–56. DOI: 10.1016/S0261-2194(00)00052-1  
613  
614 84. Pinheiro VR, Seixas CDS, Godoy CV, Soares RM, Oliveira MCN, Almeida AMR.  
615 Development of *Sclerotium rolfsii* sclerotia on soybean, corn, and wheat straw, under  
616 different soil temperatures and moisture contents. Pesquisa Agropecuária Brasileira.  
617 2010;45(3):332-334. Portuguese. DOI: 10.1590/S0100-204X2010000300014  
618  
619 85. Prahbu HV, Hiramoth PC. Bioefficacy of fungicides against collar rot of cotton caused by  
620 *Sclerotium rolfsii* Sacc. Karnataka Journal of Agricultural Sciences. 2003;16(4):576- 579.  
621 DOI:  
622  
623 86. Cilliers AJ, Pretorius ZA, Van Wyk PS. Integrated control of *Sclerotium rolfsii* on  
624 groundnut in South Africa. Journal of Phytopathology. 2003;151(5):249-258. DOI:  
625 10.1046/j.1439-0434.2003.00715.x  
626  
627 87. Menten JO, Moraes MHD. Seed Treatment: history, types, characteristics and benefits.  
628 Informativo Abrates. 2010;20(3):52-71. Portuguese. Accessed 14 Jun 2019. Available:  
629 <http://www.abrates.org.br/images/stories/informativos/v20n3/minicurso03.pdf>.  
630  
631 88. Santos FC. Scarification, chemical treatment, seed coating and seed storage of  
632 *Brachiaria brizanta* cultivar Marandu. Thesis (Doctor degree); Federal University of Lavras;  
633 2009. Portuguese. Accessed 15 Jun 2019. Available:  
634 [http://repositorio.ufla.br/jspui/bitstream/1/3099/1/TESE\\_Escarifica%C3%A7%C3%A3o%2C%20tratamento%20qu%C3%ADmico%2C%20revestimento%20e%20armazenamento%20de%20sementes%20de%20Brachiaria%20brizantha%20cultivar%20Marandu.pdf](http://repositorio.ufla.br/jspui/bitstream/1/3099/1/TESE_Escarifica%C3%A7%C3%A3o%2C%20tratamento%20qu%C3%ADmico%2C%20revestimento%20e%20armazenamento%20de%20sementes%20de%20Brachiaria%20brizantha%20cultivar%20Marandu.pdf).  
635  
636

- 637 89. Cruz, CJ. Corn Farming. Sete Lagoas: Embrapa; 2007. Portuguese.  
638
- 639 90. Rossetto, R; Santiago, AD. Other Diseases. Brasília: Embrapa. Portuguese. Accessed 03  
640 Jul 2019. Available: [http://www.agencia.cnptia.embrapa.br/gestor/cana-de-](http://www.agencia.cnptia.embrapa.br/gestor/cana-de-acucar/arvore/CONTAG01_81_22122006154841.html)  
641 [acucar/arvore/CONTAG01\\_81\\_22122006154841.html](http://www.agencia.cnptia.embrapa.br/gestor/cana-de-acucar/arvore/CONTAG01_81_22122006154841.html).  
642
- 643 91. Mendes, MAS.; Urben, AF. Related Fungi in Plants in Brazil. Brasília: Embrapa. Accessed  
644 03 Jul 2019.  
645 Available: <http://pragawall.cenargen.embrapa.br/aiqweb/michtml/fgbanco01.asp>.  
646
- 647 92. Barbosa, FR. Conidial fungi associated with decaying leaves of *Clusia melchiori gleason*  
648 and *C. nemorosa* G. Mey in fragment of Atlantic Forest, Bahia, Brazil. Dissertation. Federal  
649 University of Pernambuco; 2007. Portuguese. Accessed 03 Jul 2019. Available:  
650 [http://www.educadores.diaadia.pr.gov.br/arquivos/File/2010/artigos\\_teses/Biologia/Dissertac-](http://www.educadores.diaadia.pr.gov.br/arquivos/File/2010/artigos_teses/Biologia/Dissertacao/fungos_conidiais.pdf)  
651 [ao/fungos\\_conidiais.pdf](http://www.educadores.diaadia.pr.gov.br/arquivos/File/2010/artigos_teses/Biologia/Dissertacao/fungos_conidiais.pdf).  
652
- 653 93. Maia, LC; Cavalcanti, MA; Gibertoni, T; Goto, BT; Melo, AMM; Baseia, IG; Silvério, ML.  
654 Fungi. Brasília: Embrapa. Portuguese. Accessed 03 Jul 2019. Available:  
655 [http://www.mma.gov.br/estruturas/chm/\\_arquivos/14\\_Biodiv\\_14\\_Cap04a.pdf](http://www.mma.gov.br/estruturas/chm/_arquivos/14_Biodiv_14_Cap04a.pdf).  
656
- 657 94. Castro, NR; Coêlho, RSB; Laranjeira, D; Couto, EF; Souza, MBR. Occurrence,  
658 inoculation methods and aggressivity of *Fusarium oxysporum* f.sp. *cubense* in *Heliconia* spp.  
659 Summa Phytopathologica, 34(2):127-130; 2008. DOI: 10.1590/S0100-54052008000200003.  
660
- 661 95. Uebel,JD. Evaluation of fungicides in the control of leaf diseases, burned grains and  
662 effect on NDVI (normalized difference vegetation index) in corn hybrids. Dissertation.  
663 University of Brasília; 2015. Portuguese. Accessed 03 Jul 2019. Available:  
664 [http://repositorio.unb.br/bitstream/10482/19019/1/2015\\_JulianoDanielUebel.pdf](http://repositorio.unb.br/bitstream/10482/19019/1/2015_JulianoDanielUebel.pdf)  
665
- 666 96. Casa, RT; Reis, EM; Blum, MMC. Quantification of damage caused by diseases in  
667 maize. In: Plant Disease Epidemiology Workshop. Annals. Viçosa: Sheet of Viçosa Ltda,  
668 2005.  
669
- 670 97. Reis, EM; Casa, RT; Bresolin, ACR. Manual of diagnosis and control of diseases of  
671 maize. Lages: Graphel, 2004. Portuguese.  
672
- 673 98. Reis, EM; reis, AC; carmona, MA. Manual of fungicides: guide for the chemical control of  
674 plant diseases. Passo Fundo: UPF Editora, 2010. Portuguese.  
675
- 676 99. Costa, R. V. Integrated management of diseases in maize. In: Agricultural production  
677 system integrated in crop/livestock integration systems: maize and soybean. Course.  
678 Goiânia: Embrapa, 2009.  
679
- 680 100. Bettiol, W.; Morandi, MAB. Biocontrol of plant diseases: use and perspectives.  
681 Jaguariúna: Embrapa, 2009. Portuguese. Accessed 03 Jul 2019. Available:  
682 <https://www.alice.cnptia.embrapa.br/bitstream/doc/579982/1/2009CL07.pdf>.