

**Role of Bamboo forest for mitigation and adaptation to Climate
Change challenges in China**

ABSTRACT

Bamboo is one of the fastest growing plants on the planet, with many attributes which make it a useful potential resource for humankind. Though having fast growth and good regeneration performance after harvesting is a unique characteristic of the specie. It enhances a high carbon storage potential particularly when the harvested culms are transformed into durable products. China has many bamboo species with distribution and area coverage's, and highly connected in using the production of bamboo resources. Its characteristics make it an ideal solution for the environmental and social consequences of tropical deforestation. The aim of this review paper is to assess the contribution of bamboo in mitigating and adapting impacts of climate change and its importance in terms of ecological and socio-economic benefits. The review summarized the role of bamboo forests towards mitigating and adapting its potential to overcome the impacts of climate change currently seen globally and particularly to China. Therefore, advancing bamboo farming systems in different levels it's advantages to reduce greenhouse gas in atmosphere and expanding bamboo forests in future under wider use and intensive management is recommended.

Key words: - Bamboo; carbon sink; climate change; greenhouse gas

1. INTRODUCTION

Bamboo is a grass type of Gramineae family and it is an important component of many forest environments. It adapts easily to a range of climatic and soil conditions, and is widely distributed in the tropical and subtropical zones approximately between 46° N and 47° S latitude, covering a total area of about 31.5 million ha, and accounted for about 0.8% of the World's total forested area in 2010 [1]. It has unique features that distinguish it from most other woody plants. For example: culms

42 that are connected by an extensive system of rhizomes, leading to emerge new culms by rapid
43 asexual reproduction [2, 3, and 4].

44 The bamboo resources are distributed in many countries in the World, majorly found in Asia, Africa,
45 and Latin America; however, the origin of most of species lie in Southeast Asia. Worldwide bamboo
46 families categorized with more than 107 genera and 1300 species [6]. Along with this all China has
47 the highest bamboo species diversity, with 39 genera and 509 species, accounting for 36% and 39%,
48 respectively, of the total bamboo genera and species with the rest of the World [6]. Besides that,
49 China's bamboo forests cover an area of 4.84 million ha in 2005, which accounts for 2.8% of China's
50 forested area and 15.4% of the World's area of bamboo [7]. Bamboo forests is the most important
51 non-wood forest product and substitute wood products. As a resource, bamboo forests are an
52 important part of eco-systems, which provide a number of basic environmental services. Bamboo
53 provides food and raw materials, reduces water erosion on slopes, regulates water flows and it act
54 like windbreak in shelterbelts which offer protection against storms [1]. In addition to this because of
55 its special root re-sprouting regeneration strategy bamboo forests it generates a good potential of
56 carbon storage mechanisms, and water and soil conservation and many more advantageous [5].

57 Many studies have shown that well managed and regularly harvested bamboo can sequester more
58 carbon than bamboo in natural state. Despite that, it can sequester more carbon than fast-growing
59 tropical and subtropical trees under comparable conditions. If bamboo forests were not managed
60 through annual harvesting practices, they would be significantly less effective in carbon sequestration.
61 The aim of this review paper is to assess the contribution of bamboo in mitigating and adapting
62 impacts of climate change and its importance in terms of ecological and socio-economic benefits

63 **2. ROLES OF FORESTS IN MITIGATING CLIMATE CHANGE**

64 Forests are one of the biggest reservoirs of carbon, so they help to keep the carbon cycle and other
65 natural processes working and help reduce climate change. However, forests can also be one of the
66 biggest sources of CO₂ emissions [8]. In addition, forests provide a wide range of ecological, social,
67 and economic benefits, in the form of goods and services to society, that are much less easier to
68 quantify. Besides that, the demand for timber and related products will require more efficient and
69 sustainable use of natural resources. Forests are the most vulnerable climate dependent systems, but
70 have also been recognized to have significant and crucial contribution to address the challenges of
71 mitigation and adaptation in tandem with the issues of livelihoods, economic growth and development.
72 However, the most recent report from the International Union of Forest Research Organizations
73 (IUFRO) indicates the gloomy picture about the future of the World forests in changed climate. While,
74 it suggests that in a warmer World, the current carbon regulating services of forests as carbon sinks
75 may be entirely lost as land ecosystems could turn into a net source of carbon dioxide later in the
76 century [9]. It plays in combating climate change impact through reducing the emissions from
77 deforestation and forest degradation has become a fundamental issue to international dialogues on
78 preventing the current global temperature increases [10]. The challenges of climate change seen in
79 affecting forest ecosystems in their structure and morphology, thus causing an implication on

80 functionality of forests in every corner of the World [11, 12]. Besides that, it is considered one of the
81 greatest threats facing humanity in the current global situations. According to the Intergovernmental
82 Panel on Climate Change (IPCC), global warming is unambiguous, with evidence towards the
83 increases of average air and ocean temperatures, which leads to aggravate melting of snow and ice
84 and sea levels [8, 13].

85 **3. BAMBOOS FOR CLIMATE CHANGE ADAPTATION**

86 Bamboos are one of the World's strongest and fastest growing woody plants capable of providing
87 ecological, economic and livelihood security to the people, distributed over ranges of climate from mild
88 temperate to tropical. Bamboo's fast growth ability to grow on varied soils and climate, renewability
89 and positive socio-economic impacts make them an excellent alternative for combating climate
90 change [14]. The high growth potential and ability to store large amounts of carbon make
91 sequestration and on the other hand, their environmental and socio-economic services can help
92 communities in developing countries to adapt to the climate change impacts. According to the
93 research result by International Network of Bamboo and Rattan (INBAR) had shown that well
94 managed bamboos could be an effective in carbon sink and perform better than other species like
95 Chinese fir and eucalyptus growing under similar conditions. Furthermore, the necessities of bamboo
96 management with period, sustainable way and selective harvesting mechanism of stem which are
97 turned into products that can hold carbon for many years. The increasing popularity of durable
98 bamboo products ensures that for the foreseeable future, productive bamboo systems can be
99 considered as an important carbon sink [15].

100 **3.1 Bamboo for Timber Demand and Climate Change**

101 The demand for timber and different agricultural products will continue to increase with the global
102 population. Instead of satisfying the increasing demand for different commodities, the global policies
103 must need to shift towards using efficient and sustainable production systems [13]. Bamboo is one of
104 an alternative resource that used to play an important contribution towards reducing the direct
105 pressure on forest resources [13, 24]. Furthermore, one of the best example in China, following the
106 logging bans of certain forests resources came into effect in 1998, while bamboo forests has been
107 used as a possible substitute to timber.

108 Magel et al [16] argues that the growth of new shoots in a bamboo plantations occur because of
109 transfer of the energy accumulated in culms through photosynthesis in the previous year. The result
110 of such growth of bamboo culms is not driven by its own carbon sequestration, but by sequestration in
111 previous seasons in other parts of the bamboo system, and such growth of new shoots is not an
112 indicator of sequestration rate. Another report by Zhou [17] show that as bamboo system requires
113 more inputs in the shooting period of young culms, which means when new shoots developed during
114 that time high growth of bamboo shoots, can be equated with a high rate of carbon sequestration. The
115 maturity period of most bamboo culms estimated between 7-10 years, approximately, after that they
116 deteriorate rapidly, releasing carbon from the aboveground biomass back into the atmosphere [18].

117 Therefore, at a natural circumstance, bamboo will reach a stable level of above ground carbon
118 relatively quickly, even though carbon accumulation through sequestration is offset by carbon release
119 through deterioration of old culms.

120 **3.2 Suitable Ecological Growing Conditions for Bamboo Species**

121 Naturally, bamboo species has a potential to grow at different altitudinal range from 0 to 4000 meter
122 above sea level. It prefers well-drained sandy loams to loamy clay types of soils originated from river
123 alluvium or underlying rock. In most of the bamboo thrive well at annual average temperatures range
124 of 8.8 – 36°C and annual rainfall of 1270 – 4050 mm. On the other hand, some bamboo species are
125 also growing under high rainfall areas, while some can tolerate limited winter frost [19]. ICFRE [20]
126 reported as one of the fastest growing species in the planet, under ideal environment; it can be
127 growing up to one meter a day. The biomass of freshly planted bamboo plantation increases rapidly
128 for the early six to eight years after which emergence and death of culms tend to become equal.

129 **4. MANAGEMENT AND HARVESTING OF BAMBOO**

130 Bamboos have well-developed rhizomes with good root systems which help to obtain strengthen
131 during their existence. Its culms mature within three to four years and naturally die after eight to ten
132 years, if not used the products [21]. The periodic removal of mature culms from each bamboo clump
133 and this cycle of removal may vary from two to four year. Thus, provide a highly renewable resource
134 with a high degree of sustainability. It makes bamboos acquiescent to sequester accumulated CO₂
135 from the atmosphere throughout the lifetime [22]. The sequestration rate of bamboo is higher during
136 the initial eight to ten years period of fast growth. Numerous studies categorize the production
137 management of bamboo management practices in different five major points: timber, shoot, pulp,
138 ornamental and water/soil conservation benefits.


139 **5. BAMBOOS FOR CLIMATE CHANGE MITIGATION**

140 Bamboo grows more rapidly than any other trees and reach to give yield within three to four years
141 after planting. Hence, it is one of the fast growing and responding well against, drought which can
142 make the species more acceptable in making evergreen environment in addition to soil and water
143 conservation, carbon storage and rehabilitation of degraded lands [23]. It offers one of the quickest
144 ways to remove vast amounts of that CO₂ from the atmosphere. Lou et al [24] report that at the age of
145 9 -10 years old in moso bamboo (*Phyllostachys pubescens*) plantation the above ground carbon stock
146 ranges between 25 to 32t ha⁻¹ in China. Furthermore, another study show that in *P. pubescens* and *P.*
147 *bambusoides* from natural forests in Japan have an aboveground carbon stock of 78.6 t ha⁻¹ and 52.3
148 t ha⁻¹, respectively [25]. A four year mixed bamboo plantation (*Bambusa vulgaris*, *B. balcooa*, and *B.*
149 *cacharensis*) in India shows that the aboveground carbon stock is about 61.05t ha⁻¹ [26]. Despite that,
150 it provides a minimum CO₂ gas and generates up to 25% more oxygen than other trees within the
151 same level. One hectare of bamboo can sequester up to 62 t of CO₂ yr⁻¹, whereas equivalent of young
152 forest sequesters 15 t of CO₂ yr⁻¹. The *Guadua* plantations in Costa Rica estimated to absorb 17 t of

153 CO₂ ha yr⁻¹ as the study showed by Janssen [27]. Another research study by INBAR states that over
154 the past 15 years, areas under bamboos in Asia grew by 10%. Studies have estimated that the
155 carbon stored in Chinese bamboo forests will increase from 727.08 Tg C in 2010 to 1,017.54 Tg C in
156 2050, which equates to an increase of nearly 40% in 40 years. This represents a significant
157 contribution to the Chinese forest carbon stock and a range that shows that policies aiming at
158 combating climate change with bamboos can indeed have significant promise [28]. For example: by
159 INBAR's, as modeling shows that a managed moso bamboo forest accumulates about 300 t of carbon
160 ha⁻¹ after 60 years. As well, it does also produce more biomass under well managed and regular
161 harvesting of mature culms. Another report by Lou et al [24], confirmed that the amount of carbon
162 sequestration between a fast growing Chinese Fir plantation and monopodial (*P. pubescens*)
163 plantation modeled for subtropical agro-ecologies in South East China and the results showed that,
164 bamboo sequestered more carbon than the Chinese Fir in the first 5 years. This might be due to rapid
165 early growth; bamboos sequester more carbon in the early years of a plantation than comparable
166 forest trees. In the other way, unmanaged bamboo stands do not store high levels of carbon, as their
167 productivity is low and the accumulated carbon returns quickly to the atmosphere as the older culms
168 decompose [28].

169 **5.1 Carbon Sequestration Potential of Bamboo**

170 Carbon sequestration is the process of capturing and storing atmospheric carbon dioxide. Naturally,
171 the performance of bamboo plantation is different comparing with other tree species since it has a fast
172 growth rate with high re-growth behavior after harvesting. Despite that, it has a high carbon storage
173 potential according to Zhou and Jiang studies [29], especially when the harvested culms are
174 transformed into durable products. The increased lifespan of durable bamboo products made possible
175 by modern technology can ensure that the sequestered carbon will not return quickly in to the
176 atmosphere, thereby prolonging the carbon storage of bamboo. According to Agarwal and his
177 colleague [30] research result in the Mid Himalayan region conducting in comparing the carbon
178 sequestration potential under different bamboo species, Monopodial species (*P. nigra*) has showed a
179 high potential to sequester carbon than other Sympodial species. This might be due to high density of
180 culms and high percent dry matter in the Himalayan region [31]. Currently in China, about 53 M ha of
181 forest plantation is there with a volume stock of 1.5 billion m³. Between 2005 and 2020, China has
182 pledged to establish more than 40 million ha of plantations, referred to as carbon sink forest [32]. As
183 plantations have been recognized as the national strategy for mitigating atmospheric CO₂, it is
184 essential to assess the potential of fast-growing and high yield plantations in carbon storage and
185 sequestration at stand, regional and national scales (Figure 1).



186
187


188 Figure 1. Contribution of bamboo on adding C sink and reducing C emission (Source: Yuen 33).

189

190 The area coverage of bamboo forests in China is about 6Mha, which stores about 780 Tg carbon,
191 accounting for 14% of total forest carbon stock in China. Beside the carbon density of bamboo forest
192 ecosystems in China, the estimated global bamboo carbon stock is about 4 Pg, accounting for 0.43%-
193 0.61% of total global forest carbon stock [33].

194 To combat climate change, bamboo should be a core development resource – providing countries
195 and development partners with a wealth of practical solutions to reduce the negative effects that
196 changing climate patterns have on millions of rural communities (Figure 2).

197



198

199 Figure 2. The five key functions of Bamboo help to mitigate/adapt the impacts of Climate Change.

200 **6. SUMMARY**

201 Currently the concern of climate change is a very serious and burning issue of global agendas. In this
 202 paper, we tried to review the contribution of bamboo forests towards mitigating the impacts of climate
 203 change and the versatility of its ecological and socioeconomic development benefits. It offers one of
 204 the quickest ways to remove huge amounts of CO₂ from the atmosphere. It minimizes CO₂ gas and
 205 generates more amount of oxygen than an equivalent stand of other tree species. Many scholars
 206 suggested that bamboo forest ecosystems provide significant services for human adaptation and
 207 development at the same time mitigate climate change impacts through carbon sequestration. Under

208 well-managed bamboo plantations, it shows an effective carbon sink and better performance than
209 other tree species growing under similar conditions. Despite this, it is a source of income in rewarding
210 the diverse requirements at small and large-scales in rural areas. Generally, this review prepared to
211 demonstrate the role of bamboo forests towards mitigating and adapting potential to overcome the
212 impacts of climate change seen in the world and particularly to China. Therefore, advancing bamboo
213 farming systems in different levels it's advantageous in reducing greenhouse gas in atmosphere and
214 expanding bamboo forests in future under wider use and intensive management is recommended.

215
216

REFERENCES

- 217 1. FAO. 2010. Global Forest Resources Assessment 2010: Main Report. Food and Agriculture
218 Organization of the United Nations, Rome.
- 219 2. Janzen D.H. 1976. Why bamboos wait so long to flower Annual? Rev. Ecol. Syst. 7(1): 347–391.
220 doi:10.1146/annurev.es.07.110176.002023.
- 221 3. Isagi Y. Kawahara T. Kamo K. and Ito H. 1997. Net production and carbon cycling in a bamboo
222 *Phyllostachys pubescens* stand. Plant Ecol. 130: 41–52. doi:10.1023/A:1009711814070.
- 223 4. Li R. Werger M.J.A. During H.J. and Zhong Z.C. 1998. Carbon and nutrient dynamics in relation to
224 growth rhythm in the giant bamboo *Phyllostachys pubescens*. Plant Soil, 201(1): 113–123.
225 doi:10.1023/A:1004322812651.
- 226 5. Lobovikov M. Lou Y.P. et al. 2009. The poor man's carbon sink. Bamboo in climate change and
227 poverty alleviation. Non-Wood Forest Products. Working Document (FAO), no 8, FAO, Rome
228 (Italy), Forestry Dept. 68p., FAO.
- 229 6. Zhu Z.H. 2001. The development of bamboo and rattan in tropical China. Beijing, China: China
230 Forestry Publishing House.
- 231 7. SFAPRC. 2006. Statistics of Forest Resources in China (1999–2003). State Forestry
232 Administration, P.R. China.
- 233 8. IPCC 2007. Solomon S. Qin D. Manning M. Chen Z. Marquis M. Averyt K.B. M. Tignor and H.L.
234 Miller (eds). The Physical Science Basis, Contribution of Working Group I to the Fourth
235 Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge and New
236 York, NY, USA, 996.
- 237 9. Seppala R. Buck A. Katila. 2009. Eds. Adaptation of forests and people to climate change: A global
238 assessment report. IUFRO World Series Vol. 22. Vienna, IUFRO.
- 239 10. Wei S.J. L. Sun S.W. Wei et al. 2013: Effects of climate changes on forest disasters and the
240 preventive measures. Journal of Catastrophology, 28(1), 36-38.
- 241 11. MoEF (India. Ministry of Environment and Forests). 2012. India Second National Communication
242 to the United Nations Framework Convention on Climate Change. New Delhi, Ministry of
243 Environment and Forests, Government of India.
- 244 12. Hal Stern. 2006. Reviewed work(s): Source: The American Statistician, Vol. 60, No. 4 (Nov.), pp.
245 328-331.
- 246 13. Eliasch, J., 2008. Climate change: financing global forests: The Eliasch Review.
- 247 14. Dransfield S. Widjala E. Eds. 1995. Bamboo Plant Resources of South-East Asia.
248 No. 7. Leiden, Backhuys Publishers. 190p.
- 249 15. INBAR. 2009. Bamboo: A strategic resource for countries to reduce the effects of climate change.
250 INBAR Policy Synthesis Report
- 251 16. Magel E. Kruse S. Lutje G. Liese W. 2005. Soluble Carbohydrates and Acid Lyertases involved in
252 the rapid growth of the developing culms in *Sasa palmate* (Bean) Camus. Bamboo Science and
253 Culture, Baton Rouge/USA 19 1, S. 23-29.
- 254 17. Zhou G.M. Jiang P.K. Mo L.F. 2009. Bamboo: a possible approach to the control of global
255 warming. International Journal of Nonlinear Sciences and Numerical Simulation, 10(5): 547-550.

- 256 18. Liese W. 2009. Bamboo as Carbon sink fact or fiction? VIII World Bamboo Congress Proceedings.
257 3:71-77.
- 258 19. FSI (Forestry Survey of India). 2011. India state of forestry report 2011. Dehradun, Forest Survey
259 of India. 286p.
- 260 20. ICFRE (India Council of Forestry, Research and Education). 1992. Bamboo. Dehradun, ICFRE.
- 261 21. Banik RL. 2008. Issues of production of bamboo planting materials-Lessons and strategies. Indian
262 For (Bamb Iss) 134(3):291-304.
- 263 22. Fu Maoyi and R.L. Banik. 1995. Bamboo, people and the management. Proceedings of the Vth
264 International Bamboo Workshop and the IV International Bamboo Congress Ubud, Bali,
265 Indonesia 10-22 June 1995. Volume I Propagation and management INBAR Technical Report
266 No.8p 18-30.
- 267 23. Terefe R. Samuel D. Sanbeto M. Daba M. 2016. Adaptation and growth performance of different
268 lowland bamboo species in Bako, West Shoa, Ethiopia. Journal of Natural Sciences Research.
269 6(9):61-65.
- 270 24. Lou Y. Li Y. Kathler B. Giles H. Zhou G. Bamboo and climate change mitigation: Comparative
271 analysis of carbon sequestration. INBAR. Technical Report No. 32. 2010; 1-20.
- 272 25. Isagi Y. 1994. Carbon stock and cycling in a bamboo *Phyllostachys bambusoides* stand. Ecol.
273 Res. 9. 47-55.
- 274 26. Nath A.J. Lai R. Das A.K. 2009. Above ground standing biomass for carbon storage in village
275 bamboos in northeast India. Bio. Bioeng 33, 1188-1196.
- 276 27. Janssen J.J.A. 1996. Building codes for bamboo housing. In Ganapathy, .M.; Janssen, J.A.;
277 Sastry, C.B. ed., Bamboo, people and the environment, Vol. 3, Engineering and utilization.
278 Proceedings of the Vth International Bamboo Workshop.
- 279 28. Kuehl Y. Yiping L. 2012. Carbon off- setting with bamboo. INBAR Working paper No. 71. Beijing,
280 INBAR.
- 281 29. Zhou G.M. and Jiang P.K. 2004. Density, storage and spatial distribution of carbon in
282 *Phyllostachys pubescens* forest. Sci. Sil. Sin. 40: 20–25.
- 283 30. Agarwal A. Purwar J.P. 2009. Evaluation of above ground biomass produced by *Dendrocalamus*
284 *asper* in North Western Himalayan region of India. In: Proceedings of VIII World Bamboo
285 Congress, 4, 91-96.
- 286 31. Agarwal A. Purwar J.P. 2012. Biomass Production and Carbon Sequestration Potential of Various
287 Bamboo Species in the Mid Himalayan Region of India. Paper Presented at World Bamboo
288 Congress.
- 289 32. Chen X. Zhang X. Zhang Y. Booth T& He X. 2009. Changes of carbon stocks in bamboo stands in
290 China during 100 years. *Forest Ecology & Management* 258: 1489–1496.
- 291 33. Yuen JQ. Fung T. Ziegler AD. 2017. Carbon stocks in bamboo ecosystems worldwide: estimates
292 and uncertainties. *For Ecol Manag* 393:113–138.