

Original Research Article

Evaluation of the Ornamental Potential of Safflower (*Carthamus tinctorius* L.)

ABSTRACT

Aims: to evaluate the ornamental potential of two safflower genotypes (*Carthamus tinctorius* L.): ICA 73, ICA 193, grown under protected environment.

Place and Duration of Study: Department of Agronomy of Federal Rural University of Pernambuco, between March and May 2017.

Methodology: The methodology addressed evaluated the performance of the two genotypes, through three stages. The first stage was carried out in a completely randomized design, being evaluated: plant height; stem diameter; leaf dentin; spinescent margin of the leaves; number of branches; number of flower buds; spinescent margin of the bracts; flowering; and flower production. The second stage was conducted in a randomized complete block design in a factorial scheme, and the following variables were evaluated: plant height; stem diameter; number of branches; number of flower buds; and flower production. Finally, the third stage used a completely randomized design in a factorial scheme and evaluated the variables: number of flowers; number of flower buds; and post-harvest durability. The analysis of variance was performed using the F test at 5% of probability and, afterwards, the regression or comparison analysis of averages by the Tukey test at 5% of probability.

Results: The ICA 73 access showed plants with high flower production and the ICA 193 exhibited plants with weak or moderate spinescent margin of the leaves and bracts, besides good uniformity of the anthesis of the flowers.

Conclusion: Both accesses showed ornamental potential, demonstrating precocity, beauty and durability of the flowers. The density of one plant was the most favorable for pot plant and cut flower. The semi-open flowers harvest point was the best for maintaining the stem quality.

Keywords: Floriculture. Precocity. Flowers. Potted Plant. Cut Flower Stems.

1. INTRODUCTION

Floriculture is a segment that stands out in the world, with significant growth, constituting an important source of income for several countries and promoting the development of productive poles. In this context, production is mainly intended for export to large consumer centers located in the European Union, the United States of America and more recently Asia and the Middle East [1,2,2].

In Brazil, in contrast to other developing countries, this sector has grown mainly towards the intern market [3]. With annual growth of around 8% per year, growing exports and significant increase in domestic consumption, the floriculture in the country became one of the most prominent segments in the agribusiness market, moving around R \$ 6.7 billion in 2016 [4].

27
28 Most of the market is supplied by plants from states in the Southeast of the country [5], while
29 other important Brazilian regions, where new floriculture poles emerge, end up with
30 **difficulties to development**, despite their natural aptitude for this sector [6].
31

32 | ~~The production of flowers~~~~The floriculture~~ consists of a dynamic sector with a constant search
33 for new products that meet the new trends. In this sense, safflower has great **potential for**
34 **this** market. The safflower (*Carthamus tinctorius*) belongs to the Asteraceae family, it is an
35 herbaceous annual plant, self-pollinated and capable of developing into various
36 | edaphoclimatic conditions [7,8,8], tolerating low water availability and high temperatures [9].
37

- Comment [REV A1]: Floriculture was used previously
- Comment [REV A2]: the production of flowers
- Comment [REV A3]:

38 The use of this herbaceous plant covers many possibilities and extends to practically every
39 part of the plant. Its **seeds exploited part in the market, being** present in food products [10],
40 | cosmetics ~~and also~~ in the composition of drugs [11]. Safflower oil may also be intended
41 for biofuel production [12], **and the bagasse used in animal feed supplements** [13]. **The**
42 **flowers allow the extraction of two dyes from their petals, a soluble and another hydrophilic**
43 **[14], and have ornamental potential as fresh or dried cut flower** [15].
44

45 The variability of the color, size and arrangement of the florets that the safflower possesses,
46 make the species very attractive to the **floriculture** market, being able to be used as
47 ornamental plants, cutting stems and confection of bouquets, with potential for planting in
48 gardens or pots and trade while fresh or when dry [16]. In Europe, the use of this
49 herbaceous plant in the **flower market is common with specific cultivars for this purpose** [17].
50

51 | Safflower cultivation in Brazil is still very limited and is restricted ~~to basically~~ attending to
52 some scientific ~~research~~~~research; moreover~~~~moreover~~, usually it covers only the production
53 of oil, thus not **exploiting the ornamental potential**. However, considering the importance that
54 the floriculture is taking in Brazilian agribusiness, the search and insertion of new products to
55 expand and meet market demand **becomes a necessity** and, in this context, investing **in the**
56 **potential of this specie is a very** promising strategy.
57

58 The objective of this work was to evaluate the ornamental potential of two safflower
59 genotypes to pot plant **and cut flower, cultivated in a protected environment, determining the**
60 **best density and harvest point**.
61

62 2. MATERIAL AND METHODS

63
64 The safflower accesses (*Carthamus tinctorius*) ICA 73 and 193 used in the study were
65 imported by the Institute of Agricultural Sciences (ICA) in agreement with the Federal
66 University of Minas Gerais (UFMG) **from** germplasm banks of India and Ethiopia, which were
67 later transferred to the Plant Breeding Program of the Federal **Rural** University of
68 Pernambuco (UFRPE) to carry out this work.
69

70 **The experiment was carried out in three stages** under greenhouse conditions in the
71 Agronomy Department of the Federal Rural University of Pernambuco - UFRPE, Recife -
72 PE, whose geographical coordinates are 8°10'52"S latitude, 34°54'47" longitude and 2m
73 altitude.
74

75 | **Sowing** was done manually in pots with 5_L capacity, filled with commercial Basaplant™
76 substrate, the depth of approximately three centimeters [18]. Irrigations were performed
77 | manually and daily, **approximately 300 ml per vase**. **No fertilization was applied**~~performed~~.
78

79 The first stage was carried out in a completely randomized design, with twenty repetitions.
80 The treatments were composed of the two safflower genotypes. Twenty vases were used for
81 each access (ICA 73 and ICA 193), where each vase represented an experimental unit,
82 totaling 40 parcels.

83
84 The Emergency Velocity Index (EVI) was calculated according to the formula of [19]:

85 $EVI = \frac{E_n}{N_n}$; Where: n = the number of normal seedlings recorded in the count "n";
86 Nn = number of days of sowing until the count "n". For this, the number of emerged plants,
87 with two open cotyledon leaves, was registered until the ninth day after sowing. The
88 percentage of germination was calculated after stabilization of the emergency, considering
89 the final number of emerged plants.

90
91 The evaluations was carried out sixty days after sowing (DAS), based on the following
92 characteristics: Plant Height (PH (cm)) - performed with ruler and corresponding to the
93 measurement of the soil to the apex of the plant; Stem Diameter (SD (cm)) - measured with
94 a digital pachymeter in the base of the stem; Leaf Dentin (LD) - classified by scale of notes:
95 absent or weak (0); moderate (5); strong (10) [20] (Figure 1); Spinescent Margin of the
96 Leaves (SML) - graded by note scale: absent or weak (0); moderate (5); strong (10) [20]
97 (Figure 2); Number of Branches per plant (NB) - obtained by counting (Figure 3); Number of
98 Flower Buds (NFB) - obtained by counting the flower buds (*capitulum*); of the Spinescent
99 Margin of the Bracts (SMB) - graded by grading scale: absent or weak (0); moderate (5);
100 strong (10) [20] (Figure 4); Flowering (FI) - number of days from sowing to beginning of
101 flowering; Flower Production (FP) - obtained by counting open inflorescences.
102



103
104 Figure 1. Leaf Dentin of Safflower: (A) Weak; (B) Moderate e (C) Strong.



105
106 Figure 2. Spinescent Margin of the Leaves of Safflower: (A) Weak; (B) Moderate; (C)
107 Strong.
108



Figure 3. Safflower plants: (A) Little branched; (B) Very branched.



Figure 4. Spinescent Margin of the Bracts of Safflower: (A) Strong; (B) Moderate; (C) Absent.

109
110
111

112
113
114


115 The second stage was conducted in a randomized block, in a 4 x 2 factorial scheme,
116 combining four plant densities per vase and two safflower access, distributed in 4 blocks.
117 Each block was composed of eight vases, each one corresponding to an experimental unit,
118 totaling 32 experimental plots. The densities were evaluated referring to: four; three; two;
119 and one plant per vase.

120
121 After reaching the phase of rosette (30 DAS), the apices of the central stem of the plants
122 were pruned through a single cut. Subsequently, 30 days after the procedure, the following
123 characteristics were evaluated: Plant height (PH (cm)); Stem Diameter (SD (cm)); Number of
124 Branches per plant (NB); Number of Flower Buds (NFB); Flower production (FP).

125
126 The third stage used a completely randomized design, in a 3 x 2 factorial scheme, combining
127 three cutting points of the stems and two safflower accesses, using 4four repetitions. Twelve

128 vases were used for each genotype, wherein each vase received two seeds and
 129 corresponded to one experimental unit, the cut-off points of the stems were: closed buds,
 130 semi-open buds, and open buds. To stimulate the development of lateral buds the apices of
 131 the central stem were pruned.

132
 133 The harvest point of the stems was done according to the respective treatments: open
 134 inflorescences, above 70% of the open florets; semi-open, 30 to 40% of open florets; and
 135 closed, 5 to 15% of the open florets (Figure 5). The harvest point was determined based on
 136 the inflorescences of each stem, and the cut was performed when half of the inflorescences
 137 presented the percentage of open florets corresponding to the treatment. The stems were
 138 cut in the basal portion, about 3 cm from the base of the plant. At laboratory, the flower stem
 139 were evaluated from ornamental characteristics as follows: Number of inflorescences (NI)
 140 Number of Close Buds (NCB); Later the flower stems were placed in containers with tap
 141 water, leaving about 5 cm from the stem base submerged. The flower stems were discarded
 142 when presented an unpleasant visual aspect, with flowers, leaves and stem darkened. The
 143 Post-Harvest Durability (PHD) was consider the number of days from stem cutting to
 144 discard.
 145



146
 147 **Figure 5. Cut-off points: (a) Closed buds; (b) Semi-open buds; (c) Open buds.**
 148
 149

150 For the analysis of variance, the effects of the treatments and the averages were considered
 151 as fixed and treated according to the statistical model for the specific designs of each
 152 experiment.

153
 154 Using the F test at the 5% probability level, were tested the significance of the mean squares
 155 and posteriorly the means were submitted to polynomial regression analysis or comparison
 156 of means by the Tukey test using the GENES program [21].
 157

158 Estimates of variance components and genetic parameters were obtained from the following

159 expressions: $\sigma_g^2 = \frac{QMG - QMR}{r}$, $h^2 = \frac{\sigma_g^2}{QMG/r}$, e $CV_g = \frac{100 \sqrt{\sigma_g^2}}{\mu_g}$, for the genetic variance
 160 among means, heritability coefficient and coefficient of genetic variation, respectively.
 161
 162

163 3. RESULTS AND DISCUSSION

164
 165 The emergence of seedlings began 4 days after the sowing and continued for two days. The
 166 accesses presented 80% and 90% of germination (% G) and 5.47 and 8.22 of Emergency
 167 Velocity Index (EVI) for ICA 193 and ICA 73, respectively.
 168

Comment [REV A4]: PLEASE COULD YOU MIND CONSIDERING SUBTITLES OF 1ST STAGE; 2ND AND 3RD STAGE????

169 Significant differences were observed between the accesses evaluated at the 5% level by
170 the F test for the analyzed characteristics, except for the diameter of the stem, evidencing
171 the existence of genetic variability (Table 1).
172

173 According to [22], the plants are classified as ornamental when they present characteristics
174 that arouse attention and interest, from their aesthetic particularities, referring to the color
175 and shape of leaves and flowers, phenological aspects, among others. Taking these
176 characteristics into consideration, the ICA 73 presented plants with the highest number of
177 branches (9.15), a high number of buds (15.9) and, mainly, high flower production (14.3). On
178 the other hand, it exhibited strong spinescent margin of the leaves and bracts (9.25). On the
179 other hand, ICA 193 access presented the most favorable points to the low and moderate
180 spinescent margin of the leaves and bracts (2.39 and 3.36, respectively). However, showed
181 low branching (5.25) and consequently, lower number of buds and flower production (7.65
182 and 6.95, respectively) (Table 2).
183

184 According to [23], the leaves and bracts margins of safflower plants are peculiarly
185 ~~spinescent, spinescent; however, however,~~ the vehemence with which this characteristic is
186 expressed in the plant varies according to the different genotypes, thus allowing the
187 selection and development of varieties that exhibit a weak or moderate character
188 expression, offering attractive materials to the floriculture market.
189

UNDER PEER REVIEW

190 Table 1. Summary of variances analysis and genetic parameters estimates for Plant Height (PH), Stem Diameter (SD), Leaf Dentin
 191 (LD), Spinescent Margin of the Leaves (SML), Number of Branches (NB), Number of Flower Buds (NFB), Spinescent Margin of the
 192 Bracts (SMB), Flowering (FI), and Flower Production (FP), Recife, 2017.

SV	DF	MS								
		PH (cm)	SD (cm)	LD	SML	NB	NFB	SMB	FI	PF
Acesses	1	2822.4	0.006	30.625	122.5	152.1	680.62	90.0	198.02	540.22
Residual	38	6665.2	0.022	5.62	15.20	8.90	36.75	13.35	22.31	28.18
F		16.09**	0.29 ^{ns}	5.44*	8.06**	17.08**	18.52**	6.74*	8.87**	19.16**
Mean		68.1	0.98	4.12	7.5	7.2	11.77	7.75	56.52	10.62
CV		19.45	15.06	57.50	51.98	41.44	51.48	47.15	8.36	49.97
σ^2_g		132.35		1.25	5.36	7.16	34.03	3.83	8.78	25.60
H ²		93.78		81.63	87.60	94.15	94.60	85.16	88.73	94.78
CV _g		16.89		27.10	30.88	37.17	48.19	25.26	28.25	47.62
CV _g /CV _e		0.87		0.47	0.59	0.90	0.94	0.53	5.24	0.95

* and ** significant at the 5% and 1% levels, respectively, of the probability by the F test and "ns" not significant by the F test

193
194
195
196
197

Table 2 - Average of Plant Height (PH), Stem Diameter (SD), Leaf Dentin (LD), Spinescent Margin of the Leaves (SML), Number of
 Branches (NB), Number of Flower Buds (NFB), Spinescent Margin of the Bracts (SMB), Flowering (FI), and Flower Production (FP),
 Recife, 2017.

Accesses	PH (cm)	SD (cm)	LD	SML	NB	NFB	SMB	FI	FP
ICA 73	59.7b	9.99a	5.0a	9.25a	9.15a	15.9a	9.25a	58.7a	14.3a
ICA 193	76.5a	9.74a	3.25b	2.39b	5.25b	7.65b	3.36b	54.3b	6.95b

198 *Means followed by the same letter do not differ by Tukey test at 5%.

199 Plants with spines have less acceptance in the market, by virtue of limiting the touch, due to
200 the possibility of promoting scratches in the skin. In this context, ICA 193 stands out for
201 having naturally weak or moderate spinescent margins, both in the leaves and in the bracts,
202 presenting viability for insertion in the market of cut flowers without resistance to acceptance.
203 In contrast, access ICA 73 needs to be submitted to breeding programs in order to
204 circumvent this limitation for its use in floriculture.

205
206 Both accesses presented precocity of flowering, and ICA 73 presented a subtle highlight in
207 relation to this phenological stage, starting its flowering about 59 DAS, while the ICA 193
208 access began about 54 DAS. According to [24], this phase starts between 60 and 100 days
209 after sowing and confirms, therefore, the precocity of the materials evaluated in this work
210 (Table 2).

211
212 No early commercial material of *Carthamus tinctorius* intended for the ornamental market is
213 reported, in addition, it is possible to notice a certain difficulty in the development of cultivars
214 of this species that present attributes of ornamental interest and initiate this phenological
215 stage early [15]. Less late varieties, commonly used, show beginning of flowering only 80
216 days after sowing [25,26, 26]. In this sense, the accesses under study have a scarce and
217 desired characteristic, offering a further differential to include these in commerce, not only
218 facilitating acceptance but demonstrating competitiveness with products already available.

219
220 Regarding plant height, ICA 73 showed lower heights than ICA 193, referring to 59.7 cm and
221 76.5 cm, respectively (Table 2). Considering also the use for cutting stem, according to [15],
222 the stem length of products with superior quality must present between 70 cm and 80 cm,
223 however, there are already commercial safflower varieties specific to the ornamental market
224 with stems from 60 cm [25]. In this way, the values demonstrated by the accesses of this
225 research, fit within the allowed for both sides.

226
227 As to leaf dentin, ICA 73 presented moderate intensity (5.0) and weak to moderate ICA 193
228 (3.25) (Table 2). In order to make arrangements, the margin of the leaf does not have a fixed
229 pattern, since even the most unusual can contribute to creative and decorative
230 combinations, including being something very desired to compose bouquets base [27].

231
232 During the experiment, ~~the plants were affected by pathogens and pests~~ pathogens and
233 ~~pests affected the plants~~, since no chemical control was performed. From the symptoms and
234 a previous microscopic analysis, the presence of *Cercospora carthami* and aphid (aphis)
235 was observed (Figures 6a and 6b). However, the inflorescences were not directly affected
236 and the characteristics of interest could be effectively evaluated. [12] reported that, safflower
237 is the target of many pathogens, including fungi, bacteria and viruses, but the first group
238 cited is the most prominent. *Cercospora carthami* is one of the fungi that commonly affect
239 the culture, causing foliar damage. Among the pests, aphids ~~are said to cause the most~~
240 ~~recurrent~~ recurrent damages, however they are less worrisome than diseases [28].

241
242 The summary of the analysis of variance for the second experiment and the estimation of the
243 main genetic parameters for the agronomic and ornamental characters evaluated in the two
244 safflower accesses are organized in Table 3. According to the results, it is possible to
245 observe a significant difference between the accesses at the level of 5% by the F test for the
246 characteristics analyzed.



Figure 6. Safflower plants: (A) Pest attack (aphids); (B) Leaf disease (*Cercospora carthami*).


Table 3. Summary of variances analysis and genetic parameters estimates for Plant Height (PH), Stem Diameter (SD), Number of Branches (NB), Number of Flower Buds (NFB), and Flower Production (FP), Recife, 2017.

SV	DF	MS				
		PH (cm)	SD (cm)	NB	NFB	PF
Blocks	3	36.68	0.03	5.54	7.10	5.78
Accesses	1	570.94*	0.51*	283.52**	264.97**	242.91**
Densities	3	325.76*	0.27*	210.51**	191.36**	188.82**
AccessesxDensities	3	30.41 ^{ns}	0.04 ^{ns}	33.31**	27.76**	27.78**
Residual	21	79.10	0.05	2.90	3.58	2.93
Mean		60.30	1.75	10.69	10.43	10.05
CV		14.74	12.61	15.94	18.14	17.04
σ_g^2		30.74	0.03	17.54	16.34	15.00
H ²		86.15	90.46	98.97	68.65	98.79
CV _g		9.19	9.71	39.17	38.73	38.55
CV _g /CV _e		0.62	0.77	2.46	2.13	2.26

* and ** significant at the 5% and 1% levels, respectively, of the probability by the F test and "ns" not significant by the F test.

Figure 7 graphically shows the behavior of the accesses as a function of the different densities of plants per vase, as well as the equations and coefficient of determination (R²) that best fit the variables studied, according to the regression analysis. All variables can be explained by the linear equation of the 1st degree, with R² values higher than 0.80.

Plant height and stem diameter were inversely proportional to plant density per vase, decreasing as the number of plants increased (Figure 7a). The diameter is an important feature because it is related to rigidity and quality of the stem, since low densities can lead to flexibility and breakage [29]. Pruning did not limit the final length of the plants, which reached values characteristic of the species.



Comment [REV A5]: THIS FIGURES ARE IN SPANISH

267
268
269
270

Figure 7. Plant Height (PH), Stem Diameter (SD), Number of Branches (NB), Number of Flower Buds (NFB), and Flower Production (FP) of ICA 73 and ICA 193 safflower accesses according to four plant densities per pot.

271 The highest values of height and diameter were reached by ICA 193 access, with values
 272 ranging from 56.75 to 70.00, and 1.71 to 2.21 (Figure 7a). In the first case, the values
 273 extrapolate the recommendation of [30] for use in vases, but is suitable for employment in
 274 other areas of social recreation. In this sense, there are already commercial safflower
 275 varieties destined for the ornamental market with heights between 60 and 80 cm, such as
 276 Orange Granade, which is highly prized for beautifying gardens [25]. In contrast, ICA 73
 277 exhibited the smallest values of height and diameter, ranging from 49.00 to 63.25 and 1.44
 278 and 1.80 (Figure 2a). Despite the low diameter values, the stems showed to be well lignified
 279 and no breaks were observed.

280
 281 The decrease due to the increase in the number of plants per vase for number of branches,
 282 buds and flower production was also observed (Figure 7b and 7c) and is in agreement with
 283 the one verified by [31], that in its work with other safflower genotypes evidenced a linear
 284 reduction proportional to the increase of the density. This result is probably linked to
 285 competition between plants for nutrients, water and light, limiting their development [29]. For
 286 use in potted plants or gardens, plants with more branches, provide a aerial part more
 287 voluminous and visually pleasing. Access ICA 73, presented the highest values for these
 288 characteristics and the best density for both genotypes, refers to 4one plant per vase.

289
 290 The high number of branches acts negatively on the uniformity of the opening of the flowers,
 291 due to the different flowering rates of the buds [29]. For ornamental plants destined to
 292 gardens, vases or other leisure areas, this particularity becomes attractive, because it makes
 293 possible that the prestige of the flowers can be realized by a greater period of time, since
 294 while the first flowers are close to senescence, others will still be at the beginning of the
 295 anthesis. On the other hand, this factor is not attractive for cutting stems.

296
 297 In Table 4, the analysis of variance of the third experiment and the estimation of the main
 298 genetic parameters for characters of ornamental importance evaluated in the two accesses
 299 of safflower. The results show a significant difference between the accesses at the 5% level
 300 by the F test for the characteristics analyzed. Table 5 shows the means of the variables that
 301 were submitted to the Tukey test, depending on the treatment factors: accesses and cut-off
 302 points.

303
 304 **Table 4. Summary of variances analysis and genetic parameters estimates for Number**
 305 **of inflorescences (NI), Number of Close Buds (NCB), Post-Harvest Durability (PHD),**
 306 **Recife, 2017.**

SV	DF	MS		
		NI	NCB	PHD
Accesses	1	54.0**	84.37**	88.17**
Cut-off Point	2	26.54**	22.79**	77.17**
AccessesxCut-off	2	13.62**	15.87**	4.67 ^{ns}
Residual	18	1.17	1.12	1.33
Mean		5.33	3.21	10.58
CV		20.03	33.06	10.91
σ_g^2		4.40	6.94	7.24
H ²		97.84	98.67	98.49
CV _g		39.34	82.10	25.42
CV _g /CV _e		1.94	2.48	2.33

307 * and ** significant at the 5% and 1% levels, respectively, of the probability by the F
 308 test and "ns" not significant by the F test.

309 The maturity of the flowers is a very decisive characteristic on the quality of the product and
 310 makes it impossible in most cases to perform a mechanized harvest [15], justifying the
 311 importance of defining the best moment for cutting the stems, as far as the anthesis of
 312 flowers is concerned. Other important information that should be considered refers to the
 313 fact that the central flower opens days before the lateral flowers, about one week, leading to
 314 visual depreciation when the other flowers are opening [32], for this reason the pinch was
 315 performed, stimulating the anther of the lateral flowers in a more uniform way.

316
 317 The number of inflorescences, number of close buds and post-harvest durability should be
 318 analyzed together and as a function of the cut-off point, allowing to establish the
 319 most appropriate combination for quality of the final product.

320
 321 The ICA 73 had a higher average number of inflorescences (10.25) and a lower number of
 322 close buds (1.75) for the cut-off point when buds were open, however, in this same
 323 treatment the lower post-harvest durability of the stems was obtained (4.25), making it
 324 impossible to cut stems of this material at this maturation level, since it does not meet an
 325 adequate number of days of product life (Table 5). At the point of semi-open buds, the
 326 number of inflorescences (5.75) was reduced by half and the number of close buds
 327 increased (6.00), showing a nearly 1:-1 ratio between flowers and buds, indicating little
 328 uniformity of flower anthesis and opening of a few buds after cutting, but exhibited longer
 329 flower durability (11.00) (Table 5). Finally, at the point of closed buds the number of
 330 inflorescences reduced even more, evidencing that some of the heads did not even develop
 331 buds and the low number of close buds also confirms this hypothesis, however, the durability
 332 of the stems was equivalent to the cutting treatment with the semi-open flowers.

333
 334 **Table 5 - Average of Number of Inflorescences (NI); Number of Close Buds (NCB);**
 335 **Post-Harvest Durability (PHD), Recife, 2017.**

	Variables					
	NI		NCB		PHD	
	ICA 73	ICA 193	ICA 73	ICA 193	ICA 73	ICA 193
Open Buds	10.25Aa	4.25Ba	1.75Ac	1.25Bc	4.25Bb	9.75Ab
Semi-open Buds	5.75Ab	4.50Ba	6.00Ab	0.50Bb	11.00Ba	13.50Aa
Closed Buds	4.50Ab	2.75Bb	7.50Aa	2.25Ba	10.75Ba	14.25Aa

336 **Means followed by the same lower letters in column and capital letters on the lines*
 337 *do not differ significantly by the Tukey test at 5%.*

338
 339 ICA 193 presented a lower average number of inflorescences when compared to ICA 73 at
 340 all cut-off points; however, it also exhibited a lower number of close buds, indicating a
 341 greater uniformity of flower anthesis. At the cut-off point with open buds, exhibited the
 342 second highest value of inflorescences (4.25), according to lower flower buds values (1.25)
 343 and lower post-harvest durability of the stems (9.75), however, this useful life is already
 344 acceptable to the market (Table 5). For the cut-off point with semi-open buds, it presented
 345 the highest number of inflorescences (4.50), although it does not differ statistically from the
 346 previous treatment for this characteristic, smaller number of close buds (0.50) and second
 347 highest number of days of stem durability (13.50). Finally, at the cut-off point with closed
 348 buds, the lowest number of inflorescences (2.75) and the highest number of floral buds
 349 (2.25) were observed, evidencing the difficulty of developing the buds after cutting the
 350 stems, also presented the greatest number of days of durability of the stems (14.25), but did
 351 not differ statistically from the previous treatment for this characteristic.

352
 353 According to [32], the most favorable cutting stage of the stems corresponds to the period in
 354 which 20% to 30% of the central florets opened, which is equivalent to the cut-off point

355 denominated in this work as semi-open buds, allowing the others to open in the hands of the
356 consumer, extending the useful life of the product. In agreement with this affirmation, it is
357 observed that ICA 193 presented the best combination of factors for this cut-off point,
358 presenting a higher number of inflorescences, lower number of close buds and greater post-
359 harvest durability. The same observation can be raised for ICA 73, but with some
360 reservations, such as the performance of a removal of the secondary and tertiary branches,
361 improving the aesthetics of the product, since despite an adequate number of inflorescences
362 and stem durability, it presented high number of close buds, or search for improvements of
363 this characteristic through an improvement program.

364
365 The flowers produced by both genotypes showed a yellow color at the beginning of the
366 anthesis, changing to orange shades soon after and presented a very attractive visual
367 aspect, with abundant beauty while fresh and even after a period of drought, offering
368 potential for introduction into the Brazilian flower market, contributing to the supply of news
369 for the sector and the consumer (Figure 8a and 8b). According to [23], the characteristics of
370 greater importance and influence on the ornamental value of safflower are attributed to the
371 color of the flowers, where the oranges and yellows stand out, along with the weak
372 spinescent margin of the leaves and bracts.

373



374
375 **Figure 8. (A) Bouquet of stems after cutting; (B) Change the color of the flower.**

376
377 The evaluated characteristics presented high estimates of the genetic parameters of
378 heritability and ratio between the coefficients of genetic and experimental variation, a very
379 favorable point in breeding programs, since it indicates in a general way that these
380 characters can be easily improved through classic methods [33] and provide favorable
381 conditions for realization of selection, allowing to obtain high genetic gain within the first
382 cycles [34].

383
384 Considering the differences evidenced between the genotypes, together with the
385 completeness they demonstrate for characteristics of ornamental interest, these genotypes
386 suggest potential for inclusion in an improvement program, in order to obtain a material that
387 groups the positive characteristics presented in both accesses. [35] point out that one of the
388 criteria for success in crossbreeding depends on the divergence between the parents,
389 parallel to the superior performance they present referring to the characteristics of interest of
390 the breeder.

391

392 **4. CONCLUSION**

393

394

395

396

397

398

399

400

401

402

REFERENCES

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

1. Romero DHS, Restrepo IME. Local competitive profile as a determining factor for the development of floriculture in Madrid (Cundinamarca). Research and reflection. 2011; 19 (2): 25-43.

2. Hortiwise. A Study on the Kenyan-Dutch Horticultural Supply Chain. The Dutch Ministry of Economic Affairs, Agriculture & Innovation The Netherlands. 2012.

3. Mitsueda NC, Costa EV; D'oliveira PS. Environmental aspects of agribusiness flowers and ornamental plants. Magazine in Agribusiness and Environment. 2011; 4 (1): 9-20.

4. Ibraflor. Brazilian Institute of Floriculture. The flower market in Brazil. 2017. Accessed: 20 December 2018. Available: <http://www.ibraflor.com/site/2017/11/04/florente-de-flores-vera-longuini/>.

5. Junqueira AH, Peetz MS. The productive sector of flowers and ornamental plants of Brazil, from 2008 to 2013: updates, balance sheets and perspectives. Brazilian Journal of Ornamental Horticulture. 2014; 20 (2): 115-120.

6. Ibraflor. Brazilian Institute of Floriculture. Sector Numbers. 2014. Accessed: 28 November 2017. Available: http://www.IBRAFLOR.com/ns_mer_interno.php.

7. Dantas CVS, Silva IB, Pereira GM, Maia JM, Lima JPMS, Macedo CEC. Influence of sanity and water deficit on seed germination of *Carthamus tinctorius* L. Brazilian Journal of Seeds. 2011; 33 (3): 574-582.

8. Santos RF, Silva MA. *Carthamus tinctorius* L.: An alternative crop for Brazil. Acta Iguazu. 2015; 4 (1): 26-35.

9. Zareie S, Mohammadi-Nejad G, Sardouie-Nasab S. Screening of Iranian safflower genotypes under water deficit and normal conditions using tolerance indices. Australian Journal of Crop Science. 2013; 7(7): 1032-1037.

10. Landau S, Friedman S, Brenner S, Bruckental I, Weinberg ZG, Ashbell G, Hen Y, Dvash L, Leshem Y. The value of safflower (*Carthamus tinctorius* L.) hay and silage grow under Mediterranean conditions as forage for dairy cattle. Livestock Production Science. 2004; 88: 263-271.

11. Asgarpanah J, Kazemivash N. Review: Phytochemistry, Pharmacology and Medicinal Properties of *Carthamus tinctorius* L. Chin J Integr Med. 2013; 19(2): 153-159.

12. Mündel, H. H.; Blackshaw, R. E.; Byers, J. R.; Huang, H. C.; Johnson, D. L.; Keon, R.; Kubik, J.; Mckenzie, R.; Otto, B.; Roth, B; Stanford, K. Safflower production on the Canadian

Comment [REV A6]: PLEASE CONSIDER IN ALL DOCUMENT 3 OR 4 REFERENCES FROM JEAI

Formatted: Font: Italic

445 prairies: revisited in 2004. Agriculture and Agri-Food Canada. Lethbridge Research Centre:
446 Lethbridge; 2004.
447
448 13. Pintão AM, Silva IF. The Truth About Saffron. In: Workshop medicinal and
449 phytotherapeutic plants in the tropics. Lisbon. Summaries ... Lisbon: IICT / CCCM, 2008.
450 Electronic Version.
451
452 14. Jadhav BA, Joshi AA. Extraction and Quantitative Estimation of Bio Active Component
453 (Yellow and Red Carthamin) from Dried Safflower Petals. Indian Journal of Science and
454 Technology. 2015;8(16)1-5.
455
456 15. Uher, J. Safflower in European floriculture: a review. In: Proceedings of the 7th
457 International Safflower Conference, Wagga Wagga, New South Wales, Australia. 2008.
458
459 16. Bradley VL, Guenther RL, Johnson RC, Hannan R M. Evaluation of safflower
460 germplasm for ornamental use. In: Perspectives on new crops and new uses (J. Janick, ed.).
461 ASHS, Press: Alexandria. 1999. 433-435.
462
463 17. Oliveira GG. *Trichoderma* spp. plant growth and biocontrol of *Sclerotinia sclerotiorum*
464 and pathogens in safflower seeds (*Carthamus tinctorius*). 2007. 79 p. Dissertation (Master in
465 Plant Production), Federal University of Santa Maria, Santa Maria, 2007.
466
467 18. Zoz T. Correlation and analysis of yield traits in grains and their components and plant
468 characteristics in safflower (*Carthamus tinctorius* L.) and castor bean (*Ricinus communis* L.).
469 2012. 54 p. Dissertation (Master's degree in Agronomy) Faculty of Agronomic Sciences-
470 UNESP; 2012.
471
472
473 19. Maguire JD. Speed of germination aid in selection and evaluation for seedling
474 emergence and vigor. Crop Science. 1962;2(2):176-77.
475
476 20. Stumpf ERT, Heiden G, Barbieri RL, Fischer SZ, Neitzke RS, Zanchet B, Grolli PR.
477 Method for evaluating the ornamental potential of native and unconventional flowers and cut
478 foliage. Brazilian Journal of Ornamental Horticulture. 2007; 13: 143-148.
479
480 21. Cruz CD. GENES - versão Windows. Viçosa- MG: Editora UFV; 2006.
481
482 22. Gonçalves MF, Melo AGC. Floristic analysis of ornamental plants implanted in Garça
483 Forest / SP. Electronic Journal of Forest Engineering. 2013; 21 (1): 12-24.
484
485 23. Pahlavani MH, Mirlohi AF, Saeidi G. Inheritance of flower colour and spininess in
486 safflower. Journal of Heredity. 2004;95:265-267.
487
488 24. Emongor V. Safflower (*Carthamus Tinctorius* L.) the underutilized and neglected crop: A
489 review. Asian Journal of Plant Science. 2010;9(6):299-306.
490
491 25. Sakata. Sakata's reliable seeds. Flower seed. Sakata seed corporation, Yokohama,
492 Japan. 2010. 87 p.
493
494 26. Geneses Seeds Ltd. Catalog Flowers. Accessed: 20 November 2017. Available:
495 http://www.genesisseeds.com/PDFs/GENESIS_CATALOG_FLOWERS.pdf.
496

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- 497 27. Morais ÉB, Castro ACR, Silva TF, Soares NS, Silva JP. Evaluation of potential use of
498 | native Anthurium foliage. Ornamental Horticulture. 2017;_23:07-14.
499
- 500 28. Coronado LM. El cultivo Del cártamo (*Carthamus tinctorius* L.) em México. Instituto
501 | nacional de investigacion esforestales, agrícolas y pecuárias. México, 2010.
502
- 503 29. Bellé RA, Rocha EK, Backes FAAL, Neuhaus M, Schwab NT. Safflower grown in
504 | different sowing dates and plant densities. Ciência Rural. 2012;_42(12).
505
- 506 30. Motos J, Oliveira MJG. Produção de crisântemos em vaso. Holambra, Flortec. 1998. 34
507 | p.
508
- 509 31. Sampaio MC. Cultivo de cártamo (*Carthamus tinctorius* L.) sob variação de adubações,
510 | densidades e épocas de plantio. 2016. 63p. Dissertação (Mestrado em Engenharia de
511 | Energia na Agricultura), Universidade Federal do Oeste do Paraná, Cascavel, 2016.
512
- 513 32. Wien HC. Cut Flower Cultural Practice Studies and Variety Trials 2012. Dept. of Hort.,
514 | Cornell Univ. Ithaca, 2012.
515
- 516 33. Fox GP, Bowman J, Kelly A, Inkerman A, Poulsen D, Henry R. Assessing for genetic and
517 | environmental effects on ruminant feed quality in barley (*Hordeum vulgare*). Euphytica.
518 | 2008;_163: 249-257.
519
- 520 34. Vencovsky R, Barriga P. Genética Biométrica no Fitomelhoramento. Ribeirão Preto:
521 | SBG; 1992.
522
- 523 35. Cruz CD, Carvalho SP, Vencovsky R. Estudos sobre divergência genética II. Eficiência
524 | da predição do comportamento de híbridos com base na divergência de progenitores.
525 | Revista Ceres.1994;_41(234):183-190.

Formatted: Font: Italic