

BIOCHEMICAL AND HISTOLOGICAL INVESTIGATIONS OF ALCOHOL ADMINISTRATION IN WISTAR RATS

ABSTRACT

Introduction: Knowledge of biochemical and histological investigation of alcohol administration in Wistar rats is critical for contemporary effort to develop animal models of alcoholism.

Materials and Methods: 20 Male Wistar rats weighing (63.50 ± 3.79 g), were divided into four groups (consisting 15 treated animals and 5 control animals) and administered with varying concentrations of ethanol (5% 15% and 40%) via gavage for a period of 28 days. Probiotic evaluations, liver biochemical enzymes and alteration in histology profile of gastrointestinal tract (GIT) and viscera organs were accessed after a period of 28 days ethanol administration.

Results and Discussion: The result of biochemical study of 40% ethanol showed a significant decrease in serum gamma glutamyl transferase (GGT), serum aspartate (AST) and Alanine amino transferase (ALT) when compared to normal study while 5% and 15% ethanol intoxicated rats are within the range with respect to the normal study. The results of probiotic evaluations such as body weight, water intake and food intake show a percentage decrease in 40% ethanol administrated rat when compared with controls. The pictorial results of liver histopathology organs that received 5% and 15% ethanol did not showed a significant degeneration in histology profile when compared to the normal study while morphology degeneration in histology profile occurred in 40% ethanol administrated rats.

Conclusion: Therefore serum aspartate (AST), gamma glutamyl transferase (GGT) and Alanine amino transferase (ALT), probiotic evaluation (body weight, food intake and water intake) coupled with histopathological investigation may be used as biomarker for the early diagnosis of ethanol toxicity in human beings.

Keywords: alcohol, biomarkers, histopathology, growth performance index

Introduction

Over the centuries, alcohol has become the most socially-accepted addictive drug worldwide (Ohkubo *et al.*, 2009). Its use antedates recorded history and may go back as far as the Neolithic age around 8000BC. Ethanol is found associated with varieties of our cultural life, various names

have been ascribed to it. Among them are:- whisky in Gallic, water of life, Sapele water, gin, ogogoro, ojuna etc has been the nomenclature of alcohol in various regions and spheres. Ethanol is the type of alcohol found in alcoholic beverages (wine, beer and spirit).

Alcohol intoxication is the term used by the toxicologist to describe the point at which alcohol depresses the central nervous system so that mood, physical and mental abilities noticeably change (Sainlan, 2008). Toxicologist used the term “alcohol intoxication to discriminate between alcohols. Intoxication is the consequence of alcohol entering the bloodstream faster than it can be metabolized by the liver, common symptoms of alcohol intoxication include slurred speech, euphoria, impaired balance, loss of muscle coordination, flushed face, dehydration, vomiting, reddened eyes, reduce inhibitions and erratic behaviour. Sufficiently high levels of blood-borne alcohol will cause coma and death from the depressive effects of alcohol upon the central nervous system (Smith *et al.*, 2005).

Although the precise mechanism of alcohol intoxication is presently unknown, but studies suggest that its passes directly from the digestive tract into the blood stream in minutes, blood transports the alcohol to all parts of the body including the brain which alter their neurons in several ways by changing their membranes as well as their ion channels enzymes and receptors. (Aguayo *et al.*, 2002). Long-term use of alcohol in excessive quantities is capable of damaging practically every organ system in the body, (Testino, 2008).

Alcohol biomarkers have important applications in medicine and public safety (Litten and Fertig, 2003). The aim of this study was to investigate the biochemical and histological effects of alcohol administration in Wistar rats.

Materials and Methods

Animal Collection

Twenty Wistar rats ((63.50± 3.79g) were obtained from breeding stock maintained in the animal house of the College of health sciences, Osun State University Osogbo main campus (UNIOSUN) and were authenticated by the farm Director UNIOSUN.

The animals collected were house in well-ventilated wired plastic metabolic cages in the animal facility of the department of Biochemistry, Federal Polytechnic Ede, and approval was obtained from the Departmental Ethical Committee on animal usage.

The rats were maintained under standard room temperature (25-26⁰C) and humidity of 65.5%. They were allowed unrestricted access to water and rat chow (Tina Livestock feeds Ltd, Oke Gada Ede Osun state Nigeria).

They were allowed to acclimatize for a period of 21days before the commencement of experiments, the weight of the animals were estimated at procurement, during acclimatization, at commencement of the experiment and every day throughout the duration of the experiments using an electronic analytical precision balance.

Experimental Design

Twenty male Wistar rats (63.50± 3.79g) were used for this study. They were divided into 4 groups of 5 rats each, 1st group served as the control administered saline, 2nd-4th groups were administered 5%, 15% and 40% ethanol respectively for 28 days by gavage i.e. intra-gastric administration. All animals had access to rat chow and water *ad libitum*.

Animal experiments complied with the ARRIVE guidelines and was carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, EU Directive 2010/63/EU for animal experiments, was strictly followed throughout the experiment.

Animal Sacrifice and Sample Collection and Preparation

On the 29th day of the experiment, the rats were fasted overnight, sacrificed via cervical dislocation. Blood samples were collected via cardiac puncture, into plain tubes, centrifuged at 3000g for 5min, and serum collected and stored at -20°C for further analysis. The liver, kidney, heart and lungs, were excised and fixed in 10% formal-saline for histopathological interpretations.

Ethical Approval

Ethical approval for the study was obtained from Ethics Review Committee of College of Medicine University of Lagos, Idi-Araba with CMUL HREC REGISTRATION NUMBER: HREC/15/04/2015

Estimation of Biochemical Parameters

The methods of Reitman and Frankel (1957) and Hammed, (2011) were used for the determination of alanine amino transferase (ALT) and aspartate amino transferase (AST) respectively, while that of Szasz, (1969), Hyder *et al.*, (2013) was used for the determination of gamma glutamyl transferase (GGT) in the serum.

Determination of volume of alcohol and route of administration

The ethyl alcohol used in this study was reagent grade 200% proof. The volume of alcohol administered to the animals was calculated using the Widmark (1981) formula modified by Bouwer, (2004). Alcohol concentrations used in this study was 10-30% to represent the three classes of alcoholic beverages commonly consumed by man. Less than 10% alcoholic beverage

content represent beers, while 10-20% represents the wine group while 30% and above represents the spirit group.

Histological Procedure and Analysis

This was done as described by Saalu *et al.*, (2008), briefly, the organs were cut on slabs about 0.5cm thick and fixed in 10% formal saline for a day after which they were transferred to 70% alcohol for dehydration the tissues were passed through 90% alcohol and chloroform for different durations before they were transferred into two changes of molten paraffin wax for 20min each in an oven at 57⁰C. Serial selections of 5mm thick were obtained from a solid block of tissues and were stained with haematoxylin and eosin stains, after which they were passed through a mixture of equal concentration of xylene and alcohol. Following clearance in xylene, the tissues were oven-dried; photomicrographs were taken with a colour digital camera mounted on a light microscope.

Statistical Analysis

The SPSS v. 25.0 computer software package (SPSS Inc. Chicago, U.S.A) and GraphPad Prism 7.0 were used for this study. The results are presented as mean \pm standard error of the mean ($X \pm SEM$).

RESULTS AND DISCUSSION

Table 1: Body weight, feed and water intake of rats administered alcohol for 28 days

Probiotic Indices	Control	5% EtOH	15% EtOH	40% EtOH
B.W (g)	85.33 ± 11.52	89.67 ± 10.04	79.33 ± 8.95	73.50 ± 6.99
H.I (ml)	90.00 ± 33.15	88.67 ± 21.06	62.00 ± 27.68	53.17 ± 10.33
F.I (g)	90.00 ± 7.46	79.33 ± 9.86	67.33 ± 16.45	37.17 ± 14.93

Data are expressed as mean ± standard error of the mean (SEM) of five normal and 15 intoxicated rats.

Key: B.W-Body Weight, H.I- H₂O Intake, F.I-Feed Intake

Table 2: Serum Activities of ALT, AST & GGT of rats administered alcohol for 28 days

Parameters	Control	5% EtOH	15% EtOH	40% EtOH
GGT(U/L)	148.16±4.86	151.19±5.11	152.36±8.44	158.36±8.44
AST(U/L)	433.09±1.66	719.70±2.40	725.45±8.79	733.03±15.95
ALT(U/L)	152.20±3.40	256.49±8.01	257.43±1.55	258.43±9.64

Data are expressed as mean ± standard error of the mean (SEM) of five normal and 15 intoxicated rats.

Key: EtOH-ethanol, ALT- alanine amino transferase, AST- aspartate amino transferase, GGT- gamma glutamyl transferase.




Plate 1: Photomicrographs of **Heart** of Alcohol Administered Rats

1A: Control rats showing normal study.

1B. Rats administered 40% ethanol also showing normal heart architecture.

Key: **A:**Control, **B:** 40% EtOH




Plate 2: Photomicrographs of **Liver** of alcohol administered rats.

2A. Well preserved liver architecture showing normal appearing portal tract.

2B. Normal study with some ghost appearance of the tubules (Acute tubular necrosis **ATN**).

Key: **A:** Control, **B:** 15% EtOH




Plate 3: Photomicrographs of **Kidney** of alcohol administered rats

3A: Normal glomeruli **G** and tubules **T** in Kidney i.e. normal study, i.e. the glomeruli **G** appears normal with obvious central vein.

3B: At 15% ethanol concentration sloughing off of cells that lines tubules, a sign of **ATN** i.e. acute tubular necrosis.

Key: **A-** Control **B-**15% etoh




Plate 4: Photomicrographs of **Lungs** of alcohol administered rats

4A: Normal lungs showing clear alveolar spaces **CAS**.

4B: At 40% ethanol administration there was intense Inflammatory Cells **IC** within the interstitium and Congestion **C**

Key: **A-** Control **B-**40% etoh

Alcohol (ethanol) may lead to hepatotoxicity; ALT, AST, and GGT are most commonly used makers of hepatocyte injury, they are more specific enzymes biomarkers in intoxication experiments Palmer, (2004).

Ethanol administrated rats recorded a significant increase in gamma-glutamyl transferase (GGT), Aspartate amino transferase (AST), Alanine amino transferase by; GGT (1.2%, 0.2% and 0.5%) AST (0.13%, 0.1% and 5.21%) and ALT (0.11%, 0.21% and 11.49%) followed the administration of various concentrations of ethanol solution (Table 2) as compared to the control group respectively.

In agreement with the present study, Chen *et al.*, (2003) observed a significant increase in AST and ALT after moderate drinkers (at least once per month, < 210g ethanol/week for men <140g ethanol/week for women). In addition, Onyesom and Anosike (2007) recorded elevation in AST and ALT in rabbit orally given 1.5g ethanol/ kg body weight as single daily dose for a continuous period of fifteen weeks. The increase in enzyme activity was mainly due to the effect of ethanol that interpolates and expands bio membranes leading to increased membrane fluidity and enzyme release (Yang *et al.*, 2005). Following the treatment with alcohol, there were significant elevations in GGT, AST and ALT (Group C, 40% Ethanol) which confirms the likely hepatotoxic effect of alcohol. This finding is in line with the report of Maher (1997).

The evaluation of liver function by measuring serum GGT, AST and ALT of alcohol intoxicated rat can be used for the study of human consumption because most of the alcohol consumed by peoples is metabolized by the liver. Therefore, the liver is constantly saddled with the responsibility of detoxifications of substances ingested. It is documented that a number of potentially dangerous by-product are generated (Maher, 1997) these by-products especially free

radicals are known to cause destruction of the liver cell hence elevation of ALT, AST and GGT Onyesom and Atakuo, (1998).

The effect of ethanol on the body weight was also assessed in this study; Table 1 showed the change in body weight of rats before and after administration of ethanol solutions. Here the weight observed in intoxicated rats recorded a significant decrease by 3.6%, 1.9% and 1.3% as compared to controls. This was in accordance with the study of Rajakrishnan *et al.*, (1997) who found out that changes in the body weight of the rat may be due to the deposition of lipids in adipose tissue and fluid accumulation in the organ.

Water intake level was also measured in this study, a notable difference was observed in alcoholic treated rat, water intake was affected at the higher concentration while a less significant increase was observed in the control group.

Food intake level were also measured in this study, significant differences in daily food intake were observed between saline – control and alcohol treated rat utilizing oral administration of alcohol, this is in line with the previous study of Callaci *et al.*, (2006) that a significant difference was noticed in between control and alcohol treated rat.

This study examined the slide of Heamatoxlin and Eosin (H&E) stained tissues of the liver, kidney, heart and lungs of all the study animals that received (5%, 15%, 40%) ethanol and the control group that received 0.9% normal saline. The major histopathological changes occurred in organs of animals that received 40% ethanol, there were minimal histopathological changes in the organ tissues of the rat received 5%, and 15% ethanol indicated that high concentration of ethanol is required to caused significant histopathology changes in the liver, kidney, heart and lungs.

In all the experimental groups that received alcohol, none of the heart tissue showed significant degeneration in histology profile when compared to the control group that received normal saline. The liver tissues of 15% alcohol showed well preserved liver architecture with normal appearance in the portal tract when compared with the liver tissues of the control, this was in agreement with (Adedapo *et al.*, 2009) who observed no abnormal features in the histopathology examination of the liver tissue. This could have been caused by low doses used in the study. The kidney tissues that received 15% ethanol showed sloughing off of cells that line tubules with some ghost appearance of the tubules causing acute tubular necrosis. This was in accordance with the finding of Kasolo *et al.*, (2011) that the kidney tissue showed expanded and congested glomeruli, mononuclear cellular infiltration which are features of mild nephritis that caused kidney tissue damage followed moderate ethanol intake.

Alcohol detoxification in the body is majorly carried out by the liver. The present study revealed that liver damage is not a consequence of alcohol concentration as any level may be cause damage in different forms. The kidney major function is basically water and electrolyte balance. Renal damage has been found to be associated with acute intoxication or chronic alcoholism (Vamvakas *et al.*, 1998; Dawodu *et al.*, 2017).

Histopathological investigation of lungs that received highest concentration of ethanol 40% showed intense inflammation cells within the interstitium and congestion. This present study was in agreement with previous study on alcohol where the acute histopathological change in lungs, kidney and liver were documented at 45% alcohol for 4weeks (Abdelgadir *et al.*, 2010).

Gradual mortality was observed following ethanol administration into rat with regard to varying concentration of ethanol. There were progressive toxic signs and symptoms which resulted in pre-terminal death. Gross pathological symptoms were observed in the rat of high concentrated

ethanol treated rat. There is an observation of gross abnormality that could be attributed to ethanol administration at the time of autopsy.

Conclusion and Recommendation

Though moderate alcohol intake have shown beneficial effect, administration of different concentrations of alcohol in this study caused different alteration to the visceral organs such as liver, kidney, heart and lungs. Varying concentration of ethanol intoxications also leads to a disturbance of certain metabolic parameters that can be used as makers for early detection of ethanol toxicity.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- Adedapo A.A., Mogbojuri O.M., and Emikpe B.O. (2009). Safety evaluations of the aqueous extract of *Moringa oleifera* in rats. *Journal of Medicinal Plants Research* **3**(8):586-591.
- Abdelgadir E.H., Ahmed R.H., Adam S.L.Y., and Husein A.M. (2010). Evaluation of toxicological activity (acute and sub chronic toxicities) of *Lawsonia innermis* seeds on Wistar rats. *Journal of Pharmacological Toxicology* **5**(7):324-333.

- Aguayo L.G., Peoples R.W., Yeh H.H., and Yevenes G.E. (2002). GABA-A receptors as molecular sites of ethanol action. Direct or Indirect actions? *Current Topics in Medicinal Chemistry* **2**(8):869-885.
- Brouwer I.G. (2004). The Widmark formula for alcohol quantification. *Journal of the South African Dental Association* **59**:427–428.
- Callaci J.J., Juknelis D., Patwardhan D., and Wezeman F.H. (2006). Binge alcohol treatment increases vertebral bone loss following ovariectomy; compensation by intermittent parathyroid hormone. *Alcohol Clinical and Experimental Research* **30**(4):665–672.
- Chen J., Conigrave K.M., Mascaskill P., *et al* (2003). On behalf of the World Health Organization and the International Society for Biomedical Research on Alcoholism Collaborative Group. Combining carbohydrate-deficient transferrin and gamma-glutamyl transferase to increase diagnostic accuracy for problem drinking. *Alcohol and Alcoholism* **38**(6):574-582.
- Dawodu O.G., Ebuehi O.A.T., and Odesanmi O.S. (2017). Comparative Safety Profiles of Pure and Alcohol Beverages in Wistar Rats. *British Journal of Medicine and Medical Research* **20**(9):1-11.
- Hammed M. A. (2011). Metabolic profile of rats after one hour of intoxication with a single oral dose of ethanol. *Journal of Pharmacology and Toxicology* **6**:158-165.
- Hyder M.A., Hassan M., Mohieldein A.H. (2013). Comparative Levels of ALT, AST, ALP and GGT in Liver associated Diseases. *European Journal of Experimental Biology* **3**(2):280-284.

- Kasolo J., Bimenya G.S., Lonzy O., and Ogwal-Okeng J.W. (2011). Sub-acute toxicity evaluation of *Moringa oleifera* leaves aqueous and ethanol extracts in Swiss Albino rats. *International Journal of Medicinal Plants Research* **1**(6):75-81.
- Litten R.Z and Fertig J. (2003). Self-report and biochemical measures of alcohol consumption. *Addiction* **98**(2):3-4.
- Maher J.J. (1997). Exploring alcohol's effects on liver function. *Alcohol Health Research World* **21**(1):5-12.
- Ohkubo T., Metoki H., and Imai Y. (2009). Alcohol Intake Circadian blood pressure variation, and stroke. *Hypertension* **53**:4-5.
- Onyesom I. and Anosike E.O. (2007). Changes in Rabbit Liver function markers after chronic exposure to ethanol. *Asian Journal of Biochemistry* **2**:337-342.
- Onyesom E.O. and Atakuo (1998). An investigation into the relationship between alcohol-induced changes in serum triacylglycerol and blood pressure. *Nigerian Journal of Biochemistry and Molecular Biology* **13**:79-83.
- Reitman S. and Frankel S. (1957). A colorimetric method for the determination of serum Glutamic oxaloacetic and glutamic pyruvic transaminases. *American Journal Clinical Pathology* **28**:56-63.
- Rajakrishnan V., Viswanathan P., and Menon V.P. (1997). Adaptation of siblings of female rats given ethanol: Effect of N – acetyl – L – cysteine. *Amino acids* **12**:323-41.
- Saalu L.C., Jewo P.I., Fadeyebi L.O., and Ikuerowo S.O. (2008). The effect of unilateral varicocele on contralateral testicular Histo – morphology in *Rattus norvegicus* *Journal of Medical Science* **8**(7):654 – 655.

- Smith C., Marks A.D., and Liberman M. (2005) Marks Basic Medical Biochemistry; A Clinical Approach, 2nd Ed Lippincott Williams & Williams, USA. p.458.
- Szasz, G. (1969). A kinetic photometric method for serum gamma glutamyl transpeptidase (GGT). *Clinical Chemistry* **15**(2):124-36.
- Testino G. (2008). Alcoholic diseases in hepato-gastroenterology: a point of view. *Hepatogastroenterology* **55**(82-83):371-7.
- Vamvakas S., Bruening T., Thomasson B., Lammert M., Baumueler A., Bolt H.M., Dekant W., Birner G., Hensler D., and Ulm K. (1998). Renal Cell Cancer correlated with occupational exposure to trichloroethene, *Journal of Cancer Research Clinical Oncology*. **126**(3):178-80.
- Yang S.C., Huang J.R., Chen C.L., Chiu M.J., and Shieh S.J. (2005). Regulation of total serum Protein; Effect of antioxidant capacity on isolated rat hepatocytes. *World Journal of Gastroenterology* **11**:7272-7276.
- Widmark E. M. P. (1981). Principles and Applications of Medicolegal Alcohol Determination. Davis, CA: Biomedical Publications. 163 pages.