

1 **Original Research Article**

2 **Assessment of Production and Utilization of Black Cumin**

3 **(*Nigella sativa*) at the Oromia Region State, Ethiopia**

4

5

6

7 **ABSTRACT**

Background and Objective: Black cumin seed for local consumption and other importance, such as oil and oil rosin for medicinal purposes, export market, crop diversification, income generation, reducing the risk of crop failure and others made Black cumin as a best alternative crop under Ethiopian smaller land holdings. The objective of this study were to examine factors affecting farmer perception of the Black cumin production importance, and assess the crop utilization purpose by smallholder farmers and its income potential for the farmers in two district of Bale zone at Oromia region in Ethiopia.

Materials and methods: The survey was conducted from January to May 2018 in two districts of Oromia region. Questionnaires, focus group discussion and field observation were used to collect data. A total of 180 Black cumin producer farmers were selected randomly from 8 woredas and their response were analyzed by using descriptive statistics and Probit model in Stata version 13.

Results: The survey result indicated that, the majority (95.56%) of the households perceived that production of Black cumin crop is important. The crop used as source of better income, medicinal crop and spice in the study area. From the total mean of agriculture income, Black cumin production contributes about 39.88% of income for the study respondents. The Probit model shows that producers perception of the importance of black cumin production were found to be statistically and significant affected by age of households, education level, availability of labor for farm activities, access to credit facilities, average income from Black cumin, and its productivity level through time. **Conclusion:** The agricultural policy should give emphases at all operational level to exploit more benefit from this crop and on the production enhancement strategies, so as to bring foreseen change in the lives of producers.

8 *Keywords: [Black cumin, perception, production, use, Tobit model, Ethiopia]*

9

10 **1. INTRODUCTION**

11 Black cumin (*Nigella sativa* L.) belongs to the family Ranunculaceae. The crop is native to

12 the Mediterranean region and it has been used for thousands of years by various cultures

13 and civilizations. It grows to 20–30 cm (7.9–12 in) height, with finely divided, linear (but not

14

15 thread-like) leaves. The flowers are delicate, and usually coloured pale blue and white, with
16 5–10 petals. The fruit is a large and inflated capsule composed of 3–7 united follicles; each
17 containing numerous seeds and the seed is used as a spice [1, 2, 3].

18

19 Black cumin is one the most revered medicinal seeds in history. Though Black cumin seeds
20 are mentioned in the Bible as well as in the words of the Prophet Mohammed, they were not
21 carefully researched until about 1550 many years ago. Since 1959, over 200 studies at
22 international universities and articles published in various journals have shown remarkable
23 results supporting its traditional uses recorded almost 1400 years ago [1,2,4]. Dioscoredes,
24 a Greek physician of the century recorded that Black seeds were taken to treat headaches,
25 nasal congestion, toothache, and intestinal worms. They were also used, he reported, as a
26 diuretic to promote menstruation and increase milk production [1, 5]. Besides, Black Cumin
27 has a long history of uses for food flavors, perfumes and medicinal values. Oil has been
28 used for bringing smell to some medicines, sterilizing of surgical operation fiber, production
29 of some veterinary and agricultural medicines and plastic components. Black Cumin seeds
30 have an aromatic odor and bitter taste. They are used as an essential ingredient in soup
31 component, sausages, cheese, cakes and candies [6].

32

33 The Ethiopian variety of cumin seed accumulate up to 50% thymol, a monocyclic phenolic
34 compound. The presence of this compound makes cumin valuable source for health care
35 Industry [7] and medicinal purposes [8]. Moreover, study on phytochemical analysis of
36 Indian and Ethiopian Black cumin seeds it was investigated antioxidant capability and
37 phenolic content are higher in Ethiopian Black cumin than the Indian origin whereas
38 phytochemical content varies in each part of the seed. Seed coats of the *Nigella sativa* are
39 rich in phytochemicals rather than cotyledon as many chemical compounds are
40 concentrated into seed coat. Hence, Black cumin confirms to be a medicinal plant rich in
41 phytochemicals [9]. In Ethiopia, it is commonly used in Amharic "*Berbere*" in which it tends

42 to reduce its hotness [10], for preparation of curries, bread, katicala [1], "Shamita" [12],
43 traditional Ethiopian stews, "Wot" and preservation of butter.

44

45 In Ethiopia, the weather makes a suitable environment for the growth of Black cumin seed.
46 In the country, lot areas in Amara, Oromiya, SNNP, and Gambiella regions are found in
47 producing the Black cumin seed. Most Ethiopian people use as house holdings spice
48 preparation. Studies, also confirms that the application of Black cumin seed for medicinal
49 purpose for internal as well as external treatment problems. Besides its medicinal
50 importance, Black cumin (*Nigella Sativa*) seed is also used for production of soap, perfumes
51 and lotions, food flavorings, food preservation, nutraceuticals and cosmoceuticals from the
52 Black cumin oil [1,4].

53

54 More recently a great deal of attention has given to the seed and oils yields of Black cumin.
55 Due to this, their consumption has thus increased and Black cumin is the second cash crop
56 exported next to ginger in Ethiopia. Ethiopian annual production of Black cumin seed is
57 18000 metric tons 2014/15 [2014/15 [13]] and the national average of Black cumin
58 productivity is 0.79 tons per hectare [14]. However, the crop is produced on fragmented land
59 and soils having long cereal cropping history where crop residues are removed for various
60 purposes without any chemical fertilizer application. Additionally, information regarding its
61 response to fertilizer is insufficient in the country. Black cumin seed shows significant
62 variations in days to flowering in the tested varieties at various locations [15]. The vast
63 majority of Ethiopia's Black cumin exports go to Arabic countries, which, together with other
64 predominantly Muslim countries, accounted in 2008 for some 98% of national exports. It is
65 uncertain how reliable this market is and whether exports can be maintained at current
66 levels. Value-adding to cumin in Ethiopia is low, with all exports being made in the form of
67 whole grain [16].

68

Moreover, the production and land coverage of Black cumin has been increasing; the productivity is still less than 300 kg per hectare. Several problems including lack of improved seed, recommended fertilizer rate, lack of knowhow on postharvest handling; improved agriculture practices and extension system, marketing system, etc. are accountable for the continued low productivity and production of Black cumin [17]. Moreover, the crop is produced on fragmented land and soils having long cereal cropping history where crop residues are removed for various purposes. Due to the increased demand of Black cumin seed for local consumption and other importance, such as oil and oil rosin for medicinal purposes, its export market, its potentiality in crop diversification, income generation and its importance to reduce the risk of crop failure and others made Black cumin as a best alternative crop under Ethiopian smaller land holdings [18].

Despite the country's favorable environmental condition for its production, its importance in the economy, the Black cumin cropping system has been given little attention to improve its production and productivity, and hence, it remained an underutilized crop. The objective of this study were to examine factors affecting farmer perception of the Black cumin production importance, and establish the current status of the crop on smallholder farming sector focusing on its general utilization purpose, and income potential for the farmers in two district of Bale zone of Oromia region in Ethiopia.


2. MATERIAL AND METHODS

2.1 Description of the Study Area

Bale zone is one of the 18 administrative zones in Oromia national regional state which is located in south-eastern Ethiopia. It has borderlines with Arsi, Guji, West and East Hararge zones as well as Somali and Southern Nations and Nationalities and Peoples' Regional States. It has 18 districts out of which nine are located in highland agro-ecology whereas the remaining nine are located in mid and lowland respectively. The area receives an average annual rainfall of 400-2500mm and min and max temp 3.50c and 35.0c and altitude

97 ranges from 300 to 4377masl. Based on the figure from [19] report Bale zone has an
98 estimated total population of 1,741,197 out of which 881,559 are male and 859,638 are
99 female.

100 Goro and Ginir are the Woredas in the Bale zone of Oromia Region of Ethiopia. Goro is
101 bordered on the Southwest by Guradamole, on the west by Berbere, on the Northwest by
102 Sinanana Dinsho, on the Northeast by Ginir, and on the Southeast by the Somali Region; it
103 is separated from Guradamole and Berbere by the Gestro River (or Weyib River). Ginir is
104 bordered on the south by the Gestro River (or Weyib River) which separates it from Goro,
105 on the west by Sinanana Dinsho, on the Northwest by Gaserana Gololcha, on the
106 Northeast by Seweyna, and on the East by Raytu.



107 **Figure1. Map of the study area.**

108 **2.2. Sampling Techniques and Sample Size**

109
110
111 In this study, a multistage sampling technique was used. In the first stage from Oromia
112 regional state Bale Zone was selected purposely based on the production potential of the
113 Black cumin crop. In the second stage, from Bale Zone two Woredas (Ginir and Goro

woredas) were selected based on production potential of Black cumin. In the third stage, eight (8) rural Kebeles were selected randomly from the existing Kebeles of the Ginir and Goro woredas. Fourthly, farm households was selected using the probability proportional to size using simple random sampling technique from selected kebles of the two Woredas. Lastly, a total of 180 households was randomly selected for the analysis this study.

2.3. Types and Method of Data Collection

Both primary and secondary data were used for this study. The primary data was collected from sample respondents through face to face interview by structured questionnaire, focus group discussion and field observation. The questionnaires include the socio-economic characteristics, institutional factors, biophysical factors etc. related with the production, marketing and utilization Black cumin. Secondary, data was collected from agricultural office, kebele administration office, books, journals and documents was also be consulted for supplementing the whole work.

2.4 Data Analysis

In order to analyses the data, both descriptive analysis and econometric analysis was employed. Econometric model was applied to examine factors affecting the farmer's perception of the importance of block cumin production on the study area. In order to provide a detailed analysis of the perception on the importance of producing Black cumin, perceived it as important or not, we applied a discrete choice Probit model for binary choice (yes, no) responses to the importance Black cumin production perception question. Probit model is a statistical probability model with two categories in the dependent variable [20]. Probit analysis is based on the cumulative normal probability distribution. The binary dependent variable y , takes on the values of zero and one [21]. The Probit analysis provides statistically significant findings of which demographics increase or decrease the probability of consumption.

140 In the binary Probit model, perceived it as important to produce was taken as 1, while not
 141 perceived it as important as 0. It is assumed that the i^{th} household obtains maximum utility; it
 142 has perceived it as important to produce rather than not to produce the crop. The probability
 143 p_i of choosing any alternative over not choosing, where Φ represents the cumulative
 144 distribution of a standard normal random variable [22]:

$$145 \quad P_i = \text{prob}[Y_i = 1|X] = \int_{-\infty}^{X_i'\beta} (2\pi)^{-1/2} \exp\left(-\frac{t^2}{2}\right) dt \quad (1)$$

$$146 \quad = \Phi(X_i'\beta) \quad (2)$$

147 The relationship between a specific variable (x_i) and the outcome of the probability is
 148 interpreted by means of the marginal effect, which accounts for the partial change in the
 149 probability. The marginal effect associated with continuous explanatory variables X_k on the
 150 probability $P(Y_i = 1 | X)$, holding the other variables constant, can be derived as follows [22]:

$$151 \quad \frac{\partial P_i}{\partial X_{ik}} = \Phi(X_i'\beta)\beta_k \quad (3)$$

152 The marginal effect on dummy variables should be estimated differently from continuous
 153 variables. Discrete changes in the predicted probabilities constitute an alternative to the
 154 marginal effect when evaluating the influence of a dummy variable. Such an effect can be
 155 derived from the following [22]:

$$156 \quad \Delta = \Phi(\bar{X}\beta, d = 1) - \Phi(\bar{X}\beta, d = 0) \quad (4)$$

157 The marginal effects provide insights into how the explanatory variables shift the probability
 158 of frequency of Black cumin production. Using the econometric software Stata 13, marginal
 159 effects were calculated for each variable while holding other variables constant at their
 160 sample mean values.

161 **3. RESULTS AND DISCUSSION**

162 **3.1. Socio-Economic Characteristics of study farmers**

163
 164 In Table 1, the survey result indicated that the mean age of the respondents was 39.68 year
 165 with its minimum 20 and followed by maximum of 78 year. This shows majority of the
 166 respondents were found in economical active age category and which inturns helps to

167 accelerate the production Black cumin in the study area. Household heads an average
 168 education level in year of schooling was 4.92. The minimum level education of respondents
 169 was zero grades and the maximum was 10+3 (diploma) and this implied that the study
 170 households was comprises both educated and non-educated categories. The survey also
 171 indicted that the maximum family size was 20 and the minimum was 3, which was with the
 172 average family size 7.53 in number. The results implies that for the production of Black
 173 cumin in study area the labour availability is not problem, since in countries like Ethiopia
 174 agricultural activities needs more labour. The average land holding of the sample
 175 respondents was 2.37 hector. This is greater than the average land holding (1.37 hector)
 176 level of the Ethiopian farmers and this turns has its good implication on increasing of
 177 production of Black cumin crop. Furthermore, the mean livestock in TLU of respondents in
 178 the study area was 4.96. This was varies from zero (minimum) to 14.83 TLU.

179

180 **Table 1. Socio-economic and demographic characteristics of the Sample households**

Variables	OBS	Mean	Std. Dev.	Min	Max
Age	180	39.68	11.74	20	78
Education	180	4.92	3.06	0	13
Family size	180	7.53	3.06	3	20
Land	180	2.37	1.13	0.66	8
TLU	180	4.96	2.87	0	14.83

181 *Sources: Own survey, 2018*

182 **3.2. Land characteristics of Black cumin producers**

183

184 Land related characteristics as important features for the farmers as whole. These
 185 characteristics of land include soil fertility status and productivity of land; that are important
 186 factors influencing agricultural production in general and Black cumin production
 187 particularity. Majority (82.22%) of the respondents responded that their land soil fertility
 188 status was medium, followed by very fertile status. Only few respondents (6.11%) said that

189 their land categorized as low fertile. Therefore, this result implies that in study area there is
 190 good opportunity to increase the Black cumin crop production activities. Also, most (75 %) of the respondents revealed that the productivity of land was declining through time and
 191 about 11.11% said that their land productivity was increasing through times. Moreover,
 192 8.33% of respondents were responded that their land productivity might increases or
 193 decrease depends of farm work activity, rain fall availability and input usage. Only few (5.56
 194 %) respondents said that there was no change in productivity of land over the time (Table 2).

196 **Table 2. Land characteristics of the sample producers of the Black cumin**

Variable		Frequency	Percentage
Soil fertility status	Low fertile	11	6.11
	Medium	148	82.22
	Very fertile	21	11.67
	Total	180	100
Productivity of land	Declining	135	75.00
	No change	10	5.56
	Increasing	20	11.11
	Increase or decrease	15	8.33
	Total	180	100

197 *Source: Own survey, 2018*

198

199 **3.3. Access to Extension, Credit and Market services for producers of Black** 200 **cumin**

201 Access to development agents for training/advice is important institutional factor which have
 202 an implication on agricultural activities. High proportions (55 %) of farmers were not visited
 203 by development agents' for advices and trainings Black cumin production activities (Table,
 204 3).This implies that, for the study farmers during the study period, little emphases was
 205 placed for extension service provided by development agents on Black cumin production

and utilization opportunity. With regard to credit access, majority (93.33%) of the sample respondents reported that they were not received credit. Only very few (6.67%) respondents were received credit for livestock animal and farm input (fertilizer and improved seed) purchasing. Market related factors like access and distance are crucial for farmers for sale and buy agricultural produce and farm inputs. The result revealed that most of (87.78 %) respondents have had an access to market. The majority of sampled respondents revealed that average distance of market from their home was 10 to 30 minute and which followed (33.89) by 31 to 60 minute. Most of the respondents also perceived that the distance of market from their residence was near and about 27.78% perceived the distance as far (Table 3).

Table 3. Access to different institutional services for the sample farmers

Variable		Frequency	Percentage
Contact by Extension agents	Yes	81	45.00
	No	99	55.00
	Total	180	100
Access to Credit	Yes	12	6.67
	No	168	93.33
	Total	180	100
Market access	Yes	158	87.78
	No	22	12.22
	Total	180	100
Average distance of market from residence	<10 min	16	8.89
	10-30 min	91	50.56
	31-60 min	57	31.67
	>60 min	16	8.89
	Total	180	100
Perceive the distance of market from your residence	Near	128	71.11
	Far	50	27.78
	Very far	2	1.11
	Total	180	100

Source: Own survey, 2018

219

220 **3.4. Black cumin producer's farm input usage and its source**

221

222 Farm input liker fertilizer and improved seed helps farmers to increase the production and
 223 productivity of the farm. The result in (Table 4) revealed that majority (73.33% and 68.33%)
 224 of the respondents were using both fertilizer and improved seed for their farm production
 225 activities. As respondents said that they get fertilizer from farmers union in the form direct
 226 purchase, and the source for improved seed were the agricultural office and farmers in the
 227 form direct purchase. However, some proportions (26.67 % and 31.67%) of the framers were
 228 not using both fertilizer and improved for their farm activities. For this case some of the
 229 farmers were responded like their land doesn't need fertilizer. Hence, this implied that there
 230 should be much emphasis on awareness creation for fertilizer and improved seed usage to
 231 increase production and productivity of the farm.

232 **Table 4. Respondents' farm input use and the source**

233

Variable		Frequency	Percentage
Fertilizer use	Yes	132	73.33
	No	48	26.67
	Total	180	100.00
Improved seed	Yes	123	68.33
	No	57	31.67
	Total	180	100.00

234 Source: Own survey, 208

235

236 **3.5. Sample Households income from Black cumin and others sources**

237

238 In the study area sample respondents engaged different activities/sources to generate their
 239 income. These activities includes farm crop production, livestock and livestock products,
 240 engagement in non- farm and off-farm activities. The major crop includes wheat, *teff*, Black
 241 cumin, barley, garlic; and livestock includes cow, oxen, sheep and donkey. The major non-
 242 farm income sources are remittance, petty trade, and hand craft. The off-farm activity
 243 includes daily labor work, renting assets and firewood sale. Accordingly, Table 5 presents
 244 the mean annual income of 2018 from these activities in ETB. The result indicated that the

mean annual income from crop was 59,348.19 ETB, livestock and its product was 1,779.22 ETB; and hence, the total agriculture mean was 59,802.97 ETB. Moreover, the mean annual income of the same year from non-farm and off-farm activities earned were 1,442.22 ETB and 747.78 ETB, respectively (Table 5).

The Table also revealed that the mean income from Black cumin production and sale was **23,666.67 (39.88%)** of the mean total income of all crop production activities. It implies that this crop alone contributes high proportion of all crop income and its producers fetch higher income. Therefore, it is good opportunity for marginal farmers to cultivate and earn more income from this crop. Thus, this confirms in the both of the study Woredas there is potential of Black cumin production, and hence this in turns pave the way for the inclusion of this crop in agricultural policy to exploit more benefit, sustainable use and production enhancement strategies. Besides, the result indicates that the higher mean annual income earned from both crop and followed by livestock and its products sales. The income from non-farm activities was found next to livestock and its products. This implies that, in the study area the major livelihood and/or income source for households was crop production activities.

Table 5. Sources and mean annual income for sample households at 2018 in ETB

Variables	Obs	Mean	Std. Dev.	Min	Max
Total Crop income	180	59,348.19	40,063.71	6,650.00	202,800.00
<i>Income from Black cumin</i>	<i>180</i>	<i>23,666.67</i> <i>(39.88 %)*</i>	<i>25,316.20</i>	<i>0</i>	<i>198,000.00</i>
Livestock and its products	180	1,779.22	3,911.06	0	16,000.00
Total agriculture income	180	59,802.97	39,961.41	6,500.00	213,000.00
Off-farm income	180	747.78	4,714.16	0	60,000.00
Non-farm income	180	1,442.22	6,602.52	0	60,000.00

265 *Source: Own survey, 2018*

266 ** Income share of black cumin from the total income of crops produced by the respondents*

267

268 **3.6. Households Perception of Black cumin importance and Usage**

269

270 Sustainable production and conservation of resources is possible if the users have
271 knowledge on the importance of the resource and give the value for the resources. In line
272 with this, study households were asked to express their perception for Black cumin
273 importance, utilization purpose of the crop, productivity and market demand of Black cumin
274 through time. The result indicates that majority (95.56%) of the households perceived that
275 production of Black cumin is important in the study area. The major reason that households
276 said producing the crop is important were since the crop is source of better income, it is
277 better in market price than other crops in kg, and most their livelihood is based on Black
278 cumin. In the focus group discussion participants expressed as the seed is used as
279 medicinal crop for common cold, headache, diarrhea, asthmatic problem and spice.
280 Moreover, they added that relatively Black cumin crop adapted with the local environment,
281 give better yield with low rain fall and better market price than other crops in terms of kg. The
282 result also shows that about 83.89% (high proportion) of the sampled households were
283 allocated their farm land for production of Black cumin during the study cropping year (2018).

284

285 Table 6, result also revealed that the majority (78.89%) of the study respondents said that
286 the production of lack cumin through time in the study area was decreasing. The major
287 reasons for production decline were rain fall scarcity, disease, pests and absence of
288 improved variety. But, some (14.44%) respondents responded that the production of Black
289 may increases or decreases depending on availability of rain fall and level of farm work. This
290 means that, if there is good rainfall in production season and better farm work the production
291 will be higher and vice-versa. Additionally, the result shows that the majority (90%) of market
292 demand for Black cumin though time in study area was declining. The respondents justified
293 that the major problem for this case were unethical connection of local traders with brokers

294 and central market, poor infrastructures and some also said that declining of production
295 amount (this lead to declining the traders coming to collect and load at farm gate/site).

296 **Table 6. Respondents' perception about Black cumin importance and rated concepts**

Variable	Frequency	Percentage
Perceive that Black cumin production is important	Yes	172
	No	8
	Total	180
Farmland allocate to Black cumin production	Yes	151
	No	29
	Total	180
Black cumin productivity through time in the study area	Increasing	12
	Decreasing	144
	Increase or decrease	26
	Total	180
Market demand for Black cumin increasing though time in study area	Yes	18
	No	162
	Total	180

297 Source: Own survey, 2018

298 **3.7. Economic Model result on Factor affecting the producer's perception**

299

300 Table 7 presents the econometric model result of factor affecting the perception of producers
301 on the importance of producing Black cumin in the study area. The maximum likelihood
302 estimates of the Probit model shows that, out of 13 explanatory variables six variables were
303 found to be statistically and significant affected the perception of producers of the
304 importance of producing Black cumin. The chi-square results revealed that likelihood ratio
305 statistics are highly significant ($P<0.001$) suggesting the model has a strong explanatory
306 power. Accordingly, the interpretation and discussion of the variables were provided as
307 follows:

308

309 **Age of Respondents:** The marginal effect (dy/dx) revealed that the age and perception of
310 producers of the importance of producing Black cumin has positive and statistically
311 significant relationship at ($P < 0.001$). As age of farmers increases by one year the
312 probability of perception of the importance of producing Black cumin will rise by 1.68 %.
313 This is plausible, since adult producers might have more know-how and experience of both
314 producing and importance of farm crops than the young producers. Study in Ethiopia has
315 indeed shown a positive relationship between number of years of experience in agriculture
316 and farmers' perception for expansion of crops [23].

317

318 **Education:** In line with the expectation, the farmer's level of education and perception of the
319 importance of producing Black cumin crop has positive and significant correlation statistically
320 at ($P=0.05$). The increases in the education attainment of producers by one grade will
321 increase the probability of perception of the importance of producing Black cumin crop by
322 2.28 %. The logical analysis behind this is that, the more producers have an educational
323 opportunity the more they have knowledge of agricultural production system than the
324 counterpart. Previous Studies of [24, 25, 26] have also found that, farmers with better
325 education have more exposure to new ideas and information, and thus have better
326 knowledge to effectively analyze and use available information, and indeed it helps to
327 decide or perceive the important crop variety production.

328

329 **Labour availability:** The result revealed as the availability of labour for farm activity
330 increases by one unit (number), the probability of perception of the importance of producing
331 Black cumin crop will rise by 11.50%. In this study, it contributes positively and significantly
332 in this study for perception of the importance of producing Black cumin crop at ($P=0.1$). The
333 rational justification for this case might be households with high number family member
334 within working age group might not face the problem of labour availability to be engaged in

335 agricultural activities than small size family member, given that that agricultural crop
336 production is more labour demanding. Similarly, [27] has found that *Nigella sativa* requires
337 extensive labor in collection and harvest as the capsules (fruit) tend to shatter at maturity
338 because its post-harvest management of the fruits usually involves their harvest, one by one,
339 by hand and dry storage till natural dehiscence.

340

341 **Aces to credit:** The survey findings indicated that access to credit for the producers and
342 perception of importance of producing Black cumin in the study area has positive and
343 significant relationship at ($P=.01$) statistical level. Marginal analysis revealed that having
344 access to credit for the farmers increases the probability of the perceiving the importance of
345 producing Black cumin by 19.43%. This is true that, for the farmers having credit facilities
346 help to buy farm inputs and run their farm activity smoothly. Previous findings also confirmed
347 that credit access facilitates purchase of inputs especially improved seed varieties and
348 inorganic fertilizers if linked to well-developed input supply and market access infrastructures
349 [28, 29, 30].

350

351 **Income from Black cumin:** As it was hypothesized the income received from Black cumin
352 production and perception of its continual production importance has positive and statistically
353 significant relation at ($P<0.001$) level. As the income gained from producing Black cumin
354 rises by one Birr (ETB) the probability of perceiving the value of crop production will rises by
355 0.06%. The possible reason here is that, the rational producers give more value for crops
356 which has more return from its farm activity. Study conducted by the same authors [18]
357 confirmed that the average income from the total farm activity and participation for Black
358 cumin conservation has positive and significant correlation.

359

360 **Productivity trend of the crop:** Similar to the expected hypothesis, productivity has
361 positive and statistically significant association with producers perception of the importance

of producing Black cumin at ($P=.01$) level. Marginal effect shows that, as amount of output from a give level of input used [productivity] increases though time the probability of perceiving the importance of producing Black cumin by farmers will increases 2.89%. Possible implication is that, If the farmers get more return from the farm activity it encourages them to invest more on that activity and hence to build positive perception for production the crop.

368

369 **Table 7: Probit model result of factor affecting the perception producers on the**
370 **importance of producing Black cumin**

Variables	Coefficients	SD	Z-value	Marginal effect (dy/dx)
Age	.0966	.0202	4.78***	.0168
Education	.1308	.0559	2.34**	.0228
Total family size	.0116	.04957	0.23	.0020
Land holding	.0492	.16130	0.31	.0085
Labour availability	.6600	.35984	1.83*	.1150
Extension services	-.2242	.14924	-1.50	-.0391
Aces to credit	1.1145	.6352	1.75*	.1943
Market distance	.2747	.1987	1.38	.0479
TLU	-.0637	.0598	-1.06	-.0111
Black cumin income	.0054	.00001	3.33***	.00062
Productivity trend	.1663	.07107	2.34**	.0289
Market demand	-.0979	.16292	-0.60	.0170
Land productivity	-.06769	.055019	-1.23	-.0118
Constant	-7.6762	1.9282	-3.98***	
Number of obs = 180	Log likelihood = -58.5191			
LR chi2(13) = 93.73	Pseudo R2 = 0.4447			
Prob > chi2 =0.0000				

371 Source: Own computation, 2018
372 ****, ** and * means statistically significant at 1%, 5% and 10% level*
373

374 **4. CONCLUSION**

375
376 The survey result indicated that the mean age of the respondents was 39.68 year, education
377
378 level in year of schooling was 4.92 and the family size was 7.53 in number. The average
379 land holding of the sample respondents was 2.37 hector and the livestock in TLU was 4.96.
380 Furthermore, from the total mean of agriculture income (59,802.97 ETB) Black cumin
381 production contributes about 39.88% for the study respondents. The majority (95.56%) of the
382 households perceived that production of Black cumin crop is important in the study area. The
383 major importance and utilization of producing Black cumin crop for study households were
384 as source of better income, better market price that other crops in kg, and their major
385 livelihood is base; and used as medicinal crop for common cold, headache, diarrhea,
386 asthmatic problem; and spice.

387
388 The finding of Probit model shows that producers perception of the importance of black
389 cumin production were found to be statistically and significant affected by age of households,
390 education level, availability of labor for farm activities, access to credit facilities, average
391 income from Black cumin, and Black cumin productivity trend through time. Hence,
392 agricultural policy should give emphases at all operational level to exploit more benefit from
393 this crop and production enhancement strategies, so as to bring foreseen change in the lives
394 of producers.

395 **ACKNOWLEDGEMENTS**

396
397 Authors are grateful to Ethiopian Biodiversity Institute for providing financial assistance and
398
399 Vehicle arrangement during the data collection of this research work.

400

401 **COMPETING INTERESTS**

402

403 "Authors have declared that no competing interests exist."

404

405 **REFERENCES**

406

- 407 1. Atta MB. Some characteristics of *Nigella* (*Nigella sativa* L.) seed cultivated in Egypt
408 and its lipid profile. Food Chem. 2003; 83:63–68.
- 409 2. Burits M, Bucar F. Antioxidant activity of *Nigella sativa* essential oil. Phytother. Res.
410 2000; 14:323–328.
- 411 3. Yazachew A. Comparative study of supercritical fluid and solvent extraction. Addis
412 Ababa. 2011; 67:61-80.
- 413 4. Takrun HRH, Dameh MAF. Study of the nutritional value of Black cumin seeds
414 (*Nigella sativa* L.). J. Sci. Agric., 1998;76: 404-410
- 415 5. Ostlund, R.E. Phytosterols in human nutrition. Ann. Rev. Nutr. 2002; 22:533–549.
- 416 6. Aminpour, Karimi. Underutilized medicinal spices. Spice India. 2004; 17 (12): 5-7.
- 417 7. Black M, Bewley D, Halmer. The Encyclopedia of seed science, technology and
418 uses. wallinoford. CAB P 7. 2005.
- 419 8. Ashraf M, Orooj A. Salt stress effects on growth, ion accumulation and seed oil
420 concentration in an arid zone traditional medicinal plant ajowan (*Trachypernum*
421 *ammi* [L.]. Journal of Arid Environments. 2006; 64: 209-220.
- 422 9. Thilakarathna RCN , et al . Phytochemical Analysis of Indian and Ethiopian Black
423 Cumin Seeds (*Nigella Sativa*) Agricultural Research & Technology. 2018; 17(1).
- 424 10. Hedberge I, Edwards S, Sileshi Nemomissa (eds). Flora of Ethiopia and Eriteria.
425 Apiaceae to Dipsaceae. The Natural Herbarium. Addis Ababa University, Addis
426 Ababa. 2003; 4 (2).
- 427 11. Jansen PCM. Spices, condiments and medicinal plants in Ethiopia. Their taxonomy
428 and agricultural significance. Addis Ababa: Center for Agricultural Publishing and
429 Documentation. 1981). pp. 111-120.

- 430 12. Mogessie A, Tetemke M. Some microbiological and nutritional properties of Borde
431 and Shamita. Traditional Ethiopian fermented beverages. Ethiop. J. Health Dev.
432 1995; 9(1): 105-110.
- 433 13. Ethiopian Investment Agency. Investment opportunity Profile for Spice Processing in
434 Ethiopia. 2015; 7: 8-14.
- 435 14. Habtewold K, Demes F, Tewodros L, Dejene B, Haimanot M, Wakjira G. Seed
436 Spices Production Guideline: Ethiopian institute of agricultural Research. 2017.
437 Available: [http:// www. publication.eiar.gov.et](http://www.publication.eiar.gov.et):
- 438 15. Ermias A. Adaptability study of Black cumin (*Nigella sativa* L.) varieties in the mid
439 and high land areas of Kaffa zone, south West Ethiopia. Agriculture, Forestry and
440 Fisheries. 2015; 6:14-17.
- 441 16. Orgut. Market Assessment Study, Ethiopian Nile Irrigation and Drainage Project,
442 Main Report and Annexes, Ministry Of Water Resources, Addis Ababa, June 2007.
- 443 17. Yousif HH. Effect of high levels of nitrogen and phosphorus fertilizer on growth,
444 yield and yield components of *Nigella sativa* L. Horticulture Department College of
445 Agriculture, Duhok University, Iraq, Mesopotamia Journal Agriculture. 2008; 36(1).
- 446 18. Dessalegn A, Wubeshet T. Economic Value of Black Cumin (*Nigella sativa* L.)
447 Conservation at Bale Zone of Oromia Region, Ethiopia; American Journal of
448 Business, Economics and Management. 2018; 6(4): 104-109.
- 449 19. Bale zone administrative and development office, Bale, Ethiopia, (2012).
- 450 20. Liao TF. Interpreting Probability Models: Logit, Probit, and Other Generalized Linear
451 Models, 101 Quantitative Applications in the Social Sciences, Sage Publications,
452 Thousand Oaks, Calif, USA, 1994.
- 453 21. Aldrich JH, Nelson FD, *Linear Probability, Logit, and Probit Models*, Sage
454 Publications, Newbury Park, Calif, USA, 1984.
- 455 22. . Greene WH, *Econometric Analysis*, Prentice Hall, 7th Edition, 2011.

- 456 23. Adimassu, Kessler. Factors affecting farmers' coping and adaptation strategies
457 to perceived trends of declining rainfall and crop productivity in the central Rift valley
458 of Ethiopia: Journal of Environ system Research. 2016; 5:13.
- 459 24. Kassie M, Teklewold H, Jaleta M, Marenja, P, Erenstein O. Understanding the
460 adoption of a portfolio of sustainable intensification practices in eastern and
461 southern Africa. Land Use Policy. 2015; 42: 400-411.
- 462 25. Knowler D, Bradshaw B. Farmers' adoption of conservation agriculture: A review
463 and synthesis of recent research. Food policy. 2007; 32(1): 25-48.
- 464 26. Prokopy L, Floress K, Klotthor-Weinkauff D, Baumgart-Getz A. Determinants of
465 agricultural best management practice adoption: Evidence from the literature.
466 Journal of Soil and Water Conservation. 2008; 63(5), 300-311.
- 467
- 468 27. Animesh KD, Aditi S, Arnab B, Aninda M, Rita P, Sonali S. Black Cumin (*Nigella*
469 *Sativa* L.)- A Review; Journal of Plant Development Sciences. 2012; 4 (1): 1-43.
- 470 28. Geta E, Bogale A, Kassa B, Elias E. Determinants of Farmers' Decision on Soil
471 Fertility Management Options for Maize Production in Southern Ethiopia. American
472 Journal of Experimental Agriculture. 2013; 3(1): 226-239.
- 473 29. Jeannin M. Agricultural innovation in Africa: from soil fertility to market integration. A
474 case study from Benin. 2012.
- 475 30. Teklewold H, Kassie M, Shiferaw B. Adoption of multiple sustainable agricultural
476 practices in rural Ethiopia. Journal of agricultural economics, 2013; 64(3), 597-623.

477
478
479
480
481