

**EFFECT OF MICRONUTRIENTS IN PRESENCE OF  
DIFFERENT LEVELS OF ORGANIC MANURE ON  
GROWTH AND YIELD OF TOMATO (*Solanum  
lycopersicum* L.)**

**ABSTRACT**

A field experiment was conducted at the Horticultural Farm of Sher-e -Bangla Agricultural University, Dhaka from October, 2017 to March, 2018 to study effect of micronutrients in presence of different level of organic manure on growth and yield of tomato. There were four combinations of micronutrients viz.  $N_0=0$  kg Zn 0 kg B/ha,  $N_1=Zn_2$  kg  $B_{1.5}$  kg/ha,  $N_2=Zn_4$  kg  $B_2$  kg/ha,  $N_3=Zn_6$ kg  $B_{2.5}$  kg/ha and four organic manure viz  $M_0=0$  ton/ha,  $M_1=Cowdung$  (15 ton/ha),  $M_2=Poultry$  manure (10 ton/ha),  $M_3=(Cowdung$  7.5 ton/ha+ Poultry manure 5 ton/ha). The experiment was laid out in a Randomized Complete Block Design with 3 replications and there were altogether 48 plots. Application of micronutrients and organic manure significantly influenced the growth, yield and size of tomato. The highest yield (66.96 t/ha) was found from treatments  $N_2$  and the lowest yield (25.69 t/ha) was obtained from treatment  $N_0$ . Due to the application of organic manure, the highest yield (50.78 t/ha) was obtained from  $M_3$  and lowest yield (39.86 t/ha) was recorded from  $M_0$ . In case of combined effect, the highest yield (76.33 t/ha) was found from treatment  $N_2M_3$  and the lowest yield (24.60 t/ha) was found from treatment  $N_0M_0$ . So, application of  $Zn_4$  kg  $B_2$  kg/ha along with Cowdung 7.5 ton+Poultry manure 5 ton/ha. was the best for growth and yield of tomato. Economic analysis revealed that  $N_2M_3$  gave the maximum benefit cost ratio (3.2). So, application of  $Zn_4$  kg  $B_2$  kg/ha along with Cowdung 7.5 ton+Poultry manure 5 ton/ha was the best for growth and yield of tomato.

*Keywords: Benefit Cost Ratio (BCR), Boron, Organic manure, Yield*

**1. INTRODUCTION**

Tomato (*Solanum lycopersicum* L) is one of the most important vegetable crops grown throughout the world including Bangladesh and belongs to the Solanaceae family. Tomato is cultivated in almost all over the country for its adaptability to wide range of soil and climate in Bangladesh [1]. Tomato ranks next to potato and sweet potato in respect of vegetable production in the world [2]. It ranks fourth in respect of production and third in respect of area in Bangladesh [3]. The largest tomato producing countries of the world are China, The United States of America, India, Egypt, Turkey, Iran, Italy, Mexico, Brazil and

Indonesia [4]. The popularity and different products of tomato are increasing day by day. Tomato is very valuable crop because of higher contents of vitamins A, B and C with Ca and carotene [5]. It is also popular for its medicinal value. Tomato is a major component in the daily diet, and can be used in making soups, pickles, ketchup, sauces, juices etc. The well rippled tomato (edible portion/100g) contains water (94.1%); energy (23 calories); Ca (1.0 gm); Mg (7.0 mg); vitamin A (1000 IU); ascorbic acid (22 mg); thiamin (0.09 mg); riboflavin (0.03 mg); niacin (0.8 mg) [6]. In Bangladesh, the yield of tomato is not enough satisfactory in compare to other tomato growing countries of the World [7]. The cultivated area under tomato in Bangladesh was 75602 acre and total production was 413610 metric tons during the year 2014-2015 [3]. The low yield of tomato in Bangladesh is due to the use of poor yielding varieties and improper cultural practices and now it is considered as one of the major problem to successful upland crop production in Bangladesh [8]. For better yield the cultivation of tomato requires proper supply of plant nutrient. Sufficient supply of nutrient can improve the yield, fruit quality, fruit size, keeping quality, colour, and taste of tomato [9]. Since, the land is limited in Bangladesh, it is important to increase yield of any crop. Though the effects of different on the yield of tomato were studied earlier, the effect of micronutrients and organic manure on growth and yield of tomato were not studied in detail so far in Bangladesh. Among the micronutrients, zinc and boron play an important role in improving the yield and quality of tomato in addition to checking various diseases and physiological disorders [10]. Zinc (Zn) is another important essential micronutrient which helps in the formation of tryptophan, a precursor of IAA responsible for growth stimulation [11] and plays a vital role in synthesis of carbonic anhydrase enzyme which helps in transport of CO<sub>2</sub> in photosynthesis [12]. Zinc deficiency causes shorter and thinner internodes, stunted growth, appearance of chlorotic flecks on the older leaves and twisting of leaf borders in upward direction and plant with abnormal features [13]. The zinc deficiency may be due to soil deficient in Zn, competition with Ca, Mn, Fe, P, to some degree K, and soil properties that influence Zn availability [14]. Boron has a pronounced effect on the production and quality of tomato. Boron is needed by the crop plants for cell division, nucleic acid synthesis, uptake of calcium and transport of carbohydrates [15]. Boron also plays an important role in flowering and fruit formation [16]. Adequate levels of Boron help to maintain leaf K levels in tomato during fruit development [17]. Boron has major influence on the plasma membrane of plant cells and ion transport and those B amendments increased Calcium, and Mg levels [18]. Boron deficiency affects the growing points of roots and youngest leaves. The leaves become wrinkled and curled with light green color. Its deficiency affects translocation of sugar, starches, nitrogen and phosphorus, synthesis of amino acids and proteins [19]. In boron deficient plants the youngest leaves become pale green, losing more color at the base than at the tip. Boron deficiency symptoms will often appear in the form of thickened wilted, or curled leaves, a thickened, cracked, or water soaked condition of petioles and stems, and discoloration, cracking or rotting of fruit, tubers or roots [20]. The improvement in quality parameters of tomato fruit due to boron application could be the result of overall growth and development of the crop [21]. To improve texture, structure, humus, color, aeration, water holding capacity and microbial activity of soil used by proper amount of organic manure such as cowdung, poultry manure. In our country, the soils of most regions have less than 1.5%, some soils even have less than one percent organic matter [22]. Organic manure has the largest effect on yield and quality of tomato. It also improves the vegetative growth, flowering and fruit set of tomato. The increase in vegetative growth of tomato could be attributed to physiological role of organic manure and its involvement in the metabolism of protein, synthesis of pectin, maintaining the water relation within the plant, resynthesis of ATP and translocation of sugar at development of the flowering and fruiting stages [15]. For better growth of tomato a large amount of organic manure are required. [23]. Cowdung also contains beneficial bacteria, which convert nutrients into easily accessible forms so they can be slowly released without burning tender plant roots. Poultry manure contains high % of N and P for the healthy growth of plants [24]. The physical properties of the soil improved by the application of poultry manure. poultry manure improves the fertility of the cultivated soil by increasing the organic matter content, water holding capacity, oxygen diffusion rate and the aggregate stability of the soils [25,26]. Large quantities of poultry manure are available especially in urban centers and it is effective source of nutrients for vegetables such as tomato [27].

## 2. MATERIAL AND METHODS

### 2.1 Experimental site

The field experiment was conducted in the Horticulture farm of Sher-e-Bangla Agricultural University, Dhaka-1207 from October, 2017 to March, 2018. The location of the experimental site was at 23.75°N latitude and 90.34°E longitudes with an elevation of 8.45 meter from the sea level.

### 2.2 Experimental Treatment

The two factor experiment consisted of four levels of micronutrients (Factor A)  $N_0=0$  kg Zn 0 kg B/ha,  $N_1=Zn_2$  kg  $B_{1.5}$  kg/ha,  $N_2=Zn_4$  kg  $B_2$  kg/ha,  $N_3=Zn_6$ kg  $B_{2.5}$  kg/ha and four levels of organic manure (Factor B).  $M_0 = 0$  ton/ha,  $M_1 =$  Cowdung (15 ton/ha),  $M_2 =$  Poultry Manure (10 ton/ha),  $M_3 =$  Cowdung 7.5 ton + Poultry Manure 5 ton/ha. The experiment was laid out in Randomized Complete Block Design (RCBD) having two factors with three replications. An area of 30.8 m x 7.6 m was divided into three equal blocks. Each block consists of 16 plots where 16 treatments were allotted randomly. There were 48 unit plots in the experiment. The size of each plot was 1.5 m x 2 m. The distance between two blocks and two plots were kept 0.4 m and 0.4 m respectively

### 2.3 Application of micronutrients and organic manure

The entire amounts of  $N_1$ . $N_2$ . $N_3$  were applied during the final land preparation. The entire amounts of  $M_1$ ,  $M_2$  and  $M_3$  were applied during the final land preparation (Table 1).

### 2.4 Statistical Analysis

The data obtained for different parameters were statistically analysed by MSTAT-C computer package. The significance of the difference among the treatment combinations means was compared by LSD test at 5% level of probability.

**Table 1. Doses of nutrients application in the main field as per treatment were applied during the final land preparation.**

Treatments	Zinc	Boron	Treatments	Cowdung	Poultry manure
$N_0$	0 kg/ha	0 kg/ha	$M_0$	0 ton/ha	0 ton/ha
$N_1$	2 kg/ha	1.5 kg/ha	$M_1$	15 ton/ha	0 ton/ha
$N_2$	4 kg/ha	2 kg/ha	$M_2$	0 ton/ha	10 ton/ha
$N_3$	6 kg/ha	2.5 kg/ha	$M_3$	7.5 ton/ha	5 ton/ha

### 2.5 Economic Analysis

The cost of production was analyzed in order to find out the most economic treatment of micronutrients and organic manure. All input cost included the cost for lease of land and interest on running capital in computing the cost of production. The interest was calculated @ 13% in simple interest rate. The market price of tomato was considered for estimating the cost and return. Analyses were done according to the procedure determining by [28]. The benefit cost ratio (BCR) was calculated as follows:

$$BCR = \text{Gross return per hectare (Tk.)} \div \text{Cost of production per hectare (Tk.)}$$

## 3. RESULTS AND DISCUSSION

### 3.1 Plant height

Plant height was measured starting from 25 days after transplanting. It was measured 20 days interval and continued up to 65 DAT. At 25 DAT, the maximum plant height (51.82 cm) was observed from  $N_2$  ( $Zn_4 B_2$  kg/ha) treatment and minimum plant height (44.13 cm) was observed from  $N_0$  (0 kg/ha) treatment.

At 45 DAT, the maximum plant height (83.42 cm) was observed from N<sub>2</sub> treatment and minimum plant height (73.42 cm) was observed from N<sub>0</sub> (0 kg/ha) treatment. At 65 DAT, the maximum plant height (95.60 cm) was observed from N<sub>2</sub> treatment and minimum plant height from N<sub>0</sub> (82.86cm) treatment in Fig 1. [29] founded the soil application of zinc sulphate and borax @ 10 and 20 kg/ha, respectively in combination with their foliar spray @ 0.5% and 0.3%, respectively where most effective in improving plant height. Marked variation was observed among treatments as to the plant height of tomato due to the application of different levels of organic manure at 25, 45 and 65 DAT (Fig 2). At 25 DAT, the maximum plant height (52.31cm) was observed from M<sub>3</sub> (Cowdung 7.5 ton+Poultry manure 5 ton/ha) treatment and control treatment (M<sub>0</sub>) gave the minimum plant height (44.20 cm) treatment. At 45 DAT, the maximum plant height (82.82cm) was observed from M<sub>3</sub> treatment M<sub>2</sub> (80.77cm) and minimum plant height M<sub>0</sub> (75.31cm) treatments. At 65 DAT, the maximum plant height (93.22cm) was observed from M<sub>3</sub> and minimum plant height (85.77cm) was observed from M<sub>0</sub> treatment. The reason for higher plant height might be explained in the way that the favorable soil condition influence of balanced uptake of nutrients, which were applied, [24] observed the similar result. The combined Effect of micronutrients and organic manure showed statistically significant variation on plant height at 25, 45 and 65 DAT (Table3). At 25 DAT, the maximum plant height (57.50cm) was obtained from N<sub>2</sub>M<sub>3</sub> (Zn<sub>4</sub>B<sub>2</sub> kg/ha with Cowdung 7.5 ton+Poultry manure 5 ton/ha) and control treatment (N<sub>0</sub>M<sub>0</sub>) treatment gave the minimum plant height (40.50 cm) which. At 45 DAT, the maximum plant height (89.20cm) was obtained from N<sub>2</sub>M<sub>3</sub>.The control treatment (N<sub>0</sub>M<sub>0</sub>) gave the maximum plant height (70.00 cm).At 65 DAT, the maximum plant height (100.00cm) was obtained from N<sub>2</sub>M<sub>3</sub> and control treatment (N<sub>0</sub>M<sub>0</sub>) gave the minimum plant height (81.25cm) which similar to N<sub>0</sub>M<sub>1</sub>(82.00cm) treatment.

### 3.2 Number of leaves per plant

No of leaves per plant of tomato varied significantly due to the application of different levels of micronutrients at 25, 45, and 65 DAT (Fig 3). At 25 DAT, the maximum number of leaves (12.10) was observed in N<sub>2</sub> (Zn<sub>4</sub> B<sub>2</sub> kg/ha) and minimum number of leaves per plant (7.83) was observed in N<sub>0</sub> (0 kg/ha) treatments. At 45 DAT, the maximum number of leaves per plant (25.31) was observed from N<sub>2</sub>. The minimum number of leaves per plant (19.15) was observed from N<sub>0</sub> treatment. At 65 DAT, the maximum number of leaves per plant (33.17) was observed from N<sub>2</sub> and minimum number leaves N<sub>0</sub> (27.58) treatment. [30] reported that application of boron significantly increased the number of leaves on tomato plant compared to control.[31] found similar result. Number of leaves per plant of tomato varied significantly due to the application of different levels of organic manure at 25, 45, and 65 DAT of (Fig 4). At 25 DAT, the maximum number of leaves (11.61) was observed from M<sub>3</sub> (Cowdung 7.5 ton +Poultry manure 5 ton/ha) and minimum number of leaves per plant (8.53) was observed from M<sub>0</sub> (0 ton/ha) treatment. At 45 DAT, the maximum number of leaves per plant (24.21) was observed in M<sub>3</sub>. The minimum number of leaves per plant (20.43) was observed from M<sub>0</sub>. At 65 DAT, the maximum poultry manure 5 ton/ha) and minimum number of leaves per plant (33.51) was observed from M<sub>2</sub> and minimum number leaves M<sub>0</sub> (27.91) .Combined effect micronutrients and organic manure showed statistically significant variation on number of leaves per plant at 25, 45 and 65 DAT (Table3). At 25 DAT, the maximum number of leaves per plant (15.10) was obtained from N<sub>2</sub>M<sub>3</sub> (Zn<sub>4</sub>B<sub>2</sub> kg/ha with Cowdung7.5 ton+Poultry manure 5 ton/ha) and control treatment (N<sub>0</sub>M<sub>0</sub>) gave the minimum number of leaves per plant (7.25). At 45 DAT, the maximum no of leaves per plant (27.95) was obtained from N<sub>2</sub>M<sub>3</sub>. .The control treatment (N<sub>0</sub>M<sub>0</sub>) gave the minimum no of leaves per plant (17.40 cm). At 65 DAT, the maximum number of leaves per plant (36.80 was obtained from N<sub>2</sub>M<sub>3</sub> .The control treatment (N<sub>0</sub>M<sub>0</sub>) gave the minimum number of leaves per plant (25.50). From the result present study it can be conducted that the treatment N<sub>2</sub>M<sub>3</sub> provided better growing condition perhaps to supply of adequate plant nutrients, resulting in the number of leaves per plant.

### 3.3 Number of branches per plant

Number of branches per plant of tomato varied significantly due to the application of different levels of micronutrients at 45 and 65 DAT (Table 2). At 45 DAT, the maximum number of branches per plant (5.06) was observed from N<sub>2</sub> (Zn<sub>4</sub> B<sub>2</sub> kg/ha) and minimum number of branches per plant (3.55) was observed in from N<sub>0</sub> (0 kg/ha) treatment. At 65 DAT, the maximum number of branches per plant (5.72)

was observed from  $N_2$  treatment. The minimum number of branches per plant (4.46) was observed from  $N_0$ , [32] stated similar findings. Number of branches per plant of tomato was found significantly influenced due to the application of different levels of organic manure at 45 and 65 DAT (Table 2). At 45 DAT, the maximum number of branches (4.77) was observed from  $M_3$  (Cowdung 7.5 ton/ha+Poultry manure 5 ton/ha) and minimum number of branches per plant (3.70) was observed in  $M_0$  (0 ton/ha) treatment. At 65 DAT, the maximum number of branches per plant (5.89) was observed in  $M_3$ . The minimum number of branches per plant (4.78) was observed from  $M_0$  treatment. The combined effect of micronutrients and organic manure showed statistically significant variation on the number of branches per plant at 45 and 65 DAT (Table 4). At 45 DAT, the maximum number of leaves per plant (6.05) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung 7.5 ton/ha+Poultry manure 5 ton/ha) and control treatment ( $N_0M_0$ ) gave the minimum number of branches per plant (3.15). At 65 DAT, the maximum number of branches per plant (7.25) was obtained from  $N_2M_3$  which. The control treatment ( $N_0M_0$ ) gave the minimum number of branches per plant (4.10), this might be due to the fact that balanced uptake and influences of nutrients which improve vegetative growth. Organic manure improved physical conditions of the soil, which increase the water holding capacity and better nutrients availability and uptake by the crop.

### 3.4 Number of flowers per plant

Remarkable differences were observed among the micronutrients application of different levels of micronutrients at 45 and 65 DAT (Table 2). At 45 DAT, the maximum number of flower per plant (20.73) was observed from  $N_2$  ( $Zn_4 B_2$  kg/ha) and minimum number of flowers per plant (10.56) was observed from  $N_0$  (0 kg/ha). At 65 DAT, the maximum number of flowers per plant (84.60) was observed from treatment  $N_2$ . The minimum number of flowers per plant (27.53) was observed in treatment  $N_0$ , [31] reported similar result. Number of flowers per plant of tomato varied significantly due to the application of different concentrations of organic manure at 45 and 65 DAT (Table 2). At 45 DAT, the maximum number of flowers per plant (18.68) was observed from  $M_3$  (Cowdung 7.5 ton/ha+Poultry manure 5 ton/ha) and minimum number flowers per plant (12.03) was observed from  $M_0$  (0 ton/ha) treatment. At 65 DAT, the maximum number of flowers per plant (65.15) was observed in treatment  $M_3$ . The minimum number of flowers per plant (39.81) was observed from treatment  $M_0$ . The combined effect of micronutrients and organic manure showed statistically significant variation on number of flower per plant at 45 and 65 DAT (Table 4). At 45 DAT, the maximum number of flowers per plant (24.00) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung 7.5 ton/ha+Poultry manure 5 ton/ha) and control treatment ( $N_0M_0$ ) gave the minimum no of flowers per plant (9.25). At 65 DAT, the maximum number of flowers per plant (98.80) was obtained from treatment  $N_2M_3$ . The control treatment ( $N_0M_0$ ) gave the minimum no of flowers per plant (20.50).





Fig. 1. Graphical representation of treatment (N<sub>0</sub> to N<sub>3</sub>) and plant height in 25, 45 and 65 DAT




Fig. 2. Graphical representation of treatment (M<sub>0</sub>-M<sub>3</sub>) and plant height in 25, 45 and 65 DAT




Fig. 3. Graphical representation of treatment (N<sub>0</sub> to N<sub>3</sub>) and no. of leaves in 25, 45 and 65 DAT

Fig. 4. Graphical representation of treatment (M<sub>0</sub>-M<sub>3</sub>) and no. of leaves in 25, 45 and 65 DAT

**Table 2: The effect of different levels of micronutrients and organic manure on number of branches per plant of tomato of at different days after transplanting**

Treatments	Number of branch per plant		No of flowers per plant	
	45 DAT	65 DAT	45 DAT	65 DAT
<b>Micronutrients</b>				
N <sub>0</sub>	3.55	4.46	10.56	27.53
N <sub>1</sub>	3.92	4.79	13.63	39.51
N <sub>2</sub>	5.06	6.51	20.73	84.60
N <sub>3</sub>	4.53	5.72	17.42	57.82
CV%	10.68	11.58	9.27	9.56
LSD (0.05)	0.25	0.25	2.14	2.21
<b>Organic manure</b>				
M <sub>0</sub>	3.70	4.78	12.03	39.81
M <sub>1</sub>	4.12	5.17	14.59	47.97
M <sub>2</sub>	4.47	5.63	17.04	56.53
M <sub>3</sub>	4.77	5.89	18.68	65.15
CV%	10.68	11.58	9.27	9.56

LSD (0.05)	0.21	0.30	1.08	2.57
------------	------	------	------	------

Means, in a column followed by same letter do not differ significantly at 5% level

$N_0=0$  kg/ha ,  $N_1=Zn_2B_{1.5}$  kg/ha,  $N_2=Zn_4B_2$  kg/ha,  $N_3=Zn_6B_{2.5}$  kg/ha,  $M_0=0$  ton/ha,  $M_1=Cowdung$  (15 ton/ha),  $M_2=Poultry\ manure$  (10 ton/ha),  $M_3=CD+PM$  (7.5+5 ton/ha)

**Table 3: The combined effect of different levels of micronutrients and organic manure on the plant height (cm), number of leaves per plant of tomato at different days after transplanting**

Treatments	Plants height cm			No of leaves/ plant		
	25 DAT	45 DAT	65 DAT	25 DAT	45 DAT	65 DAT
$N_0M_0$	40.50	70.00	81.25	7.25	17.40	25.50
$N_0M_1$	43.30	72.70	82.00	7.60	18.20	26.90
$N_0M_2$	45.25	74.50	82.80	8.05	20.15	27.45
$N_0M_3$	47.50	76.50	85.40	8.45	20.85	30.50
$N_1M_0$	44.00	74.00	85.30	8.35	18.90	27.50
$N_1M_1$	45.90	76.80	86.00	9.45	21.60	29.30
$N_1M_2$	47.95	79.90	87.75	10.05	22.50	32.00
$N_1M_3$	50.05	80.80	90.75	10.50	22.95	32.45
$N_2M_0$	46.55	78.00	90.25	9.50	23.20	29.60
$N_2M_1$	50.00	81.20	94.30	11.30	24.30	32.10
$N_2M_2$	53.25	85.30	97.35	12.50	25.80	34.20
$N_2M_3$	57.50	89.20	100.50	15.10	27.95	36.80
$N_3M_0$	45.75	79.25	86.30	9.05	22.15	29.05
$N_3M_1$	47.50	80.10	88.75	10.40	23.10	30.30
$N_3M_2$	51.80	83.40	91.45	12.10	24.00	33.20
$N_3M_3$	54.20	84.80	96.50	12.40	25.10	34.30
CV%	7.41	8.67	8.25	9.98	8.16	10.97
LSD (0.05)	3.82	3.56	1.31	0.76	4.69	4.04

Means, in a column followed by same letter do not differ significantly at 5% level

$N_0=0$  kg/ha ,  $N_1=Zn_2B_{1.5}$  kg/ha,  $N_2=Zn_4B_2$  kg/ha,  $N_3=Zn_6B_{2.5}$  kg/ha,  $M_0=0$  ton/ha,  $M_1=Cowdung$  (15 ton/ha),  $M_2=Poultry\ manure$  (10 ton/ha),  $M_3=CD+PM$  (7.5+5 ton/ha)

### 3.5 Number of fruits per plant

Significant variation was noted regarding the number of fruits per plant of tomato due to the application of different levels of micronutrients at 45 and 65 DAT (Table5). AT 45 DAT, the maximum number of fruit per plant (5.61) was observed from  $N_2$  ( $Zn_4B_2$  kg/ha) and minimum number of fruits per plant (1.18) was observed from  $N_0$  (0 kg/ha) treatment .At 65 DAT, the maximum number of fruits per plant (34.80) was observed from treatment  $N_2$ .The minimum number of fruits per plant (17.68) was observed from treatment  $N_0$ , [31] reported the similar result. The number of fruits per plant of tomato varied significantly due to the application of different concentrations of organic manure at 45 and 65 DAT (Table 5). At 45 DAT, the maximum number of fruits per plant (3.79) was observed from  $M_3$  (Cowdung7.5 ton+Poultry manure 5 ton/ha) which was statistically similar to  $M_2$  (3.47) and minimum number fruits per plant (2.65) treatment was observed in  $M_0$  (0 ton/ha). At 65 DAT, the maximum number of fruit per plant (26.57) was observed from treatment  $M_0$  (control). [33] reported similar result. The combined effect of micronutrients and organic manure showed statistically significant variation on number of fruit per plant at 45 and 65 DAT (Table 6). At 45 DAT, the maximum number of fruit per plant (6.75) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5 ton+Poultry manure 5 ton/ha) and control treatment ( $N_0M_0$ ) gave the minimum number of fruits per plant (0.95). At 65 DAT, the maximum number of fruit per plant (36.80) was obtained from  $N_2M_3$  treatment. The control treatment ( $N_0M_0$ ) gave the minimum number of fruits per plant (15.70).

### 3.7 Brix%

Brix% of tomato varied significantly due to the application of different levels of micronutrients (Table 4). The highest Brix% of fruit (5.86) was observed from N<sub>2</sub> (Zn<sub>4</sub> B<sub>2</sub> kg/ha) and lowest brix% of fruit (3.61) was observed from N<sub>0</sub> (0 kg/ha) treatment. Brix% of tomato varied significantly due to the application of different levels of organic manure (Table 5). The highest brix% of fruit (6.55) was observed in M<sub>3</sub> (Cowdung 7.5 ton+Poultry manure 5 ton/ha) and lowest brix% fruit (2.88) was observed from M<sub>0</sub> (0 ton/ha). The combined effect of micronutrients and organic manure showed statistically significant variation on brix% of fruit (Table 6). The highest brix% of fruit (8.60) was obtained from N<sub>2</sub>M<sub>3</sub> (Zn<sub>4</sub>B<sub>2</sub>) kg/ha With Cowdung 7.5 ton+Poultry manure 5 ton/ha) and control treatment (N<sub>0</sub>M<sub>0</sub>) gave the lowest brix% of fruit (2.55).

**Table 4: The combined effect of different levels of micronutrients and organic manure on number of branches, no of flowers per plant of tomato at different days after transplanting**

Treatments	No of branches		No of flowers per plant	
	45 DAT	65 DAT	45 DAT	65 DAT
N0M0	3.15	4.1	9.25	20.5
N0M1	3.45	4.3	9.55	24.75
N0M2	3.7	4.6	10.5	31
N0M3	3.9	4.85	12.95	33.9
N1M0	3.7	4.25	9.74	31.75
N1M1	3.9	4.75	12.34	36.63
N1M2	4	5.05	15.17	40
N1M3	4.1	5.14	17.27	49.68
N2M0	4.05	5.7	16.05	64.5
N2M1	4.7	6.25	19.3	80.6
N2M2	5.45	6.85	23.6	94.5
N2M3	6.05	7.25	24	98.8
N3M0	3.9	5.1	13.1	42.5
N3M1	4.45	5.4	17.2	49.9
N3M2	4.75	6.05	18.9	60.65
N3M3	5.05	6.35	20.5	78.25
CV%	10.68	11.58	9.27	9.56
LSD (0.05)	0.51	0.59	4.28	4.42

Means, in a column followed by same letter do not differ significantly at 5% level

N<sub>0</sub>=0 kg/ha , N<sub>1</sub>=Zn<sub>2</sub>B<sub>1.5</sub> kg/ha, N<sub>2</sub>=Zn<sub>4</sub>B<sub>2</sub> kg/ha, N<sub>3</sub>=Zn<sub>6</sub>B<sub>2.5</sub> kg/ha, M<sub>0</sub>=0 ton/ha, M<sub>1</sub>=Cowdung (15 ton/ha), M<sub>2</sub>=Poultry manure (10 ton/ha), M<sub>3</sub>=CD+PM (7.5+5 ton/ha)

### 3.8 Dry matter contents in fruit (%)

Diverse variation was seen as to the dry matter content of tomato due to the application of different levels of micronutrients (Table 5). The highest dry matter of fruit (5.95) was observed from N<sub>2</sub> (Zn<sub>4</sub> B<sub>2</sub> kg/ha) treatment and lowest dry matter of fruit (2.31) was observed from N<sub>0</sub> (0 kg/ha) treatment. [34] reported similar result. Dry matter content of tomato varied significantly due to the application of different concentrations of organic manure Table 4. The highest dry matter of fruit (4.77) was observed in M<sub>3</sub> (Cowdung 7.5 ton+Poultry manure 5 ton/ha) and lowest dry matter of fruit (3.42) was observed from M<sub>0</sub> (0 ton/ha) treatment. The combined effect of micronutrients and organic manure showed statistically

significant variation on dry matter of fruit Table 6. The highest dry matter of fruit (7.00) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5 ton+Poultry manure 5 ton/ha) and control treatment ( $N_0M_0$ ) gave the lowest dry matter of fruit (2.05).

### 3.9. Fruit yield per plot (kg)

The yield of fruits per plot differs noticeably as to different levels of micronutrients (Fig 5). The highest yield/plot (20.09 kg) was observed from  $N_2$  ( $Zn_4 B_2$  kg/ha) and lowest yield/plot (7.70kg) was observed from  $N_0$  (0 kg/ha) treatment. Fruit yield per plot of tomato varied significantly due to the application of different levels of organic manure (Fig 6). The highest yield/plot (15.23 kg) was observed from  $M_3$  (Cowdung 7.5 ton+Poultry manure 5 ton/ha) and lowest yield/plot (11.95kg) was observed from  $M_0$  (0 ton/ha) treatment. The combined effect micronutrients and organic manure showed statistically significant variation on yield/plot (Table7). The highest yield /plot (22.90 kg) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5 ton+Poultry manure ton/ha), on the other hand, control treatment ( $N_0M_0$ ) gave the lowest yield/plot (7.38kg). Fruit yield per plant significantly affected as to combinidly application of micronutrients and organic manure. It has observed that the minimum fruit yield per plant was found from without micronutrients and organic manure (Table 7). The possible reason for higher fruit yield per plant might be due to higher number of fruits per plant, bigger fruit size and fruit weight. Dube *et al.* (2004) found the highest tomato yield with the soil application of micronutrients.

### 3.10 Fruit yield per hectare

Yield of tomato per hectare varied significantly due to the application of different levels of micronutrients (Fig 7) .The highest yield/hectare (66.96 ton/ha) was observed from  $N_2$  ( $Zn_4 B_2$  kg/ha ) and lowest yield/ha (25.69 ton/ha) was observed from  $N_0$  (0 kg/ha) treatment. [35,36] reported similar result Fruit yield of tomato per hectare varied significantly due to the application of different levels of organic manure (Fig 8). The highest yield/hectare (50.78 ton/ha) was observed from  $M_3$  (Cowdung7.5+Poultry manure 5 ton/ha) and lowest yield/hectare (39.86 ton/ha) was observed from  $M_0$  (0 ton/ha). The Combined effect of micronutrients and organic manure showed statistically significant variation on yield/ha (Table 7). The highest yield /hectare (76.33 ton/ha) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5+Poultry manure5 ton/ha) and control treatment ( $N_0M_0$ ) gave the lowest yield/hectare (24.60 ton/ha).

### 3.11 Gross return

In the combination of micronutrients and organic manure showed different gross return under the trial. The highest gross return (Tk. 725136/ha) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5 ton+Poultry manure 5 ton/ha) On the other hand, the lowest gross return (Tk. 233700/ha) was calculated from control treatment ( $N_0M_0$ ) (Table 7).

### 3.12 Net return

In case of net return, different treatment combinations showed different results. The highest (Tk. 499128/ha) net return were obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5+Poultry manure 5 ton/ha). The lowest net return (Tk. 8989/ha) was obtained from control treatment ( $N_0M_0$ ) (Table 7).

### 3.13 Benefit cost ratio (BCR)

The combination of micronutrients and organic manure for benefit cost ratio was different in all treatment combination (Table 7). The highest benefit cost ratio (3.2) was obtained from  $N_2M_3$  ( $Zn_4B_2$  kg/ha with Cowdung7.5 ton+Poultry manure 5 ton/ha) and whereas the lowest benefit cost ratio (1.04) was obtained from control treatment ( $N_0M_0$ ). From the economic point of view, it is apparent that  $N_2M_3$  treatment combination ( $Zn_4B_2$  kg/ha with Cowdung7.5+Poultry manure 5 ton/ha) was the most profitable than rest of the treatment combinations under the study.

**Table 5: The effect of different levels of micronutrients and organic manure on the number of fruits per plant, individual fruit weights, brix%, Dry matter contents in fruit % of tomato at different days after transplanting**

Treatments	Number of fruits	Individual	Brix%	Dry matter
------------	------------------	------------	-------	------------

	per plant		fruit weight(g)		contents in fruit (%)
	45 DAT	65 DAT			
<b>Micronutrients</b>					
N <sub>0</sub>	1.18	17.68	85.76	3.61	2.31
N <sub>1</sub>	2.47	21.75	92.04	4.42	3.24
N <sub>2</sub>	5.61	34.80	104.71	5.86	5.95
N <sub>3</sub>	3.71	27.01	100.66	5.45	4.85
CV%	9.37	10.42	9.31	7.17	11.74
LSD(0.05)	0.48	2.02	2.01	0.46	0.15
<b>Organic manure</b>					
M <sub>0</sub>	2.65	23.90	86.74	2.88	3.42
M <sub>1</sub>	3.07	25.04	92.48	3.68	3.80
M <sub>2</sub>	3.47	25.72	98.96	6.22	4.36
M <sub>3</sub>	3.79	26.57	105.00	6.55	4.77
CV%	9.37	10.42	9.31	7.17	11.74
LSD(0.05)	0.43	0.64	2.67	0.43	0.22




Fig. 5. Graphical representation of yield/plot for treatment (N<sub>0</sub> to N<sub>3</sub>)




Fig. 6. Graphical representation of yield/plot for treatment (M<sub>0</sub> to M<sub>3</sub>)




Fig. 7. Graphical representation of yield (Ton/ha)




Fig. 8. Graphical representation of yield (Ton/ha)

for treatment (N<sub>0</sub> to N<sub>3</sub>)

for treatment (M<sub>0</sub> to M<sub>3</sub>)

**Table 6: The combined effect of different levels of micronutrients and organic manure on the number of fruits per plant, individual fruit weights, brix%, Dry matter contents in fruit % of tomato at different days after transplanting**

Treatments	Number of fruits per plant		Individual fruit weight(g)	Brix%	Dry matter contents in fruit (%)
	45 DAT	65 DAT			
N <sub>0</sub> M <sub>0</sub>	0.95	15.70	78.95	2.55	2.05
N <sub>0</sub> M <sub>1</sub>	1.01	17.82	83.00	3.50	2.18
N <sub>0</sub> M <sub>2</sub>	1.39	18.10	87.45	4.05	2.37
N <sub>0</sub> M <sub>3</sub>	1.40	19.10	93.65	4.35	2.64
N <sub>1</sub> M <sub>0</sub>	1.99	20.40	84.20	2.70	2.79
N <sub>1</sub> M <sub>1</sub>	2.47	21.75	90.15	3.75	2.95
N <sub>1</sub> M <sub>2</sub>	2.50	22.25	94.85	4.75	3.40
N <sub>1</sub> M <sub>3</sub>	2.95	22.60	98.95	6.50	3.85
N <sub>2</sub> M <sub>0</sub>	4.65	33.18	93.70	3.05	5.00
N <sub>2</sub> M <sub>1</sub>	5.10	33.94	99.95	3.75	5.60
N <sub>2</sub> M <sub>2</sub>	5.95	35.30	107.80	8.05	6.20
N <sub>2</sub> M <sub>3</sub>	6.75	36.80	117.40	8.60	7.00
N <sub>3</sub> M <sub>0</sub>	3.04	26.35	90.10	3.25	3.85
N <sub>3</sub> M <sub>1</sub>	3.70	26.65	96.80	3.75	4.50
N <sub>3</sub> M <sub>2</sub>	4.05	27.25	105.75	8.00	5.47
N <sub>3</sub> M <sub>3</sub>	4.08	27.80	110.00	6.75	5.60
CV%	9.37	10.42	9.31	7.17	11.74
LSD (0.05)	0.97	2.04	2.03	0.93	0.31

**Table 7: The combined effect of different levels of micronutrients and organic manure on yield per plot and yield ton per hectare of tomato at different days after transplanting and return of tomato cultivation as influenced by micronutrients and organic manure**

Treatments	Yield per plot (kg)	Yield ton per ha	Gross return (Tk./ha)	Net return (Tk./ha)	Benefit cost ratio
N <sub>0</sub> M <sub>0</sub>	7.38	24.60	233700	8989	1.04
N <sub>0</sub> M <sub>1</sub>	7.55	25.16	239020	12521	1.05
N <sub>0</sub> M <sub>2</sub>	7.75	25.83	245385	37961	1.18
N <sub>0</sub> M <sub>3</sub>	8.15	27.16	258020	41059	1.25
N <sub>1</sub> M <sub>0</sub>	9.25	30.85	293075	100363	1.52
N <sub>1</sub> M <sub>1</sub>	11.65	38.81	368695	137373	1.59
N <sub>1</sub> M <sub>2</sub>	12.45	41.50	394250	182003	1.85
N <sub>1</sub> M <sub>3</sub>	13.15	43.83	416385	218856	2.11
N <sub>2</sub> M <sub>0</sub>	18.15	60.49	574655	344793	2.50
N <sub>2</sub> M <sub>1</sub>	19.13	63.76	605720	370174	2.57
N <sub>2</sub> M <sub>2</sub>	20.18	67.28	639160	422689	2.95
N <sub>2</sub> M <sub>3</sub>	22.90	76.33	725136	499128	3.20
N <sub>3</sub> M <sub>0</sub>	13.05	43.49	413155	183625	1.80
N <sub>3</sub> M <sub>1</sub>	14.15	47.16	448020	208251	1.86
N <sub>3</sub> M <sub>2</sub>	14.70	49.00	465500	244806	2.10
N <sub>3</sub> M <sub>3</sub>	16.75	55.83	530385	324678	2.57
CV%	12.87	12.43	-	-	-
LSD (0.05)	1.01	2.39	-	-	-

Means, in a column followed by same letter do not differ significantly at 5% level

Here, N<sub>0</sub>=0 kg/ha, N<sub>1</sub>=Zn<sub>2</sub>B<sub>1.5</sub> kg/ha, N<sub>2</sub>=Zn<sub>4</sub>B<sub>2</sub> kg/ha, N<sub>3</sub>=Zn<sub>6</sub>B<sub>2.5</sub> kg/ha, M<sub>0</sub>=0 ton/ha, M<sub>1</sub>= Cowdung (15 ton/ha), M<sub>2</sub>=Poultry manure (10 ton/ha), M<sub>3</sub>=CD+PM (7.5+5 ton/ha), Sale of marketable tomato @ Tk. 9,500/ton

#### 4. CONCLUSION

Both crop yield and economic benefit of crop are important for the crop production. According to the results of the present experiment, it may be concluded that efficient production of tomato is increased by the application of micronutrients and organic manure. Thus, the combined application of micronutrients and organic manure may be helpful for higher and better qualitative tomato production in considering crop productivity and economic return of tomato. On the basis of benefit cost ratio, it may be suggested that a combination of 4 kg zinc and 2 kg boron per hectare with cowdung 7.5 ton/ha+ poultry manure 5ton/ha gave maximum and profitable yield of the tomato.

#### REFERENCES

1. Ahmed, K. 1995. Phul Phal O Shak-Sabjee. 5th Edition, 414 Senpara, Parbata, Mirpur, Dhaka, Bangladesh.
2. Hossain, M.E., Alam, M.J., Hakim, M.A., Amanullah, A.S.M. and Ahsanullah, A.S.M. 2010. An Assessment of Physicochemical Properties of Some Tomato Genotypes and Varieties Grown At Rangpur. Bangladesh Res. Pub. J., 4(3): 135-243.
3. BBS. 2015. Year book of Agricultural statistics of Bangladesh. 27th edition. Bangladesh Bureau of Statistics. Statistics Division, Ministry of planning. Govt of the people Republic of Bangladesh. P-304.

4. FAO. 2002. FAO Production Year Book. Food and Agricultural Organization of the United Nations, Rome 00100, Italy.
5. Bose, T. K. and Som, M. G. 1990. Vegetable Crops in India. Naya Prakash, Calcutta Six, India. pp. 687-691.
6. Mac Gillivray, K.1961. Nutritional status of winter tomato. *Vegetable Science*. 16(2): 119-124
7. Aditya, T. L., Rahman, L., Alam M. S. and Ghoseh , A. K. 1997. Correlation and path co-efficient analysis in tomato. *Bangladesh J. Agril. Sci.*, 26(1):119-122.
8. Islam, J. K. and Noor, S. P.1982. Deficiency of soil nutrient is now considered as one of the major constraints to successful upland crop production in Bangladesh. *Pakistan J. Sci. Res.*, 34 (3-4): 113-119.
9. Shukla, V. and Naik, L.B. 1993. Agro-techniques for Solanaceous Vegetables. In: *Vegetable Crops: Part-I, Advances in Horticulture*. Vol. 5, Eds: K. L. Chadha and G. Kalloo, Maihotra Publishing House, New Delhi, India . p.371
10. Magalhaes, J.R. I., Solwa Dc,C.E.W.L and Monnerat. P.H.1980. Levels and methods of boron application in tomatoes. *Pesquisa Agropecuria Brasilesia* 10 (2): 153-157 [Cited from Hort. Abstr. 50(4): 2031, 19811.
11. Mallick, M.F.R. and Muthukrishnan, C.R.1980. Effect of micro nutrients on tomato (*Lycopersicon esculentum* Mill.), Effect on flowering, fruit-set and yield. *South Indian Hort.* 28 (1): 14-20.
12. Alloway, B.J. 2008. Fundamental aspects of zinc in soils and plants. *Zinc in Soils and crop nutrition*. 2nd Edi, Published by IZA and IFA, Brussels, Belgium and Paris, France, pp: 30-52.
13. Passam, C.H., Ioannis, C., Karapanos, J., Penelope, Bebeli and Savvas, D. 2007. A review of recent research on tomato nutrition, breeding and post-harvest technology with reference to fruit quality. *Eur. J. Plant Sci. Biotechnol.* 1(1): 121.
14. Srivastava, A.K. and Singh,S. 2003. Soil-plant nutrient limits in relation to optimum fruit yield of sweet orange (*Citrus sinensis* Osbeck) cultivar Mosambi. *Indian. J. Agric. Sci.* 73(4): 209-211.
15. Bose, U.S. and Tripathi, S.K.1996. Effect of micronutrients on growth, yield and quality of tomato cv. Pusa Ruby. *Crop. Res. Hisar*, 12(1): 61-64.
16. Nonnecke, I.B.L.1989. *Vegetable Production*. Avi Book Publishers. New York, USA. pp.200-229.
17. Sperry, W.J. 1995. Influence of boron and potassium on quality, yield, and nutrition of fresh-market tomato. PhD dissertation. N.C. State Univ., Raleigh, N.C.
18. Blevins, D.G., Reinbott, T.M. and Boyce, P.J. 1993. Foliar fertilization of soybeans with boron and magnesium. *Plant Physiol.* p. 1-6.
19. Stanley, D.W., Bourne, M.C., Stone A.P. and Wismer. W.V. 1995. Low temperature blanching effects of chemistry, firmness and structure of canned green beans and carrots. *Food Sci.* 60(5): 327-333.
20. Tisdale, S.L., Nelson,W.L. and Beaton.J.D. 1985. *Soil fertility and fertilizers*. pp: 754.
21. Naresh, B. 2002. Response of foliar application of boron on vegetative growth, fruit yield and quality of tomato var. Pusa-Ruby. *Indian J. of Hilly Farming*. 15(1):109-112.
22. BARC.1997. *Fertilizer Recommendation Guide*. Bangladesh Agricultural Research Council. Farmgate, Dhaka-1215. pp. 1-72.
23. Opena, R.T., Kuo, C.C. and Yoon, J.Y.1988. Breeding and Seed Production of Chinese cabbage in the Tropics and Subtropics. *T Ozbun, L., ech. Bul.*, 17, AVRDC. p. 97.
24. Ewulo, B. S., Ojeniyi.S. O. and Akanni, D. A.2008. Effect of poultry manure on selected soil physical and chemical properties, growth, yield and nutrient status of tomato. *African Journal of Agricultural Research* Vol. 3 (9), pp. 612-616,
25. Mahimairaja, S., Bolan, N.S. and Hedley, M.J. (1995) Agronomic effectiveness of poultry manure composts. *Communications in Soil Science and Plant Analysis* 26: 18431861.
26. Adeli, A., Tewolde, H., Sistani, K.R. and Rowe, D.E. 2009. Broiler litter fertilization and cropping system impacts on soil properties. 110: 1304-1310.
27. Adediran, J.A., Taiwo, L.B. and Sobulo, R.A. 2003. Organic wastes and their effect on tomato (*Lycopersicon esculentum*) yield. *Afr. Soils* 33: 99-116.
28. Alam, M.S., Iqbal, T.M.T., Ami, M. and Gaffar, M.A. 1989. Kamishitattic Fasaler Utpadan O Unnayan (In Bengali). T. M. Jubair Bin Iqbal, Sirajganj. pp. 231-239.

29. Dube, B. K., S. Pratima and C. Chatterjee. 2004. Effects of boron and zinc on the yield and quality of tomato. *Indian J. Hort.* 61(1): 48-52. Hunter, A. H. 1980.
30. Oyinlola, E.Y. 2004. Response of irrigated tomatoes to boron fertilizer on growth and nutrient concentration. *Nigerian J. Soil Res.* 5(2): 62-68.
31. Ejaz, M., Rahman, S., Waqas, R., Manan, A., Imran, M. and Bukhari, M.A. 2011. Combined efficacy of macro-nutrients and micro-nutrients as foliar application on growth and yield of tomato. *Int. J. Agro Vet. & Medic. Sci.* 5(3): 327-335.
32. Amrachandra, S. and Verma, B.K. 2003. Effect of boron and calcium on plant growth and seed yield of tomato. *JNKVV Res. J. of India.* 37(2): 13-14.
33. Solaiman, A. and Rabbani, M.M. 2006. Effects of NPKS and cowdung on growth and yield of tomato. *Bulletin of the Institute of Tropical Agriculture, Kyushu University.* 29: 31-37.
34. Salam, M.A., Siddique, M.A. and Rahim, M.A. 2010. Quality of tomato (*Lycopersicon esculentum* mill.) As influenced by boron and zinc under different levels of NPK fertilizers. *Bangladesh J. Agric. Res.* 35(3):475-488.
35. Gurmani, A.R., Andaleep, R. and Khan, S. 2012. Biochemical attributes, plant growth and yield of tomato as affected by boron application rates. *J. Agric. Res.* 48(2):451-455.
36. Hossein, M. 2008. Effect of zinc and boron on the growth and yield of tomato. M.S. Thesis. Department of Agricultural Chemistry, BAU, Mymensingh.