

1
2
3
4
5
6
7
8
9

Phenylthiocarbamide Taste Perception among Patients with Type 2 Diabetes Mellitus

ABSTRACT

Background: Taste perception has been associated with some diseases, disorders and treatment medications. **Aim:** To determine whether PTC taste blindness was associated with type 2 Diabetes Mellitus (DM) and possible relationship between intake of treatment medications and PTC taste sensitivity. **Methodology:** The study participants consisted of 100 type 2 DM patients on treatment (group 1) and 100 newly diagnosed type 2 DM patients not on drugs treatment (group 2). Apparently healthy individuals served as control (group 3). Informed consent was obtained from each participant at the commencement of the study. Tasters and non-tasters were determined using phenylthiocarbamide (PTC) taste strips (0.0143 mg/strip).

Results: In group 1, 66% were non-tasters; group 2 60% were non-tasters while 37% in group 3 were non-tasters. Phenylthiocarbamide taste perception varied significantly among the 3 groups studied ($p < 0.001$). Non-tasters of PTC in groups 1 and 2 were not significantly different ($p = 0.38$). Non-tasters of PTC in groups 1 and 2 ($p < 0.001$; OR 3.30 and $p = 0.001$; OR 2.55 respectively) were significantly higher than non-tasters in the control (group 3). **Conclusion:** This study shows that inability to taste PTC is associated with type 2 DM. However, intake of DM treatment medications does not appear to have any significant influence on PTC taste sensitivity.

10 **Keywords:** *Diabetes Mellitus, Phenylthiocarbamide Taste Perception, Tasters, Non-tasters*

11
12
13

14 **1. INTRODUCTION**

15

16 Diabetes mellitus (DM) is a chronic disorder which poses a major health challenge to
17 humans. About 422 million people were reported to have diabetes globally [1]. It can be
18 classified into 4 types namely; type 1 which is more common in children and adolescents
19 than in adults is an autoimmune disease where the body forms antibodies against its beta
20 cells of islet of Langerhans in the pancreas; type 2 which is associated with adults,
21 characterized by peripheral insulin resistance and inadequate insulin secretion by the
22 pancreas; Secondary DM is caused by another disease or disorder and lastly, gestational
23 DM caused by pregnancy.

24 Type 2 DM is reported to make up about 90% of all cases of DM [2, 3]. There are reports of
25 higher prevalence of type 2 diabetes in men compared to women which has been associated
26 with sex-related differences in visceral fat mass [4]. The disorder can be asymptomatic in an
27 individual for many months and years.

28 Taste has influence on one's choice of food. It allows one to choose the food one likes most.
29 Some diseases such as liver diseases, tumour and lifestyle such as consumption of alcohol
30 together with the use of drugs, head trauma, upper respiratory tract infections and exposure
31 to toxic substances have been reported to significantly influence taste [5-7]. It is thought that
32 understanding factors related to taste perception will provide opportunity to evaluate the
33 feeding behaviour of patients with chronic diseases [8].

34 Phenylthiocarbamide taste sensitivity is correlated strongly with the ability to taste other
35 naturally occurring bitter substances [9, 10]. Bitter taste perception occurs through bitter
36 taste receptors located on the surface of taste cells of the tongue [11] and is thought to be a
37 conserved chemical sense in mammals against the ingestion of naturally toxic substances
38 [12]. Taste sensitivity impairment may make an individual to ingest greater quantities of
39 substances which in turn may adversely tamper with the health of the individual. A number of

40 previous studies had been carried out on relationship between diabetes and PTC taste
41 perception. Some of these studies reported positive interactions between inability to taste
42 PTC and DM [13, 14]; others reported lack of an association between PTC taste blindness
43 and DM [15].

44 In Nigeria, PTC taste perception has been studied in relation to some common diseases
45 such as malaria, tuberculosis and HIV infection [16-18]. Type 2 DM is also quite common in
46 Nigeria but to us, there is no known investigation that has related PTC taste perception with
47 DM. Therefore, this study was carried out to determine whether there was any association
48 between type 2 DM and PTC taste perception and to ascertain whether the taste perception
49 was influenced by the intake of DM treatment medications.

50

51 **2. METHODOLOGY**

52

53 This study was carried out in Osogbo and Ogbomoso, Southwestern Nigeria. Participants
54 were drawn from patients attending diabetes clinics of Ladoké Akintola University of
55 Technology (LAUTECH) Teaching Hospitals in Osogbo and Ogbomoso, Nigeria. A total of
56 300 individuals participated in this study. The study participants consisted of 100 type 2 DM
57 patients who had been diagnosed for not less than six months and on metformin treatment
58 (group 1), 100 newly diagnosed type 2 DM patients not on drugs treatment (group 2) and
59 100 apparently healthy individuals as control (group 3). Informed consent was obtained from
60 each participant at the commencement of the study after explaining the essence and
61 procedure of the test. The criteria for diagnosis of DM included fasting blood glucose test: \geq
62 126 mg/dl (7.0 mmol/l). Two fasting glucose measurements \geq 7.0 mmol/l (126 mg/dl) were
63 considered diagnostic for diabetes mellitus. Patients who had other health conditions in
64 addition to diabetes were excluded from the study. Ethical approval for this study was
65 obtained from the Ethical Committee of the College of Health Sciences, Ladoké Akintola
66 University of Technology (LAUTECH), Osogbo.

67 Phenylthiocarbamide taste strips (0.0143 mg of PTC/strip) used were obtained from Carolina
68 Biological Supply Company, North Carolina, USA. Each participant was given a PTC taste
69 strip and a filter paper (as control) and was asked to put each on their tongue and allow to
70 be soaked in their saliva before describing their perception to each strip. Taste description of
71 each participant was recorded. Questionnaire was administered to each participant to obtain
72 relevant information such as age, sex, the drug being received for those on diabetes
73 medication and the like. Data were analysed using percentages. Differences in percentages
74 were tested by Chi-square test. A p-value of < 0.05 was considered to be significant.

75 **3. RESULTS**

76
77 Results from the study (Table1) showed that the distribution of the participants across the
78 three groups with respect to age ($\chi^2 = 0.20$, df = 2, p = 0.90) and sex ($\chi^2 = 0.51$, df = 2, p =
79 0.77) were not statistically significantly different.

80 In group1 (> 6 months DM patients), 47% were males (15% tasters plus 32% non-tasters)
81 while 53 were females (19% tasters plus 34% non-tasters). Also, of the 100 newly diagnosed
82 diabetic patients (group 2), 45% were males (18% tasters plus 27% non-tasters) and 55%
83 were females (22% tasters plus 33% non-tasters). In addition, of the 100 control subjects,
84 42 were males (26% tasters plus 16% non-tasters) and 58 were females (37 tasters plus 21
85 non-tasters). Phenylthiocarbamide taste perception varied significantly among the 3 groups
86 both in males ($\chi^2 = 8.54$, df = 2, p = 0.01) and in females ($\chi^2 = 10.29$, df = 2, p = 0.01).
87 Further Chi-Square tests showed that differences observed in the male groups were
88 between > 6 months DM group and controls ($\chi^2 = 8.03$, df = 1, p = 0.005) and between
89 newly diagnosed DM group and controls ($\chi^2 = 4.17$, df = 1, p = 0.04). Similarly, the
90 differences observed in the female groups were between group 1 and controls ($\chi^2 = 8.65$, df
91 = 1, p = 0.003) and between group 2 and controls ($\chi^2 = 6.41$, df = 1, p = 0.01).

92 Also, the distributions of the study participants with respect to PTC taste perception are
93 given in Table 1. Sixty-six percent (66%) of the diabetic group of > 6 months were non-
94 tasters, 60% of the newly diagnosed diabetic group were non-tasters while 37% of the

95 control group were non-tasters. Phenylthiocarbamide taste perception varied significantly
 96 among the three groups ($\chi^2 = 18.89$, $df = 2$, $p < 0.001$). Further Chi-Square tests showed
 97 significant differences between the diabetic group on medication and control group ($\chi^2 =$
 98 16.84 , $df = 1$, $p < 0.001$; OR 3.30, CI 1.86 – 5.85) and between the newly diagnosed diabetic
 99 group and control group ($\chi^2 = 10.59$, $df = 1$, $p = 0.001$; OR 2.55, CI 1.45 – 4.47). There was
 100 no significant difference in taste sensitivity between groups 1 and 2 ($\chi^2 = 0.77$, $df = 1$, $p =$
 101 0.38).

102

103 **Table 1: Distribution of the Study Participants by Age, Sex and Phenylthiocarbamide**
 104 **Taste Perception**
 105

Variable	DM Patients Group 1 n=100	DM Patients Group 2 n=100	Non-DM Subjects Group 3 n=100	p
Age (years)				0.90
36-45	15	17	18	
46-55	33	35	32	
≥56	52	48	50	
Sex				0.77
Male	47(15T; 32NT)	45(18T; 27NT)	42(26T;16NT)	0.01
Female	53(19T; 34NT)	55(22T; 33NT)	58(37T;21NT)	0.01
PTC Tasting				<0.001
Taster	34	40	63	
Non-Taster	66	60	37	

106 DM: Diabetes Mellitus T: Taster; NT: Non-taster. Whole figures are in percentages
 107

108

109

110 4. DISCUSSION

111

112 In this study, diabetes patients were more likely to be non-tasters of PTC than non-diabetes
 113 individuals. This is in line with the studies of some other researchers who had reported that
 114 inability to taste PTC or PTC taste blindness was associated with diabetes mellitus [13, 14].

115 The observation that non-taster status was significantly associated with diabetes in this

116 study could suggest that the gene for PTC might directly or indirectly interact with that of
117 diabetes to confer susceptibility to DM individuals. Polymorphism in TAS2R38 had been
118 linked with differences in ingestive behaviour of tasters and non-tasters which might be
119 associated the development of pre-diabetes and type 2 DM [19].

120 This study showed that the use of metformin did not influence the association reported since
121 there was no significant difference with respect to taste blindness between the participants
122 on medication and the newly diagnosed diabetic patients. This implied that unlike in HIV
123 infected persons where medication had been reported to alter taste [18, 20]; taste alteration
124 induced by medication in diabetes was insignificant.

125 It had been reported that taste perception appeared to regulate food consumption and had
126 also been linked with circulating metabolic hormones [21]. Bhatia and Sharma [22] reported
127 a decrease in palatability of glucose solution between tasters and non-tasters. Elevated
128 blood glucose levels resulted in a concentration dependent impairment of taste perception in
129 type 2 DM patients due to adaptation of the sensory cell to increased blood glucose [23].

130 Dais et al. [24] observed that the average thresholds to detect sweet taste were higher for
131 diabetic patients compared to non-diabetes showing a decreased or loss of sensitivity in
132 diabetics. This loss of sensitivity might contribute to increase in sugar consumption among
133 diabetics. Loss of taste perception in individuals with type 2 DM had been related to
134 hyposalivation, xerostomia and low salivary flow [25]. Wang et al [21] reported higher levels
135 of TNF-alpha, IGF-1 and leptin in tasters than in non-tasters and a positive correlation
136 between plasma glucose level and body mass index in non-tasters. The deficiency or
137 absence of taste interfered with salivation and maturation of the taste buds, causing changes
138 in the perception of taste [26].

139

140

141 **4. CONCLUSION**

142

143 We conclude that PTC taste blindness is significantly associated with type 2 DM and that

144 DM medication has no significant influence on PTC taste sensitivity.

145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO (World Health Organisation). Global Report on Diabetes. 2016. <http://who.int/bitstream/10665/204871/1/9789241565257>
2. Chen L, Magliano DJ, Zimmet PZ. The worldwide epidemiology of type 2 diabetes mellitus: present and future perspectives. *Nature Rev Endocrinol*. 2014; 8: 228-236.
3. Baynest HW. Classification, Pathophysiology, Diagnosis and Management of Diabetes Mellitus. *J Diabetes Metabol*. 2015. 6. 5: DOI: 10.4172/2155 – 6156.1000541.
4. Nordstrom A, Hadrevi J, Olsson T, Franks PW, Nordstrom P. Higher Prevalence of Type 2 Diabetes in Men than in Women is Associated With Differences in Visceral Fat Mass. *J Clin Endocrinol and Metabol*. 2016; 101(10): 3740-3746.
5. Hummel T, Landis BN, Hüttenbrink KB. Smell and taste disorders. *GMS Curr Top Otorhinolaryngol Head Neck Surg.*, 2011; 10: 04.
6. Lampuré A, Schlich P, Deglaire A, Castetbon K, Peneau S, Hercberg S, Mejean C. Sociodemographic, psychological, and lifestyle characteristics are associated with a liking for salty and sweet tastes in french adults. *J Nutr*. 2015; 145(3): 587-594.
7. Silva CS, Dias VR, Almeida JA, Brazil JM, Santos RA, Milagres MP. Effect of heavy consumption of alcoholic beverages on the perception of sweet and salty taste. *Alcohol*, 2016; 51(3): 302-306.
8. Loper HB, Sala ML, Dotson C, Steinle N. Taste perception, associated hormonal modulation, and nutrient intake. *Nutr Rev*. 2015; 73(2): 83–91.
9. Tepper BJ. 6-n-propylthiouracil: A genetic marker for taste, with implications for food preference and dietary habits. *Am J Hum Genet*. 1998; 63: 1271-1276.
10. Tepper BJ, Koelliker Y, Zhao L, Ullrich NV, Lanzara C, d'Adamo P, Ferrara A, Ulivi S, Esposito L., Gasparini P. Variation in the bitter-taste receptor gene TAS2R38 and adiposity in a genetically isolated population in Southern Italy. *Obesity*. 2008; 16(10): 2289-2295.
11. Adler E, Hoon MA, Mueller KL, Chandrashekar J, Ryba NJ, Zuker CS. A novel family of mammalian taste receptors. *Cell*. 2000; 100: 693–702.
12. Ueda T, Ugawa S, Yamamura H, Imaizumi Y, Shimada S. Functional interaction between T2R taste receptors and G-protein alpha subunits expressed in taste receptor cells. *J Neurosci*. 2003; 23: 7376–7380.

- 198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
13. Ali S.G, Azad-Khan AK, Mahtab H, Khan AR, Muhibullah M. Association of phenylthiocarbamide taste sensitivity with diabetes mellitus in Bangladesh. *Hum Hered.* 1994; 44: 14-17.
 14. Baruah M, Joshi A. Comparative Study of Taste Sensitivity to Phenylthiocarbamide in Normoglycaemics and in Diabetes Mellitus Type 2. *Int J Pharm Bio Sci.* 2013; 4(4): 727-730.
 15. Sriram K. Balaraman VT. Usha J. The association between taste sensitivity to PTC and Diabetes Mellitus. *Indian J Med Res.*, 1975; 63(3): 390-395.
 16. Igbeneghu C, Owoeye Y, Akanni EO. Association between phenylthiocarbamide (PTC) taste perception and falciparum malaria infection in Osogbo, Southwestern Nigeria. *Ann. Res. Rev. Biol.*, 2014; 4 (14): 2295-2301.
 17. Igbeneghu C, Gabriel BA, Onuegbu JA, Olisekodiaka JM, Adesiyun AA. 2016. Phenylthiocarbamide (PTC) Taste Perception among Pulmonary Tuberculosis Patients in Southwest Nigeria. *Schl J App Med Sci.* 2016; 4 (6F): 2248-2251.
 18. Igbeneghu C, Oluwatunbi TB, Aina OA, Olisekodiaka MJ. 2017. Phenylthiocarbamide Taste Perception among HIV-infected Patients on Highly Active Antiretroviral Therapy in Southwestern Nigeria. *Bri J Med Med Res.* 2017; 19 (10): 1-7.
 19. Dotson CD, Shaw HL, Mitchell BD, Munger SD, Steinle NI. Variation in the gene TAS2R38 is associated with the eating behaviour disinhibition in older Amish women. *Appetite.* 2010; 54(1): 93-99.
 20. Mattes RD. The chemical senses and nutrition in aging: challenging old assumptions. *J Am Diet Assoc.* 2002; 102 (2): 192-196.
 21. Wang R, van Keeken NM, Siddiqui S, Dijkman LM, Maudsley S, Derval D, van Dam PS, Martin B. Higher TNF-alpha, IGF-I and Leptin Levels are Found in Tasters than Non-tasters. *Front Endocrinol. (Lausanne)* 2014; 5: 125.
 22. Bhatia S, Sharma KN. Taste impairment for glucose in diabetes PTC tasters and non-tasters. *Diabetes Res Clin Pract.* 1991; 12(3): 193-199.
 23. Bustos-Saldana R, Alfaro-Rodriguez M, Solis-Ruiz Mde L, Trujillo-Hernandez B, Pacheco-Carrasco M, Vazquez-Jimenez C, Calis-de la Rosa Ade J. Taste sensitivity diminution in hyperglycemic type 2 diabetics patients. *Rev Med Inst Mex Seguro Soc.* 2009; 47(5):483-488.
 24. Dias VR, Brazil JM, Almeida JAR, Silva CS, Milagres MP. Evaluation of the sensory perception of sweat taste in people with diabetes mellitus type 2. *Rev Rene.* 2016; 17(4): 483-489.
 25. Lopez-Pintor RM, Casarias E, Gonzalez-Serrano J, Serrano J, Ramirez L, de Arriba L, Hernandez G. Xerostomia, Hyposalivation and Salivary flow in Diabetes Patients. *J Diabetes Res.* 2016; ID 4372852, 15 pp
 26. Negrato CA, Tarzia O. 2010. Buccal alterations in diabetes mellitus. *Diabetol Metab Syndr.* 2010; 15; 2: 3.