

**Prevalence and associated factors of urinary schistosomiasis among basic school children in the Akyemansa District, Ghana**

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

---

**ABSTRACT**

**Background:** Globally, urinary schistosomiasis has devastating implications on school children. It predisposes them to dysuria, haematuria among others which can negatively influence their academic performance. The study determined the prevalence and associated risk factors of urinary schistosomiasis among basic school children in the Akyemansa district.

**Materials and Methods:** A simple randomized cross-sectional study was conducted among 504 basic school children whose urine samples were collected. Each urine sample was microscopically examined for the presence or absence of ova. *Schistosoma haematobium* (SH) ova were quantified as light or heavy.

**Results:** Prevalence of SH infection among school children in Akyemansa District was 10.32% [95% CI: 7.80 -13.31%]. Out of 52 participants who were infected, 69.2% had light infection while the rest had heavy infection. Female participants were less likely to be infected with SH than males [OR=0.47; 95% CI: 0.23-0.97], children who do not stay by the river/stream were also less likely to be infected with SH compared to those who lived near waterbodies [OR=0.35; 95% CI: 0.17-0.72]. Additionally, participants who did not play around water bodies were also less likely to be infected with SH compared to those who did [OR=0.17; 95% CI= 0.04-0.71; p=0.015]. However, inhabitants of Kotokuom were more likely to be infected compared to those in Pawuda [OR=8.54; 95% CI: 1.91-38.27; p=0.005]

**Conclusion:** The prevalence of urinary schistosomiasis among basic school children in the Akyemansa district was found to be 10.3% [95% CI: 7.80 -13.31%]. Gender, staying around river/ stream, playing at river/ stream and inhabitant of participants were significantly associated with the prevalence of *Schistosoma haematobium* infection. The study therefore recommends that periodic drug administration and a comprehensive intervention strategy should be designed and implemented to reduce schistosomiasis prevalence.

*Keywords: Urinary Schistosomiasis, School Children, Akyemansa District.*

**1. INTRODUCTION**

Schistosomiasis has been shortlisted among the world's major public health challenges with marked incidence in about 77 developing countries in the tropics and subtropics resulting in increased morbidities and mortalities (Bruun & Aagaard-Hansen, 2008; Thétiot-Laurent, J., Robert, & Meunier, 2013). It has been put on record that about 249 million people are

23 infected globally with an alarming 779 million people at risk across the globe (Thétiot-  
24 Laurent, J., Robert, & Meunier, 2013).

25 Human schistosomiasis is a parasitic infection caused by various species of *Schistosoma*,  
26 namely, *Schistosoma haematobium* (*S. haematobium*), *S. mansoni*, *S. japonicum*, *S.*  
27 *mekongi*, and *S. intercalatum* (Gryseels, Polman, Clerinx, & Kestens, 2006). In 2015, a  
28 report by the Centre for disease Control (CDC) revealed that about 224 million malignant  
29 effects of schistosomiasis are seen annually in sub-Saharan Africa with an estimated  
30 280,000 deaths, largely among rural inhabitants. A study by van der Werf, et al., (2003)  
31 depicted *S. haematobium* infection as the cause of 70 million cases of hematuria, 32 million  
32 cases of dysuria, 18 million cases of bladder-wall pathology and 10 million cases of major  
33 hydronephrosis in sub-Saharan Africa of which Ghana is no exception. The infection which is  
34 crucial among children has been linked with complications such as nutritional deficiencies,  
35 growth retardation, cognitive function impairment, decreasing physical activity, school  
36 performance, and work capacity and productivity (Stephenson, 1993).

37 Urinary schistosomiasis is endemic in Ghana with widespread distribution across the country  
38 causing considerable morbidities. Its transmission is by fresh water snail (host) that belongs  
39 to the genus *Bulinus* and water contacts through human activities, particularly with children  
40 between the ages of 4 and 16 years, the most vulnerable and heavily infected year group.  
41 This makes it difficult to control in relation to children since children who live by water bodies  
42 always love to play with water most of the time. Reports from studies conducted by Aryeetey  
43 et al., (2000), have depicted a range between 54.8 and 60.0% urinary schistosomiasis  
44 prevalence in southern Ghana, with increasing infection rates with age and a peak in the 10-  
45 19year category, and decreasing rate with increasing age. These figures reflect the need to  
46 monitor and control the rate at which children get infected with urinary schistosomiasis.  
47 Therefore, the risk and re-emergence of urinary schistosomiasis is attributable to the range  
48 of snail habitats that are promoted by water development schemes.

49 In ensuring appropriate interventions against schistosomiasis, information on the distribution  
50 and associated risk factors of the disease in different transmission settings remains a  
51 prerequisite. Furthermore, extensive mapping programme is required to facilitate scaling-up  
52 of the much needed preventive chemotherapy intervention in rural communities. This study  
53 therefore determined the prevalence and associated risk factors of *S. haematobium* infection  
54 among basic school children in the Akyemansa district, Eastern region, Ghana.

55

## 56 **2. MATERIAL AND METHODS**

57

### 58 **2.1 Study Design/ Eligibility Criteria**

59 A simple randomized cross-sectional study was conducted in the Akyemansa district in  
60 January, 2014. A total of five hundred and four (504) basic school children were recruited  
61 from six communities within the Akyemansa district. These communities were Pawudu,  
62 Adubiase, Adjobue, Abenase, Nyamebekyere and Kotokuom. The ages of these participants  
63 ranged from seven (7) to eighteen (18) years. Structured questionnaire was administered to  
64 obtain information on participants' socio-demographic characteristics, predisposing factors  
65 and knowledge on urinary schistosomiasis. Basic school children on chemotherapeutic  
66 agents for at most one week prior to the day of sampling and those below seven years and  
67 above eighteen years were excluded from the study.

### 68 **2.2 Sample collection and processing**

## 69 2.2.1 Urine specimen collection

70 Each participant was given a well labeled sterile urine container to collect random urine  
71 sample, these samples were received on the same day between 12 noon and 2 pm. During  
72 the urine collection, the participants were directed to empty their bladder completely into the  
73 container after some minimal exercise. After collection, the urine samples were assembled  
74 and kept in cold boxes with ice packs and transported to the laboratory attached to the  
75 Ayirebi Health Centre for immediate parasitological investigation.

## 76 2.2.2 Sample analysis

77 All the urine samples collected were taken through microscopy.

### 78 2.2.2.1 Microscopy

79 10ml of each urine sample was measured with a graduated measuring cylinder into a test  
80 tube and centrifuged at 1000rpm for 5 minutes to obtain the sediment and the supernatant.  
81 The supernatant was poured away and the sediment was re-suspended. A drop of the re-  
82 suspended sediment was then placed on a clean glass slide, cover slipped and examined  
83 using x10 and later x40 objective lenses of light microscope (Olympus American  
84 Clinical/Education CX® light microscope) to detect the presence or absence of *Schistosoma*  
85 *haematobium* egg.

### 86 2.2.2.2 Egg counting

87 Egg count was performed on each urine sample containing *Schistosoma haematobium* ova  
88 to rank the intensity of infection. Urinary schistosomiasis infection was categorized as light or  
89 heavy depending on the number of *Schistosoma haematobium* ova per 10ml of urine. Light  
90 infection represented less than fifty (50) *Schistosoma haematobium* ova per 10ml of urine  
91 while heavy infection meant more than fifty (50) *Schistosoma haematobium* ova per 10ml of  
92 urine.

## 93 2.3 Statistical analysis

94 Data collected was captured into Microsoft excel 2010 and analyzed using SPSS version  
95 22.0 for statistical significance. Crude odds ratios, 95% confidence intervals and p-values  
96 were calculated using univariate logistic regression to describe the relationship between  
97 dependent and independent variables where p-value <0.05 was regarded as statistical  
98 significant.

99

## 100 3. RESULTS AND DISCUSSION

101

### 102 3.1 RESULTS

103

104 **Table 1** below shows the general characteristics of the study population. A total of 504  
105 participants comprising 343 (68.1%) males and 161 (31.9%) females were included in the  
106 study. Majority, 264 (52.4%) of the respondents were aged 11-14 years. Among the  
107 respondents, 321 (63.7%) indicated that borehole was their source of drinking water. Most  
108 311 (61.7%) of the respondents stayed by a river/stream. Out of the 504 respondents  
109 interviewed, 416 (82.5%) played with a waterbody where they lived and 495 (98.2%) swam  
110 in the waterbody (river/stream). Majority 476 (94.4%) indicated that they do not urinate in the

111 river/stream. Also, 343 (68.1%) indicated that they were knowledgeable on SH infection. Out  
 112 of the 504 respondents, 141 (28%) had ever suffered from SH infection.

113 **Table 1: General Characteristics of Study Population**

| <b>Variable</b>                      | <b>Male<br/>(n = 343)</b> | <b>Female<br/>(n = 161)</b> | <b>Total<br/>(n = 504)</b> |
|--------------------------------------|---------------------------|-----------------------------|----------------------------|
| <b>Age range</b>                     |                           |                             |                            |
| 7-10                                 | 117(60.9)                 | 75(39.1)                    | 192(38.1)                  |
| 11-14                                | 197(74.6)                 | 67(25.4)                    | 264(52.4)                  |
| 15-18                                | 29(60.4)                  | 19(39.6)                    | 48(9.5)                    |
| <b>Source of Drinking Water</b>      |                           |                             |                            |
| Pipe Borne Water                     | 1(33.3)                   | 2(66.7)                     | 3(0.6)                     |
| Well                                 | 48(55.8)                  | 38(44.2)                    | 86(17.1)                   |
| River/Stream                         | 64(68.1)                  | 30(31.9)                    | 94(18.7)                   |
| Borehole                             | 230(71.7)                 | 91(28.3)                    | 321(63.7)                  |
| <b>Staying around River/Stream</b>   |                           |                             |                            |
| Yes                                  | 208(66.9))                | 103(33.1)                   | 311(61.7)                  |
| No                                   | 135(70.0)                 | 58(30.0)                    | 193(38.3)                  |
| <b>Playing at River/Stream Bank</b>  |                           |                             |                            |
| Yes                                  | 297(71.4)                 | 119(28.6)                   | 416(82.5)                  |
| No                                   | 46(52.3)                  | 42(47.7)                    | 88(17.5)                   |
| <b>Swimming in River/Stream</b>      |                           |                             |                            |
| Yes                                  | 339(68.5)                 | 156(31.5)                   | 495(98.2)                  |
| No                                   | 4(44.4)                   | 5(55.6)                     | 9(1.8)                     |
| <b>Urinating in River/Stream</b>     |                           |                             |                            |
| Yes                                  | 19(67.9)                  | 9(32.1)                     | 28(5.6)                    |
| No                                   | 324(68.1)                 | 152(31.9)                   | 476(94.4)                  |
| <b>Knowledge of Schistosomiasis</b>  |                           |                             |                            |
| Yes                                  | 253(73.8)                 | 90(26.2)                    | 343(68.1)                  |
| No                                   | 90(55.9)                  | 71(44.1)                    | 161(31.9)                  |
| <b>Ever Suffered Schistosomiasis</b> |                           |                             |                            |
| Yes                                  | 126(89.4)                 | 15(10.6)                    | 141(28.0)                  |
| No                                   | 217(59.8)                 | 146(40.2)                   | 363(72.0)                  |

114

115 **Table 2** shows the prevalence of urinary schistosomiasis stratified by demographic data. Out  
 116 of the 504 participants, 52 representing 10.3% tested positive for urinary schistosomiasis. A  
 117 higher prevalence (12.2%) of SH was found among males. Those who used river/stream as  
 118 their major source of drinking water had a highest prevalence (11.7%) while those who

119 drank pipe borne water had no prevalence. A higher prevalence, (13.5), (12.0%), (10.5%)  
 120 and (10.7%) of SH was found among those stayed in river/stream, played in river/stream,  
 121 swam in river/stream and did not urinate in river/stream respectively. The highest prevalence  
 122 (26.3%) of SH was recorded in Kotokoum while no prevalence was found among  
 123 inhabitants of Adubiase. A greater prevalence, (11.4%) was also found among participants  
 124 who had knowledge of schistosomiasis. Gender, staying around river/ stream, playing at  
 125 river/stream and inhabitant of school children were significantly associated with the  
 126 prevalence of *Schistosoma haematobium* infection. Females had 53% reduced odds of SH  
 127 infections compared to their male counterparts [OR=0.47; 95% CI= 0.23-0.97; p=0.042].  
 128 Those who did not stay around water bodies had 65% reduced odds of SH infections  
 129 compared to those who did [OR=0.35; 95% CI= 0.17-0.72; p=0.004]. Furthermore, those  
 130 who did not play around water bodies had 83% reduced odds of SH infections compared to  
 131 those who did [OR=0.17; 95% CI= 0.04-0.71; p=0.015]. Participants who lived in Kotokuom  
 132 had 8.54 times increased odds of SH infections compared to those in Pawudu [OR=8.54;  
 133 95% CI= 1.91-38.27; p=0.005].

134 **Table 2: Prevalence of SH Infection Stratified by Demographic Data**

| <b>Variable</b>                     | <b>Positive<br/>(n = 52)</b> | <b>Total<br/>(n = 504)</b> | <b>OR (95% CI)</b> | <b>p-value</b> |
|-------------------------------------|------------------------------|----------------------------|--------------------|----------------|
| <b>Age range</b>                    |                              |                            |                    |                |
| 7-10                                | 17(8.9)                      | 192(38.1)                  | 1                  |                |
| 11-14                               | 32(12.1)                     | 264(52.4)                  | 1.42 (0.76 – 2.64) | 0.268          |
| 15-18                               | 3(6.2)                       | 48(9.5)                    | 0.68 (0.19 – 2.44) | 0.561          |
| <b>Gender</b>                       |                              |                            |                    |                |
| Male                                | 42(12.2)                     | 343(68.1)                  | 1                  |                |
| Female                              | 10(6.2)                      | 161(31.9)                  | 0.47 (0.23 – 0.97) | <b>0.042</b>   |
| <b>Source of Drinking Water</b>     |                              |                            |                    |                |
| Pipe Borne Water                    | 0(0.0)                       | 3(0.6)                     | 1                  |                |
| Well                                | 8(9.3)                       | 86(17.1)                   | 0.31 (0.03 – 3.32) | 0.331          |
| River/Stream                        | 11(11.7)                     | 94(18.7)                   | 0.40 (0.04 – 4.16) | 0.442          |
| Borehole                            | 33(10.3)                     | 321(63.7)                  | 0.34 (0.03 – 3.40) | 0.361          |
| <b>Staying around River/Stream</b>  |                              |                            |                    |                |
| Yes                                 | 42(13.5)                     | 311(61.7)                  | 1                  |                |
| No                                  | 10(5.2)                      | 193(38.3)                  | 0.35 (0.17 – 0.72) | <b>0.004</b>   |
| <b>Playing at River/Stream Bank</b> |                              |                            |                    |                |
| Yes                                 | 50(12.0)                     | 416(82.5)                  | 1                  |                |
| No                                  | 2(2.3)                       | 88(17.5)                   | 0.17 (0.04 – 0.71) | <b>0.015</b>   |

**Swimming in River/Stream**

|     |          |           |                    |       |
|-----|----------|-----------|--------------------|-------|
| Yes | 52(10.5) | 495(98.2) | 1                  |       |
| No  | 0(0.0)   | 9(1.8)    | 0.94 (0.12 – 7.62) | 0.959 |

**Urinating in River/Stream**

|     |          |           |                     |       |
|-----|----------|-----------|---------------------|-------|
| Yes | 1(3.6)   | 28(5.6)   | 1                   |       |
| No  | 51(10.7) | 476(94.4) | 3.24 (0.43 – 24.35) | 0.253 |

**Knowledge of Schistosomiasis**

|     |          |           |                    |       |
|-----|----------|-----------|--------------------|-------|
| Yes | 39(11.4) | 343(68.1) | 1                  |       |
| No  | 13(8.1)  | 161(31.9) | 0.68 (0.35 – 1.32) | 0.259 |

**Communities**

|              |           |            |                     |              |
|--------------|-----------|------------|---------------------|--------------|
| Pawudu       | 2 (4.0)   | 50 (9.92)  | 1                   |              |
| Adubiase     | 0 (0.0)   | 50 (9.9)   | 0.48 (0.04 – 5.47)  | 0.554        |
| Adjobue      | 7 (5.6)   | 126 (25.0) | 1.41 (0.28 – 7.04)  | 0.674        |
| Abenase      | 13 (9.8)  | 133 (25.0) | 2.60 (0.57 – 11.96) | 0.220        |
| Nyamebekyere | 9 (13.9)  | 65 (12.90) | 3.86 (0.79 – 18.72) | 0.094        |
| Kotokuom     | 21 (26.3) | 80 (15.87) | 8.54 (1.91 – 38.27) | <b>0.005</b> |

135

136 **Table 3** shows the intensity of SH infection stratified by Demographic data. In relation to the  
 137 52 (10.3%) participants who tested positive for SH, 36 (69.2%) had light infection and 16  
 138 (30.8%) had heavy infection. A higher prevalence, (33.3%) and (37.5%) of heavy infection  
 139 was found among males and 11-14 year old participants correspondingly while there was  
 140 no prevalence of heavy infection among 15-18 year olds. A greater prevalence, (33.3%),  
 141 (35.7%), (32.0%), (30.8%) and (31.4%) of heavy intensity SH was found among used  
 142 borehole as source of drinking water, stayed around river/stream, played in river/stream,  
 143 swam in river/stream and did not urinate in river/stream respectively. The topmost  
 144 prevalence (42.9%) of heavy intensity SH was recorded in Adjobue while no prevalence  
 145 was found among inhabitants of Pawuda. A greater prevalence, (38.5%) of heavy intensity  
 146 SH was found among participants who had knowledge of schistosomiasis while a higher  
 147 prevalence (43.3%) was also found among those who had suffered from the infection before.

148 **Table 3: Intensity of SH Infection Stratified by Demographic Data**

| <b>Variable</b>  | <b>Light<br/>(n = 36)</b> | <b>Heavy<br/>(n = 16)</b> | <b>Total<br/>(n = 52)</b> |
|------------------|---------------------------|---------------------------|---------------------------|
| <b>Age range</b> |                           |                           |                           |
| 7-10             | 13(76.5)                  | 4(23.5)                   | 17(32.7)                  |
| 11-14            | 20(62.5)                  | 12(37.5)                  | 32(61.5)                  |

|                                      |           |          |           |
|--------------------------------------|-----------|----------|-----------|
| 15-18                                | 3(100.0)  | 0(0.0)   | 3(5.8)    |
| <b>Gender</b>                        |           |          |           |
| Male                                 | 28(66.7)  | 14(33.3) | 42(80.8)  |
| Female                               | 8(80.0)   | 2(20.0)  | 10(19.2)  |
| <b>Source of Drinking Water</b>      |           |          |           |
| Well                                 | 6(75.0)   | 2(25.0)  | 8(15.4)   |
| River/Stream                         | 8(72.7)   | 3(27.3)  | 11(21.2)  |
| Borehole                             | 22(66.7)  | 11(33.3) | 33(63.5)  |
| <b>Staying around River/Stream</b>   |           |          |           |
| Yes                                  | 27(64.3)  | 15(35.7) | 42(80.8)  |
| No                                   | 9(90.0)   | 1(10.0)  | 10(19.2)  |
| <b>Playing at River/Stream Bank</b>  |           |          |           |
| Yes                                  | 34(68.0)  | 16(32.0) | 50(96.2)  |
| No                                   | 2(100.0)  | 0(0.0)   | 2(3.8)    |
| <b>Swimming in River/Stream</b>      |           |          |           |
| Yes                                  | 36(69.2)  | 16(30.8) | 52(100.0) |
| No                                   | 0(0.0)    | 0(0.0)   | 0(0.0)    |
| <b>Urinating in River/Stream</b>     |           |          |           |
| Yes                                  | 1(100.0)  | 0(0.0)   | 1(1.9)    |
| No                                   | 35(68.6)  | 16(31.4) | 51(52.6)  |
| <b>Knowledge of Schistosomiasis</b>  |           |          |           |
| Yes                                  | 24(61.5)  | 15(38.5) | 39(75.0)  |
| No                                   | 12(92.3)  | 1(7.7)   | 13(25.0)  |
| <b>Ever Suffered Schistosomiasis</b> |           |          |           |
| Yes                                  | 17(56.7)  | 13(43.3) | 30(57.7)  |
| No                                   | 19(86.4)  | 3(13.6)  | 22(42.3)  |
| <b>Communities</b>                   |           |          |           |
| Pawudu                               | 2 (100.0) | 0 (0.0)  | 2 (3.8)   |
| Adubiase                             | 0 (0.0)   | 0 (0.0)  | 0 (0.0)   |
| Adjobue                              | 4 (57.1)  | 3 (42.9) | 7(13.5)   |
| Abenase                              | 10 (76.9) | 3 (23.1) | 13 (25.0) |

|              |           |          |           |
|--------------|-----------|----------|-----------|
| Nyamebekyere | 7 (77.8)  | 2 (22.2) | 9 (17.3)  |
| Kotokuom     | 13 (61.9) | 8 (38.1) | 21 (40.4) |

149

150

151 **3.2 DISCUSSION**

152 The overall prevalence of urinary schistosomiasis was found to be 10.32% [95% CI: 7.80 -  
 153 13.31%]. This prevalence is slightly higher compared to that reported in other studies; (8.6%)  
 154 in Central Sudan and (7.7%) in Burkina Faso (Ahmed, Abbas, Mansour, Gasim, & Adam,  
 155 2012; Touré, et al., 2008). On the contrary, prevalence rate reported in this study was lower  
 156 than that reported in other similar studies (Nkenazong, Nijioku, & Asonganyi, 2013; Augusto,  
 157 Magnussen, Kristensen, Appleton, & Vennervald, 2009). A study by Augusto et al., (2009) in  
 158 Mozambique reported a prevalence of 51.7%. Also, a recent study by Nkenazong, Nijioku, &  
 159 Asonganyi, (2013), in Barombi Focus, South West Cameroon, reported a prevalence of  
 160 13.7%. The inconsistency in the above prevalence rates can be attributed to differences in  
 161 study periods, sample sizes, participants recruited for the study and geographical location.

162

163 Additionally, the present study found a statistically significant association between gender  
 164 and the prevalence of *Schistosoma haematobium* infection. Further, the prevalence of SH  
 165 infection and the intensity of heavy SH infection were higher in males than in females.  
 166 Collaborating the findings of this study with other studies, our study was parallel to studies  
 167 conducted by Benwat et al., (2012) in Langai plateau. Similarly, the high prevalence of heavy  
 168 SH infection common among males than females in our study is similar to the observation  
 169 made by Nkenazong, Nijioku, & Asonganyi, (2013), in South-West Cameroon. Activities such  
 170 as swimming and playing in water bodies which bring people in contact with water could be a  
 171 possible cause of the variation in the prevalence of SH infection and the intensity of SH  
 172 infection among males and females. In the transmission of urinary schistosomiasis, the risk  
 173 of infection varies directly with frequency of contact with contaminated water (Jordan &  
 174 Webbe, 1999). Secondary to the frequency of contact, the time of the day, duration of  
 175 exposure and the area of the skin exposed also account for the risk of infection (Jordan and  
 176 Webbe, 1999). Generally, more males often engage in swimming than females which leads  
 177 to the higher prevalence of SH infection in males. Females mostly fetch water from the rivers  
 178 and streams for washing and cooking at home. Hence, the females have less contact to  
 179 water than males and this accounted for the low prevalence of SH infection in females. As  
 180 proposed by Remoue *et al.*, (2001), females have stronger immune response to  
 181 schistosomiasis infection than males because the females are known to produce more  
 182 specific IgA, TGF- $\beta$  and Interleukin-10 (IL-10) to the infection. Hence, there is higher  
 183 intensity of heavy infection in males than in females.

184

185 In relation to the 52 positive SH ova cases, 16 (30.8%) were heavy SH infections and 36  
 186 (69.2%) were light infections. Hence, light infection was more prevalent than heavy infection.  
 187 The higher prevalence of light infection is in conformity with an earlier study conducted which  
 188 investigated the prevalence of urinary schistosomiasis among Nigerien school children  
 189 (Tohon, et al., 2008). The low prevalence of heavy infection is important because the  
 190 severity of morbidity caused by urinary schistosomiasis is directly related to the intensity of  
 191 infection. Therefore, the heavy infection causes more severe morbidity and the light infection  
 192 causes less severe morbidity (King and Dangerfield-Cha, 2008).

193



194 Knowledge of schistosomiasis ( $P=0.037$ ) and having suffered from schistosomiasis infection  
195 in the past ( $P= 0.022$ ) showed significant association with the intensity of SH infection. The  
196 age group 11-14 years had the highest prevalence of SH infection followed by 7-10 years  
197 and 15-18 years. The intensity of heavy infection was highest among the age group 11-14  
198 years whilst the intensity of light SH infection was predominantly high among the age group  
199 15-18. The findings in our study in relation to the prevalence and intensity of SH infection  
200 among the age groups of the participants is in keeping with studies conducted among basic  
201 school children in Kigogo administrative ward of the Kinondoni district of Dare-es-Salam  
202 (Ndyomygyenyi *et al.*, 2001) and two peri-urban villages in the South- Western state of  
203 Osun, Nigeria (Ugbomoiko *et al.*, 2010) where the participants aged 10-14 years had the  
204 highest prevalence and intensities of infection than those in the younger or older age groups  
205 studied. These findings, may be attributed to the swimming or playing behavior of  
206 participants aged 11-14 years in infested river or stream, resulting in the high urinary  
207 schistosomiasis infections recorded among these subjects. Also, lower immunity at age 11-  
208 14 years may be another contributing factor to their susceptibility. The prevalence of  
209 infection among the age group 15-18 years could have resulted from prolong exposure to  
210 infested water bodies.

211

212 The study also found a significant statistical association between age distribution, source of  
213 drinking water, staying around river or stream, playing at river or stream bank, swimming in  
214 river or stream, knowledge of schistosomiasis and SH ova with the communities in which the  
215 study was carried out. The prevalence of SH ova or urinary schistosomiasis was 26.3% in  
216 Kotokuom, 13.8% in Nyamebikyere, 9.8% in Abenase, 5.6% in Adjobue and 4% in Pawudu.  
217 No participant from Adubiase tested positive to SH ova. Hence, Kotokuom recorded the  
218 highest prevalence of urinary schistosomiasis in the district with zero prevalence in  
219 Adubiase. These findings are in conformity with a study carried out in the Ga district by  
220 Aboagye and Edoh (2009), which revealed a high prevalence rate of SH infection in Mahem  
221 (58%) which is closer to the Weija dam than a prevalence rate of 49% in Galilea. Also, the  
222 present study is in agreement with a study carried out in Kumasi comparing the prevalence  
223 of urinary schistosomiasis in Kumasi South hospital (surrounded by Atonsu, Gyinyase and  
224 Ahinsan communities) and Animwa Medical Centre (surrounded by Boadi, Ayeduase,  
225 Emena, Appiadu and Kokoben communities) which revealed prevalence of 40.2% and 4.5%  
226 respectively (Tay, Amankwa, & Gbedema, 2011). The variations in the prevalence rates  
227 within the communities could be due to differences in the number of freshwater bodies  
228 infested with the snail intermediate host (*Bulinus* species) of the parasite as well as the  
229 behavioural patterns of the communities (Tay, Amankwa, & Gbedema, 2011). The highest  
230 prevalence of urinary schistosomiasis in Kotokuom may be due to unsanitary conditions,  
231 unavailability of clean household water, proximity of the town to infested water bodies,  
232 ignorance of the knowledge of schistosomiasis and anthropological activities such as  
233 swimming, fishing, fetching and washing in water infested with the snail intermediate hosts of  
234 the parasite. The zero prevalence in Adubiase may be attributed to the non-availability of the  
235 intermediate water snail of the parasite secondary to good sanitary conditions in the  
236 community. The prevalence of 13.8% observed in Nyamebikyere may be as results of low  
237 standard of living and over reliance on river or stream as source of drinking water. Although  
238 the Adjobue community is traversed by a river suspected to be infested with the intermediate  
239 water snail of the parasite, the annual initiative community outreach programme undertaken  
240 by the University of Cape Coast School of Medical Sciences (UCCSMS) to educate  
241 inhabitants about diseases of public health interest in rural areas including Adjobue  
242 community may have contributed to the low prevalence (5.6%) of urinary schistosomiasis  
243 recorded in Adjobue. The prevalence of 9.8% recorded in Abenase is secondary to  
244 predominance of agricultural activities in the community whereas the prevalence of 4% in  
245 Pawudu may be due to scarcity of the intermediate water snail for transmission.

246

247 **CONCLUSION AND RECOMMENDATIONS**

248

249 The prevalence of urinary schistosomiasis among basic school children in the Akyemansa  
250 district was found to be 10.3% [95% CI: 7.80 -13.31%]. Gender, staying around river/  
251 stream, playing at river/ stream and inhabitant of participants were significantly associated  
252 with the prevalence of *Schistosoma haematobium* infection. Periodic drug administration is  
253 recommended and a comprehensive intervention strategy should be designed and  
254 implemented to reduce schistosomiasis prevalence.

255

256 **LIMITATION**

257

258 Real-time PCR assay approach would have been more sensitive in detecting *Schistosoma*  
259 *haematobium* ova compared to the light microscopy which was the diagnostic technique  
260 employed in the study.

261

262

263 **COMPETING INTERESTS**

264

265 The authors hereby declare that there are no competing interests regarding the publication  
266 of this article.

267

268

269 **CONSENT (WHERE EVER APPLICABLE)**

270

271 Written informed consent was obtained from participants before the study took place and  
272 these documents are preserved by the authors.

273

274

275 **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

276

277 Ethical clearance was sought from the management of Ayirebi Health Centre before the  
278 study was commenced. In addition, study participants gave informed consent after thorough  
279 explanation of the rationale of the study has been given.

280

281 **REFERENCES**

282

283 1. Ahmed, A., Abbas, H., Mansour, F., Gasim, G., & Adam, I. (2012). *Schistosoma*  
284 *haematobium* infections among school children in central Sudan one year after treatment  
285 with praziquantel. *Parasite and vectors*, 5, 108.

286

287 2. Aryeetey, M. E., Wagatsuma, Y., Yeboah, G., Asante, M., Mensah, G., Nkrumah, F. K.,  
288 & Kojima, S. (2000). Urinary schistosomiasis in southern Ghana: Prevalence and morbidity  
289 assessment in three (defined) rural areas drained by the Densu river. *Parasitol Parasitology*  
290 *International*, 49(2), 155–163. doi:doi:10.1016/s1383-5769(00)00044-1

291

292 3. Benwat, M., Ogbonna, C., Daboer, J., Chingle, M., Envuladu, E., Audu, S., & Lar, L.  
293 (2012). Prevalence of urinary schistosomiasis in school-aged children in Langai Plateau  
294 state: Pre and Post- intervention. *Niger J Med.*, 21(2).

295

- 296 4. Bruun, B., & Aagaard-Hansen, J. (2008). The social context of schistosomiasis and its  
297 control. . *Geneva: WHO*.  
298
- 299 5. CDC. (2015, February 12). Retrieved from The burden of schistosomiasis.: Retrieved  
300 from CDC: [http://www.cdc.gov/globalhealth/ntd/diseases/schisto\\_burden.html](http://www.cdc.gov/globalhealth/ntd/diseases/schisto_burden.html).  
301  
302
- 303 6. Gryseels, B., Polman, K., Clerinx, J., & Kestens, L. (2006). *The Lancet*, 368 (no. 9541,  
304 8), pp. 1106–111.  
305
- 306 7. Jordan, P., Webbe, G., & Sturrock, R. (1999). *Epidemiology in Human schistosomiasis* .  
307 Wallingford, UK: CAB International.  
308  
309
- 310 8. King, C., & Dangerfield-Cha, M. (2008). The unacknowledged impact of schistosomiasis.  
311 *Chronic Illn*, 4, 65-79.  
312
- 313 9. Ndyomugenyi, R., & Minjas, J. (2001). Urinary Schistosomiasis in school children in  
314 Dar-es- Salam, Tanzania, and the factors influencing its transmission. *Ann Trop Med*  
315 *Parasitol*, 95(7), 697-706.  
316
- 317 10. Nkenazong, L., Nijioku, F., & Asonganyi, T. (2013). Two years impact of single  
318 Praziquantel treatment on urinary schistosomiasis in the Barombi Kotto Focus, South West  
319 Cameroon. *J. Parasitol. Vector Biol.*, 5(6), 83-89.  
320
- 321 11. Remoue, F., To Van, D., Schacht, A. M., Picquet, M., Garraud, O., Vercruysse, J., . . .  
322 Riveau, G. (2013). Gender-dependent specific immune response during chronic human  
323 Schistosomiasis haematobia. *Clin Exp Immunol*, 124, 62-68.  
324
- 325 12. Stephenson, L. (1993). The impact of schistosomiasis on human nutrition. *Parasitology*,  
326 107, 107–23.  
327
- 328 13. Tay, S. C., Amankwa, R., & Gbedema, S. Y. (2011). Prevalence of Schistosoma  
329 Haematobium infection in Ghana: a retrospective case study in Kumasi. *Int J Parasitol Res*,  
330 3(2), 48-52.  
331
- 332 14. Thétiot-Laurent, S., J., B., Robert, A., & Meunier, B. (2013). Schistosomiasis  
333 Chemotherapy. *Angew Chem Int Ed Engl*, 52(31), 7936-56. doi:doi:  
334 10.1002/anie.201208390  
335
- 336 15. Tohon, Z., Mainassara, H., Garba, A., Ibrahim, M. L., Mahamane, A., Bosqué- Oliva, E.,  
337 & al., e. (2008). Controlling schistosomiasis: Significant decrease of anaemia prevalence  
338 one year after a single dose of Praziquantel in Nigerien School children.  
339
- 340 16. Touré, S., Zhang, Y., Bosqué-Oliva, E., Ouedraogo, A., Koukounari, A., Gabrielli, A., &  
341 et al. (2008). Two-year impact of single praziquantel treatment on infection in the national  
342 control programme on schistosomiasis in Burkina Faso. *Bull World organ*, 86, 780-787.  
343
- 344 17. Ugbomoiko, U., Ofoezie, I., Okoye, I., & Heukelbach, J. (2010). Factors associated with  
345 urinary schistosomiasis in two peri-urban communities in South- western Nigeria . *Ann Trop*  
346 *Med Parasitol.*, 104(5), 409-19.van der Werf, M., de Vlas, S. B., Looman, C., Nagelkerke, N.,

347 & Habbema, J. (2003). Quantification of clinical morbidity associated with schistosome  
348 infection in sub-Saharan Africa. *Acta Trop.*, 86(2–3), 125–39.

349

350 18. Augusto, G., Magnussen, P., Kristensen, T., Appleton, C., & Vennervald, B. (2009). The  
351 influence of transmission season on parasitological cure rates and intensity of infection after  
352 praziquantel treatment of *Schistosoma haematobium*-infected School children in  
353 Mozambique. *Parasitology*, 136, 1771-1779.