

**EFFECTS OF ORGANIC AND INORGANIC FERTILIZER ON THE EARLY
GROWTH RESPONSE OF *Azelia Africana***

ABSTRACT

An experiment was conducted in the nursery of Department of Forestry Technology at the Federal College of Forestry Ibadan, Oyo state, Nigeria to determine the effect of organic and inorganic fertilizers on early growth response of *Azelia africana*. Seedlings were collected from Forestry Research Institute of Nigeria, Ibadan, Oyo State and were transplanted into 2kg perforated polythene pots. The experiment was laid out in a completely randomized design with five treatments replicated three times. The treatments were: 20 t/ha water hyacinth compost (T1), 20 t/ha water hyacinth + poultry manure compost (T2), 20 t/ha poultry manure (T3) and 50 kg/ha N:P:K 20:10:10 (T4) and T5 - control (no fertilizer application). The compost was applied two (2) weeks before planting while NPK 20:10:10 was applied two 2 weeks after planting. The experiment was monitored for eight (8) weeks after transplanting (WAT) while growth parameters were measured. The results of the study showed that application of fertilizers gave significant ($p=0.05$) increase in plant height (cm), stem diameter (mm), leaf production, and leaf area (cm^2) of *A. africana*. Plant height ranged from 41.43 cm in the control to 47.96 cm in the pots where 20 t/ha water hyacinth compost was applied. Stem diameter also increased appreciably across treatments, while leaf production ranged from 9 in the control treatment to 14 in the pots with 20 t/ha poultry manure. This result suggested that the incorporation of organic and inorganic fertilizers increased productivity of *Azelia africana* and hence application of 20 t/ha water hyacinth compost may be recommended for the production of *Azelia africana* especially in the study area.

Keywords: Water hyacinth compost; *Azelia africana*; NPK 20:10:10

27

28

INTRODUCTION

29 Forests and trees perform various functions in the ecosystem namely; aesthetics, provision of
30 food and medicine, provision of shelter to wildlife and hygienic purpose (Agbogidi and
31 Eshegbeyi, 2008). It is universally accepted that forests and trees carry out a fundamental role
32 in soil and water resources conservation (Broadhead and Leslie, 2007; Hamilton, 2008). As
33 population density increases and land for food production expands due to agricultural
34 activities and urbanization, natural forests became degraded (Salim and Ullsten, 1999). The
35 degradation has led to the disappearance of most species including agro forest trees and
36 causing difficulty in growing some seedlings including *Afzelia africana* (Nwoboshi, 1985;
37 Keay, 1989; Etukudo, 2000; Ezenwaka *et al.*, 2004). *Afzelia africana* is a leguminous tree
38 found in the humid and dry forest savannah borders or semi-deciduous forest (Keay, 1985). It
39 is used for soil conservation and improvement (Agbogidi and Onomeregbor, 2007). *Afzelia*
40 *africana* is a timber species with high forage, economic and pharmacological values. Its
41 leaves are harvested for grazing during the dry season. The high demand for *A. africana*
42 leaves, seeds, roots and barks for various uses has resulted in corresponding increase in the
43 exploitation at such a rate that sustainability of this natural resource cannot be guaranteed
44 (Mtambalika *et al.*, 2014; Palgreave, 2002). Documented reports on the cultivation and
45 seedling growth of this multipurpose tree known commonly as African mahogany are scarce
46 (Okeke, 1996; Burkill, 1999; Etukudo, 2000; Agbogidi *et al.*, 2008). If the benefits derivable
47 from *A. africana* must continue especially for the future generations, there is the need to
48 stimulate farmers' interest in the cultivation of *A. africana* thereby helping to reduce poverty,
49 helping in conservation role as well as to boost the source of revenue for the government. In
50 the same vein, information on the domestication of the plant seeds and seedlings are in piece
51 meal due mainly to the poor nature of the soil caused by human influences. Consequently, the

52 need to acquaint farmers with the most successful soil or manures that could enhance the
53 growth of *A. africana* seedlings cannot be overemphasized. This study was aimed at
54 investigating the effect of organic and inorganic fertilizers on the seedling establishment of
55 *Azelia africana* with a view to recommend the best fertilizer to *A. africana* growers
56 especially at the nursery stage and to multiply this multi-purpose species that nature has
57 bequeathed to mankind.

58 MATERIALS AND METHODS

59 The experiment was carried out in 2018 at nursery site of the Department of Forestry
60 Technology, Federal College of Forestry Jericho Ibadan (Latitude 07⁰ 27¹N and longitude 03⁰
61 53¹E), Ibadan, Nigeria (FRIN Meteorological Station, 2018). The annual rainfall is 1250 mm
62 with a bimodal pattern and has a minimum temperature of 21.9 °C and maximum temperature
63 of 35.5 °C. The experiment was laid out in a Complete Randomized Design with five
64 treatments replicated three times. The treatments were: 20 t/ha water hyacinth compost (T1),
65 20 t/ha water hyacinth + poultry manure compost (T2), 20 t/ha poultry manure (T3) and 50
66 kg/ha N:P:K 20:10:10 (T4) and T5 - control (no fertilizer application). The compost was
67 applied two (2) weeks before planting. Eight week old seedlings of *A. africana* were
68 collected from Forestry Research Institute of Nigeria (FRIN), Ibadan, Oyo State, and potted
69 into 2kg polythene pots filled with different media treatments, watered and allowed to
70 stabilize for two weeks before the commencement of growth assessment. Plant height was
71 measured with a meter rule at the distance from soil level to terminal bud. Leaf production
72 was determined by counting. Stem diameter at the collar was measured with venier caliper.
73 Data collected were analysed statistically using Genstat Software Package and were subjected
74 to analysis of variance. Means were separated using Duncan's multiple range test (DMRT) at
75 5% level of significance.

76

77 **RESULTS**

78 Table 1: Pre-planting soil physical and chemical properties of the experimental site

Soil parameters	Content in soil
pH (H ₂ O)	5.8
Organic carbon (g kg ⁻¹)	9.0
Total nitrogen (g kg ⁻¹)	1.0
Available phosphorus(mg kg ⁻¹)	6.0
Exchangeable cations (cmol kg ⁻¹)	
Ca	1.32
Mg	0.25
K	0.11
Na	0.31
Extractable micronutrient (mg kg ⁻¹)	
Mn	302.0
Fe	265.0
Cu	3.39
Zn	1.2
Exchangeable Acidity (cmol kg ⁻¹)	0.40
Particle size distribution (g kg ⁻¹)	
Sand	838
Silt	54
Clay	108
Textural class	Sandy loam
Bulk density (g cm ⁻¹)	1.62
Saturated hydraulic conductivity (cm hr ⁻¹)	12.4

79

80

81

82

83

84 **Table 2: Chemical properties of poultry manure, water hyacinth compost and water**
 85 **hyacinth + poultry manure compost used.**


Parameter	Poultry manure	Water hyacinth compost	Water hyacinth + Poultry manure compost
pH (H ₂ O)	6.8	5.89	5.76
Organic carbon (%)	22.94	31.92	32.11
Total Nitrogen (%)	3.30	2.6	1.73
C:N	6.8	12.52	43.99
Phosphorus (%)	0.83	1.24	1.55
Potassium (%)	1.80	0.47	0.80
Calcium (%)	2.56	1.6	1.80
Magnesium (%)	1.58	4.30	0.92

86

87 The physical and chemical properties of the soil (0 – 15 cm depth) at the experimental site
 88 before planting is as presented in Table 1. The soil is sandy loam, slightly acidic, high bulk
 89 density (1.62 g/cm³) and has been classified as an Alfisol (Smyth and Montgomery, 1962)
 90 with its distinctive characteristics. The data in Table 1 further confirms this assertion and also
 91 reveals that the soils are moderate in zinc, low in potassium (0.11 cmol/kg), organic carbon
 92 (9.0 g/kg), total nitrogen (1.0 g/kg) and phosphorous (6.0 mg/kg). Saturated hydraulic
 93 conductivity value of 12.4 cm hr⁻¹ indicated a well drained soil.

94 The chemical composition of water hyacinth compost, poultry manure and water
 95 hyacinth + poultry manure compost used is as presented in Table 2. The pH of all the organic
 96 fertilizers were all acidic with the water hyacinth + poultry manure compost having the most
 97 acidic pH of 5.76. Organic carbon, C/N and phosphorus were highest in the mixture of water


98 hyacinth + poultry manure compost. Poultry manure was highest in nitrogen, potassium and
 99 calcium content



100

101 Figure 1: Growth pattern of number of leaves of *Afzelia africana* seedlings as influenced by
 102 fertilizers

103



104

105 Figure 2: Effects of fertilizers on number of leaves of *Afzelia africana* seedlings

106


107

108

109 **Effect of organic and inorganic fertilizers on number of leaves of *Azelia africana***
110 **seedling**

111 Figure 1 showed the growth pattern of leaf production of *Azelia africana* seedlings as
112 influenced by the application of organic and inorganic fertilizers. Results revealed that mean
113 leaf production of *Azelia Africana* increased across the study period. T3 (20t/ha poultry
114 manure) recorded the highest mean number of leaves across the weeks while T5 (top soil
115 only) had the least mean number of leaves all through the weeks.


116 The effect of organic and inorganic fertilizers on the number of leaves of *Azelia africana* is
117 as presented in Figure 2. Leaf production increased appreciably across treatments and differs
118 significantly ($p=0.05$) throughout the experiment. Leaf production ranged from 9 in the
119 control medium to 14 in the pots where 50 kg/ha NPK 20:10:10 was applied. The highest
120 mean leaf production of *Azelia africana* was observed in T3 with 14 and it was closely
121 followed by T1, T2, and T4 with 13 and the least was found in T5 (control) with
122 9



123
124 T1 (20t/ha water hyacinth) T2 (20t/ha water hyacinth + poultry manure compost)
125 T3 (20t/ha poultry manure) T4 (50kg/ha NPK 20:10:10) T5 (control – no fertilizer application)
126
127
128

129
130
131

Figure 3: Growth pattern of height of *Azelia africana* seedlings as influenced by fertilizers



132

133 Figure 4: Effects of fertilizers on plant height of *Azelia africana* seedlings

134
135
136
137


138 **Effect of organic and inorganic fertilizers on plant height of *Azelia africana* seedling**

139 Results in Figure 3 showed the growth pattern of plant height of *Azelia africana* seedlings,
140 the control experiment followed similar trend with the growth pattern of number of leaves
141 and performed poorly in comparison to others in terms of the number of leaves produced. The
142 highest mean height of *Azelia africana* was observed in T1 (20t/ha water hyacinth compost)
143 with T2 (20t/ha water hyacinth + poultry manure compost) closely followed and T5 (control)
144 performed least. The effect of organic and inorganic fertilizers on the plant height of *Azelia*
145 *africana* is as presented in Figure 4. Plant height increased appreciably across treatments but
146 did not differ significantly ($p=0.05$) throughout the experiment. 20 t/ha water hyacinth

147 compost (47.96 cm) had highest plant height followed by 20 t/ha water hyacinth + poultry
 148 manure compost (47.89 cm) and least by control pot (41.43 cm).

149


150



151

152 Figure 5: Growth pattern of stem diameter of *Azelia africana* seedlings as influenced by
 153 fertilizers

154



155

156 Figure 6: Effects of fertilizers on stem diameter of *Azelia africana* seedlings

157 **Effect of organic and inorganic fertilizers on stem diameter of *Afzelia africana* seedling**

158 Results in figure 5 showed the growth pattern of stem diameter of *Afzelia africana* seedlings,
159 the control experiment performed well at the beginning but poorly at the end of the eighth
160 week, with T1 (20t/ha water hyacinth) performing best.

161 The effect of organic and inorganic fertilizers on the stem diameter of *Afzelia africana* is
162 presented in Figure 6. Stem diameter increased appreciably across treatments but did not
163 differ significantly ($p=0.05$) throughout the experiment. 20 t/ha water hyacinth compost (0.81
164 mm) had highest plant height followed by 20 t/ha water hyacinth + poultry manure compost
165 (0.76 mm) and least by control plot (0.62 mm).

167 **DISCUSSION**

168 The major factor affecting plant growth in the tropics is the nutrient deficiency in tropical
169 soils resulting from degraded farmland. The low levels of nitrogen, phosphorus, and organic
170 carbon observed in the experimental soil indicated that the soil had a low fertility status. The
171 value obtained for N, P, K and organic C is below the critical range (Adeoye and Agboola,
172 1985, Akinrinde, *et al.*, 2005), thus indicating poor soil fertility not suitable without the
173 addition of external input for planting *Afzelia africana*. The chemical composition of water
174 hyacinth compost, poultry manure and water hyacinth + poultry manure compost used in the
175 experiment was relatively high in major elements (N, P, K, Ca and Mg). The organic carbon
176 content of all the organic fertilizers were less than the values obtained for the composts of
177 *Azadirachta indica*, *Albizia lebbek* and *Khaya senegalensis* by Daldoum and Hammad
178 (2015).

179 The application of the various fertilizers increased the growth of *A. Africana* this agreed with
180 the findings of Uddin (2014) where organic fertilizers enhanced the seedling growth of some
181 leguminous agroforestry species. This could result from the nutritional benefits of organic

182 and inorganic fertilizers which include improvement of soil fertility. The result obtained from
183 the plant height showed that water hyacinth compost significantly induced the shoot growth,
184 leaf production and stem diameter of *A.africana* seedlings. This is in support with Razaq, *et*
185 *al.* (2017); Talkah (2015) and Cuesta (2010) that reported that plant height and number of
186 leaves of plants treated with water hyacinth compost had been used and showed showed
187 better performance than control. This result is also in line with the results by Lata (2013) that
188 experimented with water hyacinth manure on *Coriandrum sativum* and revealed positive
189 response with increase in manure rates. This was also supported by study done by Osoro, *et*
190 *al.*, (2014) and Aboul-Enein *et al.* (2011) who advocated that water hyacinth has good N, P,
191 K absorbing capacity from water and thus can be used as a good source of compost material
192 to serve as fertilizer in soil with poor amount of N, P, K and C values. Water hyacinth which
193 used to be tagged as waste and nuisance to aquatic environment can be converted to compost
194 for fertilizing plants at the nursery stage in order to improve early growth. This might replace
195 use of expensive, scarce and environmentally hazardous inorganic fertilizers in forest nursery
196 work.

197

198 **CONCLUSION**

199 Organic and inorganic fertilizer had effect on the growth of *Afzelia Africana*. It could be
200 observed from the result obtained that there were increases in plant height, stem diameter and
201 number of leaves. However, there was poor performance throughout the assessment period in
202 control treatment when compared to other treatments in terms of the number of leaves
203 produced per plant. Seedlings with 50 kg/ha N:P:K 20:10:10 (T4) performed best at week
204 eight though it was poor at the early stage of the experiment. Seedlings with 20 t/ha poultry
205 manure (T3) performed well from the beginning of the experiment up to the penultimate
206 week to the end of the experiment. Application of 20 t/ha water hyacinth compost gave the

207 highest plant height. Stem diameter also increased appreciably across treatments. Therefore,
208 it can be concluded and recommended that 20 t/ha water hyacinth compost can be used by
209 farmers to increase the growth of *Azizelia africana*.

210

211 REFERENCES

212 Aboul-Enein A, Al-Abd A, Shalaby E, Abul- Ela F, Nasr-Allah A, Mahmoud A. (2011).
213 *Eichhornia crassipes* (Mart) solms, Plant Signaling & Behavior. 2011;6(6):834-836.

214

215 Adeoye, G. O., and Agboola, A. A., (1985). Critical level for soil pH, available P, k, Zn and
216 maize ear leaf content for P, Cu and Mn in sedimentary soils of Southwestern Nigeria. 6: 65 –
217 71.

218

219 Agbogidi O. M, Onomerebor V. A (2007). Enhancing germination in seeds of *Azizelia*
220 *africana* (Sm. Expers.). In: Olufajo, OO, Omokore DF, Akpa GN Sanni SA (ed.).
221 *Proceedings of the 41st Annual Conference of the Agricultural Society of Nigeria* held in
222 Zaria, Kaduna State, between 22nd and 26th October, 2007. Pp 260-263.

223

224 Agbogidi OM, Onomerebor VA, Avwevughwar OI (2008). Eco Physiological effects of
225 crude oil pollution *Azizelia africana* (Sm. Ex pers.) In: Ezike, K.N.N., Osakwe, I.I., Ekwu,
226 L.G., Utobo, E. and Mba, C.N. (eds.). *Proceedings of the 42nd Annual Conference of the*
227 *Agricultural Society of Nigeria* held at the Ebonyi State University, Abakakili, Ebonyi state,
228 between 19th and 23rd October, 2008. Pp 365-369.

229

230 Agbogidi, O. M. and Eshegbeyi, O. F. (2008). Forestry development for a safe environment.
231 In: Onyekwelu J.C. Adekunle, V.A.J.and Oke, D.O. (ed.). *Proceedings of the First National*
232 *Conference of the Forests and Forest Products Society* held at the Federal University of
233 Technology, Akure, Ondo State between 16th and 18th of April, 2008. Pp 95-98

234

235 Akinrinde, E. A., Obigbesan, J. A., and Onanuga, O. A., (2005). Extractable micronutrient in
236 relation to physiochemical properties of soil in four ecological zones in Nigeria. *Global*
237 *Journal of Pure and Applied science* 2. (11), 197-200.

238

239 Broadhead, J., Leslie, R., 2007. Coastal protection in the aftermath of the Indian Ocean
240 tsunami: what role for forests and trees? In: Proceedings of an FAO Regional Technical
241 Workshop, Khao Lak, Thailand, 28–31 August 2006, FAO, Thailand.

242

243 Burkill HM (1999). The useful plants of West Africa. Vol. Calesal piniaceae Royal botanic
244 Gardens. Longman publishing U.K.

245

246 Cuesta B, Villar-Salvador P, Puertolas J, F. Jacobs D, M. Rey Benayas J. (2010). Why do
247 large, nitrogen rich seedlings better resist stressful transplanting conditions? A physiological
248 analysis in two functionally contrasting Mediterranean forest species. Forest ecology and
249 management [Internet].;260:71-78. Available from:
250 <http://80.24.165.149/webproduccion/PDFs/10ART 10.PDF>

251

252 Daldoum D.M.A. and Hammad G.H.(2015): Performance of *Acacia senegal* (L.) wild
253 seedlings growth under some tree manures and NPK fertilizers in nursery Site. *Journal of*
254 *Forest and Environmental Science* Vol. 31, No. 4, pp. 303-311

255

256 Etukudo I. (2000). Forests; our divine treasure. Dorand publishers, Uyo.

257

258 Ezenwaka J, Ekeke B. A., Oyebade B. A. and Anyadiegwe AO (2004). Identification and
259 evolution of some threatened indigenous medical plants of the tropical rainforest of Eastern
260 Nigeria. *J. Conserv.* 2 (122): 131-141. F.P.L.G. Ibadan.

261 Hamilton, L.S. (2008). Forests and water – a thematic study prepared in the framework of the
262 Global Forest Resources Assessment 2005. Forestry Paper #155. Rome, Italy. 78 p.

263

264

265 Keay R. W. J. (1989). Trees of Nigeria: a revised version of Nigeria trees vol. 1 & 2.

266

267 Lata, N. (2013) . The impact of water hyacinth manure on growth attributes and yields in
268 Coriandrum. *IOSR Journal Of Enviromental Science, Toxicology and Food*
269 *Technology.*;5(3):04-07.

270

271 Mtambalika K., Munthali C., Gondwe D. and Missanjo E.(2014): Effect of seed size of
272 *Azalia quanzensis* on germination and seedling growth. *International Journal for Forestry*
273 *Research.* 2014;5.

274

275 Nwoboshi L. C. (1985). Teak (*Tectona grandis* Linn. F.) ten year after thinning. *Nig.J.*
276 *Forestry* 15:76-79.

277

278 Okeke R. E. (1996). Comparative anatomy of trees in Nigeria: *Azalia africana* species.

279

280 Osoro N. (2014). Effects of Water Hyacinth (*Eichhornia crassipes* [Mart.] Solms) Compost
281 on Growth and Yield Parameters of Maize (*Zea mays*). *British Journal of Applied Science &*
282 *Technology*.;4(4):617-633.

283

284

285 Palgreave K.C. (2002), *Trees of Southern Africa*. (3rd Ed). Revised by Coates MC. Struik
286 Publishers, Cape Town.

287

288 Razaq M., Zhang P. and Sheng H. (2017). Influence of nitrogen and phosphorus on the
289 growth and root morphology of Acer mono. *PLoS one*. 12(2):8.

290

291 Salim E. and Ullsten O. (1999). *Our forest: our future*. A paper of the world commission on
292 forest and sustainable development, Cambridge Press, U.K.

293

294 Smyth, A. J., and Montgomery, R. F., (1962). *Soil and landuse in central western Nigeria*.
295 Government of Western Nigeria, Ibadan, Nigeria.

296

297 Talkah A. (2015). Effect of Organic Fertilizer Water Hyacinth on the Growth and Production
298 Plant Taro (*Colocasia esculenta* L.). *Journal of Environment and Earth Sciences* [Internet].;5
299 (22): 5. Available from
300 :<http://www.iiste.org/Journals/index.php/JESS/article/view/27595/28309>.

301

302 Uddin M.B., Mukul S.A., Arfin Khan M.A.S. and Hossain M.K. (2012). Effects of organic
303 manure on seedlings growth and nodulation capabilities of five popular leguminous
304 agroforestry tree components of Bangladesh. *Journal of Forestry Science* 28 (4): 212 - 219

305

306 .

307