

# Effect of Different Moisture Harvesting Techniques on Seedling Survivals and Growth of Trees in degraded lands of southern Tigray

## Abstract

Tree planting on degraded lands play a key role in forest rehabilitation processes through afforestation and/or reforestation. Moisture harvesting structures (MHSs) has significant impact on seedling survivals at degraded lands. The objectives of this study were to investigate the impact of water harvesting techniques on seedling survival and growth performance of trees. Field experiments were conducted for two rainy seasons in southern Tigray, Atsela watershed. The experimental design followed was the split plot design. The MHSs as main plot used were eye-brow basin (EBs), micro trench (MTs), improved pit (IPs) and as control normal pit (NPs). The tree species grown as subplots were *Eucalyptus camaldulensis*, *Grevillea robusta*, *Olea europaea* and *Cupressus lusitanica*. The four tree species were planted by using seedlings. The tree survival rate, height, crown width (CW) and root collar diameter (RCD) of the four tree species were measured every six months after transplanting. The result shows that MHSs were significant in tree seedling height, CW and RCD but not in tree survival rate. Tree seedling height and CW grown in EBs was significantly higher than grown in MTs, IPs and NP ( $p \leq 0.05$ ). RCD of tree seedling was higher grown in EBs than NP (control) ( $p \leq 0.05$ ). The interaction of tree species seedlings and MHSs shows that those seedlings grown on MHSs were significantly thicker, taller and more survived than those grown on the NPs (control) ( $p \leq 0.05$ ). So based on the experiments, it is concluded that MHSs particularly the eyebrow basin (EBs) was considered as the most appropriate planting pit. So, further demonstration of eyebrow basin tree planting should be carried out.

## Introduction

More than 60 % of Tigray region have a mountainous topography and it aggravates to land degradation (1). The severity of soil erosion in the region is the result of the mountainous and hilly topography, erratic rainfall, and low degree of vegetation cover (2). In many parts of Tigray, soil erosion has made cultivation of old farmland impossible. Farmers have been forced to constantly cultivate new and more marginal areas.

Deforestation in the highlands of Ethiopia started already 2000 years ago (3). Forest plantation on degraded lands can play a key role in harmonizing long-term forest ecosystem rehabilitation process (4). Forest resources in Ethiopia in general, particularly in Tigray have experienced so much pressure due to increasing need for wood products and conversion to agriculture (5). The trend in the region today is to protect the remaining natural forests for their various social, economic and Environmental values. To strike the balance between the two interests, afforestation/ reforestation is very important. According to different scholars, “afforestation”, “reforestation” is used to distinguish new planted forests. The term “afforestation” is used in describing forests established artificially on land that previously did not carry forest for at least 50 years whereas, reforestation is activity of planting on forest exists area to replace or enrich the previous one (6). The purpose can be wood production or protection under the ownerships of the private sector, individual farmers, the community, or the state.

Afforestation/reforestation is the common approach of rehabilitation on degraded lands (2,7–10). Hence, millions of tree seedlings have been planted by different afforestation/reforestation programs in order to provide a basis for environmental improvement and increase the forest cover of the degraded lands. As cited by (11) the national average for tree seedling survival is less than 20%.

In most cases, the afforestation/reforestation are suffering from multiple environmental factors like limited water availability, free grazing, lack of proper management, premature cutting by peasants and inadequate care (9,10,12–14). Limited water availability is the primary factor controlling plant establishment and growth in the degraded lands (7–10,12,15–17). Therefore, to insure success of establishment of tree species MHSs can be used to collect rain water in areas close to the trees. Among the widely used micro catchment MHSs are eye-brow basin (EBs), micro trench (MTs), improved pit (IPs) and as control normal pit (NPs (10,18).

So, the hypothesis of the research was;

1) Hypothesis about the WH techniques

HO: The growth performance and survival of tree is the same under the four types of WH techniques

HA: The growth performance and survival of tree is different under the four types of WH techniques

2) Hypothesis about the tree species

HO: The growth performance and survival of the four tree species is the same.

HA: The growth performance and survival of the four tree species is different

3) Hypothesis about the interaction effect

HO: The difference in growth performance and survival among the four tree species does not depend on types of WH techniques

HA: The difference in growth performance and survival among the four tree species depend on types of WH techniques


Thus, the main aim of the study was to investigate the possible contribution of MHSs in improving survival rate of tree seedlings and to estimate their effects on the performance of the seedling.

The specific objective of this study was; 1). To evaluate the effect of different MHSs on growth performance and survival rates of tree species seedlings 2). To determine the interaction effect of MHSs and tree species seedlings 3). To evaluate the four tree species seedling performance in the in degraded lands/ moisture stressed areas of Southern Tigrai, Atsela watershed, Northern Ethiopia.

## Materials and methods

### Description of the study area

The study was conducted in Southern Tigray, Atsela watershed, Northern Ethiopia



( Figure 1). It is located 702 km north of Addis Ababa and about 90 km south of the Tigray Regional capital state Mekelle. The agro ecology of the experimental site is classified as high land with an average temperature and annual rain fall of 15.8 degree centigrade and 570.2 mm respectively. The altitude of the district ranges from 2,907-2,938m.a.s.l.The occurrence of rain is highly variable in the study area, and rain is not evenly distributed throughout the months when it rains. The annual mean precipitation ranges from 238-939 mm.




Figure 1: Study area

### Species selection

Four woody species were carefully selected for experimentation based on the preference of the community's multi-criteria decision approach taking into account indicators of ecological suitability, socio-economical functions, protection functions and root characteristics. The selected species have the following characteristics based on (19–21):

#### I. *Eucalyptus camaldulensis* Labill.

*Eucalyptus camaldulensis* is from the Myrtaceae family. It has been planted in Africa since around 1900, it does well in semi-arid regions and tolerates a long dry season as well as some salinity. It does well in deep silt or clay soil in Dry and Moist Kolla agroclimatic zones in Tigray, 200–2,800 m. *Eucalyptus camaldulensis* is very suitable for fuel, construction and soil conservation. In local medicine, the steam from boiled leaves is inhaled to relieve the common cold and other bronchial problems. Steam from eucalyptus is believed to kill disease-causing bacteria and microbes in sick rooms. It is used as a steam inhalant for chest congestion. Eucalyptus is rarely used internally.

#### *Grevillea robusta* R. Br.

*Grevillea robusta* is from Proteaceae family with medium-sized to tall tree up to 30 m high, usually less than 10 m in Ethiopia. A very successful Australian tree planted and widely used in dry, moist and wet Weyna Dega and Dega agroclimatic zones, 0–3,000 m. It used as firewood, charcoal, timber (furniture), poles, fodder (leaves), bee forage, shade, ornamental, soil conservation and windbreak.

***Olea europaea* L. subsp. *cuspidata***

It is from oleaceae family with a height of 10-15 m. It is widely distributed in dry forest in east Africa and Ethiopia. It reaches southern Africa, it is best in good forest soil, but hardy and drought resistant once established, even in poor soils. The species found in moist and Wet Weyna Dega and lower Dega agroclimatic zones in all regions, 1,400-3,100 m. In Ethiopia, the leaves, twigs and wood of the African olive are used to fumigate pots for milk and for the local beverages, Tella and Tej. Twigs are used as toothbrushes (tooth sticks) and hard wood for carving. In addition, it used for firewood, charcoal, timber (furniture, floors, panelling, walking sticks), poles, posts, medicine (stem, bark, leaves) and bee forage.

***Juniperus procera* Hochst. ex Endl.**

*Juniperus procera* (Cupressaceae) an erect evergreen tree grow up to 40 m tall, trunk in old trees up to 1 m in diameter. It is valuable timber tree indigenous to Ethiopia and eastern Africa highland forests. *Juniperus* is very common plant generally in the northern and northern central parts of Ethiopia and particularly in Tigray about 1100 to 3500 m altitudes. Widely used for firewood, timber, poles, posts, medicine (bark, leaves, twigs, buds), shade, ornamental and windbreak.

**Moisture harvesting structures (MHSs)**

The effective MHSs which are familiar in the watersheds of Tigray region for rehabilitation of degraded land were adopted for the experiment from ministry of agriculture and rural development. According to (18) the specification and design of the structures we used were as follows.

**EYEBROW BASINS (EBs):** are larger circular and stone faced structures for tree and other species planting. They are effective in low rainfall areas to

grow trees and harvest moisture. The technical standards constructed were;

Size: 2.5 m diameter;

- Stone riser with 0.25 m depth of the foundation;
- Height 0.5 m;
- Stone riser sealed with soil excavated from water collection area;
- Water collection area: dug behind the plantation pit: 1 m width x 1 m length x 25 cm depth (lower side);
- Three plantation pits of 40cm depth x 40cm diameter dug between riser and water collection area as in the left figures (2,3 and 4) shown.

**MICROTRENCHES (MTRs):** are rectangular and deep pits constructed along the contours. The technical standards constructed and design we follow was;

- Size of the trench: 1.5 length x 0.4 m width x 0.5 m depth (downside);
- Trenches were provided with a small and low tie in the middle to regulate water flow (15 cm width);
- Trees were not planted in the middle of the trench but in front of it;
- Plantation pit 40cm depth x 40 cm width.

**IMPROVED PITS (IPs):** Was square shaped water collection pits constructed along the contours with a plantation pit in front of the main water storage pit.

- Dimension: 0.60m length x 0.60 width x 0,5 m depth
- Planting pit 40cm depth x 40 cm width, were prepared in front of the square shaped water collection pit

**NORMAL PIT (NPs):** are circular and deep pits constructed along the contour..

- Size: 40cm depth x 40 cm width

## **Experimental design and method of implementation**

### **Experimental design**

Four tree species were established in four MHSs in a split-plot design. There were 16 treatments combination of two factors (Four MHSs \* Four Tree species = 16 treatments) Table 1. Each treatment combination was tried on 3 replications and thus a total of 48 plots (Four MHSs \* Four Tree species\* Three replication= 48 plots). MHSs were the main plot factor in this experimental

design, and tree species were randomly assigned to one of four MHSs. The tree species were planted by using seedlings.

Table 1. The treatments of the experiment

No.	Type of pit	Tree species	Treatments (Code)
1	EYEBROW BASINS (EBs)	Eucalyptus camaldulensis (E )	EBsE
		Grevillea robusta (G)	EBsG
		Olea europaea (O)	EBsO
		Cupressus lusitanica (C )	EBsC
2	MICROTRENCHES (MTRs)	Eucalyptus camaldulensis (E )	MTRsE
		Grevillea robusta (G)	MTRsG
		Olea europaea (O)	MTRsO
		Cupressus lusitanica (C )	MTRsC
3	IMPROVED PITS (IPs)	Eucalyptus camaldulensis (E )	IPsE
		Grevillea robusta (G)	IPsG
		Olea europaea (O)	IPsO
		Cupressus lusitanica (C )	IPsC
4	NORMAL PIT (NPs)	Eucalyptus camaldulensis (E )	NPE
		Grevillea robusta (G)	NPG
		Olea europaea (O)	NPO
		Cupressus lusitanica (C )	NPC

### Data collection

The data collected were growth variables of tree species like 1<sup>st</sup> tree survival rate (December, 2016) in %; 2<sup>nd</sup> survival rate (May, 2016) in %; 3<sup>rd</sup> survival rate (December, 2017) in %; Tree height (H) in cm; Root collar diameter (RCD @<sub>10cm</sub> stamp height) in cm and Crown width (CW) in cm were taken.



## Statistical analysis

Results were analyzed using R software. Analysis of variance (ANOVA) was made to determine the significance of variation between the tree species and MHSs. Tukey Multiple Range Test was used to compare mean values of various growth variables in each treatment.

## Results and discussions

### Response of tree growth to different MHSs

The result of this study showed that 1<sup>st</sup> survival, 2<sup>nd</sup> survival and 3<sup>rd</sup> survival rates of tree seedlings grown in different MHSs were not varied significantly ( $p \geq 0.05$ ). However, it was significantly different ( $p \leq 0.05$ ) for the growth variables tree seedling height, root collar diameter (RCD) and crown width (CW) Table 2. Tree height and CW was significantly higher in EBs than MTs, IPs and NP. In addition, RCD was higher in EBs than NP (control) and not varied significantly with MTs and IPs. Not differences were shown in tree height grown in MTs, IPs and NP.

Even though the MHSs doesn't affect the survival of tree seedling, those structures that can conserve water and soil like EBs has a positive and significant on growth variables. This finding confirms, these structures could solve the moisture stress that commonly limits growth, survival and distribution of tree seedlings (22). The results of this study clearly indicated that the rehabilitation of degraded land and ensuring the survival of tree seedling is possible using MHSs. However, the implementation of various MHSs depends on local rainfall characteristics, construction materials and site conditions. MHSs have proved to be a valuable tool, especially in degraded lands to establish trees and to allow reforestation.

Similar results were shown in different studies of MHSs. According to (23) the impact of MHSs is high and showed an 8 times increase in total biomass compared with the normal pit and also proved that there was an increase in tree height by 20%. Similarly other study showed that *Zizyphus maurutuana* growth rates in the Jodhpur province of India were from 25-33% higher in shallow one meter(24). The results from the semiarid loess region of China (25) also showed that water harvesting treatments had a prominent effect on the growth characteristics of *T. ramosissima*.

Table 2: The effect of different MHSs on growth variables of tree species

Growth	Moisture harvesting structures (MHSs) (Mean $\pm$ (SE))	P-
--------	---	----

variables	EBs	MTs	IPs	NP (control)	Value
1 <sup>st</sup> survival (%)	91.67 ± (5.98)	97.22 ± (2.78)	88.89 ± (6.27)	91.67 ± (5.98)	NS
2 <sup>nd</sup> survival (%)	63.89 ± (13.27)	63.92 ± (11.19)	55.56 ± (9.48)	41.67 ± (10.95)	NS
3 <sup>rd</sup> survival (%)	63.89 ± (13.27)	63.89 ± (11.21)	55.56 ± (9.48)	33.33 ± (10.05)	NS
Height (cm)	75.50 ± (17.73) <sup>a</sup>	34.83 ± (8.16) <sup>b</sup>	42.08 ± (8.19) <sup>b</sup>	27.33 ± (8.1) <sup>b</sup>	0.002
RCD (cm)	1.36 ± (0.27) <sup>a</sup>	0.95 ± (0.14) <sup>ab</sup>	1.01 ± (0.14) <sup>ab</sup>	0.63 ± (0.16) <sup>b</sup>	0.023
CW (cm)	70.54 ± (13.88) <sup>a</sup>	43.17 ± (8.58) <sup>b</sup>	38.50 ± (8.72) <sup>bc</sup>	18.58 ± (5.89) <sup>c</sup>	0.001

**Note:** EBs (eyebrow basin), (MTs) micro trench, (IPs) improved pit and (NPs) normal pit as control. Similar letter in the row shows not significant difference and different letters indicate significance differences, NS: not significant difference between moisture harvesting structures (MHSs) at  $p \leq 0.05$

### The interaction effect of tree species and MHSs

The interaction effect in Table 3 shows that those seedlings grown on MHSs were significantly thicker, taller and more survived than those grown on the NPs (control). The 2<sup>nd</sup> survival, 3<sup>rd</sup> survival and height of *Grevillea robusta* seedlings planted in EBs MHSs were significantly higher than the same species planted in MTs, IPs and NPs ( $p \leq 0.05$ ). In addition, RCD and CW were higher in EBs MHSs compare to the control (NPs).

The survival rate of *Eucalyptus globulus* was not varied with the different MHSs. It could be the ability of the species to survive to the different MHSs. However, height was significantly higher in MHSs of EBs and MTs than IPs and NPs ( $p \leq 0.05$ ). Furthermore, RCD and CW growth variables were significantly higher in *Eucalyptus globulus* seedlings grown in EBs MHSs than seedlings planted in MTs, IPs and NPs ( $p \leq 0.05$ ). These results agreed with the results of (10) who showed EBs showed a significantly higher survival rate, RCD and DBH in growing trees seedlings.

Whereas, the two indigenous tree species i.e. *Olea europaea* and *Juniperus procera* tree species were not varied significantly with the different MHSs. It could be the late response of the tree seedlings to the growth variables.

Table 3 The Interaction effect of different *MHSs* on seedling survival and growth performance tree species

Tree species	Growth Variables	Moisture harvesting structures (MHSs)				<i>P</i> -value
		EBs	MTs	IPs	NPs	
<i>Grevillea robusta</i>	1 <sup>st</sup> survival (%)	100	100	88.89 (11.11)	100	NS
	2 <sup>nd</sup> survival (%)	100 <sup>a</sup>	55.56 (11.11) <sup>b</sup>	44.44 (11.11) <sup>b</sup>	55.55 (11.11) <sup>b</sup>	0.015
	3 <sup>rd</sup> survival (%)	100 <sup>a</sup>	55.56 (11.11) <sup>b</sup>	44.44 (11.11) <sup>b</sup>	55.56 (11.11) <sup>b</sup>	0.015
	Height (cm)	70 (9.6) <sup>a</sup>	24 (6) <sup>b</sup>	37.67 (5.24) <sup>b</sup>	35.33 (7.06) <sup>b</sup>	0.01
	RCD (cm)	1.7 (0.2) <sup>a</sup>	1.17 (0.17) <sup>ab</sup>	1.13 (0.07) <sup>ab</sup>	1.07 (0.07) <sup>b</sup>	0.041
	CW (cm)	91.33 (9.49) <sup>a</sup>	57 (16.29) <sup>a</sup>	28 (9.5) <sup>b</sup>	21.67 (6.11) <sup>b</sup>	0.006
<i>Eucalyptus globulus</i>	1 <sup>st</sup> survival (%)	100	100	77.78 (22.22)	100	NS
	2 <sup>nd</sup> survival (%)	100	66.67 (33.33)	44.45 (22.22)	44.44 (29.4)	NS
	3 <sup>rd</sup> survival (%)	100	66.67 (33.33)	44.45 (22.22)	11.11 (11.11)	NS
	Height (cm)	155 (17.35) <sup>a</sup>	56 (28.48) <sup>ab</sup>	51.33 (25.69) <sup>b</sup>	24.33 (24.33) <sup>b</sup>	0.023
	RCD (cm)	2.27 (0.18) <sup>a</sup>	0.8 (0.42) <sup>b</sup>	0.83 (0.42) <sup>b</sup>	0.33 (0.33) <sup>c</sup>	0.022
	CW (cm)	113.33 (12.39) <sup>a</sup>	48 (25.15) <sup>b</sup>	47.67 (28.32) <sup>b</sup>	13.33 (13.09) <sup>c</sup>	0.05
<i>Olea europaea</i>	1 <sup>st</sup> survival (%)	77.78 (22.22)	88.89 (11.11)	88.89 (11.11)	77.78 (22.22)	NS
	2 <sup>nd</sup> survival (%)	11.11 (11.11)	33.44 (19.15)	55.56 (50.92)	11.11 (19.24)	NS
	3 <sup>rd</sup> survival (%)	11.11 (19.24)	33.33 (33.33)	55.56 (29.4)	11.11 (11.11)	NS
	Height (cm)	12.33 (12.33)	15.67 (7.88)	17.33 (8.69)	8.67 (8.67)	NS
	RCD (cm)	0.43 (0.43)	0.67 (0.33)	0.67 (0.33)	0.33 (0.33)	NS
	CW (cm)	14.33 (14.33)	13 (7)	16 (8.72)	5 (5)	NS
<i>Juniperus procera</i>	1 <sup>st</sup> survival (%)	88.89 (11.11)	100	100	88.89 (11.11)	NS
	2 <sup>nd</sup> survival (%)	44.43 (29.4)	100	77.77 (11.11)	55.55 (29.4)	NS

Tree species	Growth Variables	Moisture harvesting structures (MHSs)				<i>P</i> -value
		EBs	MTs	IPs	NPs	
	3 <sup>rd</sup> survival (%)	44.44 (29.4)	100	77.78 (11.11)	55.56 (29.4)	NS
	Height (cm)	64.67 (33.79)	43.67 (7.12)	62 (12.49)	41 (20.52)	NS
	RCD (cm)	1.03 (0.58)	1.17 (0.09)	1.4 (0.1)	0.77 (0.39)	NS
	CW (cm)	63.17 (32.37)	54.67 (6.77)	62.33 (8.45)	34.33 (17.7)	NS

**Note:** EBs (eyebrow basin), (MTs) micro trench, (IPs) improved pit and (NPs) normal pit as control. Similar letter in the row shows not significant difference and different letters indicate significance differences, NS: not significant difference between moisture harvesting structures (MHSs) at  $p \leq 0.05$

## Response of tree species under different MHT in the site

Species differed significantly in all growth variables, except the 1<sup>st</sup> survival % in the study site. *Olea europaea* had significantly lower survival in the watershed when statistically compared to *Juniperus procera* and *grevillea robusta* in the 3<sup>rd</sup> survival rate. Similarly, height and CW of *Olea europaea* was significantly lower from *Eucalyptus globulus*, *Grevillea robusta* and *Juniperus procera* tree seedlings. This could be the morphological characteristics of the species.

Table 4 Response of tree species in different MHSs for the different growth Variables

Growth Variables	Tree seedling species				P-Value
	<i>Eucalyptus globulus</i>	<i>Juniperus procera</i>	<i>Olea europaea</i>	<i>Grevillea robusta</i>	
1 <sup>st</sup> survival (%)	94.44 ± (5.56)	94.44 ± (3.75)	83.33 ± (7.68)	97.22 ± (2.78)	NS
2 <sup>nd</sup> survival (%)	63.89 ± (12.62) <sup>ab</sup>	69.44 ± (11.21) <sup>a</sup>	27.81 ± (9.90) <sup>b</sup>	63.89 ± (7.63) <sup>ab</sup>	0.019
3 <sup>rd</sup> survival (%)	55.56 ± (13.19) <sup>ab</sup>	69.44 ± (11.21) <sup>a</sup>	27.78 ± (9.91) <sup>b</sup>	63.89 ± (7.63) <sup>a</sup>	0.015
Height (cm)	71.66 ± (18.2) <sup>a</sup>	52.83 ± (9.52) <sup>a</sup>	13.50 ± (4.19) <sup>b</sup>	41.75 ± (6) <sup>ab</sup>	<0.001
RCD (cm)	1.06 ± (0.26) <sup>ab</sup>	1.09 ± (0.17) <sup>ab</sup>	0.53 ± (0.16) <sup>b</sup>	1.27 ± (0.1) <sup>a</sup>	0.014
CW (cm)	55.58 ± (14.11) <sup>a</sup>	53.63 ± (8.91) <sup>a</sup>	12.08 ± (4.22) <sup>b</sup>	49.50 ± (9.49) <sup>a</sup>	0.001

**Note:** EBs (eyebrow basin), (MTs) micro trench, (IPs) improved pit and (NPs) normal pit as control. Similar letter in the row shows not significant difference and different letters indicate significance differences, NS: not significant difference between moisture harvesting structures (MHSs) at  $p \leq 0.05$

## Conclusion and recommendation

The study results revealed that MHSs are verified as effective structures in producing well survived and grown trees species. The growth variables of the tree species planted in the EBs is better than MTs, IPs and NPs. Tree species planted in MTs and IPs were also the most appropriate planting pit than NP. The MHSs shows great potential in increasing tree survival and growth performance due to helping to harvest rainwater and protecting them. However, the use

MHSs for native tree species (*Juniperus procera* and *Olea europaea*) didn't affect their growth performance and survivals.

So, expanding MHSs is the most appropriate afforestation method particularly for degraded area. More elaborative studies are required with more representative locations and in different soil and agro ecology.

## References

1. BoAaNR. Tigray Region Bureau of Agriculture and Rural Development 2017 Annual Reports, Northern Ethiopia. Mekelle; 2017.
2. Kjell E, Tor-Gunnar V, Yibabe T, Mitiku H. SOIL CONSERVATION IN TIGRAY, . 2002.
3. Hamilton AC. An upper Pleistocene pollen diagram from highland Ethiopia. In: International Quaternary Congress. 1977. p. 193.
4. Sharma D, Sunderraj S. Species selection for improving disturbed habitats in Western India. *Curr Sci*. 2005;462–7.
5. Sileshi GW. A critical review of forest biomass estimation models, common mistakes and corrective measures. *For Ecol Manage* [Internet]. 2014;329:237–54. Available from: <http://dx.doi.org/10.1016/j.foreco.2014.06.026>
6. Yitebitu M, Eshetu Z, Nune S. Ethiopian forest resources: current status and future management options in view of access to carbon incomes. 2010;(October):9–11.
7. Dastorani MT, Kouhzad B, Sepehr A, Talebi A. The role of rainwater harvesting on providing vegetation water requirement in arid and semi-arid regions. *Eur Water*. 2017;58(2017):521–7.
8. Hani M, Raed A, Khaled A, Jemal Y. Effect of Three Water Harvesting Techniques on Forage Shrub and Natural Vegetation in the Badia of Jordan. *Int J Bot* [Internet]. 2011;7(3):230–6. Available from: <http://docsdrive.com/pdfs/ansinet/ijb/2011/230-236.pdf>
9. Singh G, Khan AU, Kumar A, Bala N, Tomar UK. Effects of rainwater harvesting and afforestation on soil properties and growth of *Embliba officinalis* while restoring degraded hills in western India. *African J Environ Sci Technol* Vol. 2012;6(August):300–11.
10. Sisay D, Tewodros A, Belete B, Gete Z. Impacts of micro-basin water harvesting structures in improving vegetative cover in degraded hillslope areas of north-east Ethiopia. *Rangel J*. 2009;31(January):259–265.

11. Badege B. Deforestation and Land Degradation on the Ethiopian Highlands: A Strategy for Physical Recovery. In: International Conference on Contemporary Development Issues in Ethiopia. 2003. p. 1–9.
12. Sona M, Abdel M. Impact of Water Harvesting Techniques on Growth of Some Indigenous Tree Species in Jebel Awlia Locality, Sudan. *Glob J Sci Front Res D Agric Vet* [Internet]. 2016;16(3). Available from: [https://globaljournals.org/GJSFR\\_Volume16/7-Impact-of-Water-Harvesting.pdf](https://globaljournals.org/GJSFR_Volume16/7-Impact-of-Water-Harvesting.pdf)
13. Temesgen G, Amare B, Hagos G. Land Degradation in Ethiopia : Causes , Impacts and Rehabilitation Techniques. *J Environ Earth Sci*. 2014;4(9).
14. Thinh V, Ralph M, Nguyen V, Tran V. Environmental Factors Affecting the Abundance and Presence of Tree Species in a Tropical Lowland Limestone and Non-limestone Forest in Ben En National Park, Vietnam. *J For Environ Sci* [Internet]. 2015;31(3):177–91. Available from: <http://dx.doi.org/10.7747/JFES.2015.31.3.177>
15. Battaglia LL, Fore SA, Sharitz R. Seedling emergence , survival and size in relation to light and water availability in two bottomland hardwood species. *J Ecol*. 2000;88:1041–50.
16. Bert R, Jean P, Jan N, Yves L, Amanuel Z, Sarah T, et al. Establishment and management of woody seedlings in gullies in a semi-arid environment (Tigray, Ethiopia). *Plant Soil*. 2009;324(1):131–56.
17. Demisachew T, Awdenegest M, Mihret D. Impact evaluation on survival of tree seedling using selected in situ rainwater harvesting methods in. *J Hortic For*. 2018;10(April):43–51.
18. Lakew D, Carucci V, Asrat W, Yitayew A (eds). *Community Based Participatory Watershed Development: A Guideline*. Ministry of Agriculture and Rural Development. Addis Ababa, Ethiopia; 2005.
19. Azene B, Tengnäs B. *Useful Trees and Shrubs of Ethiopia: Identification, Propagation, and Management for 17 Agroclimatic Zones*. 2007;566. Available from: <http://books.google.com.et/books?id=15UfAQAAIAAJ>
20. Ermias D. *Natural Database for Africa(NDA)* [Internet]. Addis Ababa, Ethiopia; 2011. Available from: <http://go.microsoft.com/fwlink/?LinkId=69157>
21. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. *Agroforestry Database: a tree*

reference and selection guide [Internet]. 2009. Available from:  
<http://www.worldagroforestry.org/af/treedb/>

22. Warren J, Meinzer F, Brooks J, Domec J. Vertical stratification of soil water storage and release dynamics in Pacific Northwest coniferous forests. *Agric For Meteorol.* 2005;130(1–2):39–58.
23. Gupta G. Rain-water management for tree planting in the Indian Desert. *J Arid Environ.* 1995;31:219–35.
24. Ojasvi P, Goyal R, Gupta J. The micro-catchment water harvesting technique for the plantation of jujube (*Zizyphus mauritiana*) in an agroforestry system under arid conditions, Division of Resources Management, Central Arid Zone Research Institute. Jodhpur 342003, India. 1999;8.
25. Xiao-Yan L, Lian-You L, Shang-Yu G, Pei-Jun S, Xue-Yong Z, Chun-Lai Z. Microcatchment water harvesting for growing *Tamarix ramosissima* in the semiarid loess region of China. *J For Ecol Manag.* 2005;214:111–7.




Fig. 2. Figure showing water collection ditch and planting pits






Fig. 3. Plantation pits of 40cm depth x 40cm diameter dug between riser and water collection Area




Fig. 4. Preparation of planting pit (40cm depth x 40 cm width) in front of the square shaped water collection pit