

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

**PERFORMANCE CHARACTERISTICS OF BROILERS FED WITH TERMITES  
(*MACROTERMES BELLICOSUS*) AS PROTEIN SUPPLEMENT IN POULTRY FEED.**

**ABSTRACT**

Eight weeks feeding trails were conducted with 120 broiler chicks at poultry production unit of the Department of Animal Science, Usmanu Danfodiyo University Sokoto. A commercial diet was compared with two other diets; fishmeal diet and termite supplemented diet. Each of the three treatments was replicated four times, with ten chicks per replicate in a Complete Randomized Design (CRD). Broilers were fed regularly throughout the period of the experiment. Results of Carcass evaluation showed significant ( $P<0.05$ ) difference in weight of broiler parts like; breast, drum stick and liver, but there was no significant ( $P>0.05$ ) difference in back, thigh, wings and gizzard in all treatment diets. Other parts; feet, head, and spleen showed significant ( $P<0.05$ ) difference. But neck, heart, lungs, intestine and crop shows no significant ( $P>0.05$ ) difference in their weights. Proximate analysis showed no significant ( $P>0.05$ ) difference in nutrient composition in all the treatment diets, only ash showed significant ( $P<0.05$ ) difference in fishmeal. Mineral composition showed significant ( $P<0.05$ ) difference in almost all treatment diets except in phosphorus, copper and zinc. It was concluded that termite supplemented diet can successfully and economically replace fishmeal as protein source in the diets of broilers without deleterious effect on their health and carcass quality.

Key words: *Macrotermes bellicosus*, Broilers, Carcass, Nutrients and mineral composition.

**INTRODUCTION**

Termites are social land dwelling insects. They are cosmopolitan and polymorphic mainly found in tropical and sub tropical areas, comprising some 21,000 to 30,000 species (Malaka, 1996, Kambhampati and Eggleton, 2000). They are further divided into seven Families namely; Mastotermitidae, Termopsidae, Kalotermitidae, Hodotermitidae, Rhinotermitidae, Semitermitidae and Termitidae (Kambhampati and Eggleton, 2000). The first six families are referred to as lower termites and the remaining family of the termites is referred to as higher termites (Myles, 2003).

Abd Rahman *et al.* (2012) stated that, insects are more promising alternative protein source to rearing animals (Allotey, 2003). To a larger extent hundreds of insect species, have been used as animal protein supplements with some more important groups including, termites,

34 grasshoppers, caterpillars, beetle grubs and adults, bees etc (Banjo *et al.*, 2006). Termites are a  
35 delicacy in the diet of some human cultures. In many cultures, termites are used as food  
36 particularly the alates type (reproductive forms) (Grimaldi and Engel, 2005, Oguwike, *et al.*,  
37 2013). The use of insects as human food and animal feed is widely spread in tropical and sub  
38 tropical countries and is the cheapest source of animal protein (Paul and Sudipta, 2011).

39 Poultry farming has expanded rapidly in developing countries in last two decades. Termites are  
40 used as complementary feed sources for poultry and they are used as feed for chicken and  
41 guinea fowl in Togo and Burkina Faso (Ravindran and Blair, 1993). In some Countries of  
42 West Africa they already have a primitive way of rearing termites on crop residues (on inverted  
43 clay pots or baskets) for food and poultry feed supplement, many developing countries termites  
44 are used as feed, locally to supply day old chicken or guinea fowls and in Togo termites are  
45 bred for this purpose (Ravindran and Blair, 1993). *Macrotermes bellicosus*, this species simply  
46 called termites in most Nigerian communities are the commonest and has high composition of  
47 mineral elements; vitamins (N'tukuyo *et al.*, 2012). Nutritive potentials and utilization of  
48 termites as poultry feed ingredients have been documented in Botswana and recommend  
49 termites as source of protein in poultry diets in poultry production (Bamphith and John, 2012).  
50 The protein content of termites has been reviewed to vary from 20.00 to 46.3 per cent  
51 (Sogbesan and Ugwumba, 2008), while (Paul and Sudipta, 2011) reported termites to have  
52 protein percentage of about 81.66 per cent and 87.33 for workers and sexual forms  
53 respectively, and they could help in maintaining acidity and water balance in the body of  
54 poultry.

55 This work incorporated termites (*Macrotermes bellicosus*) in poultry feed as protein  
56 supplement, collected in four agricultural zones of Kebbi State, North-Western Nigeria.

## 57 **MATERIALS AND METHODS**

58 The study on feeding trial of broilers was conducted at poultry production Unit of Department  
59 of Animal science, Usmanu Danfodiyo University Sokoto, which is located at state Veterinary  
60 Centre along Aliyu Jodi road, and Sokoto lies between latitudes 12<sup>0</sup> and 13<sup>0</sup> 05'N and  
61 longitudes 4<sup>0</sup> 8' and 6<sup>0</sup> 4'E (Mamman *et al.*, 2000).

62 Termites (*Macrotermes bellicosus*) were collected including workers, soldiers in selected areas  
63 both dry land and wet land alike. Mounds were excavated using diggers, spade, shovels and  
64 hoes causing termites to come out. They were collected into plastic containers and sacs as in  
65 method reported by N'tukuyoh *et al.* (2012). Termite samples were randomly collected from  
66 mounds in different locations within the study area. They were collected with sand and  
67 separation of termites was done in the Laboratory by flotation methods as reported by N'tukuyo  
68 *et al.* (2012). Mound soils were transferred into buckets separately and water was poured into  
69 them and floating termites were skimmed using a sieve (2.0 mm), sandy water was poured into  
70 a sieve (2.00 mm) for further separation of the sunken termites. After separating them from  
71 sand, they were then washed with clean water and sun-dried. They were ground into powder  
72 and package in plastic containers and kept in the laboratory until the time of used.

73 Day old broilers were obtained from Sokoto market. The birds were sourced from a  
74 commercial hatchery Ibadan, Oyo State; they were purchased according to the number of  
75 treatments and replicates. A total of one hundred and twenty birds were used for this study.  
76 They were divided into three treatments and four replicates, ten birds per replicate; they were  
77 further divided into three dietary groups. Experimental birds were randomly selected as in the  
78 method of (Sogbesan and Ugwumba, 2008 and Oguwike *et al.*, 2013). They were fed with three  
79 different diets, that is termite-protein supplemented meal, fishmeal and commercial feeds as  
80 control.

81 Poultry house was well cleaned and sprayed with fumigants and disinfectants to avoid infection  
82 of various diseases. This was done before the arrival of the birds and the house was partitioned  
83 into pens according to the replicates. Experimental birds were kept for three days after arrival  
84 to take care of stress. Within these three days, they were administered with anti stress drugs  
85 after which they were weighed and allocated to their replicate groups. Each group was  
86 replicated four times. Vaccines were administered according to the routine; antibiotics and  
87 Coccidiostats were administered according to recommendations of Roberts (1998).  
88 Experimental birds were kept in a cross ventilation of open side walls house and were kept on  
89 deep litter house.

90 At the end of experiment that is at 8<sup>th</sup> week, six birds from each treatment (three birds per  
 91 replicate) were taken at random tagged for carcass analysis. They were weighed separately and  
 92 slaughtered Plucked carcass was dissected and eviscerated. The head, breast and internal organs  
 93 (i.e. intestine, crop, gizzard, liver, lungs, heart, spleen) were weighed separately, eviscerated  
 94 carcass was then cut into parts (i.e. back, breast, thigh, drum stick, neck, wings, feet) weighed  
 95 separately.

96 The results obtained were tested for significance of differences between treatments that is  
 97 carcass evaluation and proximate composition using Start View Statistical Analysis Software  
 98 (SAS. 2002)

## 99 Results

100 **Table 1: Carcass evaluation of experimental broilers (Primal cuts)**

101 102 103 104 105 106 107 108	Parameters						
	Treatments	Breast	Back	Drum stick	Thigh	Wings	Liver
Commercial diet	410.43 <sup>a</sup>	159.95 <sup>a</sup>	152.10 <sup>a</sup>	160.73 <sup>a</sup>	127.10 <sup>a</sup>	95.98 <sup>a</sup>	41.88 <sup>a</sup>
Fishmeal diet	348.60 <sup>b</sup>	152.20 <sup>a</sup>	117.43 <sup>b</sup>	134.75 <sup>a</sup>	104.20 <sup>a</sup>	34.78 <sup>b</sup>	36.13 <sup>a</sup>
Termite diet	269.35 <sup>b</sup>	149.98 <sup>a</sup>	132.73 <sup>a</sup>	147.13 <sup>a</sup>	101.85 <sup>a</sup>	36.13 <sup>b</sup>	36.98 <sup>a</sup>
<b>SEM</b>	<b>9.479</b>	<b>5.935</b>	<b>2.66</b>	<b>3.209</b>	<b>2.076</b>	<b>0.735</b>	<b>0.796</b>

109 Means along the same column with similar superscripts are not significantly (P>0.05) different  
 110 from one another

111 **SEM** = Standard error of means

114 **Table 2: Carcass evaluation of the experimental broilers (Other cuts)**

115 116 117	Parameters							
	Treatments	Feet	Head	Neck	Heart	Lungs	Spleen	Intestine
Commercial diet	70.45 <sup>a</sup>	47.59 <sup>a</sup>	65.33 <sup>a</sup>	7.93 <sup>a</sup>	8.83 <sup>a</sup>	2.60 <sup>a</sup>	113.15 <sup>a</sup>	9.05 <sup>a</sup>

118

119 Fishmeal diet 52.93<sup>b</sup> 38.65<sup>b</sup> 49.70<sup>a</sup> 6.80<sup>a</sup> 7.90<sup>a</sup> 1.75<sup>b</sup> 95.65<sup>a</sup> 8.25<sup>a</sup>

120

121 Termite diet 55.58<sup>b</sup> 40.30<sup>b</sup> 51.15<sup>a</sup> 7.73<sup>a</sup> 7.63<sup>a</sup> 1.75<sup>b</sup> 121.65<sup>a</sup> 7.95<sup>a</sup>

122 **SEM 0.815 0.674 1.340 0.101 0.209 0.065 2.942 0.174**

123 Means along the same column with similar superscripts are not significantly (P>0.05) different  
124 from one another

125 SEM = Standard error of means

126

127 **Table 3: Nutrient composition of experimental broilers (percentage)**

128	Parameters						
	129	Carbohydrate	Protein	Fat/Lipids	Moisture	Fibre	Ash
130	Commercial diet	44.22 <sup>a</sup>	34.56 <sup>a</sup>	9.67 <sup>a</sup>	11.38 <sup>a</sup>	2.30 <sup>b</sup>	8.42 <sup>b</sup>
131							
132	Fishmeal diet	50.38 <sup>a</sup>	33.01 <sup>a</sup>	10.17 <sup>a</sup>	11.17 <sup>a</sup>	3.92 <sup>a</sup>	12.90 <sup>a</sup>
133							
134	Termite supplemented diet	47.76 <sup>a</sup>	29.04 <sup>a</sup>	10.63 <sup>a</sup>	12.96 <sup>a</sup>	2.43 <sup>a</sup>	10.17 <sup>b</sup>
135	<b>SEM</b>	<b>0.819</b>	<b>0.754</b>	<b>0.245</b>	<b>0.160</b>	<b>0.052</b>	<b>0.193</b>

136 Means along the same column with similar superscripts are not significantly (P>0.05) different  
137 from one another

138 SEM = Standard error of means

139

140 **Table 4: Mineral composition, sodium, potassium, magnesium, phosphorus, zinc, copper**  
141 **and iron of experimental broilers (mg)**

142	Parameters								
	143	Na	K	Ca	Mg	P	Zn	Cu	Fe
144	Commercial diet	127.08 <sup>b</sup>	155.00 <sup>b</sup>	1.467 <sup>b</sup>	1.158 <sup>b</sup>	5.408 <sup>a</sup>	0.034 <sup>a</sup>	0.081 <sup>a</sup>	2.357 <sup>a</sup>

145

146	Fishmeal diet	160.31 <sup>a</sup>	221.38 <sup>a</sup>	1.877 <sup>a</sup>	1.104 <sup>b</sup>	5.868 <sup>a</sup>	0.007 <sup>b</sup>	0.075 <sup>a</sup>	2.767 <sup>a</sup>
147									
148	Termite supplemented diet	163.75 <sup>a</sup>	182.08 <sup>a</sup>	1.254 <sup>b</sup>	1.588 <sup>a</sup>	5.605 <sup>a</sup>	0.002 <sup>b</sup>	0.072 <sup>a</sup>	1.955 <sup>a</sup>
149	<b>SEM</b>	<b>2.456</b>	<b>3.986</b>	<b>0.030</b>	<b>0.023</b>	<b>0.109</b>	<b>0.002</b>	<b>0.005</b>	<b>0.119</b>

150 Means along the same column with similar superscripts are not significantly (P>0.05) different  
 151 from one another

152 **SEM** = Standard error of means

153

## 154 Discussion and Conclusion

155 Species of termites were collected in selected sites of the study areas. *Macrotermes bellicosus* that  
 156 include workers, soldiers and queens were collected, collected termites were washed very well  
 157 with clean water and sun dried and ground into powder which was used for feed formulation.

158 Weight of broiler parts fed with commercial feed was significant (P<0.05) higher compared to  
 159 fishmeal and termite supplemented diet. However, weights of back, thigh, wings and gizzard were  
 160 not significantly (P>0.05) different in all treatments. Drum stick weight was similar in commercial  
 161 and termite supplemented diet, results showed that feeds were relatively consumed by broilers in  
 162 all treatment groups. This was similar to Paul and Sudipta (2011), who reported in his findings  
 163 that insects are used as animal feed in tropical and sub tropical regions and are the cheapest source  
 164 of animal protein. Bamphith and John (2012) also reported that nutritive potentials and utilization  
 165 of termites as poultry ingredients have been in use and documented. Results of carcass evaluation  
 166 followed the same trend of performance of experimental broilers. The results of this finding  
 167 demonstrated no significant (P>0.05) difference of the experimental broilers placed on different  
 168 diets, there seems to be relationship of weight of broiler parts, which agrees with Kwari *et al.*  
 169 (2008) findings who reported that, carcass and organs of measurement of broilers fed with  
 170 different meals most cases followed the same trend.

171 Nutrients and mineral composition of broilers was also observed at finisher stage in all treatment  
 172 groups. Result obtained showed no significant (P>0.05). This was similar with Bamphith and John  
 173 (2012) who reported that insects are capable of replacing fishmeal completely for growing

174 chickens. Agomuo (2011) also reported that, termites (*Macrotermes bellicosus*) a reproductive  
175 have good composition of moisture, crude protein, crude fibre, crude lipid, ash, carbohydrate and  
176 energy.

177 Conclusively, termites can serve as a good protein supplement in poultry feed, which can replace  
178 not only fishmeal but can also replace animal protein, soybean meal and can perform very well.

179

## 180 REFERENCE

181 Abd Rahman Jabir, M. D., Razak, S. A. and Vikineswary, S. (2012). “Nutrient potential and  
182 utilization of super worm (*Zophobas Mario*) meal in the diet of Nile tilapia (*Oreochromis*  
183 *niloticus*): Juvenile”. *African Journal of Biotechnology*, 11 (24): 6592 – 6598

184 Allotey, J. (2003). “Utilization of useful insects as food; improved Nutrition for Africa”. *Africa*  
185 *Journal of food, Agriculture, Nutrition and Development*. 3 (2): 1-8.

186 Bamphith, T. and John, C. M. (2012). Termites and Earthworms as potential alternative sources  
187 of protein for poultry. *Tiroesele and moreke, IJAVM*. 6(5): 368-376 Doi: 10. 5455/ijvms.  
188 174

189 Banjo, A.D., Lawal, O.A and Songonuga, E. A. (2006). “The nutritional Value of fourteen  
190 species of edible insects in South Western Nigeria”. *African Journal of Biotechnology*, 5:  
191 298-301

192 Grimaldi, D. and Engel, M.S. (2005). *Evolution of insects*. Cambridge University Press. ISBN 0-  
193 521-821149-5

194  
195 Kambhampati, S and Eggleton, P. (2000). Taxonomy and Phylogeny of Termites, In: Termites  
196 Evolution, Sociality, Symbiosis, Biology, Abe, T., Bigness, D. E. and Hu, M. (Ed).  
197 Kluver Academic Publishers, Netherlands, 1-23pp

198  
199 Mamman, A.B., J. O. Oyebanji and S. W. Peters (2000). In: Nigeria A people united, a future  
200 assured (survey) Calaba, Gabumo publishing Company. Pp 6-7

201 Malaka, S. L. O. (1996). *Termites in West Africa*. 1<sup>st</sup> Edition University of Lagos press, USA.,  
202 165pp

203  
204 Myles, T.G. (2003). Phylogeny of Taxonomy of the isopteran. [Internet], University of  
205 Toronto. Available from <http://www.utoronto.ca/forest/termite/speclist.htm> accessed

206 22<sup>nd</sup> february, 2016.

207

208 N'tukuyoh, A. I., Udiong, D. S., Ikpe, E. and Akpakpan, A. E. (2012). "Evaluation of Nutritional  
209 value of Termites (*Macrotermes Bellicosus*): Soldeirs, Workers and Queen in Niger Delta  
210 Region of Nigeria". *International Journal of food Nutrition and safety*, 1(2): 60-65.

211 Oguwike, F. N., Ebede, S. O., Offor, C. C. and Onubeze, D. P. M. (2013). "Evaluation of effects  
212 of consumption of termites (*Macrotermes nigeriensis*) on Biochemical and  
213 Heamatological profile of made albino Wister rates". *Greeener journal of medical  
214 Science*. 3 (9)1: 314-318

215 Paul, D, and Sudipta, D. (2011). "Nutrient content of sexual and worker forms of subterranean  
216 termites *Reticulitermes spp*". *Indian Journal of Traditional Knowledge*, 10: 505-507

217 Revindran, V., Blair, R. (1993). Food resource for poultry production in Asia and the pacific.  
218 *World's poultry science journal*. 49: 219-235

219  
220 Robberts, A. S. (1998). Broiler Management in warm Climate, Animal Science Annual Technical  
221 Bulletin 1: 40  
222

223 SAS. (2002). Stat View Statistical Package.English Version, SAS Inc. Newyork

224 Sogbesan, A. O. and Ugwumba, A. A. A. (2008). "Nutritional evaluation of termites  
225 (*Macrotermes sublyalinus*) meal as animal protein supplements in the diets of  
226 *Heterobranchus longifilis* (Valenciennes, 1940) Fingerlings". *Turkish Journal of fish and  
227 aquatic science*, 8: 149-157.

228

229

230

231