

2

3 **Theoretical Investigation into the Change in**

4 **the Number of Water Molecules in Solvent**

5 **Inaccessible Region of an Enzyme and**

6 **Enzyme-Substrate Complex**

7 **ABSTRACT**

8 **Background:** There may be dry enzymes, but water remains indispensable for the catalytic action of enzymes. There is not as much interest in how the presence of a drug such as aspirin and a psychoactive compound such as ethanol may affect the water-mediated role of the enzyme.

Objectives: The objectives of this research are: 1) To assess the changes in the number of water molecules interacting with the enzyme-substrate complex and the solvent inaccessible region of a protein, 2) to determine the free energy difference due to preferential solvation and hydration, and 3) to re-examine theoretical issues in literature and relate them to the interpretation of the results.

Methods: A major theoretical research and minor experimentation using Bernfeld method.

Results and discussion: The presence of ethanol/aspirin alone yielded only dehydration of the osmolyte inaccessible region and the enzyme substrate complex (*ES*). There was positive free energy difference ($\Delta\Delta G$) if the equilibrium constant for hydration change ($K_{eq}(1)$) > the equilibrium constant for folding-unfolding transition ($K_{eq}(3)$); it is negative where $K_{eq}(3)$ > $K_{eq}(1)$. Analysis of various models made them valuable for the interpretation of result

for feature application.

Conclusion

The change in the number of water molecules in an osmolyte inaccessible region of the enzyme and those interacting with the ES may be either positive or negative due respectively to sucrose and ethanol/aspirin. The spontaneity of two processes, hydration and folding-unfolding transition, the free energy difference, differs. The model for water stripping, preferential interaction concept, and the KBI for osmolation and hydration can guide the interpretation of the effects of any cosolute.

9 *Keywords: Porcine pancreatic alpha amylase; change of Gibbs free energy; change of the*
10 *number of water molecules; enzyme-substrate complex; osmolyte-inaccessible region of*
11 *enzyme; cosolutes; Kirkwood-Buff Integrals(KBI).*

12
13

1. INTRODUCTION

14 There are quite a lot of controversies surrounding the hydration of biomolecules. The
15 hydration of biomolecules is not in doubt but the effect of such hydration on internal
16 dynamics of the biomolecules is commonly of general interest to scientists [1]. However, this
17 does not exclude intermolecular dynamics needed for contact with each other or with other
18 solution components otherwise the needed contact for whatever transformation may not
19 occur; hence the proposition that enzymes must diffuse towards the substrate to align itself
20 with it to achieve a catalytic orientation [2]. This is notwithstanding current trend in the
21 development of immobilised enzymes, from amylase family. It must however, be made clear
22 that it is very impossible to digest polysaccharide without hydration of both substrate and
23 enzyme. A lot of interest has been shown in immobilised enzymes [3] for different reasons.
24 In those studies concern has been shown for the need for hydration, its purpose and effect
25 on the kinetic and thermodynamic stability of the enzyme. But there seem not to be much
26 interest on how the presence of drug such as aspirin and psychoactive compound such as
27 ethanol may affect the role of all kinds of hydration of the enzyme.

28 Some enzymes are known to possess conserved water molecules as part of the
29 structure of the enzyme's active site suggesting they play an important function in the active
30 site stability, flexibility, ligand coordination and residue positioning, hence their guided
31 evolutionary conservation [4]. Nuclear Magnetic resonance spectroscopy (NMR) analysis of
32 the hydration process indicates that the onset of catalytic activity is a direct consequence of
33 an increase in enzyme's (lysozyme's) conformational flexibility; it has been suggested that
34 this increased flexibility may be due, in part, to the reduced interaction of charged and / or
35 polar amino acid residues within the enzyme molecule caused by water's ability to effect
36 dielectric screening [5]. Yet there is objection against total reliance on flexibility for function
37 considering that enzyme activity can occur at very low hydration levels, coupled with
38 reduction in protein's flexibility; this according to [6] calls for a rethink regarding the dynamic
39 requirement for enzyme activity and stability.

40 In this research, the changes in the number of water molecules interacting with the
41 enzyme via its enzyme-substrate complex due to the presence of the additives that appears
42 not to feature very prominently in literature have become the concern of this research. In this
43 regard the view by Damien is relevant. Citing other workers, Damien et al [7] posits that
44 water strongly influences the structure and function of biomolecules within it. According to
45 them [7] the most relevant interactions are hydrogen bonds, a mainly local type of weak
46 bonding among water molecules which also exist between water and the polar or ionic
47 groups of the biomolecule; this is besides other long-range Coulomb forces, between
48 formally charged groups of the biomolecule and hydrophobic forces; the latter is relevant for
49 the aggregation of hydrophobic moieties and/or protein folding. It is obvious that the
50 relevance of water is accomplished through various forms of interaction that cannot preclude
51 interaction energy and solution structure in the presence of additives in particular.

52 It should be realised that the presence of cosolvent or cosolute can alter the effect of
53 aqueous solvent on the structure and function of the enzyme. The thermodynamic and
54 activation parameters in terms of energy associated with *ES* may not remain the same in the

55 presence of cosolvents, otherwise called osmolytes. The description of the interaction
56 requires mathematical models that will be briefly addressed in theoretical section while a
57 detailed qualitative aspect of theory is to be addressed in the discussion section as part of
58 interpretational goal. The objectives of this research are 1) To assess the changes in the
59 number of water molecules interacting with enzyme-substrate complex and solvent
60 inaccessible region of a protein, 2) determine the free energy difference due to preferential
61 solvation and hydration and 3) reexamine theoretical issues in literature and relate same to
62 the interpretation of results.

63 2. THEORY

64 To begin with there is need to state that the major motivation of this section is the
65 need to establish a justifiable theoretical background that can enhance the quality and
66 perhaps, the validity and serve as a basis for the generation and possibly the interpretation
67 of result. This section has two parts viz: The review of the derived equation related to
68 difference in interaction free energy and the changes in the number of water molecules
69 interacting with the enzyme substrate complex ($[ES]$); the second part is concerned with the
70 change in the number of water molecules in osmolyte-inaccessible regions. The equation [8]
71 adopted as in previous publication [9] in the quantitative determination of pair-wise solute-
72 solute interaction parameter is as follows:

$$73 \quad \ln[k_{(m_3)}/k_{(m_3=0)}] = \frac{2[g_{cx} - g_c^\#]m_3}{RTm_0^2} - N\phi M_1 m_3 \quad (1)$$

74 where $k_{(m_3)}$ is the (pseudo –) first – order rate constant in a reaction mixture containing co –
75 solute whose concentration is m_3 and $K_{(m_3=0)}$ is the rate constant in the absence of the co –
76 solute; R and T are the molar gas constant and thermodynamic temperature; m_0 is the
77 (hypothetical) ideal reference state and it is equal to 1 mol/kg; $g_{cx} - g_c^\#$ is the difference in
78 interaction Gibbs free energies between the co –solute c and the reactants β (and by
79 extension substrate and a biochemical catalyst) on one hand and the activated complex $\#$ on
80 the other hand; M_1 , ϕ , N and m_3 are the molar mass of water, practical osmotic coefficient for

81 the aqueous solution, the number of water molecules, and the molarity of the added cosolute
 82 respectively [8]. The equation seems to represent another way of expressing preferential
 83 interaction, a thermodynamic phenomenon applicable to multi-component solution. In the
 84 original equation by Buurma *et al* [8],

$$85 \quad RT \ln (k_{(m_3)}/k_{(m_3=0)}) = \Delta G(c)m_3 - N\phi RT M_1 m_3 \quad (2)$$

86 Where, R is the universal gas constant.

87 Thus,

$$88 \quad G(c)_{m_3} = \frac{2[g_{cx} - g_c^\#]m_3}{m_0^2} \quad (3)$$

89 The most important function of the enzyme is the lowering of activation energy and
 90 free energy of activation. Enzyme – substrate cannot proceed to product without initial
 91 activation which however occurs at a lower energy cost. Previous research attempted to
 92 apply this concept of pair-wise solution component interaction to biological system such as
 93 enzyme catalysed reaction in the presence of cosolute [9]. Here a more straight forward
 94 approach is further adopted to achieve similar result. If assay is at very high enzyme
 95 concentration, and if the substrate is not soluble, and if the raw insoluble starch was the
 96 substrate as in this research, a situation that satisfies the condition for reverse quasi steady
 97 state approximation (rQSSA) [10], then the equilibrium dissociation constant of the substrate
 98 from the complex given as $K_s = k_{-1}/k_1$ where k_{-1} and k_1 are the rate constant for the
 99 dissociation of enzyme-substrate complex (ES) and the 2nd order rate constant for the
 100 formation of the ES respectively, should be the case. But the concept is also applicable to a
 101 situation where the substrate concentration is very high such that $\frac{[E_0]}{K_M + [S_0]} \ll 1$
 102 ($[E_0]$, K_M , and $[S_0]$ are the total concentration of enzyme, is the Michaelis-Menten constant,
 103 and concentration of the substrate) as to satisfy the condition for standard QSSA (sQSSA)
 104 [10]. This takes the form $K_M = \frac{k_{-1} + k_2}{k_1}$ where k_2 is the rate constant for product formation and
 105 release. However, the key issue is that the rate constants for the dissociation of ES can be
 106 expressed respectively as

107
$$k_{-1} = K_s k_1 \quad (4)$$

108
$$k_{-1} = K_M k_1 - k_2 \quad (5)$$

109 In this research Eq. (4) unlike Eq. (5) does not present any issue because $\ln[k_{-1(m_3)}/$
 110 $\ln k_{-1(m_3=0)}]$ will always eliminate the need for the molar mass of the substrate. Besides,

111
$$k_1 = - \frac{\partial \ln([S](t)/[S]_0)}{\partial t [E_0]} \quad (6)$$

112 Where the concentration of enzyme ($[E_0]$) assayed is held constant or fixed while the
 113 concentration of the substrate in time $t = 0$ is $[S]_0$, and $[S](t)$ is the concentration of the
 114 substrate in time, t .


115
$$\ln(k_{s(m_3=0)}/(k_{s(m_3)})) = (\Delta G_{m_3} - \Delta G_{-1 m_3})/RT + M m_3 (\varphi_{-1} n_{-1} - n\varphi) \quad (7)$$

116

117
$$= \frac{\Delta \Delta G_{m_3}}{RT} + M m_3 \varphi \Delta n \quad (8)$$

118 It is not in doubt that Michaelis-Menten (MM) constant is a sum of equilibrium

119 constants. This is to say that it is given as $K_M = \frac{k_{-1}}{k_1} + \frac{k_2}{k_1}$; this implies that



121 The variable, S_{FR} as explained elsewhere [11], is the fragment of the polysaccharide left after
 122 a given catalytic cycle; no single polysaccharide is totally hydrolysed by an appropriate
 123 hydrolase.

124 The change in the number of water molecules in osmolyte-inaccessible regions,
 125 ΔN_w , is given by the slope of line relating $\ln(K_{eq})$ and the osmolyte concentration as follows
 126 [12].

127
$$\ln K_{eq} = - \frac{\Delta N_w [\text{Osmolal}]}{55.56} \quad (9b)$$

128 Leading to Eq. (9b) from the perspective of osmolyte-inaccessible regions is simply
 129 reaffirmation of the principle of preferential exclusion anchored on Kirkwood-Buff theory
 130 [KBT] of solution structure that has been popularised in recent papers [13, 14]. The
 131 theoretical interest arises from what appears to be a common ground for Eq. (2), Eq. (8), and

132 Eq. (9) in that the number of water molecules for different purposes can be calculated from
133 all equations, one from the slope (Eq. (9)) and the other from intercept of either Eq. (2) or
134 Eq. (8). Equation (8) which arises from theoretical exposition of Buurma *et al* [8] represented
135 the first time observation was made of the appearance of variable - concentration of cosolute
136 - in two places as an independent variable in an equation.

137 Soluble polar organic substances called osmolytes may be excluded from the
138 protein surface domain on account of their inability to penetrate protein's inner region. This
139 issue is important in the light of the fact that solvent accessibility change plays a critical role
140 in protein misfolding and aggregation, the culprit for several neurodegenerative diseases,
141 including amyotrophic lateral sclerosis (ALS) [15]. Furthermore, solvent accessibility may be
142 part of the structural environment of amino acids in the protein that might influence the
143 function-structural (mechanical) and catalytic in nature-of any of such amino acids [16].
144 Directly or indirectly, this important issues may have prompted research in this issue of
145 inaccessible core of the protein as exemplified in very recent research outcome which
146 showed that the solvent-inaccessible cores of the three classes of proteins are equally
147 densely packed [17]; this constitute steric hindrance to the penetration of relatively large
148 organic osmolytes. This may have promoted excess flexibility that caused increasing velocity
149 of hydrolysis with higher concentration of ethanol. One must not fail to point out that
150 osmophobic concept [18] has been advanced as basis for the action that compels a protein
151 to fold due to exclusion of such osmolyte which exist in nature from protein back bone.

152 On account of the issues raised in the text, there is need to recall that preferential
153 osmolation, either negative or positive, can yield (re) folding and unfolding as the case may
154 be leading to equilibrium state if a two-state model is assumed. Hence, the equilibrium
155 constants (K_{eq}) defined mathematically and given below are of paramount relevance to a
156 system in near dynamic equilibrium.

157
$$K_{eq(3)} = \frac{[U]}{[N]} \quad (10)$$

158 Equation (10) is adapted from the work by Pace [19] which the author restate as $K_{\text{eq}(3)} =$
159 $U/1 - U$ where in this case U and $1-U$ denotes fraction of unfolded protein molecular
160 population and fraction of folded protein respectively. Equation (10) or its alternative is
161 expressible in two ways in accordance to whether or not the observed catalytic activity of the
162 enzyme in the presence of osmolyte is greater than the same activity in the absence of the
163 osmolyte. The equations which are applicable to the effects arising from the presence of an
164 osmolyte are to be stated in method subsection. But there is also preferential hydration and
165 dehydration due to preferential exclusion and binding of appropriate osmolyte respectively.
166 This creates directional aqueous molecular motion to and from the protein's surface domain
167 leading to an equilibrium system described by the second equation of equilibrium constant
168 given elsewhere [20] as

$$169 \quad K_{\text{eq}(1)} = \exp\left(-\frac{\ln a_1 C_1}{C_3} \Delta\Gamma_{23}\right) \quad (11)$$

170 Where $\Delta\Gamma_{23}$ is the change in preferential interaction by either binding or relative exclusion of
171 an osmolyte; C_1 and C_3 are molar concentrations of water and osmolyte respectively; a_1 is
172 the activity of water in aqueous solution of osmolyte. Meanwhile the equation of preferential
173 interaction [14] is given as

$$174 \quad \Delta\Gamma_{23} = \frac{\ln K_{\text{eq}(3)}}{\ln a_3} \quad (12)$$

175 The emergence of Eq. (12), as in previous publication (14), is as a result of the proposition
176 that a parameter cannot be a devise-based measurable quantity (without definite or finite
177 magnitude) as well as a constant quantity. A measurable quantity is an extensive
178 thermodynamic parameter and, if a given ratio is always constant regardless of the
179 magnitudes of the compared parameters, it becomes an intensive thermodynamic quantity.
180 The report at the web site, en.Wikipedia.org (<https://www.en.Wikipedia.org>) shows that
181 Richard C. Tolman was the author who first introduced the concept of extensive and
182 intensive quantities.

183 **3. MATERIALS AND METHODS**

184 **3.1 Material**

185 As stated elsewhere[9], the materials are The chemicals used were: The chemicals
186 used were: Sucrose (St Lious France); raw (native) potato starch (Sigma Chemicals Co, USA);
187 ethanol, hydrochloric acid and sodium chloride (BDH Chemical Ltd, Poole England); 3,5-
188 dinitrosalicylic acid (DNA) (Lab Tech Chemicals, India); Tris (Kiran Light Laboratories, USA);
189 porcine pancreatic alpha amylase (EC 3.2.1.1) (Sigma, Adrich, USA); all other chemicals were of
190 analytical grade and solutions were made in distilled water. Aspirin was purchased from CP
191 Pharmaceuticals Ltd, Ash road North, Wrexham, LL 13 9UF, and U.K

192 **3.2 Equipment**

193 *pH* meter (tester) from Hanna Instruments, Mauritius; electronic weighing machine
194 from Wensar Weighing Scale Ltd, Chennai; Centrifuge, 300D model from China; 721/722
195 visible spectrophotometer from Spectrum Instruments Co Ltd, China.

196 **3.3 Method**

197 As stated elsewhere [20], 0.01 g of PPA was dissolved in 20 mL of distilled water to give
198 500 µg/L while potato starch was prepared by dissolving 1 g in tris-HCl buffer (aq.) buffer (90
199 mL), 5 mL, 6% (W/W), NaCl (aq.) and 5 mL distilled water to give 1 g/ 100 mL. Approximate
200 dilutions were carried out for the determination of K_M and v_{max} at 37⁰C and pH of 7.4 by
201 Lineweaver-Burk plot [21]. As in previous investigation [14] a method adopted for the
202 determination of velocity (v) where $C_3 \rightarrow 0$ is a re-modification of the same equation found in
203 literature [22]. It may appear theoretical but that is the essence of this research, a combination of
204 a major theory and minor experimentation. Increasing v with increasing C_3 , demands a plot of log
205 v versus C_3 which gives an intercept, being an extrapolated velocity ($v_{C_3 \rightarrow 0}$) of hydrolysis as C_3
206 $\rightarrow 0$.

207
$$\log v = \log v_{C_3 \rightarrow \infty} - \beta [C_3]. \quad (13)$$

208 Equation (13) is for the increasing v , while for the decreasing case, it is given as

209
$$\log v = \log v_{C_3 \rightarrow \infty} - \beta / [C_3]. \quad (14)$$

210 However, in order to obviate the effect of outliers, linear regression line was allowed to link
211 the lowest point with the highest point for the determination of the minimum v as $C_3 \rightarrow 0$.
212 Assay for the generation of velocities of the hydrolysis of starch is according to Bernfeld
213 method [23].

214

215 The equation (Eq. (15)) below is explored for the purpose of comparing the transition state
216 energies of two different equilibrium systems dehydration/hydration and
217 osmolation/exclusion equilibria due to the presence of osmolytes or cosolvents; it is
218 therefore, restated as

$$219 \quad \Delta\Delta G_{ES}^{\#} = -RT \ln \frac{K_{eq(1)}}{K_{eq(3)}} \quad (15)$$

220 The equilibrium constant $K_{eq(1)}$ is determined by substituting relevant parameters into Eq.
221 (11); $K_{eq(3)}$ is determined by exploring either Eq. (16) or Eq. (17) below.

$$222 \quad K_{eq(3)} = \frac{V_N - V_{OBS}}{V_{OBS} - V_{MIN}} \quad (16)$$

223 Where $V_N > V_{OBS} > V_{MIN}$ and the subscripts, N, OBS, MIN are respectively, catalytic activity
224 of native enzyme, observed activity of treated enzyme and minimum activity of treated
225 enzyme.

$$226 \quad K_{eq(3)} = \frac{V_N - V_{MIN}}{V_{MAX} - V_N} \quad (17)$$

227 The issues that led to the emergence of Eq. (16) and Eq. (17) were addressed in part
228 elsewhere [24]. Further details are currently in manuscript under preparation.

229 **3.4 STATISTICAL ANALYSIS**


230 The velocities of hydrolysis were determined in triplicates. The mean values were
231 used to determine the equilibrium constant. Microsoft Excel (2007) was used to plot the
232 dependent variable versus independent variable.

233 **4. RESULTS AND DISCUSSION**


234 Before analysis and discussion of results, there is a need to review Eq. (2) and Eq.
235 (8). A careful examination of the equations reveals that the slope and intercept may not be

236 positive or negative; the values depend on the magnitude of the ratio given as $K_{(m_3)}/K_{(m_3=0)}$.
237 If $K_{(m_3=0)} < K_{(m_3)}$, with increasing values of $K_{(m_3)}$, the value of the ratio should be decreasing
238 from smaller negative values to larger negative values, such that a plot of $\ln(K_{(m_3)}/K_{(m_3=0)})$
239 versus m_3 should give a negative slope and definitely a negative intercept. It is also
240 probable that $\frac{K_{(m_3)}}{K_{(m_3=0)}} > 1$ such that any plot may give a positive slope or correlation and
241 either a positive intercept or intercept which is negative but much smaller in magnitude.
242 Therefore, characteristics such as the magnitudes and signs of the slope and intercept of a
243 straight line from the plot express the type of change in the number of water molecules,
244 which is either net hydration (positive) or net dehydration (negative). These account for the
245 shapes of various curves shown in Figure 1 through 6. To show the direction of shift in the
246 hydration process in the equilibrium, $E + S \rightleftharpoons ES$, a plot of Δn versus m_3 was carried out.
247 The result (Fig. 1) shows that there was a decreasing trend in the change in the number of
248 water molecules interacting with the ES . The observed trend is due to the effect of aspirin.
249 The decreasing trend along the positive axis suggests that there was a decrease in
250 hydration due to the effect of aspirin alone (Fig. 1). It is a loss-dehydration- the magnitude of
251 which showed a decreasing trend, progressing towards net hydration (Fig. 2) due to the
252 effect of the second cosolute, sucrose. In this case, the variation of the change in the
253 number of water molecules with the molar concentrations of sucrose showed mixed trend.
254 With a lower concentration of the cosolute (1.55 mmol/L and 0.73 mmol/L) – aspirin – there
255 was an increasing trend unlike with higher concentration of the same cosolute, due perhaps,
256 to the effect of the 2nd cosolute (sucrose) in the reaction mixture (Fig. 3).


257



258
 259 **Fig. 1. Variation of the change in the number of water molecules interacting with the**
 260 **enzyme- substrate complex with different concentration of aspirin. n is the number of**
 261 **water molecules.**
 262



263
 264
 265
 266 **Fig. 2. Variation of the change in the number of water molecules interacting with**
 267 **enzyme substrate complex with different [Aspirin] at different fixed concentration of**
 268 **sucrose whose concentration ranges between 3.60-57.53 mmol/L. n and SUC denote**
 269 **the number of water molecules and sucrose.**
 270



272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288


289

290

291

Fig. 3. Variation of the change in the number of water molecules interacting with the enzyme-substrate complex with different concentration of sucrose (SUC) at different fixed concentration of aspirin. $n(1)$ and ASP denote the number of water molecules and aspirin respectively.

Like the trend observed with the effect of aspirin there is also a decreasing trend in the positive values of Δn with increasing concentration of ethanol (Fig.4). Variation with different concentrations of ethanol exhibited similar trend observed for the variation of Δn with molar concentration of aspirin (Fig.5). With a mixture of ethanol and sucrose, there was, as was the case with the effect of a mixture of aspirin and sucrose, a mixed trend in the variation of Δn with molar concentrations of sucrose (Fig. 6). All these observation notwithstanding, it is rather difficult to suggest why such observations cannot be mere coincidence taking into account the effect of high degree of improvisation in the conduct of the experiment. It is not an overemphasis to opine that ethanol is totally different from aspirin; while the former is essentially psychoactive, the latter is a well known non-steroidal anti-inflammatory drug [25, 26], and both have adverse effects on intestinal brush border membranes that could compromise the biological function of brush border membrane enzymes and transporters respectively.



292

293

294

295


296

297

298


Fig. 4. Variation of the change in the number of water molecules interacting with enzyme substrate complex with different concentration of ethanol. $n(1)$ and ETH denote the number of water molecules and ethanol respectively.

299
300



301
302
303
304
305
306

Fig. 5. Variation of the change in the number of water molecules interacting with enzyme- substrate complex with different [Ethanol] at different fixed concentration of sucrose. *SUC* and *n(1)* denote sucrose and number of water molecules respectively.



307
308
309
310
311
312
313
314
315

Fig. 6. Variation of the change in the number of water molecules interacting with enzyme-substrate complex with different [Sucrose] at different fixed concentration of ethanol. *ETH* and *n(1)* denote ethanol and number of water molecules respectively.

Table 1. Change in the number of water molecules in osmolyte-inaccessible regions due to ethanol-sucrose mixture

As function of sucrose concentration			
[Ethanol]/mol/L	1.247	3.228	5.279
ΔN_w	-601.667	615.05	3809.19

r^2	0.817		0.533		0.766	
As function of ethanol concentration						
[Sucrose]/mmol/L	0.00	3.57	7.14	14.29	28.57	57.14
ΔN_W	-34.56	-50.06	-69.26	-36.00	-9.06	-21.11
r^2	0.855	1 (2dpts)	0.954	0.943	0.948	0.599

316 *Changes in the number of water molecules (ΔN_W) are calculated as the product of slope and*
317 *55.56; the slope may be obtained from the plot of $\ln K_{eq}$ versus [cosolute] at a fixed*
318 *concentration of the 2nd cosolute.*

319
320 The change in the number of water molecules (ΔN_W) on osmolyte inaccessible
321 region as a function of sucrose concentration is similar to the exclusion of aqueous solvent
322 or dehydration with lower concentration of ethanol unlike with higher concentration ethanol in
323 which there was hydration (Table 1). But as function of the concentration of ethanol, there
324 was irregular trend couple with a case of dehydration similar to result obtained in only
325 ethanol treated enzyme (sucrose concentration = 0) (Table 1). This is not unexpected
326 considering ethanol as a fluidising and water-stripping agent.

327

328 **Table 2. Change in the number of water molecules in osmolyte-inaccessible regions**
329 **due to aspirin-sucrose mixture**

As function of sucrose concentration						
[Aspirin]/mmol/L	0.76		3.05		6.10	
ΔN_W	2042.86		4380.01		6261.61	
r^2	0.978		0.965		1 (2 dpts)	
As function of aspirin concentration						
[Sucrose]/mmol/L	0.00	7.19	14.38	28.76	57.75	
ΔN_W	-40625.47	-80839.8	-83895.60	-52854.23	-44925.82	
r^2	0.865	1 (2dpts)	1 (2 dpts)	0.993	0.982	

330 *Changes in the number of water molecules (ΔN_W) are calculated as the product of slope and*
331 *55.56; the slope may be obtained from the plot of $\ln K_{eq}$ versus [cosolute] at a fixed*
332 *concentration of the 2nd cosolute; dpt stands for data point.*

333
334 The change in the number of water molecules as a function of sucrose concentration
335 showed increasing trend unlike such change as a function of aspirin concentration in which
336 there was irregular trend and negative in sign as to imply dehydration (Table 2) similar to the
337 result obtained due to the effect of aspirin alone. This may imply that aspirin like ethanol has
338 water-stripping properties.

339 The difference in free energies between two thermodynamic processes
 340 dehydration/hydration and osmolation/exclusion arising from the effect of cosolutes and
 341 water are recorded in Tables (3a), (3b), (4a), and (4b). There is need to state that the data
 342 generated is not an outcome of high precision measurement as a result of improvisation.
 343 This leaves room for further research using state-of-the-art facilities while the current data
 344 remains purely illustrative of the fact and principle enunciated in this research. Usually, a
 345 spontaneous process is one in which the free energy is relatively large and negative in sign.
 346 The effect of ethanol and aspirin separately alone, yielded a mixed result of negative and
 347 positive free energies as shown in Tables (3a) and (4a) respectively. The negative difference
 348 in free energy occurred with higher concentration of the cosolutes. With a mixture of ethanol
 349 and sucrose (Table 3b) and a mixture of aspirin and sucrose (Table 4b), the negative values
 350 occurred with higher concentration of ethanol and aspirin. What one can deduce is that
 351 positive $\Delta\Delta G$ occurs if $K_{eq}(1) > K_{eq}(3)$ and as such (de) hydration is more spontaneous. On
 352 the other hand if $K_{eq}(1) < K_{eq}(3)$, a negative $\Delta\Delta G$ may be given with the result that,
 353 osmolation/exclusion is more spontaneous.

354 **Table 3a. Difference in free energies between dehydration/hydration and**
 355 **osmolation/exclusion with only ethanol.**

[Ethanol]/mol/L	1.247	2.398	3.228	4.311	5.279
$\Delta\Delta G$ /kJ/mol	18.68	0.96	-0.38	-0.36	-0.28

356 *$\Delta\Delta G$ is the difference in free energy.*

357

358 **Table 3b. Difference in free energies between (de)hydration and (de) osmolation with a**
 359 **mixture of ethanol and sucrose**

[Ethanol]/mol/L	[Sucrose]/mmol/L				
	3.57	7.14	14.29	28.57	57.14
	$\Delta\Delta G$ /kJ/mol				
1.247	3.53	4.22	2.43	2.42	1.81
3.228	-1.92	-0.49	0.12	0.28	0.64
5.279	-	-11.35	-4.84	-0.95	4.44

360 *$\Delta\Delta G$ is the difference in free energy.*

361

362

Table 4a. Difference in free energies between dehydration/dehydration and
osmolation/exclusion with only aspirin.

[Aspirin]/mmol/L	0.76	1.53	3.05	4.58	6.10
$\Delta\Delta G$ /kJ/mol	2.63	2.49	1.84	-2.85	-9.55

363 *$\Delta\Delta G$ is the difference in free energy.*

364

365

366

Table 4b. Difference in free energies between dehydration/hydration and
osmolation/exclusion with a mixture of aspirin and sucrose

[Aspirin]/mol/L	[Sucrose]/mmol/L			
	7.19	14.38	28.57	57.14
	$\Delta\Delta G$			
0.76	6.03	7.39	9.11	9.80
3.05	-4.29	-3.61	0.74	1.49
6.10	-	-	-7.76	-5.50

367 $\Delta\Delta G$ is the difference in free energy.

368 The results obtained so far are significant because of the biological role of water,
369 and, there are a lot of theoretical basis for them. Such theoretical foundation or basis
370 broadens the scope for further research. Beginning from what is known is the fact that
371 proteins are strongly hydrated in aqueous medium. The density of water molecules close to
372 the protein surface due to effect of polar and non-polar groups is as high as 1.25 g/mL within
373 3-4.25 Å of protein surface, mainly as result of large number of water molecules that are
374 3.75 Å from non-polar atoms; within 2.5 Å of the protein surface there is a small increase in
375 density of water molecules due to electrostriction around the polar groups; but 3-4.5 Å from
376 the surface, there is a slight decrease in density [27]. Water molecules are clustered
377 perpendicular to the protein surface while in the parallel direction to the protein surface the
378 water molecules are more disperse [27]. This means that given suitable pH, an enzyme
379 exhibits a level of hydration needed for function. For a particular group, the fraction of time
380 when a water protein hydrogen bond is formed otherwise called hydrogen bond probability
381 (P_{hb}) is strongly dependent on protein accessible surface area (ASA). The lower the latter,
382 the higher the entropic barrier (cost) that should be paid to significantly reduce the flux of
383 water molecules on the protein surface hydration site where H-bond is expected [28]. In the
384 same vein, Ooi & Oobatake [29] also posited that each atomic group interacts with water in
385 proportion to its water-ASA. The effect of the presence of chaotropes is of major concern as
386 it has been observed that more polar organic solvents (tetrahydrofuran and acetonitrile)
387 replace mobile and weakly bound water molecules in the active site and leave primarily the
388 tightly bound water in that region [30].

389 In this research, aspirin and ethanol decreased the velocities of amylolysis of raw
390 potato starch. These velocities under the influence of aspirin and ethanol are respectively
391 21-74 U/mL and 38 – 61 U/mL, all being < the velocity of amylolysis (97 U/mL) by the
392 untreated enzyme. This means that the entropic cost of fixing water of hydration increased
393 as a consequence. This can be elucidated via the equation made popular by Petukhov [28].
394 The equation is

$$395 \quad \Delta G_{\text{hb}} = -RT \ln \frac{P_{\text{hb}}}{1 - P_{\text{hb}}} \quad (18)$$

396 Equation (18) suggests that as $P_{\text{hb}} \rightarrow 1$, the value of the free energy of hydration tends
397 toward higher negative magnitude as an expression of feasibility or spontaneity of hydration.

$$398 \quad P_{\text{hb}} = \frac{\exp\left(-\frac{\Delta G_{\text{hb}}}{RT}\right)}{1 + \exp\left(-\frac{\Delta G_{\text{hb}}}{RT}\right)} \quad (19)$$

399 Hence if $P_{\text{hb}} \rightarrow 1$ (or if it's equivalent 100 $P_{\text{hb}} \rightarrow 100\%$), the entropic cost should tend to
400 minimum. It is important to point out that the value of P_{hb} may be a function of the fraction of
401 water population that can form H-bond with 4 water molecules and 3 water molecules both of
402 which are a function of the prevailing temperature in a equation given according to Petukhov
403 [28] as

$$404 \quad P_{\text{hb}} = 100(4X_1 + 3X_2)/4 \quad (20)$$

405 Where X_1 and X_2 are respectively the fraction of water that can H-bond with 4 and 3
406 molecules of water. As this research shows, the presence of sucrose seems to have
407 opposed the effect of aspirin and ethanol. As such it is expected that P_{hb} may have
408 increased as a result of the effect of sucrose.

409 The role of water, or the effect of hydration, has its theoretical foundation that
410 enhances the interpretation of results. It has been reported that “the hydration environment
411 of a protein significantly affects its dynamics. This is why changes in the number of water in
412 cosolvent inaccessible site of the protein have become very important because such can
413 affect enzyme function. A positive change indicates that there may have been hydration and
414 negative change means the opposite. Such changes may not have been possible if there

415 was no initial hydration and preferential interaction with molecules of water. Although the
416 method adopted by Mitchell and Litman [12] and Buurma [8] are different they have a
417 common ground for addressing the issue of hydration changes. This is the case because
418 osmolyte inaccessible region of the protein may accommodate the active site. The active site
419 is either located within the protein's inner part or at locations close to the surface domain of
420 the enzyme. Hence changes in the number of water molecules in an osmolyte inaccessible
421 region cannot totally exclude the active site. This is the case, if one recalls that *ES* complex
422 is the result of complex formation between substrate and active site of the enzyme.
423 Therefore, there could be changes in the number of water molecules interacting with *ES*.

424 In literature, following the application of osmotic stress, is the observation that
425 protein-DNA complex can be hydrated with measurable volume changes. The sign of the
426 changes of the number of water molecules interacting with the protein and *ES* as well as
427 osmolyte inaccessible region indicates the occurrence of either hydration or dehydration. As
428 in this research such change occurs when a cosolute is introduced into the medium. It has
429 been observed that the catalytic activity of lyophilised oxidative enzyme was lower when
430 directly suspended in organic solvents containing little water than when they are introduced
431 into the same largely nonaqueous media by first dissolving them in water and then diluting
432 with anhydrous solvents [31]. Despite the need for water for maximum catalytic activity of
433 enzymes, an obvious paradox exists to the effect that, some enzymes (subtilisin and alpha-
434 chymotrypsin) showed a 100 billion-fold enhancement in nonpolar solvent like octane with
435 just an amount of water much less than needed to form a monolayer [32]. This is attributed
436 to an increase in the kinetic barrier (activation energy) needed to be overcome in order to
437 transform from native to unfolded conformation [32]. This should not be surprising because
438 unlike polar solvents, *e.g.* ethanol and polar solute, *e.g.* aspirin in this research, that have
439 water-stripping power, octane does not being hydrophobic.


440 It is quite obvious that infinitesimal amount of water is needed to trigger catalytic
441 action as to imply that water may be described as a *prima facie* example of an inorganic

442 catalyst. Highly structured water molecules are needed around the protein surface as part of
443 efficient chemistry of the protein by which they promote the protein's three dimensional (3-D)
444 structures [33]. According to Csermely [34], water molecules within the region of perturbative
445 influence of the enzyme provides the environment by which fluctuating changes in hydrogen
446 bond could occur as a necessary requirement for protein flexibility, structural
447 rearrangements leading to conformational transitions needed for catalytic functions. This
448 very much in agreement with the observed hydration induced conformation and dynamic
449 changes which are completed just before the onset of enzyme biological function [35]. It
450 goes to confirm that an increased rigidity in the protein at low hydration can be reversed
451 when water is added to the dry enzyme leading to a "loosening up" or increase in flexibility
452 [35]. Protein flexibility means inter-domain and catalytic site mobility made possible waters of
453 hydration. The deduction one can make is that polar solvent like ethanol as in this research
454 displaces the weakly bound structural water molecules and preferential water of hydration
455 leading to alteration and distortion in the catalytic conformational transition needed for
456 function that culminate in lower velocity of amylolysis.


457 However, the hydration effects are strongly dependent on both temperature and
458 hydration. At cryogenic temperatures, hydration stiffens protein structure because of the
459 hydrogen-bond interaction, whereas at physiological temperatures, hydration softens the
460 structure through the activation of anharmonic motion"[1]. The hydration water dynamics and
461 their dynamical coupling with the protein are presumed to be essential for protein dynamics
462 and biological function [1]. The protein dynamics in question is actually intra-molecular
463 dynamics needed for conformational flexibility for function. According to Chaplin
464 (www1.lsbu.ac.uk), proteins are characterized by conformational flexibility, which entails a
465 wide range of hydration states, in a state of dynamic equilibrium, facilitated by the ease of
466 hydration. The ease of hydration is dependent on the activity of the surrounding water
467 molecules. The enzymatic function of the enzyme is dependent on the position of the
468 equilibrium, $es=cs$ (where es and cs mean the expanded state and compact state of water

469 respectively) around the protein; the es is also called the Ih-type with lower density-the low
470 density water (LDW) while cs is called II-type with higher density-the high density water
471 (HDW). The LDW and HDW are respectively more ordered and less ordered. Thus an
472 intermediate mixture of nonionic kosmotropes and nonionic chaotropes such as sucrose and
473 aspirin/ethanol respectively as in this research can enhance biological activity of the
474 enzyme: It is neither an excessive rigidity nor an over flexibility of the structure of the protein
475 that can enhance the function of an enzyme.

476 The effect of a mixture of protecting and destabilising cosolute as observed in this
477 research has its theoretical foundation. Beginning with aspirin and ethanol, the theory is
478 described as preferential interaction by osmolation or by binding and by exclusion. The
479 former, according to Timasheff [36], leads to water stripping according to the equation:



481 Where, L is the ligand otherwise called cosolute. Citing his previous paper, Timasheff [36]
482 posits that “the reference state is the protein dissolved in water, in which it is fully hydrated.
483 Therefore, in a binary solvent, the binding of the nonaqueous solvent component to any
484 locus must displace water, *i.e.*, binding is an exchange reaction” [36]. Unfortunately there is
485 no equation for exclusion of ligand as at this moment. Nonetheless, the following equation
486 may serve this purpose.



488 Equation (22) (which reflects only the expulsion of water) symbolically shows that when an
489 aqueous solution of a protein is introduced into a stabilising cosolute, the macromolecule will
490 not be totally free from the molecules of the cosolute. Hence β is the small portion of L
491 (stabilising osmolyte in this case) that binds while $L - \beta$ is the vast amount of L that is
492 excluded given that β is equal to the amount of water displaced. This could have given rise
493 to $L + E \cdot nH_2O \rightleftharpoons E \cdot (n - \beta)H_2O \cdot \beta + \beta H_2O + L - \beta$. However, one must not overlook the effect
494 of osmotic stress that might be created when any cosolute is excluded leaving the
495 surrounding of the protein more concentrated as to create concentration gradient. This can

496 compel loosely bound water molecules to depart the protein into the bulk; this may also be
497 interpreted as a translational entropy gain of the aqueous solvent that drives re-folding [37].
498 This is in agreement with the view that osmotic pressure controls the activity of water in an
499 aqueous compartment inaccessible to neutral solutes (osmolytes). The osmotic stress
500 created then induces the release of bound water from macromolecules into bulk solvent.
501 Macromolecular conformations are thus shifted toward the state with the smallest volume,
502 which is the state with the least amount of bound water [38, 39]. The folded state promoted
503 by stabilisers such as sucrose in this research has smaller volume. This is another evidence
504 of the importance of hydration, be it water of hydration or water of preferential hydration.
505 Osmotic stress will always occur when there is the presence of a stabiliser in particular.

506 Further evidence is the observation about individual, internal water molecules that
507 may be reactants in a catalyzed reaction and/or may be integral parts of a protein structure,
508 providing stereospecific interactions; the correlation between hydration and increased
509 activity means that it is likely that the observed hydration-induced dynamical changes may
510 facilitate activity [6]. However, the presence of amyolytic activities, in the absence of intra
511 molecular motion, indicates that the motions are not an absolute requirement; this seems to
512 imply that if enzyme activity can occur at very low hydration levels, and if at these levels
513 protein flexibility is reduced, then the dynamic requirement for enzyme activity and stability
514 may be questionable [6]. Here one must strongly disagree on account of this research
515 outcome and on the basis of common sense. The lower amyolytic activity of ethanol/aspirin
516 treated-enzyme suggest that the structural water within the protein core and active site in
517 particular may be weakly intact, but other catalytically supportive mobile water molecules
518 may have been stripped off. In other words the ubiquitous surface hydration shell has
519 influence on protein dynamics and function such that if adversely affected by the chaotropes
520 [6], a reduction in amyolytic activity becomes inevitable. This clearly explains the decrease
521 in the velocity of amyolysis for ethanol/aspirin treated enzyme. Scholars who are only
522 interested in balanced diet may consume the usual diet thrice without water for two days but

523 with implication of being inflicted with indigestion and constipation despite the fact that ab
524 initio, the gastrointestinal tract is not dry. No matter the length of time no reaction can be
525 noticed in a dry mixture of enzyme and its substrate.

526 This can be accounted for in terms of Kirkwood-Buff theory (KBT) of solution
527 structure which states that the average structure of all solutions [40] is given by radial
528 distribution function ($g_{\alpha\beta}(r)$) between two species, namely, α and β . The term, radial
529 distribution function, is a measure of the deviation from the random distribution of particles of
530 type β from a central particle of type α as a function of the distance (r) from the central
531 particle [40]. A positive or negative deviation of $g_{\alpha\beta}(r)$ (also known as pair correlation
532 function) from unity, at a certain distance corresponds to excess or deficit of β at the
533 indicated distance from the particle designated as ' α '. The issue remains the combined
534 effect of aspirin/ethanol and sucrose.

535 According to Bolen and Baskakov [41], the osmophobic effect of osmolyte is a vital
536 property that is beneficial to life, being the capacity for an unfavourable interaction between
537 the osmolyte/cosolute and peptide backbone. In the same vein, Baskakov and Bolen [42]
538 opined that the osmophobic effect of stabilisers on the peptide back bone made the unfolded
539 state of protein in osmolyte solution very unfavourable relative to the folded state; therefore,
540 it was the strongly destabilising effect of stabilisers such as sucrose on the unfolded state as
541 in this research, that forces the enzyme to refold. From the perspective of thermodynamic
542 stability, Bolen and Baskakov [41], see solvophobic action which Schellman [43] and Rösger
543 *et al* [40] called excluded volume action, as a factor which raised the free energy of the
544 denatured state, shifting the equilibrium in favour of the native state. In this research sucrose
545 is a well known stabiliser which acts by preferential exclusion. On account of this sucrose
546 was able to enhance the amylolytic velocities of sucrose treated-enzyme in a reaction
547 mixture containing aspirin (3.052 mmol/L) and ethanol (3.228 mol/L): The velocities ranges
548 from 132-140 and 116-136 U/mL respectively. These values were higher than values

549 obtained for the untreated native enzyme (97 U/mL), only ethanol-treated (102 U/mL) and
550 only aspirin treated (69 U/mL) enzyme.

551 The phenomena of solvophobic and solvophilic effect are the root cause of the
552 change of biological function-either an increase or decrease in the velocity of catalytic action
553 for instance as noted in this research. Osmolytes (as cosolvents/cosolute) may be
554 solvophobic or solvophilic (preferential exclusion or osmolation *i.e.* preferential interaction by
555 binding) which causes respectively refolding and unfolding; this presupposes changes in the
556 volume or 3-D structure of the macromolecule. This needs interpretational analysis based on
557 what Rösgen *et al* called inverse KBT. It is usually a context between solvation and hydration
558 change expressed via the KB integrals (KBIs). From the point of view of preferential
559 hydration integral, the following equation is inevitable. The partial molar volume of the
560 protein is in contention. Thus,

$$561 \quad \Delta_N^D(G_{pw}) = -\Delta_N^D V_{prot} + \frac{m\phi_{os}}{RT} \quad (23)$$

562 Where $\Delta_N^D V_{prot}$ is the partial molar volume of the protein, ϕ_{os} is the volume fraction of the
563 osmolyte, G_{pw} is the KBI for hydration, m is the short form of m -value, the capacity of
564 osmolyte to cause conformational change and Δ_N^D means folded to unfolded transition. For
565 the ideal case as may be applicable to dilute solution of sucrose, positive m -value for the
566 protecting osmolyte, should be such that $\Delta_N^D(G_{pw})$ may be positive in sign as to imply an
567 increase in the number of water molecules around the protein. This view is premised on the
568 fact that the (re)folded state has smaller hydrodynamic radius than the unfolded which is
569 also more hydrated [40]. The implication of this premise is that $\Delta_N^D V_{prot}$ being small, implies
570 that its negative magnitude ($-\Delta_N^D V_{prot}$) may be small. The outcome is that the right hand side
571 (RHS) may be large and positive. It must be made clear that $\Delta_N^D(G_{pw})$ needs to be
572 determined but it remains outside the scope of this research. This view explains the effect of
573 sucrose which promotes initial preferential hydration of the enzyme before other

574 physicochemical events such osmotic stress effect due to concentration gradient created by
575 the excluded osmolyte.

576 The equation for the osmolation case is given as

577
$$\Delta_N^D(G_{po}) = -\Delta_N^D V_{prot} - \frac{m(1-\phi_{os})}{RT} \quad (24)$$

578 In this research, aspirin was noticed to have reduced the velocity of amylolysis as to imply
579 that the enzyme was destabilised. In this case, the m -value may be large and negative; there
580 may be a positive value of $\Delta_N^D(G_{po})$. This is to say the number of water molecules around the
581 protein decreases while there is a relative excess of the osmolyte around the protein surface
582 domain including some molecule that may have penetrated into the protein 3-D structure.

583 A very important deduction one can make is that hydration and dehydration are
584 merely precedent to the initial events, preferential exclusion and binding respectively.
585 Otherwise, upon unfolding due to osmolation, maximum hydration may occur as expected
586 for the unfolded protein [40]; this is clearly the case because if there is excess of the
587 osmolyte on the protein surface, osmotic gradient should be created towards the protein.
588 Diffusion of water towards the unfolded should occur, making available enough water
589 molecules for the hydration of exposed polar groups. This is without prejudice to the initial
590 displacement of weakly bound water by the binding of the osmolyte. On the other hand
591 translational entropy gain of departing water from hitherto hydrated protein due to excluded
592 osmolyte compels the protein to (re)fold. As presented in literature [40] the first order case
593 which seem to be applicable to highly concentrated osmolyte/cosolute, requires the
594 introduction of apparent hydrated molar volume of the former as follows: Equation (23), by
595 so doing, is transformed to

596
$$\Delta_N^D(G_{pw}) = -\Delta_N^D V_{prot} + \frac{m(1-C_3 V_1)\phi_{os}}{RT} \quad (25)$$

597 Equation (25) enables the determination of the integral for hydration at none destabilising
598 concentration of the stabilising osmolyte as long as $1 > C_3 V_1$ and $V_1 \ll 1$. This equation is
599 reserved for feature investigation in which the concentration range of sucrose may be 0.25 -

600 1.25 mol/L. For the purpose of discussion Eq. (25) reminds one of the high molar
601 concentrations of ethanol explored in this research whose effect requires another equation
602 slightly different from Eq. (25). The equation is

$$603 \quad \Delta_N^D(G_{po}) = -\Delta_N^D V_{prot} - \frac{m(1-\phi_{os})(1-C_3V_1)}{RT} \quad (26)$$

604 Since ϕ_{os} is the volume fraction of cosolvent (or rather mole fraction which covers non-
605 solvent and solvents, e.g. sucrose and ethanol respectively), and its value being < 1 means
606 that $1 - \phi_{os}$ is always > 0 . The implication is that for the osmolation (positive preferential
607 interaction parameter) case $1 - C_3V_1$ should also be > 0 . This explains the osmolation (and
608 its effect) whereby $\Delta_N^D(G_{po})$ needs to be positive due to the binding of ethanol alone and only
609 aspirin in separate assays. Osmolation leads to unfolding and consequently, a decrease in
610 the amyolytic action of the enzyme as observed. The question that needs to be answered is,
611 what means can be applied for the determination of V_1 ? The issue of interest is always the
612 hydration changes linked either to the *ES* or cosolvent inaccessible region of the protein.
613 This is despite objection against total reliance on flexibility for function considering that some
614 enzyme activity can occur at very low hydration levels, coupled with a reduction in protein's
615 flexibility. On the contrary, Poole [35] observed that hydration induced conformation and
616 dynamic changes are completed just before the onset of enzyme activity which occurs
617 before all polar groups are hydrated. There was confirmatory evidence via increased alpha –
618 helicity that leads to increased rigidity in the protein at low hydration (dry); this led to the
619 deduction that when water is added to the dry enzyme a “loosening up” or increase in
620 flexibility occurs around a threshold of hydration [35]. It appears therefore, that it is an
621 excessive flexibility that leads to total unfolding due to the action of destabilisers that
622 reduces the biological function of the protein as observed in this research.

623 In summary there may be changes in the number of water interacting with *ES*, be it
624 negative or positive. The change in the number of water molecules interacting with the *ES* as
625 a function of ethanol/aspirin concentration indicates dehydration more so with a lower

626 concentration of sucrose. Thus ethanol is destabilising. The change as a function of sucrose
627 concentration with different concentration of ethanol/aspirin shows mixed trend, increasing
628 hydration with lower fixed concentration of ethanol/aspirin and decreasing with higher fixed
629 concentration of ethanol/aspirin. Thus sucrose promotes hydration being a protecting
630 osmolyte.

631 Generally, the change in the number of water molecules (ΔN_W) in an osmolyte
632 inaccessible region of the enzyme as a function of sucrose concentration with different fixed
633 concentration of aspirin/ethanol is positive as to imply hydration. Perhaps, the increasing
634 solubility of raw starch in increasing concentration of ethanol may presumably account for
635 the negative ΔN_W with lower concentration of ethanol. The values of ΔN_W as a function of
636 aspirin/ethanol concentration with different fixed concentration of sucrose are negative as to
637 imply dehydration peculiar to osmolation by destabilising cosolute.

638 The spontaneity of the processes, folding to unfolding transition and accompanying
639 hydration changes, has been illustrated with the quantification of the free energy difference;
640 in line with the approach, the results shows that *ab initio* the equilibrium constant for
641 hydration change ($K_{eq}(1)$) may be < or > equilibrium constant ($K_{eq}(3)$) for folding to unfolding
642 transition. A positive free energy difference means that hydration change is more
643 spontaneous than folding transition which may be attributable to the effect of sucrose. The
644 converse is the case with ethanol/aspirin in which the free energy difference is negative
645 ($K_{eq}(3) > K_{eq}(1)$).

646 For the purpose of interpretation, theories in literature were adopted for the
647 elucidation of results. The model for water stripping effect of aspirin/ethanol, preferential
648 interaction concept and the KBT for KBI for osmolation and hydration guided the
649 interpretation of the root basis of the effects of the cosolutes.

650 5. CONCLUSION

651 The change in the number of water molecules in an osmolyte inaccessible region of
652 the enzyme and those interacting with the ES may be either positive or negative due

653 respectively to sucrose and ethanol/aspirin. The spontaneity of two processes, hydration and
654 folding-unfolding transition, the free energy difference, differs. The mathematical model for
655 water stripping, preferential interaction concept, and the KBI for osmolation and hydration
656 can guide the interpretation of the effects of any cosolute.

657

658 **COMPETING INTERESTS DISCLAIMER:**

659 **Authors have declared that no competing interests exist. The products used**
660 **for this research are commonly and predominantly use products in our area of**
661 **research and country. There is absolutely no conflict of interest between the**
662 **authors and producers of the products because we do not intend to use these**
663 **products as an avenue for any litigation but for the advancement of**
664 **knowledge. Also, the research was not funded by the producing company**
665 **rather it was funded by personal efforts of the authors.**

666

667 **COMPETING INTERESTS**

668 There is no competing interest.

669 **REFERENCES**

- 670 1. Nakagawa H, Kitao JA, Kataoka M. Hydration affects both harmonic and
671 anharmonic nature of protein dynamics. Biophys. J. 2008; 95(6): 2916–2923.
- 672 2. Bernazzani P. Structural changes associated with interactions between starch and
673 particles of TiO₂ ZnSe. J. Chem. Biochem. Mol. Biol. 2008; 2(1) 1 – 13.
- 674 3. Gangadharan D, Nampoothiri KM, Sivaramakrishnan S, Pandey A. Immobilized
675 bacterial α-amylase for effective hydrolysis of raw and soluble starch. Food Res. Int.
676 2009; 42: 436-442.
- 677 4. DR. Knight J, Hamelberg D, McCammon AJ, Kothary R. The role of conserved
678 water molecules in the catalytic domain of protein kinases. 2009; 76(3): 527-535.
- 679 5. Affleck R, Xu Z-F, Suzawa V, Focht K, Clark DS, Enzymatic catalysis and dynamics
680 in low-water environments. Biochemistry. 1992; 89: 1100 -1104.

- 681 6. Kurkal V, Daniel RM, Finney JL, Tehei M, Dunn RV, Smith JC. Enzyme Activity and
682 Flexibility at Very Low Hydration. *Biophys. J.* 2005; 89: 1282–1287.
- 683 7. Laage D, Elsaesser T, Hynes JT. Water dynamics in the hydration shells of
684 biomolecules. *Chem. Rev.* 2017; 117:10694-10725.
- 685 8. Buurma, NJ, Pas Torello L, Blandermer JM, Engberts JBFN. Kinetic evidence for
686 hydrophobically stabilized encounter complexes formed by hydrophobic esters in
687 aqueous solutions containing monohydric alcohols. *J. Am. chem. Soc.* 2001; 123:
688 11848 – 11853.
- 689 9. Udema II, Onigbinde AO. Effect of interacting organic co-solutes with enzyme
690 substrate complex on the hydrolysis of raw soluble starch with alpha-amylase:
691 Theory and experimentation. *Adv. Res.* 2016; 7(1): 1-19.
- 692 10. Schnell S, Maini PK. Enzyme kinetics at high enzyme concentration. *Bull. Math. Biol.*
693 2000; 62: 483–499.
- 694 11. Udema II. Substrate mass conservation in enzyme catalyzed amylolytic activity. *Int.*
695 *J. Biochem. Res. Rev.* 2017; 18 (1): 1-10.
- 696 12. Mitchell DC, Litman DT. Effect of ethanol and osmotic stress on receptor
697 conformation reduced water activity amplifies the effect of ethanol on Metarhodopsin
698 II formation. *J. Biol. Chem.* 2000; 275 (8): 5355-5360.
- 699 13. Sirotkin VA, Kuchierskaya AA. Alpha-Chymotrypsin in water-ethanol mixtures: Effect
700 of preferential interactions. *Chem. Phys. Lett.* 2017; 689: 156-161.
- 701 14. Udema II, Onigbinde AO. Basic Kirkwood –Buff theory of solution structure and
702 appropriate application of Wyman linkage equation to biochemical phenomena.
703 *Asian J. Phys. Chem. Sci.* 2019; 7(1):1-14.
- 704 15. Sheng Y, Capri J, Waring A, Valentine JS, Whitelegge J. Exposure of solvent-
705 inaccessible regions in the amyloidogenic protein human SOD1 determined by
706 hydroxyl radical foot printing. *J. Am. Soc. Mass Spectrum.* 2019; 30 (2): 218-226.

- 707 16. Blundell TL, Gong S. Structural and functional restraints on the occurrence of single
708 amino acid variations in human proteins. *PLoS one*. 2010; 5(2): e9186, 1-12.
- 709 17. Gains JC, Acebes S, Virueta A, Butter M, Regan L, O' Hern CS. Comparing side
710 chain packing in soluble proteins, protein-protein interfaces, and transmembrane
711 proteins. *Proteins*. 2018; 85 (5):581-591.
- 712 18. Bolen DW, Baskakov IV. The osmophobic effect: Natural selection of a
713 thermodynamic force in protein folding. *J. Mol. Biol.* 2001; 310 (5): 955-963.
- 714 19. Pace CN. Measuring and increasing protein stability. *Trends Biotechnol.*1990; 8: 93
715 – 98.
- 716 20. Udemá II, Onigbinde AO. Activity coefficient of solution components and salts as
717 special osmolyte from Kirkwood-Buff theoretical perspective. *Asian Res. Biochem.*
718 2019; 4(3): 1-20.
- 719 21. Lineweaver H, Burk D. The determination of enzyme dissociation constants. *J. Am.*
720 *Chem. Soc.* 1934; 56: 658–666.
- 721 22. Kramer RM, Shende VR, Motl N, Pace N, Scholtz JM. Toward a molecular
722 understanding of protein solubility: Increased negative surface charge correlates
723 with increased solubility. *Biophys. J.* 2012; 102: 1907–1915.
- 724 23. Bernfeld P. Amylases, alpha and beta. *Methods. Enzymol.* 1955;1:149–152.
- 725 726 24. Udemá II. The effect of additives and temperature on the velocity of hydrolysis of raw
727 starch with human salivary α - amylase. *J. Biochem. Res. Rev.* 2016; 10(2): 1-17.
- 728 25. Sanyal NS, Kaushal N. Effect of two non-steroidal anti-inflammatory drugs, aspirin
729 and nimesulide on the G-glucose transport and disaccharide hydrolases in the
730 intestinal brush border membrane *Pharmacol Rep* 2005; 57: 833-838.
- 731 26. de Piña MZ, Saldaña-Balmori Y, Hernández-Tobias A, Piña E. Nonsteroidal
732 anti-inflammatory drugs lower ethanol-mediated liver increase in lipids and
733 thiobarbituric acid reactive substances. *Alcohol Clin Exp Res* 1994; 17(6):1228-
734 12232.

- 735 27. Levitt M, Sharon R. Accurate simulation of protein dynamics in solution. Proc. Nat.
736 Acad. Sci. U.S.A. 1988; 85: 7557-7561.
- 737 28. Petukhov M, Rychkov G, Firsov L., Serrano L. H-bonding in protein hydration
738 revisited. Protein Sci. 2004; 13 (8): 22120-2129.
- 739 29. Ooi T, Oobatake M. Prediction of the thermodynamics of protein unfolding: The
740 helix-coil transition of poly(L-alanine). Biochemistry. 1991; 88: 974-975.
- 741 30. Yang L, Dordick JS, Garde S. Hydration in nonaqueous media is consistent with
742 solvent dependence of its activity. Biophys. J. 2004; (87): 812-821.
- 743 31. Dai L, Klibanov AM. Striking activation of oxidative enzymes suspended in
744 nonaqueous media. Proc. Nat. Acad. Sci. U.S.A. 1999; 96: 9475 - 9478.
- 745 32. Zaks A, Klibanov AM. The effect of water on enzyme action in organic media. J. Biol.
746 Chem. 1988; 263 (17): 8017 - 8021.
- 747 33. Pal SKJ, Zewail AH. Biological water at the protein surface, dynamical solvation
748 probed directly with femtosecond resolution. Proc. Nat. Acad. Sci. U.S.A. U.S.A.
749 2001; 94 (4):1763-1768.
- 750 34. Csermely, P. (2001). Water and cellular folding process. Cell. Mol. Biol. 47 (5): 1-9.
- 751 35. Poole P. Hydration and enzymatic activity. J. Phys. Colloq. 1984; 45(c7): 249-253.
- 752 36. Timasheff SN. Protein solvent preferential interaction, protein hydration, and the
753 modulation of biochemical reactions by solvent components. Biochemistry. 2002; 99(15):
754 9721-9726.
- 755 37. Harano Y, Kinoshita M. Translational-entropy gain of the solvent upon protein
756 folding. Biophys. J. 2005; 89: 2701-2710.
- 757 38. Robinson CR, Sligar SG. Molecular recognition mediated by bound water: A
758 mechanism for star activity of the restriction endonuclease *EcoRI*. J.Mol. Biol. 1993;
759 234(2): 302-306.
- 760 39. Lynch TW, Sligar SG. Macromolecular hydration changes associated with *BamHI*
761 binding and catalysis. J. Biol. Chem. 2000; 275 (39): 30561-30565.

- 762 40. Rösger J, Pettit MB, Bolen DW. Protein folding, stability, and solvation structure in
763 osmolyte solution. *Biophys. J.* 2005; 89: 2988–2997.
- 764 41. Bolen DW, Baskakov IV. The osmophobic effect: Natural selection of a
765 thermodynamic force in protein folding. *J. Mol. Biol.* 2001; 310 (5): 955-963.
- 766 42. Baskakov I. and Bolen DW. Forcing thermodynamically unfolded proteins to fold
767 (communication). *J. Biol. Chem.* 1998; 273 (9): 1-5.
- 768 43. Schellman JA. Protein stability in mixed solvents: A balance of contact interaction
769 and excluded volume. *Biophys J.* 2003; 85:108 -105.

770

771

772

773

774

775

776

777

778

779

780

781

782