

***In Vitro* Analysis of Antimicrobial Susceptibility of *Xanthomonas axonopodis* pv. *punicae* and Comparative Study of Protein Profiling under Biotic Stress**

Abstract

Bacterial blight of pomegranate caused by *X. axonopodis* pv. *Punicae* (*XAP*) assumed epidemic form and resulted in economic burden on farmers. In the current study the pathogen infected samples were collected and the isolated *XAP* was identity and confirmed through the morphological, biochemical characterization and Pathogenicity test. Bacterium was reisolated from infected plant to prove Koch's postulates. Efficacy of different chemicals and oils were tested by disc diffusion assay and turbidometrically. Bronopol 3000 ppm (25.6 ± 1.6 mm) and Clove oil (18.0 ± 0.7 mm) formed highest zone of inhibition Turbidometri showed the highest O.D. (0.908 nm) by Copper oxy chloride and Neem oil showed maximum inhibition of growth with O.D. (0.842 nm). Biotic stress (pathogen) induced protein response was studies by using SDS-PAGE method after protein extraction from *XAP*, healthy *P. granatum* L. and infected *P. granatum* L. The protein banding pattern showed the unique band no. 2 (Mol.Wt.66000 Da) in infected *P. granatum* L. as compared to the banding pattern of *XAP* and healthy *P. granatum* L. The over expressed protein due to biotic stress could be useful as a marker for detection of the disease at the early stage and for control of the diseases after knowing the biochemical significance of the protein.

Keywords: Pomegranate, bacterial blight, *Xanthomonas axonopodis* pv. *punicae*, *In vitro* efficacy, Antimicrobial activity, SDS-PAGE.

Introduction

Pomegranate (*Punica granatum* L.) belongs to the family Lythraceae, is an ancient, medical fruit and cash crop of Maharashtra. It is also grown in commercially in other state of india viz. Karnataka, Andhra Pradesh, Rajasthan, Gujarat, and Tamil Nadu (Manjula et al., 2009). India is one of the major producers of pomegranate in the world with average total production of 8 lakh tons per annum.

Many diseases occurred on the pomegranate plant out of them bacterial disease caused by the *Xanthomonas axonopodis* pv. *punicae* (Xap) strongly damaged the pomegranate production, including large economic losses to the Indian growers (Mondal et al. 2009). Bacterial disease cause substantial loss to the productivity of major crop plants. Unlike fungal disease, bacterial disease not control effectively through chemical methods.

34 The symptoms of the bacterial blight were observed on the leaves stems and fruits as
35 small (2 to 5 mm) in size, irregular, prominent water soaked spots, which later become black
36 often leads to the breaking of branches and cracking of fruits, reducing to the market value of
37 the fruit (Rangaswamy, 1962; Manjula and khan 2002).

38 It is gram negative, aerobic rods belonging to the family *Pseudomonodaceae*. The
39 pathogen causes disease actively at high temperature (up to 32°C) and high rainfall and
40 humidity and p^H at 7.0. The pathogen causes destruction of seedlings and adult plants and thus
41 results in yield reduction. The aim of the study was to screen *in vitro* a range of antimicrobial
42 agents and their efficacy against Xap and protein profiling; in laboratory conditions.

43 **Material and methods**

44 **Collection and isolation of disease sample:**

45 The samples were collected from two different districts of Maharashtra viz.
46 Ahmednagar and pune as per typical symptoms showed by plant externally. The fresh
47 infected leaves samples were used for isolation employing tissue isolation method by using
48 Nutrient Agar medium (Antre et.al. (2015)).

49 **Purification and Maintenance of bacterial culture:**

50 The respective bacterial cultures were maintained on NA medium at room
51 temperature by adopting subsequent subculturing at periodical, regular intervals. Three days
52 old cultures were used for further studies (Kanwar et.al. (1976)).

53 **Identification of the pathogen:**

54 The identification of the pathogen involved in causing of oily spot in pomegranate
55 was determined by conducting studies on its morphological (Ryan *et.al.* (2011); Kishun R *et*
56 *al.* (1993)) and biochemical (Duche *et.al.*(2015); Giri *et.al.*(2011)) features of the pathogen as
57 per standard microbiological procedures.

58 **Pathogenicity test:**

59 Veinlets of some leaves were injected with 48 h old bacterial suspension with the help
60 of sterile Dispo Van syringe. Some plant leaves were inoculated by injuring the leaves with
61 sharp needle and by cotton swabbing the bacterial suspension on injured leaves. The plant
62 was kept for incubation. After 10 – 12 days observations made for typical symptoms of
63 bacterial blight on leaves, the organism was reisolated from artificially inoculated leaves of
64 Pomegranate plants showing typical symptoms of disease. The re-isolation was carried out on
65 NAS medium and the growth of organism was observed for next 2-3 days (Popovic *et al.*
66 (2013); Abdo-Hasan *et al.* (2008)).

67 ***In vitro* efficacy of different chemicals and plant derived oils against Xap by using paper** 68 **disc and turbidimetric method:**

In vitro efficacy of different chemicals and oils was checked against *Xanthomonas axonopodis* pv. *Punicae* by using paper disc and turbidimetric method (Copper oxy chloride: 2500ppm, 1000ppm, Bronopol: 3000ppm, 1000ppm, Bordeaux mixture: 3000ppm, 1000ppm, Bactomycin: 1200ppm, 1000ppm, Clove oil: 100%, Nilgiri oil: 100%, Neem oil: 100%). (Kizil et al.(2005), Pawar et al. (2014)).

Isolation of protein from healthy and infected pomegranate leaves:

Kumar et al. (2015) standardized the protein isolation protocol from plant leaves.

Isolation of protein from *Xanthomonas axonopodis* pv. *punicae*:

Bioera's teaching kit (2017) described the protein extraction from bacteria and clear result was observed.

Results and discussion:

Collection and isolation of diseased sample:

Bacterial blight infected samples were collected from Ahmednagar and Pune districts of Maharashtra, associated bacterium *Xanthomonas axonopodis* pv.*punicae* was isolated. Purified culture was maintained on NSA slants and coded as Xap I, Xap II respectively. (Bradbury et al. (1970)).

Identification of the pathogen:

Morphological and biochemical characteristic viz., shape, colony colour, Gram reaction, starch hydrolysis, gelatin liquefaction, indole production, acid and gas production, KOH, catalase test were performed and confirmed to identity the bacterium as *Xanthomonas axonopodis* pv. *punicae* and following table 1 showed the morphological and biochemical test. (Gamangatti et al. (2013),Bhardwaj et al. (2014)).

Pathogenicity test:

Pathogenicity test of isolates was confirmed by injecting bacterial solution to disease free pomegranate plant. Occurrence symptoms of bacterial blight symptoms on plant leaf confirmed pathogenicity test of isolates (Fig. 1). Bacterium was reisolated from infected plant to prove Koch's postulates. (Yenjerappa et al. (2009)).

***In vitro* efficacy of different chemicals and plant derived oils against Xap by using paper disc method:**

In order to assess the efficacy of chemicals and oils against two isolates of *Xanthomonas axonopodis* pv. *Punicae* an experiment was conducted and the evaluation was made by paper disc method. The data presented in (Table 2) revealed the significant differences among the different treatment. In the chemical treatment Bronopol were significantly superior in inhibiting the growth of bacteria (18.6 ± 0.7 mm, 25.6 ± 1.6 mm) against *Xap*. Among the oils Clove oil showed maximum zone of inhibition 18.3 ± 0.7 mm (Fig. 2). (Meerts and Fadavi et al. (2009)).

***In vitro* efficacy of different chemicals and plant derived oils against Xap by using turbidimetric method:**

Efficacy different chemicals and oils was tested turbidometrically. After 24 hours of incubation list O.D. 0.713 nm was exhibited by Bronopol(3000 ppm), followed by Copper oxy chloride (3000 ppm) with O.D. 0.908 nm and Bordeaux (3000 ppm) 0.834 nm respectively whereas Bronopol (1000 ppm) with O.D 0.219 nm showed minimum inhibition of growth and Among the oils tested Neem oil showed maximum inhibition of growth with O.D. 0.842 nm, followed by Clove oil with O.D. 0.810 nm, Nilgiri oil showed minimum inhibition of growth among oils showed in following table 3.

Protein isolation from Healthy pomegranate leaves, infected pomegranate leaves and *Xap I*, *Xap II* Bacteria:

Data analysis of Healthy pomegranate leaves, infected pomegranate leaves and Bacteria:

Following data analysis table shows the actual position of bands on the gel as compared to the standard marker on the basis of molecular weights provided by BioEra's Teaching Kit.

The protein banding pattern showed the unique band no. 2 (Mol.Wt.66000 Da) in infected *P. granatum L.* as compared to the banding pattern of *X. axonopodis* pv.*punicae* and healthy *P. granatum L.* (Fig. 3) (Walker J.M. (2005)).

126 Conclusion:

127 In the current study the pathogen infected samples identified as infected with *XAP*
 128 was through morphological, biochemical characterization and Pathogenicity test. Efficacy of
 129 different chemicals and oils were tested by disc diffusion assay and turbidometrically.
 130 Bronopol 3000 ppm (25.6 ± 1.6 mm) formed highest zone of inhibition and Clove oil formed
 131 highest zone of inhibition (18.0 ± 0.7 mm). Turbidometri showed the highest O.D. (0.908 nm)
 132 by Copper oxy chloride and Neem oil showed maximum inhibition of growth with O.D.
 133 (0.842 nm).

134 Biotic stress (pathogen) induced protein response was studies by using SDS-PAGE
 135 method after protein extraction from *XAP*, healthy *P. granatum L.* and infected *P. granatum*
 136 *L.* The protein banding pattern showed the unique band no. 2 (Mol.Wt.66000 Da) in infected
 137 *P. granatum L.* as compared to the banding pattern of *X. axonopodis pv. punicae* and healthy
 138 *P. granatum L.* The overexpressed protein due to biotic stress could be useful as a marker for
 139 detection of the disease at the early stage and for control of the diseases after knowing the
 140 biochemical significance of the protein.

141 **Table1. Morphological and Biochemical characteristics of *Xanthomonas axonopodis***
 142 ***pv. punicae* isolates.**

Sr. No.	Morphological and Biochemical test	<i>Xap I</i>	<i>Xap II</i>
1.	Colony color	Yellow	Yellow
2.	Gram's reaction	-ve	-ve
3.	Starch hydrolysis	+++	++
4.	Indole production	++	+
5.	Catalase test	+++	+++
6.	KOH test	++	+
7.	Gelatin liquefaction	++	++
8.	Acid production	++	+

144 - Negative, + Positive, Varying degree of + ve reaction (+ poor, ++ moderate, +++ strong)

145 **Table 2 revealed the significant differences among the different treatment by paper disc**
 146 **method.**

Sr. No.	Treatment	Concentration	Zone of inhibition produced by Xap I	Zone of inhibition produced by Xap II
T ₀	Control(+ve) water	10µl	00	00
T ₁	Control(-ve) Ampicillin	15µl	17.9±0.6	27.7±0.7
T ₂	Bronopol	1000 ppm	10.8±0.4	18.6±0.7
T ₃	Bordeaux	1000 ppm	10.2±0.2	9.3±0.1
T ₄	COC	1000 ppm	12.1±0.3	11.9±1.5
T ₅	Bactomycin	1000 ppm	10.1±0.3	13.2±0.9
T ₆	Bronopol	3000 ppm	11.6±0.4	25.6±1.6
T ₇	Bordeaux	3000 ppm	9.7±0.7	8.1±.02
T ₈	COC	2500 ppm	7.1±1.0	9.2±0.4
T ₉	Bactomycin	1200 ppm	11.2±0.4	15.3±0.2
T ₁₀	Clove oil	100%	15.9±0.9	18.3±0.7
T ₁₁	Neem oil	100%	13.4±1.6	7.3±1.1
T ₁₂	Nilgiri oil	100%	8.3±0.9	10.6±0.4

147

148 **Table 3 Efficacy of chemicals and oils against *Xanthomonas axonopodis pv***
 149 ***punicae* by turbidometric method:**

150

Sr.No.	Treatment	Concentration	O.D. of culture	O.D. after 24 hrs at 620 nm
T ₀	Control(+ve)Nutrient Broth	1 ml		00
T ₁	Control (-ve) antibiotic	500 µl		0.324
T ₂	Bronopol	1000 ppm		0.219
T ₃	Bordeaux	1000 ppm		0.337
T ₄	COC	1000 ppm		0.410

T ₅	Bronopol	3000 ppm	0.998	0.713
T ₆	Bordeaux	3000 ppm		0.834
T ₇	COC	3000 ppm		0.908
T ₈	Clove oil	100%		0.810
T ₉	Neem oil	100%		0.842
T ₁₀	Nilgiri oil	100%		0.392

151 **Table4. Data analysis of Healthy pomegranate leaves, infected pomegranate leaves and**
152 **bacteria**

Sr. No.	Standard Marker (Band no)	Molecular weight	Healthy plant (Band no)	Infected plant (Band no)	<i>Xap I</i> proteins (Band no)	<i>Xap II</i> proteins (Band no)
1.	1	97,400	Absent	Absent	Absent	Absent
2.	2	66,000	Absent	Present	Absent	Absent
3.	3	43,000	Present	Present	Absent	Absent
4.	4	29,000	Present	Present	Absent	Absent
5.	5	20,100	Present	Present	Present	Present
6.	6	14,300	Present	Present	Present	Present

153

154 **Fig. 1 (a) Symptoms of bacterial blight observed after 10-12 days of inoculation (b)**
155 **Reisolated *Xanthomonas axonopodis* pv. *punicae* after pathogenicity test**

156



157


158

(a)




(b)


Fig. 2 Efficacy of different chemicals and oils against *Xanthomonas axonopodis* pv. *Punicae* by Paper disc method.



a) Bactomycin




b) Bordeaux



168

169


c) Bronopol



170

171

d) Copper oxychloride




172

173

e) Oils

174

175 **Fig 3(a) Isolated proteins of 1) Healthy, 2) Marker, 3) Infected leaves (b) 1) *Xap I*, 2)**
 176 **Marker and 3) *XapII* Bacteria (*Xanthomonas axonopodis* pv.*punicae*).**



177
 178 **References:**

- 179 Abdo-Hasan,M., Khalil, H., Debis, B., & MirAli, N. (2008). Molecular characterization of
 180 Syrian races of *Xanthomonas axonopodis* pv.*malvacearum*. *Journal of plant*
 181 *Pathology*, 431-439.
- 182 Antre, S.H., Gadhe, S.K., Abhang, P.B. and Autade, R.H., In vitro Efficacy of Different
 183 Chemicals, Botanicals and Bioagent Against *Xanthomonas axonopodis* pv. *punicae*,
 184 *Int. J. Pure App. Biosci.* 4(3): 112-118 (2016). doi: [http://dx.doi.org/10.18782/2320-](http://dx.doi.org/10.18782/2320-7051.2310)
 185 7051.2310
- 186 Bhardwaj, N.R., Upadhyay, V., & Nagar,m. (2014). Biochemical characterization of
 187 *xanthomonas axonopodis* pv..*citri*, causal agent of citrus canker.
- 188 BioEra Life Sciences Pvt. Ltd., Pune. (2017). *Teaching Kit, Total Protein Extract Kit (From*
 189 *Bacteria)* [Brochure]. Pune, Maharashtra: Author. Catalog number BTK 130934.
- 190 Bradbury, J. F. (1970). Isolation and preliminary study of bacteria from plants.*PANS Pest*
 191 *Articles & News Summaries*, 16(4), 632-637.
- 192 Duche, T. R., Omoigui, L., & Iheukwumere, C. C. (2015). Variations among *Xanthomonas*
 193 *axonopodis* pv.*vignicola* isolates in Benue State, Nigeria. *International Journal of*
 194 *Innovation and Scientific Research*, 13(1), 271-278.
- 195 Gamangatti, R. B., & Patil, M. B. (2013). Biochemical studies on *Xanthomonas axonopodis*
 196 pv..*punicae* causing bacterial blight of pomegranate. *International Journal of Plant*
 197 *Protection*,6(2), 401-404.

- 198 Giri M.S., Prasanthi., Kulkarni, S., Benagi, V.I. and Hegde, Y.R. 2011. Biochemical and
199 molecular variability among *Xanthomonas axonopodis* pv. *Punicae* strains, the
200 pathogen of pomegranate bacterial blight. *Indian phytopathol.* 64(1): 1-4.
- 201 Kanwar, Z. S.(1976). A note on bacterial disease of Pomegranate (*Punica granatum* L.) in
202 Haryana.*Haryana J. Horti. Sci.*, 5 : 177-180.
- 203 Kishun R (1993). Bacterial diseases of fruits. In: Chadha KL, Parek OP (Eds) *Advances in*
204 *Horticulture-Fruit crops* (Vol 3), Malhotra Publishing House, New Delhi, India, pp
205 1389-1404.
- 206 Kizil, S., Uyar, F., & Sagir, A. (2005).Antibacterial activities of some essential oils against
207 plant pathogens.*Asian Journal of Plant Sciences*.
- 208 Kumar Suresh, Vinutha T, Archana Singh, R.D.Rai et.al., 2015 *training manual Plantomics-*
209 *Emerging Tools and Techniques for Crop improvement*.
- 210 Manjula N., Sheikh M.K., Teggi M.Y., 2009. Bacterial blight management practice adopted
211 by farmers ISHS ACTA Horticulture 890: International symposium on pomegranate
212 and minor-including Mediterranean-Fruits.
- 213 Manjula, C.P. And Khan, A.N.A., 2002. Incidence of bacterial blight of pomegranate (*Punica*
214 *granatum* L.) in Karnataka. Paper presented at the Annu. Meet. Symp. Plant disease
215 scenario in southern India, held at Bangalore, December, 19-21.
- 216 Meerts and Fadavi et al.,(2009). In vitro control of bacterial blight causing organism
217 (*Xanthomonas axonopodis* pv. *Punicae*) of pomegranate journal of environmental
218 research and development, 8(3a), 568.
- 219 Mondal, K.K., Sharma, J. 20019. Bacterial blight: an emerging threat to pomegranate export.
220 *Ind. Farm.*, 59: 22-23.
- 221 Pawar, M., kadam, M., & Nirichan, B. K. (2014). in vitro control of bacterial blight causing
222 organism (*xanthomonas axonopodis* pv.. *punicae*) of pomegranate. *journals of*
223 *environmental research and development*, 8(3a), 568.
- 224 Popovic, T., Josic, D., Starovic, M., Milovanovic, P., Dolovac, N., Postic, D., & Stankovic,
225 S. (2013). Phenotypic and genotypic characterization of *Xanthomonas campestris*

- 226 strains isolated from cabbage, kale and broccoli. *Archives of Biological Science*, 65,
227 585-593.
- 228 Rangaswamy, G., 1962 Pomegranate. In: Bacterial plant disease in India. Asia publication
229 house, Bombay, pp.830.
- 230 Ryan, R. P., Vorhölter, F. J., Potnis, N., Jones, J. B., Van Sluys, M. A., Bogdanove, A. J., &
231 Dow, J. M. (2011). Pathogenomics of *Xanthomonas*: understanding bacterium–plant
232 interactions. *Nature Reviews Microbiology*, 9(5), 344-355.
- 233 Walker, J. M. (2005). Electrophoretic techniques. In *Principles and Techniques of*
234 *Biochemistry and Molecular Biology* (6th ed., pp. 449-485). New York: Cambridge
235 University Press.
- 236 Walker, J. M. (2005). Protein structure, purification, characterization and functional analysis.
237 In *Principles and Techniques of Biochemistry and Molecular Biology* (6th ed., pp.
238 349-405). New York: Cambridge University Press.
- 239 Yenjerappa, S. T., (2009). Epidemiology and management of bacterial blight of pomegranate
240 caused by *Xanthomonas axonopodis* pv. *punicae* (Hingorani and Singh) Vauterin *et al*
241 .Ph. D.Thesis, Univ.Agri. Sci. Dharwad (India).147.