

Short Research Article

Response of Heat Tolerant Variety (Kufri Surya) of Potato (*Solanum tuberosum*) Under Different Levels of Nitrogen

Abstract

Keeping in view, the deficiency of detailed information on adoption of heat tolerant potato (*Solanum tuberosum*) variety 'Kufri Surya' in Terai Agro-Climatic situation of West Bengal, the field experiment was conducted at the Instructional farm of Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal during the rabi season of 2016 to study the effect on heat tolerant variety (Kufri Surya) with different nitrogen levels. Experiment was laid out in a Split-plot design taking two varieties 'Kufri Jyoti' and 'Kufri Surya' as main plot with six different levels of nitrogen of 0 kg N ha⁻¹, 50 kg N ha⁻¹, 100 kg N ha⁻¹, 150 kg N ha⁻¹, 200 kg N ha⁻¹ and 250 kg N ha⁻¹ as subplot. Results of the experiment showed that the higher values of the growth attributes like dry matter accumulation, leaf area index, in all the sampling dates of experimentation was recorded with 100 kg N ha⁻¹. Owing to the higher leaf area index and dry matter accumulation in shoot, tuber yield was recorded highest from the treatment having 100 kg N ha⁻¹ (28.46 t ha⁻¹).

Key words: Potato, Nitrogen, Kufri Surya, Kufri Jyoti, Heat Tolerant Variety.

23 Introduction

24 Potato (*Solanum tuberosum* L.) is the third most important food crop in the
25 world after rice and wheat in terms of human consumption. India ranks as the world's
26 2nd largest potato producing nation after China. Production in India is about 48.52
27 million tonnes (Government of India, 2018) of which 26% are produced by West
28 Bengal itself. Potato is a cool season long day crop. High temperatures and long days
29 favour assimilate partitioning to the above ground vegetative parts, as a result, above
30 ground bio-mass and plant height is increased and tuber yield is reduced (Wolf *et al.*,
31 1990). Potato gives good yield at day temperature of 30-35 °C. But if night temperature
32 go beyond 22 °C, there will be little tuberization even when day temperature is 25-27
33 °C. Due to intense climate change the favourable temperatures for its growth is
34 increased at its later stages hampering the tuberization. On this context, whether the
35 heat tolerant variety 'Kufri Surya' could perform better than check variety 'KufriJyoti'
36 was brought into notice from the experiment. Nitrogen is beneficial for the tuber quality,
37 dry matter production, size of tubers etc. More application of nitrogen fertilizers can
38 increase size of tubers and hence the yield but there is a particular dose limit up to
39 which it will show positive results; beyond that limit the application of nitrogen
40 fertilizer will not increase the yield but rather it would be harmful because of deposition
41 of nitrogen in tubers in the form of nitrates (Mohammad and Mohammadreza, 2012)
42 which is not at all favourable for human consumption and moreover excessive
43 application can cause environmental pollution. The use of low N results in reduction in
44 yield of potato. Judicious use of balanced dose of fertilizers is very critical for higher
45 tuber yield. Keeping the above ideas in view, a field experiment was undertaken to
46 study the effect of different levels of Nitrogen on growth and yield of potato cultivars
47 'Kufri Surya' with check variety of 'Kufri Jyoti' and to study the effect of different
48 levels of nitrogen on net photosynthesis rate, stomatal conductance rate, transpiration
49 rate of potato.

50

51 Materials and Methods

52 A field experiment was conducted to study the effect of different doses of
53 nitrogen on two different varieties of potato that is, 'KufriJyoti' and heat tolerant
54 variety 'Kufri Surya' at Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Bihar,
55 West Bengal during *rabi* season of 2016. The farm is situated at 26°12'78"N latitude
56 and 89°24'55" E longitude at an elevation of 43 meters above mean sea level. The
57 climatic zone where the farm is situated is in *Terai* zone which is subtropical in nature
58 having its prominent characteristics of very high rainfall, high humidity and a
59 prolonged winter season. The average rainfall of this zone varies from 2000-3000 mm.
60 The soil of the experimental field was sandy loam in texture, a true representative of the
61 *terai* region of West Bengal with a pH of 5.6. The experiment was carried out in split
62 plot design with two varieties of potato 'Kufri Surya' and 'KufriJyoti' as main plots and
63 six nitrogen levels as subplots i.e., 0 kg N ha⁻¹, 50 kg N ha⁻¹, 100 kg N ha⁻¹, 150 kg N ha⁻¹,
64 200 kg N ha⁻¹ and 250 kg N ha⁻¹. The experiment had three replications with a plot size
65 of 5m x 3.45 m and a spacing of 45 cm x 15 cm. The crop was planted on 26th
66 November of 2016.

67 Healthy cut tubers were selected each having two-three eyes weighing 25-40
 68 gm. The seed tubers were treated to protect them from an attack of fungal diseases
 69 when planted in the field. So, before planting, the seed tubers were dipped in solution of
 70 Sprint @ 2.5 g lit.⁻¹+ streptomycin (Plantamycin) @ 2.5 g lit.⁻¹of water for 15 minutes
 71 and then they were dried in shade to protect it from direct sunlight prior to planting.
 72 Farmyard manure was applied on the field @ 5t ha⁻¹ at the time of final land
 73 preparation. The different doses of nitrogen were 0, 50, 100, 150, 200 and 250 Kg N ha⁻¹
 74 ¹+ 100 Kg P₂O₅ha⁻¹+100 Kg K₂O ha⁻¹were given respective plots. Out of these doses
 75 1/3rd of nitrogen and full dose of P₂O₅ and full dose of K₂O were applied as basal at the
 76 time of planting of tubers. The rest half of the 2/3rd nitrogen was given in two equal
 77 splits, one as first top dressing at 21 DAP (days after planting) and the second split of
 78 Nitrogen was applied at second top dressing at 41 DAP. The fertilizers were applied by
 79 broadcasting method in the form of Urea, Single Super Phosphate (SSP) and Muriate of
 80 Potash (MOP) as the sources of N, P₂O₅ and K₂O respectively. Two irrigations were
 81 given to the crop. First irrigation was given at 22 DAP after first top dressing and
 82 earthing up. Second irrigation was given at 44 DAP after second top dressing. Before
 83 10 days of harvesting of the crop dehauling was done.

84 The growth attributes like number of haulms per plant, leaf area index (LAI),
 85 crop growth rate(CGR),net assimilation rate(NAR) were recorded at 20,40, 60 and 80
 86 DAP(days after planting). The photosynthetic and transpiration parameters like net
 87 photosynthesis rate, leaf stomatal conductance, transpiration rate were recorded at 20,
 88 40,60DAP as 80 DAP the leaves starts showing senescence. Though the photosynthetic
 89 and transpiration parameters were recorded by the instrument CI -340 Handheld
 90 Photosynthesis system but the parameters can be also calculated by the following
 91 formula.The Net Photosynthesis Rate (Pn) (micro mol m⁻² s⁻¹) was calculated by the
 92 following formula:

93
$$P_n = -W \times (C_0 - C_i) = - \frac{V \times P}{T_a \times A} \times (C_0 - C_i)$$

96 Where, C₀(C_i): outlet (inlet) CO₂ concentration (ppm or micro mol/ mol)
 97 W = Mass flow rate per leaf area, V = Leaf chamber volume, T_a = air temperature (K), P
 98 = Atmospheric pressure (bar) ,A = Leaf area (cm²).

99 The Transpiration Rate (E) (milimol m⁻² s⁻¹) was calculated by:

100
 101
$$E = \frac{e_0 - e_i}{P - e_0} \times W \times 10^3$$

104
$$e_0 = hr_0 \times es100^{-1}$$

 105
$$e_i = hr_i \times es100^{-1}$$

106 Where, e₀(e_i): Outlet (inlet) water vapour (bar) ,W = Mass flow rate per leaf area ,T_a =
 107 air temperature (K) ,P = Atmospheric pressure (bar) ,hr₀ (hr_i) = Outlet (inlet) relative
 108 humidity (%).

109
 110

111
112

W

113 Leaf Stomatal Conductance= $\frac{W}{e_0 - e_i} \times \frac{p - e_0}{p} \times 1000$

114
$$\frac{e_{\text{leaf}} - e_0}{e_0 - e_i} \times \frac{p - e_0}{p}$$

115

116 Where, e_{leaf} = saturated water vapour at leaf temperature (bar) W = Mass flow
117 rate per leaf area, P = Atmospheric pressure (bar) , e_0 (e_i) : outlet (inlet) water vapour
118 (bar) , R_b = leaf boundary layer resistance ($\text{m}^2\text{s mol}^{-1}$) $0.3\text{m}^2\text{s mol}^{-1}$ is used. Observation
119 were done at 20,40,60 DAP.

120 The data collected from the field and laboratory experiments were subjected to
121 statistical analysis with appropriate design and treatment variations were tested for
122 significance by F-test (Cochran and Cox, 1955). The standard error of mean and critical
123 difference is indicated in the tables. For determination of critical difference at 5% level
124 of significance Fisher and Yates (1963) table was consulted. The statistical analysis was
125 evaluated by SPSS software.

126

127

128 Results and Discussion

129 *Effect on growth attributes and yield:*

130 Among the two varieties 'Kufri Surya' and 'KufriJyoti' both of them have given
131 statistically similar results for most of the growth attributes at 20 and 80 DAP. This was
132 because at 20 DAP the plants are yet to be developed because of its early stages of
133 growth and at 80 DAP the plants started showing senescence symptoms of stems and
134 leaves. But quite significant differences were observed when the observations were
135 taken at 40 and 60 DAP because it was the peak period of vegetative growth of the
136 potato crop. In case of number of haulms per plant at any stages of the crop there were
137 no significant differences among the varieties. At 40 and 60 DAP it was observed that
138 'KufriJyoti' had performed 29 % better than 'Kufri Surya' at 40DAP and 24 % better at
139 60DAP. This can be pertained to 'KufriJyoti' having higher leaf area than 'Kufri Surya'
140 whose leaves are narrower in shape resulting in lesser leaf area. Dry matter
141 accumulation at 40 DAP was 19% more in 'KufriJyoti' because of more leaf area
142 index, for which photosynthesis was more resulting in better accumulation of
143 photosynthates and at 60 DAP Kufri Surya (303.13 g m^{-2}) performed better than
144 KufriJyoti (279.53 g m^{-2}). This was due to mild attack of *Phomaspp.onKufriJyoti* at
145 60DAP for which growth of the plant was hampered. In case of crop growth rate
146 KufriJyoti performed better at 20- 40 DAP ($7.315 \text{ g m}^{-2} \text{ day}^{-1}$) and at 40-60 DAP Kufri
147 Surya ($8.569 \text{ g m}^{-2} \text{ day}^{-1}$) gave maximum crop growth rate. Crop growth rate was
148 hampered for KufriJyoti due to the same reason for which the dry matter accumulation
149 was less at 60DAP. Net assimilation rate which is the amount of dry matter produced in

150 gram per unit area of leaf per day was found significant at 40-60 DAP for both the
151 varieties due to its peak period of growth in which Kufri Surya has performed 65%
152 better than KufriJyoti at 40-60 DAP as it was resistant to pathogen attack. Though
153 KufriJyoti (25.80t ha⁻¹) was mildly affected by *Phoma* spp. at 60DAP, timely control
154 measures had helped immensely to revert back its negative effects on yield and hence
155 had shown better yield compared to Kufri Surya(19.76 t ha⁻¹) . The reasons might be
156 due to bigger size and weight of tubers per plant in case of KufriJyoti.

157 Nitrogen is a very essential nutrient for growth of plants because its an
158 important constituent of key photosynthetic enzyme RuBPCarboxygenase/ oxygenase.
159 Total sugar accumulation in leaves and tubers are positively influenced by nitrogen
160 application. Total sugar increased with the rate of N-fertilizer application. The higher
161 sugar content was due to higher photosynthetic rate, which is enhanced due to
162 enzymatic activity. Increase in nitrogen levels increases the carbohydrate production by
163 more number of chlorophylls. But there is a limit of nitrogen application beyond which
164 if nitrogen fertilizers are added the plants won't show a positive result. It was observed
165 that number of haulms increased linearly with increase in dose of nitrogen since
166 nitrogen has a positive role in increase in vegetative growth of plant. There was
167 significant difference for most of the growth attributes in all the stages among the
168 various nitrogen levels except 20 DAP because of early stages of growth. So, in 40
169 DAP maximum number of haulms were observed in the treatment of 200 kg N ha⁻¹
170 (3.35) and in 60 DAP for 250 kg N ha⁻¹ (5.75). For dry matter accumulation 100 kg N
171 ha⁻¹ was found optimum for the maximum dry matter production at 40 DAP(171.46 g
172 m⁻²) and 60 DAP (322.25 g m⁻²). These results were in accord with the findings of
173 Sharma et al (1991). This might be assigned to LAI at 40 DAP and 60 DAP having the
174 highest value for 100 kg N ha⁻¹ as it was optimum amount nitrogen required for
175 enlargement of leaves resulting in production of more photosynthates . Crop growth
176 rate among the various nitrogen levels for both 20-40 DAP and 40-60 DAP were
177 statistically at par with each other. There was no significant difference among the
178 various nitrogen levels in 20-40 DAP except 40-60 DAP. Maximum net assimilation
179 rate (g m⁻² day⁻¹) was observed at 200 kg N ha⁻¹ in 40-60DAP(1.087). Highest yield was
180 obtained at 100 kg N ha⁻¹ due to better tuber development at the optimum level.

181

182 **Table 1.**Effect of Variety and Nitrogen levels on number of haulms per plant, LAI and dry matter accumulation of plant.

TREATMENTS	Number of haulms per plant			Leaf Area Index(LAI)				Dry matter accumulation(g m ⁻²)			
	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP	80 DAP	20 DAP	40 DAP	60 DAP	80 DAP
V ₁	2.50	2.87	4.47	0.15	3.98	4.46	2.20	8.26	150.97	279.53	274.55
V ₂	2.38	2.93	5.00	0.14	3.08	3.60	2.10	8.68	131.75	303.13	272.66
SEm(±)	0.01	0.03	0.59	0.05	0.12	0.02	0.09	0.24	2.83	1.18	.94
CD(0.05)	NS	NS	NS	NS	0.72	0.10	NS	NS	17.24	7.18	NS
Nitrogen levels											
N ₀	2.15	2.25	3.40	0.11	3.04	3.39	1.78	7.21	112.78	218.01	218.33
N ₁	2.40	2.65	4.30	0.12	3.50	3.77	1.93	8.25	136.15	290.72	261.76
N ₂	2.70	2.90	5.05	0.13	4.40	4.70	2.45	9.53	171.46	322.25	327.76
N ₃	2.30	3.05	5.00	0.19	4.01	4.18	2.69	8.54	154.28	321.60	306.71
N ₄	2.75	3.35	4.90	0.17	3.10	4.53	1.97	9.15	128.46	297.72	263.63
N ₅	2.35	3.20	5.75	0.16	3.14	3.60	2.11	8.15	145.03	297.66	263.44
SEm(±)	0.17	0.13	0.20	0.01	0.14	0.10	0.06	0.41	3.88	7.34	5.75
CD(0.05)	NS	0.39	0.58	NS	0.41	0.30	NS	NS	11.46	21.65	16.96

V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha⁻¹, N1-50 kg ha⁻¹, N2-100 kg ha⁻¹, N3- 150 kg ha⁻¹,N4-200 kg ha⁻¹, N5-250 kg ha⁻¹.

183

184

185

186 **Table 2.**Effect of Variety and Nitrogen levels on Crop Growth Rate, Net Assimilation Rate and Yield of crop.

TREATMENTS	Crop growth rate (g m ⁻²)		Net assimilation rate(g m ⁻² day ⁻¹)		Yield(t ha ⁻¹)
	Variety	20- 40 DAP	40-60 DAP	20-40 DAP	
V ₁	7.135	6.428	2.734	0.679	25.80
V ₂	6.153	8.569	2.825	1.120	19.76
SEm(±)	0.130	0.083	0.057	0.045	0.19
CD(0.05)	0.789	0.503	NS	0.142	1.17
Nitrogen levels					
N ₀	5.278	5.262	2.458	0.641	9.28
N ₁	6.395	7.729	2.832	0.789	20.45
N ₂	8.116	6.313	2.861	0.985	28.46
N ₃	7.287	8.366	2.954	0.905	26.78
N ₄	5.947	9.690	2.555	1.087	26.08
N ₅	6.844	7.632	3.016	0.991	25.63
SEm(±)	7.135	6.428	0.082	0.061	0.45
CD(0.05)	NS	NS	NS	0.179	1.33

187 V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha⁻¹, N1-50 kg ha⁻¹, N2-100 kg ha⁻¹, N3- 150 kg ha⁻¹,N4-200 kg ha⁻¹, N5-250 kg ha⁻¹.

188 ***Effect on photosynthetic parameters:***

189 There were significant differences among the varieties, maximum net
190 photosynthesis rate was observed in 20 DAP (7.7 micro mol m⁻² s⁻¹), 40 DAP
191 (14.77 micro mol m⁻² s⁻¹) in KufriJyoti and in 60 DAP in Kufri Surya (10.88 micro mol
192 m⁻² s⁻¹) which might be due to the possible reason of attack of pathogen on KufriJyoti
193 hampering its healthy leaf growth and hence the photosynthetic activity. Both the
194 varieties were statistically at par with each other with respect to transpiration
195 rate. Maximum stomatal conductance rate was in Kufri Surya (269.42 millimol m⁻² s⁻¹)
196 at 20 DAP, in KufriJyoti (368.55 millimol m⁻² s⁻¹) at 40 DAP which might be due to
197 faster development of leaves of Kufri Surya at 20 DAP and better development of
198 leaves and number of stomata in KufriJyoti at 40DAP.

199 Significant differences were observed between different nitrogen levels for
200 various photosynthetic parameters. All the photosynthetic characters have been
201 recorded maximum at 150 kg N ha⁻¹. Stomatal conductance rate is the rate at which
202 carbon dioxide is uptaken and water vapour is released through stomata. Nitrogen plays
203 an important role in stomatal conductance by cell expansion and altering the cation and
204 anion concentration of cytoplasmic environment which can actually change the stomatal
205 conductance rate (Nasabet al.2014). More nitrogen application also increases the leaf
206 growth and hence the number of stomata increasing the stomatal conductance rate.
207 Highest stomatal observations were found at 150 kg nitrogen ha⁻¹, for all the stages with
208 a maximum of 395.33 millimol m⁻² s⁻¹ at 60 DAP because this was the optimum dose
209 above which no such effect was seen. Since stomatal conductance rate is closely related
210 with transpiration rate, maximum transpiration rate similarly observed at 150 kg N ha⁻¹
211 at all stages of growth. Net photosynthetic rate may be assigned to the possible reasons
212 of larger number of chlorophyll and stomata due to optimum doses of nitrogen.

213 From the above experiment it can be concluded that Kufri Surya didn't perform
214 better than check variety KufriJyoti because the high temperature at which Kufri Surya
215 might have shown better performance with respect to yield than KufriJyoti due to its
216 heat tolerant characteristics which was not obtained. So the experiment requires
217 repetition for further studies in Terai region of West Bengal.

218

219

220

221

222

223

224

225

226

227 **Table 3.**Effect of Variety and Nitrogen levels on photosynthetic parameters.

TREATMENTS	Stomatal conductance rate (millimol m ⁻² s ⁻¹)			Transpiration rate (millimol m ⁻² s ⁻¹)			Net Photosynthesis rate (micro mol m ⁻² s ⁻¹)		
	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP
V ₁	235.43	368.55	355.63	1.06	2.80	2.85	7.77	14.17	9.19
V ₂	269.42	343.33	365.55	0.99	3.19	2.95	5.66	12.64	10.88
SEm(±)	1.44	2.79	0.58	0.07	0.05	0.05	0.04	0.02	0.23
CD(0.05)	8.76	16.98	NS	NS	0.30	NS	0.27	0.10	1.42
N ₀	220.60	315.27	355.89	0.72	3.18	2.04	4.29	11.94	7.97
N ₁	229.09	333.97	353.43	1.01	2.44	2.54	5.45	13.14	8.77
N ₂	244.39	365.07	387.76	1.04	2.94	3.04	7.46	13.60	9.34
N ₃	292.51	389.69	395.33	1.33	3.68	3.54	10.15	15.05	12.52
N ₄	258.91	371.73	349.98	1.02	2.85	3.13	7.16	13.07	11.49
N ₅	269.06	359.92	321.16	1.04	2.87	3.09	5.77	13.65	10.14
SEm(±)	2.49	3.96	10.02	0.09	0.10	0.09	0.21	0.16	0.15
CD(0.05)	7.35	11.68	29.57	0.26	0.29	0.26	0.62	0.47	0.44

247

V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha⁻¹, N1-50 kg ha⁻¹, N2-100 kg ha⁻¹, N3- 150 kg ha⁻¹,N4-200 kg ha⁻¹, N5-250 kg ha⁻¹.

248 **REFERENCES**

249 Cochran, W.G. and Cox, G.M. 1957.(2nd Edition).*Experimental Design*.John Wiley and Sons.
250 New York.Pp: 615.

251 Mohammad, V. and Mohammadreza, N.2012.The effect of various nitrogen fertilizer amounts
252 of yield and nitrate accumulation in tubers of two potato cultivars in cold–IV regions of
253 Isfahan (Iran). *International Journal of Agriculture and Crop Sciences* .**4(22)**: 1688-
254 1691.

255 Nasab H.M. , Siadat S.A., Naderi A., Lack S.andModhej A.2014.The effects of Drought Stress
256 and Nitrogen Levels on Yield , Stomatal Conductance and Temperature Stabily of
257 Rapeseed (Canola) Genotypes. *Advances in Environmental Biology*.8(10).1239-1247.

258 Sharma, B.D. and Sharma, U.C. 1991. Dry matter accumulation behaviour of different potato
259 cultivars (*Solanumtuberosum* L.).*Indian J Hill Farming*.**4(1)**: 11-14.

260 Wolf, S.A., Marani and Rudich, J. 1990.Effect of temperature and photoperiod on assimilate
261 partitioning in potato plants. *Annals of Botany*.**66**:513-520.

262

263

UNDER PEER REVIEW