

1 **Evaluation of Different Nutritional and Soil Sources on the Early Growth of**  
2 ***Moringa oleifera* (Lam)**

3  
4 **ABSTRACT**

5 In any plantation establishment programme their must be adequate number of healthy  
6 seedlings and these depend on the nutrition and care given to them at the nursery stage.  
7 Pot experiment was conducted to assess the response of *Moringa oleifera* (Lam)  
8 seedlings to different soil and nutritional sources at nursery stage. One hundred seeds  
9 were obtained from Centre for Environmental Renewable Resources Research and  
10 Development (CENRAD) Ibadan, Nigeria and sown in germination trays. Seed  
11 germination was completed between 10-15 days, 36 uniformly growing seedlings were  
12 transplanted into polythene pots of size 29×25cm and were filled with different soil  
13 sources at 500gm (arable) soil, forest reserve soil and natural forest mixed with  
14 nutritional sources of the same ratio (10gm) i.e. poultry manure, cow dung and N.P.K  
15 15.15.15. The experiment was factorial arranged in a completely randomized design.  
16 Result showed that pots with natural forest soil mixed with 10gm N.P.K ( $s_3f_3$ ) produced  
17 seedlings with highest value In plant height (76.30cm), stem diameter (3.47mm) and  
18 number of branches (10.00) which was significantly different ( $p < 0.01$ ) from the other  
19 treatments. The least value in plant height (30.70cm) stem diameter (1.50mm) and  
20 number of branches (3.3) were recorded in pots with forest reserve soil mixed with cow  
21 dung ( $s_2f_2$ ). Interaction effect of soil and nutritional sources were significant for all the  
22 growth parameters assessed ( $p < 0.01$ ). Natural forest soils treated with N.P.K 15.15.15  
23 should be employed in raising *Moringa oleifera* seedlings at nursery stage for optimum  
24 performance.

25 **Key Words:** *Moringa oleifera*, Growth, Nutritional sources, Soil

26 **INTRODUCTION**

27 The forest plays an important role in protecting the soil, ameliorating the environment  
28 and protecting water resources. Non-timber forest products are very essential in urban  
29 and rural life, under which *Moringa oleifera* (Lam) belong (Leone *et al.*, 2015). *Moringa*  
30 *oleifera* is the most widely cultivated species of the genus *Moringa*, is the only genus in  
31 the family Moringaceae. English common names include: Moringa, (Olson, 2010)  
32 drumstick tree (from the appearance of the long, slender, triangular seed pods),  
33 horseradish tree (from the taste of the roots, which resembles horseradish, ben oil tree, or  
34 benzoil tree (from the oil which is derived from the seeds). It is a fast-growing, drought-  
35 resistance tree, native to the southern foothills of the Himalayas in the northwestern  
36 Indian, and widely cultivated in tropical and subtropical areas where its young seed pods  
37 and leaves are used as vegetables. It can also be used for water purification and washing,  
38 and is sometimes used in herbal medicine (Torondel, *et al.*, 2014).

39 It has become a clear issue that man cannot sufficiently sustain its existence without  
40 adequately improving the level of food and fiber production as raw material for industrial  
41 uses. Most tropical soils are deficient in nitrogen and other macronutrients and uptake of  
42 these limited quantities of nutrients by plant roots from litters is difficult (Jose, 2003).  
43 Nitrogen allows plants to produce proteins needed to build living tissues for green stems,  
44 leaves and strong roots, phosphorus helps move energy throughout the plant while  
45 potassium aids plants in adapting sugars needed in growth. Fertilization is the only way  
46 to supply nutrients within a short period of time. Adegbidi et al, (2003) reported that the  
47 effects of the mixed use of chemical fertilizer and organic matter on the growth of trees  
48 and soil fertility vary substantially according to the fertilizer amounts and the organic  
49 manure characteristics. The need to investigate the response of *Moringa oleifera* to  
50 different ratios of inorganic and organic fertilizer application on soil sources is essential  
51 as this will determine its optimum growth performance at the nursery stage. The objective  
52 of the study was to investigate the effect of different nutritional and soil sources on early  
53 growth of *M. oleifera* so as to find the optimum dose of fertizer for raising quality  
54 seedlings.

## 55 MATERIALS AND METHOD

56 The experiment was carried out at the West African Hardwood Improvement Project  
57 (WAHIP) of the Forestry Research Institute of Nigeria (FRIN), Ibadan (Latitude 7°39'13"  
58 and longitude 3°8'28"E. the Institute is situated at Jericho Hills in Ibadan North West  
59 Local Government Area of Oyo State. The climate of the area is tropically dominated by  
60 rainfall pattern ranging between 1400 - 1500mm, average temperature is 30°C. It has 2  
61 distinct seasons rainy season (April – October) and dry season (November – March)  
62 (FRIN, 2015). The black polythene pots were purchased from CENRAD, Ibadan. Cow  
63 dung and poultry manure were collected from Federal College of Forestry Teaching and  
64 Research Farm Ibadan. N. P. K 15.15.15 fertilizer was obtained from Centre for  
65 Environmental Renewable Resources and Management Development (CENRAD),  
66 Jericho, Ibadan. The natural forest soil was collected from FRIN arboretum, forest  
67 reserve soil was collected from *Pinus caribea* Morelet plantation FRIN while the  
68 cultivated soil used was collected from Federal College of Forestry Ibadan farm. One  
69 hundred *Moringa oleifera* seeds were obtained from CENRAD and sown in germination  
70 trays filled with sterilized river sand.

71 Cow dung and poultry manure were dried, crushed and sieved with 2mm sieve while the  
72 soil samples were also sieved. The same ratio of cow dung, poultry manure and fertilizer  
73 (N.P.K. 15.15.15) were measured in grams (10gm) and mixed with the soil samples, each  
74 treatment contained the same level of organic and inorganic fertilizers. 10gm of organic  
75 fertilizer, (cow dung and poultry manure of same ratio) and 10gm of N.P.K. 15.15.15  
76 were weighed in the soil laboratory of Forestry Research Institute of Nigeria. 10mg of  
77 cow dung and poultry manure were applied on 36 seedlings i.e. 18 seedlings contained  
78 10gm of cow dung and 18 seedlings contained 10gm of poultry manure, while 18  
79 seedlings contained 10gm of N.P.K. 15.15.15. Eighteen seedlings were used as control. A  
80 total of 72 seedlings were transplanted after four weeks into polythene pots of 29×25cm  
81 size filled with the different potting mixtures. Watering of the seedlings was done once

82 daily. Seedling height (cm), stem diameter (mm) and number of branches were assessed  
 83 after 4weeks for 12 weeks.

84 **Table 1:** Laboratory Analysis of Organic and Inorganic Fertilizer

85 <b>Sample Code</b>	%N	%P	%K
86 Poultry manure	3.47	1.18	1.38
87 Cow dung	4.06	0.33	0.77
88 N.P.K. 15.15.15	15.00	15.00	15.00

89 **Treatment Combination**

- 90 Where; (1) (i)  $S_1f_0$  - Arable soil without fertilizer (control)  
 91 (ii)  $S_1f_1$  - Arable soil with poultry manure  
 92 (iii)  $S_1f_2$  - Arable soil with cowdung  
 93 (iv)  $S_1f_3$  - Arable soil with N.P.K (15.15.15)  
 94 (2) (i)  $S_2f_0$  - Forest reserve soil without fertilizer (control)  
 95 (ii)  $S_2f_1$  - Forest reserve soil with poultry manure  
 96 (iii)  $S_2f_2$  - Forest reserve soil with cowdung  
 97 (iv)  $S_2f_3$  - Forest reserve soil with N.P.K. (15.15.15)  
 98 (3) (i)  $S_3f_0$  - Natural Forest soil without fertilizer (control)  
 99 (ii)  $S_3f_1$  - Natural Forest soil with poultry manure  
 100 (iii)  $S_3f_2$  - Natural Forest soil with cowdung  
 101 (iv)  $S_3f_3$  - Natural forest soil with N.P.K (15.15.15)

102 **Data Analysis**

103 Analysis of variance was used to analyze the data obtained while least significant  
 104 difference (LSD) was used to separate the means where significant.

105 **RESULTS AND DISCUSSION**

106 **Table2:** Effect of treatment on stem diameter (mm) of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
-----------	------	------	------	------	-------	-------

S <sub>1</sub> F <sub>0</sub>	1.90	2.06	2.00	2.03	2.17	3.19
S <sub>1</sub> F <sub>1</sub>	1.93	2.07	1.97	1.97	2.17	3.20
S <sub>1</sub> F <sub>2</sub>	2.03	2.10	2.00	2.03	2.20	2.84
S <sub>1</sub> F <sub>3</sub>	2.07	2.20	2.40	2.63	3.17	4.28
S <sub>2</sub> F <sub>0</sub>	1.97	2.03	0.63	0.63	0.63	1.45
S <sub>2</sub> F <sub>1</sub>	2.10	2.13	1.97	1.93	2.00	3.17
S <sub>2</sub> F <sub>2</sub>	2.07	1.40	1.40	1.43	1.50	2.42
S <sub>2</sub> F <sub>3</sub>	2.03	2.03	0.00	0.00	0.00	0.00
S <sub>3</sub> F <sub>0</sub>	1.90	2.03	1.40	3.80	1.77	2.85
S <sub>3</sub> F <sub>1</sub>	2.03	2.07	2.13	2.17	2.60	3.00
S <sub>3</sub> F <sub>2</sub>	1.90	2.03	2.17	2.22	2.60	3.00
S <sub>3</sub> F <sub>3</sub>	1.93	2.00	2.07	2.47	3.47	4.53
Mean	1.99	2.01	1.68	1.94	2.02	2.83
Significance	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
Se±	0.1023	0.1023	0.1023	0.1023	0.1023	0.1023
LSD	0.9105	0.9105	0.9105	0.9105	0.9105	0.9105
C.V%	5.3	5.3	5.3	5.3	5.3	5.3

107 **Table 2:** Effect of treatment on stem diameter (mm) of *Moringa oleifera* seedlings  
108 Where: - WAT= Week after transplanting\*\*\* significantly difference (p<0.001).

109 Table 2 shows that there was no significant difference in stem diameter of the seedlings  
110 subjected to various treatments at P<0.001 while N.P.K. 15.15.15 in natural forest soil  
111 (S<sub>3</sub>F<sub>3</sub>) had the highest stem diameter of 4.53mm. There was a continuous increase in stem  
112 diameter across the weeks. This was followed by N.P.K. 15.15.15 in arable soil that had  
113 value of 4.28mm against cowdung in forest reserve soil which had the lowest stem  
114 diameter of 1.5mm across the weeks. This confirms the findings of Waheed et al. (2001)  
115 for *Camelia sinensis* that nitrogen containing fertilizers such as N. P.K had a significant  
116 effect on seedling growth parameters. Almeida (1997) had earlier reported that N.P.K  
117 fertilizer gave a positive response in seedling growth of cashew. This also supports the  
118 findings of Larcheveque et al. (2011) that chemical fertilizers promote higher growth and  
119 root development compared to livestock organic manure in a Poplar plantation.

**Table 3:** Effect of soil mixture on the height (cm) of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
S <sub>1</sub> F <sub>0</sub>	32.40	34.50	36.40	37.40	40.00	43.04
S <sub>1</sub> F <sub>1</sub>	28.40	37.33	41.33	42.33	46.00	54.25
S <sub>1</sub> F <sub>2</sub>	32.00	36.67	39.50	34.17	46.67	55.80
S <sub>1</sub> F <sub>3</sub>	40.17	41.33	46.00	54.33	66.67	74.55
S <sub>2</sub> F <sub>0</sub>	32.33	31.67	11.00	11.67	12.33	15.61
S <sub>2</sub> F <sub>1</sub>	34.73	33.73	35.17	37.83	42.50	48.72
S <sub>2</sub> F <sub>2</sub>	24.83	20.67	28.77	30.00	30.67	35.74
S <sub>2</sub> F <sub>3</sub>	26.33	34.33	0.00	0.00	0.00	0.00
S <sub>3</sub> F <sub>0</sub>	28.50	31.50	32.17	35.00	36.00	39.50
S <sub>3</sub> F <sub>1</sub>	34.83	35.57	46.50	50.33	52.33	60.28
S <sub>3</sub> F <sub>2</sub>	34.50	34.17	42.33	49.33	54.67	64.45
S <sub>3</sub> F <sub>3</sub>	30.90	32.33	43.33	58.00	73.33	89.67
Mean	31.66	33.81	3.38	37.45	41.74	48.47
significance	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
Se±	3.638	3.638	3.638	3.638	3.638	3.638
LSD	0.9105	0.9105	0.9105	0.9105	0.9105	0.9105
C.V%	5.3	5.3	5.3	5.3	5.3	5.3

121 Where:- WAT = Week after transplanting\*\*\* significantly difference (<0.001).

122 Table 3 revealed that there was no significant difference in the height of Moringa  
 123 seedlings among the treatments at P<0.001. N.P.K. 15.15.15 in natural forest soil had a  
 124 height of 89.67cm (P<0.001), this was achieved due to the increase in the growth rate  
 125 (height) of Moringa across the weeks. This was followed by N.P.K. 15.15.15 in arable  
 126 soil that had a value of 74.55cm, the trend was also maintained across the weeks while  
 127 cowdung in forest reserve soil had the lowest height of 35.74cm. Similar positive results  
 128 has been reported by (Hoque, *et al.*, 2004) seedling growth was enhanced significantly  
 129 with the application of N.P.K fertilizer. Tree seedlings need nutrients to grow, nitrogen  
 130 for lots of green leaves, phosphorus for new tissues particularly the roots and potassium  
 131 for seedling vigour.

132

**Table 4:** Effect of soil mixture on number of branches of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
S <sub>1</sub> F <sub>0</sub>	6.00	6.00	5.67	6.33	6.67	6.67
S <sub>1</sub> F <sub>1</sub>	7.00	6.00	6.33	6.00	6.00	6.00
S <sub>1</sub> F <sub>2</sub>	5.67	4.67	6.00	7.33	6.67	7.52
S <sub>1</sub> F <sub>3</sub>	4.00	6.00	8.67	9.67	9.67	10.23
S <sub>2</sub> F <sub>0</sub>	5.00	2.00	1.33	1.33	1.33	1.33
S <sub>2</sub> F <sub>1</sub>	6.33	3.67	4.00	5.00	5.33	6.50
S <sub>2</sub> F <sub>2</sub>	5.67	3.00	3.00	3.67	3.33	3.33
S <sub>2</sub> F <sub>3</sub>	5.67	4.33	0.00	0.00	0.00	4.33
S <sub>3</sub> F <sub>0</sub>	5.00	3.67	4.67	4.67	4.00	4.00
S <sub>3</sub> F <sub>1</sub>	6.67	6.33	7.33	6.67	6.67	6.67
S <sub>3</sub> F <sub>2</sub>	6.33	7.00	7.00	8.67	8.00	8.00
S <sub>3</sub> F <sub>3</sub>	6.33	6.33	8.67	10.33	10.00	11.00
Mean	5.81	4.91	5.58	6.17	6.00	6.33
significance	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
Se±	0.570	0.570	0.570	0.570	0.570	0.570
LSD	2.830	2.830	2.830	2.830	2.830	2.830
C.V%	10.3	10.3	10.3	10.3	10.3	10.3

133 Where: WAT= week after transplanting\*\*\* significantly different (P &lt; 0.001)

134 Table 4 shows that there were significant differences in number of branches of Moringa  
135 seedlings among the treatments at P<0.001. N.P.K. 15.15.15 in natural forest soil (S<sub>3</sub>F<sub>3</sub>)  
136 had the highest value of 10.0. This was achieved due to continuous production in number  
137 of branches across the weeks. This was followed by N.P.K. 15.15.15 in arable soil that  
138 had the value of 9.7 which was also maintained across the weeks while cowdung in forest  
139 reserve soil (S<sub>2</sub>F<sub>2</sub>) had the lowest value of 3.33 across the weeks. This supports the  
140 findings of Jaenicke (1999) who stated that cowdung contains 0.3% Nitrogen, 0.2%  
141 phosphoric acid and 0.1- 0.5% while Ajay, (2017) also reported that cowdung is not as  
142 rich in nitrogen as many other types of fertilizers. He reported that it has about 8%  
143 nitrogen, 2% phosphorus and 1% potassium. These nutrients are also slowly infused into  
144 the soil.

145

146

147

148 **Conclusion**

149 Based on the findings of these studies, seedlings raised with N.P.K. 15.15.15 mixed with  
150 natural forest soil had the highest stem diameter, height and number of branches. Also,  
151 there were significant differences in seedlings height, stem diameter and branches at  
152  $P < 0.001$ , interaction effect among soil and nutritional sources was also significant.

153 **Recommendation**

154 Natural forest soil mixed with N.P.K. 15:15:15 should be used in raising seedlings of  
155 *Moringa oleifera* for optimum growth performance since the seedlings presented most  
156 noticeable positive influence on seedling growth in the nursery.

157

158

159 **References**

160 Adegbidi, H.G.; Briggs, R.D.; White, E.H.; Abrahamson, L.P.; Volk, T.A. Effects of  
161 organic amendments and slow release  $N_2$  fertilizer on willow stem biomass  
162 production and soil chemical characteristics. **Biomass Bioenergy** v.25,  
163 no.4, p. 389-398, 2003.

164 Ajay, M.; R. Effective use of cowdung manure for healthy plant growth.  
165 **International Journal of Advanced Research and Development** v, 2 5,  
166 no. p 218-221, 2017.

167 Almeida, A.;G.; Almeida, F. C. G. Meneses,J. J. Effect of N, P and K fertilization on  
168 productivity in dwarf early air-layered Cashew under irrigation.**Turri alba**  
169 v. 44 no.3, p. 168-178,1997.

170 FRIN-Forestry Research Institute of Nigeria. Annual MEtereological Report. 2013.

171 Hoque, A. T.; M. R.; Hossian, M.K.; Mohiuddin, M.; Hoque, M. M. (2004). Effect of  
172 inorganic fertilizers on initial growth performance of *Michelia champaca*  
173 Linn. Seedlings Nursery. **J.Biol.Sci** v 4 p. 489-497.

174 Jaenicke,H. Good tree nursery practices: practical guidelines for research nurseries.  
175 Nairobi ICRAF, 1999

176 Jose, L.M (2003). Nitrogen and phosphorus resorption in tree of neotropical rainforest.  
177 **J.Trop Ecol** v, 19 p.465-468.

178 Larcheveque, M.; Desroches,A.;Larocque, G.R. Comparison of manure bcompost and  
179 mineral fertilizer for hybrid poplar plantation establishment on boreal clay  
180 soils . **Ann. For, Sci** v. 68, no.4, p 849-860

- 181 Leone, A.; Spada, A.; Battezzati, A.; Schiraldi, A.; Aristil, J.; Bertoli, S. "Cultivation,  
182 Genetic, Ethnopharmacology, Phytochemistry and Pharmacology of  
183 *Moringa oleifera* Leaves: An Overview" .**Int J Mol Sci**.v, 16 no.6: 12.  
184 p791~835. 2015.
- 185 Olson, M.;E. Flora of North America Committee, ed.flora summary :Moringaceae  
186 Drumstick Family. **Flora of North America North of Mexico**.7.New York  
187 and Oxford.p 167-169.
- 188 Torondel, B.; Opare, D.; Brandberg, B.; Cobb, E.; Cairncross, S. (2014). "Efficacy of  
189 *Moringa oleifera* leaf powder as a hand- washing product: A crossover  
190 controlled study among healthy volunteers". **BMC Complementary and  
191 Alternative Medicine**.v,14 no, 57. doi : 10.1186/1472-6882-14-57 .2014.
- 192 Waheed, A.; Jamal, A.; Hamad,; F.S; Ahmad, N. Growth of tea seedlings affected by  
193 different levels and application of N.P.K Urea. **Online Journal of  
194 Biological Sciences** v, 1 no.3.p101-102.2001.

195

196